





3 1761 04382 0943

UNIV. OF
TORONTO
LIBRARY









GENERA INSECTORUM

FASC. CLXXXIV-CLXXXV



Zool
Insecta
W

5

GENERA INSECTORUM

DIRIGÉS PAR

P. WYTSMAN

"



FASCICULES CLXXXIV-CLXXXV

*364191
16. 3. 39*

184. Lepidoptera Heterocera. Fam. Gelechiadæ, par E. MEYRICK.
185. Diptera. Fam. Empididæ, par A. L. MELANDER.

BRUXELLES
LOUIS DESMET - VERTENEUIL
IMPRIMEUR - ÉDITEUR
1925-1927

QL
468
W87
Fasc. 184-185

**LEPIDOPTERA HETEROCEA
FAM. GELECHIADÆ**

LEPIDOPTERA HETEROCHERA

FAM. GELECHIADÆ

by E. MEYRICK

WITH 5 COLOURED PLATES

General Characters. — Head usually smooth or with appressed scales, sidetuffs sometimes raised, only in *Panicotricha* with erect rough hairs on crown; ocelli small, posterior or rarely inferior; tongue usually developed. Antennae varying from $2/3$ to $1\frac{1}{2}$, in ♂ usually simple or shortly ciliated, but sometimes with long fasciculate ciliations or even pectinated (*Ptilothyris*), basal joint more or less elongate, usually without pecten (but pecten present in several independent genera). Labial palpi normally long or very long, recurved, sickle-shaped, acute, variably scaled, rarely very short or aborted (*Amblypalpis* and allies), terminal joint in ♂ sometimes aborted. Maxillary palpi short or very short, scaled, appressed or rarely porrected (*Lacistodes*). Anterior legs normally long, slender; posterior tibiae usually with rough scales or long hairs above. Forewings moderate in earlier but becoming narrow in higher forms; 1b normally furcate, 2 from or towards angle, rarely remote, 7 and 8 stalked or coincident (only in *Anomoxena* separate by obsolescence of stalk), 11 usually from middle. Hindwings usually 1 or over 1, but sometimes under 1, usually trapezoidal, with termen slightly or strongly sinuate or abruptly emarginate beneath apex or even with deep excavation, apex sometimes strongly or even excessively produced, but sometimes elongate-ovate without terminal sinuation, cilia 1/6-6, cubital pecten sometimes present (groups 4 and 7); cell complete but transverse vein sometimes obsolescent, 3 and 4 originally connate but in higher forms often parallel, 5 parallel or approximated to 4 or when 4 is absent connate with 3, 6 and 7 normally connate or stalked, yet sometimes remote and parallel (*Protolechia* and allies), 8 normally appressed or connected with middle of upper margin of cell.

Larva with prolegs on segments 7-10 and 13, rarely rudimentary; rarely with clothing of dense long hairs (*Hygroplasta*); feeding in more or less web, the more primitive forms on lichens, moss, and dry vegetable or seldom animal refuse, those more advanced on phanerogamous plants and trees, in spun shoots, rolled leaves, within stems or roots or seed-heads or sometimes mining in leaves, occasionally living in a portable case, and in general displaying flexibility of habit and adaptability.

LEPIDOPTERA HETEROCEA

Pupa with first four segments of abdomen fixed; sometimes with rather prominent scattered hairs (group 8); usually in slight cocoon amongst refuse or on food-plant.

* * *

The conception of the family *Gelechiidae* is apparently due to Stainton, but (as noted in my remarks on the *Oecophoridae*) he did not discriminate properly between these two families, their limits being defined by myself in 1883. Subsequent discoveries of material have shown that (as indicated in the family characters above) there is really not a single character to which an exception does not occur; as knowledge widens, all boundaries become less absolute. I am however still satisfied that the *Gelechiidae* and *Oecophoridae* do represent two natural associations or lines of development, arising from a common base and not intermixing at any other point, but in order to distinguish the two in practice it is necessary to consider the whole of the characters in conjunction, and not to argue from single points. There are in fact many points of distinction between the two families, and the failure of one can always be corrected by the inspection of the rest. In *Anomoxena* a character (common to both families) fails which might have been thought (and probably really is) difficult of modification, and to which no other exception is known amongst the whole of the 700 genera of the two families, yet this is in a highly specialised genus far up the line of ascent; as is easily seen from the rest of its characters. Great caution is therefore advisable in accepting the statements of those whose limited experience recommends faith in any particular characters whatsoever.

I have divided the family into nine groups which I consider undoubtedly natural, but I have not ventured to attach names to these groups, because they are not properly definable; there are in all cases structural characters which serve as guides to their identification, but there are so many exceptions that it would only be misleading to draw up definitions. When familiar with these groups, however, it is generally possible to say at once of a new genus to which group it should be referred, and thus they are really serviceable as mental conceptions, in view of the large size of the family, and contribute materially to its comprehension.

The family as a whole must be regarded as a rather modern development, which has succeeded in adapting itself to a great variety of situations. In New Zealand it is very scantily represented, obviously only by three or four types that have found their way there in recent times. Elsewhere it is generally abundant (not in oceanic islands), but with different types predominant in different regions; thus in Europe and North America the *Gelechia* type predominates, in India the *Lecithocera* type, in South America the *Compsolechia* type, in Australia the *Protolechia* type; these groups are therefore more recent in origin than the period since which these main regions have been disconnected. These distinctions are very pronounced; but owing to the obscure appearance and secretive habits of many of these insects, there are undoubtedly a large proportion still awaiting discovery.

In the present revision I have classified 391 genera and 3542 species.

KEY TO THE GENERA

- | | |
|---|---------------------------------|
| 1. Forewings with naked glandular spot beneath costa towards base | 64. Genus CLISTOTHYRIS, Zeller. |
| Forewings without such spot | 2. |
| 2. Forewings with 7 and 8 separate | 33. Genus ANOMOXENA, Meyrick. |
| Forewings with 7 and 8 stalked or coincident | 3. |
| 3. Forewings with 7 to termen or apex, or 7 (or 8 or 9) absent | 4. |
| Forewings with 7 to costa, 7-9 present | 198. |
| 4. Forewings with 6 to costa | 5. |
| Forewings with 6 to apex or termen | 6. |

5. Forewings with 2 and 3 stalked	146. Genus EUNOMARCHA, Meyrick.
Forewings with 2 and 3 remote	126. Genus SIMONEURA, Walsingham.
6. Antennae about 1 or over 1	7.
Antennae not over 5/6	47.
7. Anterior coxae and middle tibiae long-haired	284. Genus TOGIA, Walker.
Anterior coxae and middle tibiae not long-haired	8.
8. Costa of hindwings with strong postmedian scale-projection	349. Genus PHILOPTILA, Meyrick.
Costa of hindwings without scale-projection	9.
9. Posterior tibiae with median tuft of rough hairscales	10.
Posterior tibiae without median tuft of rough hairscales	17.
10. Forewings with 7 absent	11.
Forewings with 7 present	13.
11. Forewings with 8 and 9 stalked	12.
Forewings with 8 and 9 separate	299. Genus ANAXYRINA, Meyrick.
12. Palpi in ♂ with terminal joint absent	300. Genus HETERODELTIS, Meyrick.
Palpi in ♂ with terminal joint developed	301. Genus CANTHONISTIS, Meyrick.
13. Forewings with 2 and 3 stalked	302. Genus HABROGENES, Meyrick.
Forewings with 2 and 3 remote	14.
14. Hindwings trapezoidal	15.
Hindwings lanceolate	16.
15. Hindwings in ♂ with 2 and 4 connate, 3 absent	298. Genus TIMYRA, Walker.
Hindwings in ♂ with 2 remote, 3 and 4 connate	297. Genus OLBOTHREPTA, Meyrick.
16. Forewings with 4 absent	295. Genus MONERISTA, Meyrick.
Forewings with 4 present	294. Genus MAGONYMPHA, Meyrick.
17. Forewings with 2 and 3 stalked or coincident (3 absent)	18.
Forewings with 2 separate	38.
18. Hindwings with 5 separate	19.
Hindwings with 5 connate or stalked with 3	32.
19. Forewings with 10 out of 7	333. Genus PHATNOTIS, Meyrick.
Forewings with 10 separate	20.
20. Hindwings with 3 and 4 remote	306. Genus TELEPHATA, Meyrick.
Hindwings with 3 and 4 connate, stalked, or coincident	21.
21. Second joint of palpi with apical tuft beneath	22.
Second joint of palpi without apical tuft beneath	24.
22. Forewings with 7 absent	262. Genus NOEZA, Walker.
Forewings with 7 present	23.
23. Antennae over 1	304. Genus FRISILIA, Walker.
Antennae under 1	257. Genus TOCMIA, Walker.
24. Antennae thick, flatly compressed throughout	305. Genus NARTHECOCEROS, Meyrick.
Antennae not markedly flattened	25.
25. Palpi in ♂ with terminal joint absent	288. Genus DOXOGENES, Meyrick.
Palpi in ♂ with terminal joint developed	26.
26. Palpi with terminal joint shorter than second	27.
Palpi with terminal joint as long as second or longer	29.
27. Forewings with 7 absent	148. Genus DREPANOTERMA, Walsingham.
Forewings with 7 present	28.

LEPIDOPTERA HETEROCHERA

28. Forewings with 7 to apex	354. Genus BRACHYERGA, Meyrick.
Forewings with 7 to termen	308. Genus SYNCATHEDRA, Meyrick.
29. Hindwings with 5 equidistant, parallel	351. Genus ADELOMORPHA, Snellen.
Hindwings with 5 nearer 4	30.
30. Forewings with 5 connate with 2 from angle	353. Genus THUBANA, Walker.
Forewings with 5 from above angle	31.
31. Hindwings in ♂ with submedian groove and hair-pencil	287. Genus DINOCHARES, Meyrick.
Hindwings in ♂ without submedian groove and hair-pencil	357. Genus LECITHOCERA, Herr.-Schäff.
32. Forewings with 3 absent	33.
Forewings with 3 present	36.
33. Forewings with 7 absent	345. Genus CROCANTHES, Meyrick.
Forewings with 7 present	34.
34. Forewings with 2 and 4 stalked	35.
Forewings with 2 and 4 separate	340. Genus COYDALLA, Walker.
35. Forewings with 10 absent	352. Genus SPHENOCRATES, Meyrick.
Forewings with 10 present	347. Genus PLACANTHES, Meyrick.
36. Forewings with 7 absent	309. Genus PSAMMORIS, Meyrick.
Forewings with 7 present	37.
37. Forewings with 11 absent	350. Genus COPROPTILIA, Snellen.
Forewings with 11 present	356. Genus SARISOPHORA, Meyrick.
38. Forewings with 7 absent	39.
Forewings with 7 present	43.
39. Hindwings with 4 absent	313. Genus PARELLIPTIS, Meyrick.
Hindwings with 4 present	40.
40. Hindwings with 3 and 4 out of 5	314. Genus DOLICHOTORMA, Meyrick.
Hindwings with 5 separate	41.
41. Forewings with 3 and 4 long-stalked or coincident	321. Genus CELETODES, Meyrick.
Forewings with 3 and 4 separate	42.
42. Palpi in ♂ with terminal joint absent	286. Genus MNESTERIA, Meyrick.
Palpi in ♂ with terminal joint developed	144. Genus MERIMNETRIA, Walsingham.
43. Hindwings with 5 separate	44.
Hindwings with 5 connate or stalked with 3	320. Genus ERIDACHTHA, Meyrick.
44. Hindwings with 4 absent	319. Genus CROCOGMA, MEYRICK.
Hindwings with 4 present	45.
45. Second joint of palpi with apical tuft beneath	282. Genus CHERSOGENES, Walsingham.
Second joint of palpi without apical tuft beneath	46.
46. Hindwings in ♂ with expansible fringe in submedian groove	292. Genus PHANOSCHISTA, Meyrick.
Hindwings in ♂ without expansible fringe in submedian groove	331. Genus HOMALOXESTIS, Meyrick.
47. Hindwings with 7 absent	327. Genus APHNOGENES, Meyrick.
Hindwings with 7 present	48.
48. Hindwings with 3 (or 4) absent (at least in ♂)	49.
Hindwings with 3 and 4 present	67.
49. Forewings with 2 and 3 separate	50.
Forewings with 2 and 3 stalked or coincident	53.
50. Second joint of palpi tufted beneath	202. Genus PILOCRATES, Meyrick.
Second joint of palpi not tufted beneath	51.

51. Forewings with 7 absent	52.
Forewings with 7 present	310. Genus PSEUDOCRATES, Meyrick.
52. Forewings with 12 sinuate and approximated to 11	325. Genus EUPRAGIA, Walsingham.
Forewings with 12 straight	322. Genus ATHRINACIA, Walsingham.
53. Costa of hindwings with projecting scale-teeth	348. Genus NOSPHISTICA, Meyrick.
Costa of hindwings without projecting scale-teeth	54.
54. Forewings with 7 absent	55.
Forewings with 7 present	56.
55. Terminal joint of palpi longer than second, antennae in ♂ pectinated	336. Genus PTILOTHYRIS, Walsingham.
Terminal joint of palpi not longer than second, antennae in ♂ not pectinated	362. Genus COPHOMANTIS, Meyrick.
56. Forewings with 7 to termen	57.
Forewings with 7 to apex	60.
57. Forewings with 5 from angle	58.
Forewings with 5 from above angle	59.
58. Second joint of palpi tufted beneath	343. Genus CYNICOSTOLA, Meyrick.
Second joint of palpi smooth	342. Genus GASMARA, Walker.
59. Forewings with 9 out of 7	334. Genus EPHARMONIA, Meyrick.
Forewings with 9 separate	344. Genus THYMBRITIS, Meyrick.
60. Hindwings with 5 separate	61.
Hindwings with 5 connate or stalked with 3	62.
61. Forewings with 3 absent	341. Genus HARMATITIS, Meyrick.
Forewings with 3 present	255. Genus MACHLOTRICA, Meyrick.
62. Palpi with terminal joint as long as second	63.
Palpi with terminal joint longer than second	66.
63. Forewings with 4 absent	363. Genus ORGANITIS, Meyrick.
Forewings with 4 present	64.
64. Forewings with 4 out of 2	360. Genus CHLOROLYCHNIS, Meyrick.
Forewings with 4 separate	65.
65. Forewings with 3 absent	359. Genus LARCOPHORA, Meyrick.
Forewings with 3 present	361. Genus PROTOLYCHNIS, Meyrick.
66. Forewings with 5 connate with 2	335. Genus SISYRODONTA, Meyrick.
Forewings with 5 not connate with 2	339. Genus DELTOPLASTIS, Meyrick.
67. Hindwings with 5 absent	68.
Hindwings with 5 present	70.
68. Forewings with 2 separate	323. Genus THRYSIGENES, Meyrick.
Forewings with 2 stalked with 3 or 4	69.
69. Forewings with 7 to termen	253. Genus STRYPHNOCOPA, Meyrick.
Forewings with 7 to apex	365. Genus PROSODARMA, Meyrick.
70. Forewings with 3 and 4 coincident	71.
Forewings with 3 and 4 not coincident	78.
71. Second joint of palpi shortly tufted beneath	199. Genus ENCOLAPTA, Meyrick.
Second joint of palpi not tufted beneath	72.
72. Hindwings with 3 and 4 remote	73.
Hindwings with 3 and 4 connate	75.

73. Basal joint of antennae with pecten	2. Genus SICERA, Chrétien.
Basal joint of antennae without pecten	74.
74. Forewings with 7 absent	102. Genus THYRSOSTOMA, Meyrick.
Forewings with 7 present	318. Genus PHARANGITIS, Meyrick.
75. Hindwings with 6 to apex	53. Genus ELASIPRORA, Meyrick.
Hindwings with 6 to termen	76.
76. Hindwings with termen rounded	115. Genus PAURONEURA, Turner.
Hindwings with termen sinuate or emarginate	77.
77. Forewings with 7 absent	105. Genus THIOTRICA, Meyrick.
Forewings with 7 present, to termen	247. Genus AULIDIOTIS, Meyrick.
78. Head above with long rough spreading hairs	388. Genus PANICOTRICA, Meyrick.
Head above with appressed scales	79.
79. Labial palpi very short	80.
Labial palpi moderate or long	81.
80. Forewings with 6 and 8 stalked	3. Genus SCLEROCECIS, Chrétien.
Forewings with 6 and 8 separate	1. Genus AMBLYPALPIS, Ragonot.
81. Basal joint of antennae with pecten	82.
Basal joint of antennae without pecten	83.
82. Hindwings with 6 and 7 parallel	11. Genus EPIDOLA, Staudinger.
Hindwings with 6 and 7 closely approximated towards base	312. Genus MNESISTEGA, Meyrick.
83. Hindwings with 3 or 4 remote	84.
Hindwings with 3 and 4 connate or seldom approximated	92.
84. Forewings with 2 and 3 stalked	85.
Forewings with 2 and 3 remote	87.
85. Forewings with 7 absent	86.
Forewings with 7 present	258. Genus SCHEMATISTIS, Meyrick.
86. Second joint of palpi rough-scaled above	385. Genus ULIARIA, Dumont
Second joint of palpi not rough-scaled above	384. Genus SYRMADAULA, Meyrick.
87. Forewings with 9 absent	13. Genus ENCENTROTIS, Meyrick.
Forewings with 9 present	88.
88. Forewings with 8 and 9 stalked	109. Genus CRAMBODOXA, Meyrick.
Forewings with 8 and 9 separate	89.
89. Terminal joint of palpi concealed	14. Genus NEVADIA, Caradja.
Terminal joint of palpi exposed	90.
90. Forewings with 6 and 8 stalked	91.
Forewings with 6 and 8 separate	92.
91. Second joint of palpi with long tuft beneath	211. Genus AXYSTOLA, Meyrick.
Second joint of palpi without tuft	108. Genus CNAPHOSTOLA, Meyrick.
92. Terminal joint of palpi longer than second	124. Genus ANTERETHISTA, Meyrick.
Terminal joint of palpi half second	10. Genus COLOPTERYX, Hofmann.
93. Hindwings with 6 and 7 remote at origin, nearly parallel	94.
Hindwings with 6 and 7 approximated or stalked or coincident	104.
94. Forewings with 7 absent	95.
Forewings with 7 present	192. Genus PHLOEOGRAPTIS, Meyrick.
95. Forewings with 9 absent	114. Genus IDIOPHANTIS, Meyrick.
Forewings with 9 present	96.

96. Forewings with 2 separate	97.
Forewings with 2 connate or stalked with 3	101.
97. Second joint of palpi smooth beneath	98.
Second joint of palpi with brush of scales beneath	162. Genus ZELOSYNE, Walsingham. (part).
98. Palpi moderate, termen of hindwings emarginate	72. Genus GUEBLA, Chrétien.
Palpi long, termen of hindwings at most sinuate	99.
99. Terminal joint of palpi shorter than second, tongue absent	149. Genus ERIPNURA, Meyrick.
Terminal joint of palpi as long as second, tongue present	100.
100. Hindwings 1, termen sinuate	69. Genus APOCRITICA, Meyrick.
Hindwings considerably over 1, termen not sinuate	116. Genus CHALINIASTIS, Meyrick.
101. Forewings with 6 to apex	178. Genus EPIBRONTIS, Meyrick.
Forewings with 6 to termen	102.
102. Terminal joint of palpi as long as second	190. Genus PANCOENIA, Meyrick.
Terminal joint of palpi shorter than second	103.
103. Terminal joint of palpi roughened, hindwings over 1	181. Genus SPHALERACTIS, Meyrick.
Terminal joint of palpi smooth, hindwings not over 1	179. Genus EPIMIMASTIS, Meyrick.
104. Forewings with 7 (or 8) absent	105.
Forewings with 7 and 8 present	168.
105. Second joint of palpi tufted beneath	106.
Second joint of palpi not tufted beneath	117.
106. Forewings with 2 and 3 stalked	107.
Forewings with 2 and 3 separate	109.
107. Terminal joint of palpi as long as second	108.
Terminal joint of palpi much shorter than second	243. Genus PROPHORULA, Meyrick.
108. Hindwings with cubital pecten	245. Genus SEMIOMERIS, Meyrick.
Hindwings without cubital pecten	263. Genus ILINGIOTIS, Meyrick (part).
109. Forewings with tufts or raised scales	110.
Forewings without tufts or raised scales	111.
110. Forewings with 3 and 4 stalked	204. Genus OESTOMORPHA, Walsingham.
Forewings with 3 and 4 approximated	203. Genus METABOLAEA, Meyrick.
111. Second joint of palpi expanded with rough hairs above towards apex	112.
Second joint of palpi not expanded with rough hairs above towards apex	114.
112. Forewings with 8 and 9 stalked	106. Genus SEMNOSTOMA, Meyrick.
Forewings with 8 and 9 separate	113.
113. Terminal joint of palpi longer than second	129. Genus HAPALONOMA, Meyrick.
Terminal joint of palpi much shorter than second	220. Genus THIOGNATHA, Meyrick.
114. Forewings with 6 and 8 stalked	229. Genus LACHNSTOLA, Meyrick.
Forewings with 6 and 8 separate	115.
115. Forewings with 3 and 4 stalked	209. Genus TORNODOXA, Meyrick.
Forewings with 3 and 4 separate	116.
116. Hindwings elongate-ovate	278. Genus HOLCOPOGON, Staudinger.
Hindwings trapezoidal	200. Genus PESSOGRAPTIS, Meyr. (part).
117. Forewings with 2 and 3 stalked	118.
Forewings with 2 and 3 normally separate	148.

118. Forewings with 6 and 8 stalked	119.
Forewings with 6 and 8 separate	121.
119. Hindwings over 1	379. Genus SCEPTEA, Walsingham.
Hindwings not over 1	120.
120. Terminal joint of palpi as long as second	378. Genus ISCHNODORIS, Meyrick.
Terminal joint of palpi shorter than second	180. Genus CRASPEDOTIS, Meyrick.
121. Second joint of palpi strongly compressed	122.
Second joint of palpi not strongly compressed	125.
122. Forewings with 9 out of 8	166. Genus SCHEMATASPIS, Meyr. (part).
Forewings with 9 separate	123.
123. Hindwings with cubital pecten	151. Genus STROBISIA, Clemens.
Hindwings without cubital pecten	124.
124. Terminal joint of palpi longer than second	153. Genus TRICYANAULA, Meyrick.
Terminal joint of palpi not longer than second	150. Genus PARELECTRA, Meyrick.
125. Antennae in ♂ with subbasal notch	126.
Antennae in ♂ without subbasal notch	127.
126. Palpi expanded with long rough hairscales	380. Genus PTILOSTONYCHIA, Walsingham.
Palpi not rough-scaled	381. Genus GLYPHIDOCERA, Walsingham.
127. Second joint of palpi with apex truncate	267. Genus PARANOEA, Walsingham.
Second joint of palpi with apex not truncate	128.
128. Forewings with 7 (or 8) and 9 stalked	129.
Forewings with 7 (or 8) and 9 separate	135.
129. Forewings with 4 out of 2	364. Genus STELECHORIS, Meyrick.
Forewings with 4 separate	130.
130. Terminal joint of palpi in ♂ expanded posteriorly with loose scales	382. Genus ANAPTILORA, Meyrick.
Terminal joint of palpi in ♂ not expanded posteriorly with loose scales	131.
131. Second joint of palpi very long, rough-scaled above	261. Genus DEIMNESTRA, Meyrick.
Second joint of palpi normal	132.
132. Forewings with 7 to apex, 8 absent	370. Genus PARALLACTIS, Meyrick.
Forewings with 7 absent, 8 to costa	133.
133. Hindwings under 1, 5 closely approximated to 4 at base	368. Genus ERYTHRIASTIS, Meyrick.
Hindwings 1 or over 1, 5 nearly parallel	134.
134. Hindwings with 5 much nearer 4 than 6	366. Genus HYGROPLASTA, Meyrick.
Hindwings with 5 not much nearer 4 than 6	369. Genus PACHNISTIS, Meyrick.
135. Second joint of palpi with rough projecting scales above	136.
Second joint of palpi without rough projecting scales above	141.
136. Hindwings with cubital pecten	137.
Hindwings without cubital pecten	138.
137. Terminal joint of palpi as long as second	260. Genus CYMOTRICA, Meyr. (part).
Terminal joint of palpi shorter than second	244. Genus NEOCHRISTA, Meyrick.
138. Terminal joint of palpi slender	139.
Terminal joint of palpi thickened, not longer than second	117. Genus HYPELICTIS, Meyr. (part).
139. Terminal joint of palpi longer than second	140.
Terminal joint of palpi shorter than second	269. Genus EUNEBRISTIS, Meyrick.

140. Forewings with 7 absent, 8 to costa	263. Genus ILINGIOTIS, Meyrick (part)
Forewings with 7 to apex, 8 absent	270. Genus ZOMEUTIS, Meyrick.
141. Hindwings with cubital pecten	142.
Hindwings without cubital pecten	143.
142. Hindwings 1	159. Genus CATOPTRISTIS, Meyrick.
Hindwings over 1	242. Genus EUPOLIS, Meyrick.
143. Terminal joint of palpi thickened and roughened anteriorly	371. Genus DEROXENA, Meyrick.
Terminal joint of palpi not thickened and roughened anteriorly	144.
144. Second joint of palpi rough anteriorly, terminal joint shorter	147. Genus SPHENOGRYPA, Meyrick.
Second joint of palpi smooth anteriorly, terminal joint nearly as long or longer	145.
145. Palpi in ♂ abnormally elongate, terminal joint with long hair-scales posteriorly	383. Genus STOEBERHINUS, Butler.
Palpi in ♂ normal	146.
146. Second joint of palpi slender, terminal joint longer	147.
Second joint of palpi rather thickened, terminal joint not longer	386. Genus AUTOSTICHA, Meyrick.
147. Terminal joint of palpi almost twice second	156. Genus SATRAPODOXA, Meyrick.
Terminal joint of palpi not almost twice second	158. Genus CHARISTICA, Meyrick (part).
148. Forewings with 3 and 4 stalked	149.
Forewings with 3 and 4 separate	150.
149. Hindwings with cubital pecten	128. Genus UNTOMIA, Busck.
Hindwings without cubital pecten	387. Genus DEMIOPHILA, Meyrick.
150. Forewings with 4 and 5 stalked or connate from angle	132. Genus MOLOPOSTOLA, Meyrick.
Forewings with 4 and 5 separate	151.
151. Terminal joint of palpi half second	152.
Terminal joint of palpi more than half second	153.
152. Hindwings with cubital pecten	250. Genus IOCHARES, Meyrick.
Hindwings without cubital pecten	208. Genus METOPLEURA, Busck.
153. Forewings with 9 out of 8	154.
Forewings with 9 separate or connate	156.
154. Hindwings with cubital pecten	137. Genus ACANTHOPHILA, Heinemann.
Hindwings without cubital pecten	155.
155. Hindwings with termen emarginate	103. Genus HIERANGELA, Meyrick.
Hindwings with termen rounded	143. Genus DESMAUCHA, Meyrick.
156. Second joint of palpi expanded at apex above, truncate	127. Genus COMMATICA, Meyrick.
Second joint of palpi not expanded nor truncate	157.
157. Terminal joint of palpi longer than second	158.
Terminal joint of palpi not longer than second	161.
158. Hindwings with cubital pecten	159.
Hindwings without cubital pecten	160.
159. Hindwings 2/3	125. Genus STAGMATURGIS, Meyrick.
Hindwings 1	135. Genus COMPOLECHIA, Meyr. (part).
160. Forewings with apex falcate	111. Genus CALLIPRORA, Meyrick.
Forewings with apex not falcate	107. Genus PLECTROCOSMA, Meyrick.
161. Forewings with 6 and 8 stalked	68. Genus LATROLOGA, Meyrick.
Forewings with 6 and 8 not stalked	162.

162. Hindwings narrower than forewings	163.
Hindwings not narrower than forewings	164.
163. Hindwings with 6 and 7 long-stalked	110. Genus POLYHYMNO, Chamb. (part).
Hindwings with 6 and 7 approximated towards base	12. Genus TIRANIMIA, Chrétien.
164. Posterior tibiae smooth	161. Genus HARPAGIDIA, Ragonot.
Posterior tibiae rough-haired above	165.
165. Hindwings with 6 and 7 well separated at base	122. Genus GLAPHYRERGA, Meyrick.
Hindwings with 6 and 7 connate	166.
166. Terminal joint of palpi as long as second	167.
Terminal joint of palpi much shorter than second	329. Genus CYMATOPLEX, Meyrick.
167. Forewings with 8 and 9 connate	328. Genus CARTERICA, Meyrick.
Forewings with 8 and 9 separate	389. Genus ENCRASIMA, Meyrick.
168. Forewings with 2 and 3 remote	169.
Forewings with 2 and 3 stalked or coincident	176.
169. Hindwings trapezoidal	170.
Hindwings elongate-ovate or ovate-lanceolate	281. Genus SYMMOCA, Hübner.
170. Forewings with 7 to apex	171.
Forewings with 7 to termen	174.
171. Tongue absent	173. Genus AREGHA, Chrétien.
Tongue developed	172.
172. Maxillary palpi porrected	376. Genus LACISTODES, Meyrick.
Maxillary palpi appressed to tongue	173.
173. Forewings with 9 and 10 stalked	330. Genus CARODISTA, Meyrick.
Forewings with 9 and 10 separate	390. Genus PROTOBATHRA, Meyrick.
174. Forewings with 8 to apex	391. Genus AMPHIGENES, Meyrick.
Forewings with 8 to costa	175.
175. Second joint of palpi strongly compressed	171. Genus AELOTROCHA, Meyrick.
Second joint of palpi not strongly compressed	377. Genus ENCOLPOTIS, Meyrick.
176. Forewings with 5 absent	307. Genus STEREMNIODES, Meyrick.
Forewings with 5 present	177.
177. Terminal joint of palpi thickened with somewhat rough scales at least at base	164. Genus ATASTHALISTIS, Meyrick.
Terminal joint of palpi not thickened with rough scales	178.
178. Second joint of palpi strongly compressed	179.
Second joint of palpi not strongly compressed	181.
179. Forewings with 9 out of 7	166. Genus SCHEMATASPIS, Meyr. (part).
Forewings with 9 separate	180.
180. Terminal joint of palpi as long as second, or longer	167. Genus ONEBALA, Walker.
Terminal joint of palpi shorter than second	168. Genus MYCONITA, Meyrick.
181. Second joint of palpi roughened or tafted beneath	182.
Second joint of palpi with appressed scales	186.
182. Forewings with 9 out of 7	373. Genus SYNDESMICA, Turner.
Forewings with 9 separate	183.
183. Second joint of palpi with dense rough projecting scales above	265. Genus HOLAXYRA, Meyrick.
Second joint of palpi without dense rough projecting scales above	184.
184. Hindwings with cubital pecten	237. Genus VAZUGADA, Walker.
Hindwings without cubital pecten	185.

185. Second joint of palpi tufted beneath	238. Genus GAESA, Walker.
Second joint of palpi not tufted beneath	283. Genus CEUTHOMADARUS, Mann.
186. Hindwings with cubital pecten	187.
Hindwings without cubital pecten	191.
187. Forewings with 9 out of 7	188.
Forewings with 9 separate	190.
188. Terminal joint of palpi shorter than second	236. Genus HYLOGRAPTIS, Meyrick.
Terminal joint of palpi not shorter than second	189.
189. Terminal joint of palpi flatly compressed	248. Genus SPHAGIOCRATES, Meyrick.
Terminal joint of palpi not flatly compressed	254. Genus MYTHOGRAPHA, Meyrick.
190. Terminal joint of palpi half second	266. Genus EPICORTHYLIS, Zeller.
Terminal joint of palpi as long as second	240. Genus MYROPHILA, Meyrick.
191. Terminal joint of palpi extremely short	272. Genus TAPHROSARIS, Meyrick.
Terminal joint of palpi not extremely short	192.
192. Forewings with 7 to apex	193.
Forewings with 7 to termen	196.
193. Hindwings with cilia 2	277. Genus EREMICA, Walsingham.
Hindwings with cilia not over 1	194.
194. Second joint of palpi triangularly expanded above towards apex	271. Genus MUSURGA, Meyrick.
Second joint of palpi not triangularly expanded above towards apex	195.
195. Terminal joint of palpi longer than second, hindwings with 7 to costa	367. Genus TORODORA, Meyrick.
Terminal joint of palpi not longer than second, hindwings with 7 to apex	374. Genus BRACHMIA Hübner.
196. Forewings with 9 out of 7	197.
Forewings with 9 separate	375. Genus APETHISTIS, Meyrick.
197. Second joint of palpi triangularly expanded with dense scales rough terminally	264. Genus SATHROGENES, Meyrick.
Second joint of palpi not triangularly expanded	372. Genus PHILARACHNIS, Meyrick.
198. Antennae I or over I	199.
Antennae under I	214.
199. Forewings with 3 and 4 absent	285. Genus TISIS, Walker.
Forewings with 3 and 4 not both absent	200.
200. Posterior tibiae with median tuft above	296. Genus HETERALCIS, Meyrick.
Posterior tibiae without median tuft above	201.
201. Forewings with 9 out of 7	202.
Forewings with 9 separate	208.
202. Forewings with 3 and 6 absent	332. Genus PROCHARISTA, Meyrick.
Forewings with 3 and 6 present	203.
203. Forewings with 2 and 3 stalked	355. Genus PERIPHORECTIS, Meyrick.
Forewings with 2 and 3 separate	204.
204. Hindwings with 5 obsolete	205.
Hindwings with 5 developed	207.
205. Forewings with 4 and 5 stalked	206.
Forewings with 4 and 5 separate	290. Genus TEUCRODOXA, Meyrick.

206. Hindwings with 6 to costa or apex	289. Genus OXYGNOSTIS, Meyrick.
Hindwings with 6 to termen	293. Genus TECHNOGRAPHA, Meyrick.
207. Hindwings with 4 and 5 stalked	315. Genus ACHORIA, Meyrick.
Hindwings with 4 and 5 separate	316. Genus NEOCORODES, Meyrick.
208. Forewings with 3 absent	311. Genus HELIANGARA, Meyrick.
Forewings with 3 present	209.
209. Forewings with 2 and 3 stalked	210.
Forewings with 2 and 3 separate	211.
210. Hindwings with 4 absent	358. Genus ASMENISTIS, Meyrick.
Hindwings with 4 present	303. Genus HYPTIASTIS, Meyrick.
211. Hindwings with 3 and 4 remote	274. Genus OECIA, Walsingham.
Hindwings with 3 and 4 connate or stalked	212.
212. Terminal joint of palpi in ♂ obsolete	291. Genus ALCIPHANES, Meyrick.
Terminal joint of palpi in ♂ developed	213.
213. Second joint of palpi with long tuft beneath	317. Genus ENTHETICA, Meyrick.
Second joint of palpi without tuft	275. Genus AMBLOMA, Walsingham.
214. Hindwings with 3 (or 4) absent (at least in ♂)	215.
Hindwings with 3 and 4 present	219.
215. Forewings with 9 out of 7	216.
Forewings with 9 separate	218.
216. Hindwings with 3 and 5 connate or stalked	217.
Hindwings with 3 and 5 remote	337. Genus IDIOPTERYX, Walsingham.
217. Terminal joint of palpi longer than second	338. Genus HYPEROCHTHA, Meyrick.
Terminal joint of palpi not longer than second	346. Genus GONAEPA, Walker.
218. Forewings with 3 absent	195. Genus PALINTROPA, Meyrick.
Forewings with 3 present	230. Genus PAPPOPHORUS, Walsingham.
219. Hindwings with 2 and 3 stalked (? ♂ only)	259. Genus PARISTHMIA, Meyrick.
Hindwings with 2 and 3 separate	220.
220. Hindwings with 3 and 4 separate	221.
Hindwings with 3 and 4 connate or stalked	280.
221. Hindwings with 5 and 6 absent	6. Genus NEALYDA, Dietz.
Hindwings with 5 and 6 not both absent	222.
222. Terminal joint of palpi with posterior scale-projection or loose scales	223.
Terminal joint of palpi without posterior scale-projection or loose scales	227.
223. Forewings with 3 and 4 stalked	90. Genus PARASTEGA, Meyrick.
Forewings with 3 and 4 separate	224.
224. Hindwings with cubital pecten	239. Genus OXYCRYPTIS, Meyrick.
Hindwings without cubital pecten	225.
225. Hindwings with 6 and 7 stalked or nearly approximated	196. Genus CHELARIA, Haworth.
Hindwings with 6 and 7 nearly parallel	226.
226. Hindwings with 5 straight, parallel to 4	38. Genus EMPEDAULA, Meyrick.
Hindwings with 5 curved, approximated	19. Genus PROSELOTIS, Meyrick.
227. Forewings with tufts of scales on surface	228.
Forewings without tufts of scales on surface	235.

228. Second joint of palpi tufted beneath	229.
Second joint of palpi not tufted beneath	232.
229. Forewings with 2 remote, 3 and 4 connate	41. Genus LEUCE, Chambers.
Forewings with 2-4 approximated	230.
230. Terminal joint of palpi stout, roughened anteriorly	42. Genus PACHYGENEIA, Meyrick.
Terminal joint of palpi slender	231.
231. Second joint of palpi with rough tuft	43. Genus LEPTOGENEIA, Meyrick.
Second joint of palpi with median and apical triangular tufts	55. Genus COLONANTHES, Meyrick.
232. Forewings with 11 absent	51. Genus AGNIPPE, Chambers.
Forewings with 11 present	233.
233. Hindwings with 6 and 7 nearly parallel	76. Genus AROGALEA, Walsingham.
Hindwings with 6 and 7 closely approximated or stalked	234.
234. Forewings with 7 and 8 out of 6	78. Genus SCHISTOPHILA, Chrétien.
Forewings with 6 separate, or out of 7 near base	81. Genus TELPHUSA, Chambers.
235. Basal joint of antennae with pecten	236.
Basal joint of antennae without pecten	239.
236. Hindwings with 6 and 7 remote	237.
Hindwings with 6 and 7 stalked	238.
237. Hindwings with apex very long-produced	5. Genus APATETRIS, Staudinger.
Hindwings with apex somewhat produced	9. Genus METANARSIA, Staudinger.
238. Forewings with 9 out of 7	4. Genus OECOCECIS, Guenée.
Forewings with 9 separate	31. Genus SITOTROGA, Heinemann.
239. Second joint of palpi with tuft or rough projecting hairs beneath	240.
Second joint of palpi without tuft or rough projecting hairs beneath	255.
240. Terminal joint of palpi (at least in ♂) very short, more or less concealed	241.
Terminal joint of palpi moderate or long, exposed	243.
241. Hindwings with 6 and 7 separate	8. Genus PARANARSIA, Ragonot.
Hindwings with 6 and 7 stalked	242.
242. Palpi very long, porrected	15. Genus CHILOSELAPHUS, Mann.
Palpi in ♂ moderate, in ♀ recurved	193. Genus ANARSIA, Zeller (part).
243. Forewings with 7 and 8 out of 6	244.
Forewings with 6 separate, or 6 and 7 out of 8	248.
244. Terminal joint of palpi twice second	214. Genus HYODECTIS, Meyrick.
Terminal joint of palpi not longer than second	245.
245. Forewings with 3 absent	29. Genus STEREOMITA, Braun.
Forewings with 3 present	246.
246. Second joint of palpi beneath with long rough spreading hairs	25. Genus ISOPHRICHTIS, Meyrick.
Second joint of palpi beneath with compact tuft of dense scales	247.
247. Hindwings with 6 and 7 connate	30. Genus PTYCERATA, Ely.
Hindwings with 6 and 7 remote	23. Genus PYCNOTOLA, Meyrick.
248. Terminal joint of palpi roughened anteriorly	249.
Terminal joint of palpi not roughened anteriorly	250.
249. Hindwings with 5 nearer 4	44. Genus PHOTODOTIS, Meyrick.
Hindwings with 5 nearer 6	28. Genus DORYCNOPA, Lower.
250. Forewings with 9 out of 7	207. Genus STACHYOSTOMA, Meyrick.
Forewings with 9 separate	251.

LEPIDOPTERA HETEROCEA

251. Hindwings with 5 nearer 4	252.
Hindwings with 5 nearer 6	253.
252. Forewings with 4 absent	48. Genus OXYLECHIA, Meyrick.
Forewings with 4 present	56. Genus COMPSOSARIS, Meyrick.
253. Forewings with 7 and 8 out of 6	254.
Forewings with 6 separate	22. Genus CATAMECES, Turner.
254. Second joint of palpi beneath with long rough spreading hairs	26. Genus PALTODORA, Meyrick.
Second joint of palpi beneath with apical tuft of dense hairscales	24. Genus MEGACRASPEDUS, Zeller.
255. Second joint of palpi extremely long, straight, prorected, with long rough hairs above	256.
Second joint of palpi not so formed	257.
256. Hindwings with cubital pecten	231. Genus RHYNCHOTONA, Meyrick.
Hindwings without cubital pecten	16. Genus TRICHEMBOLA, Meyrick.
257. Second joint of palpi with rough projecting scales above towards apex	258.
Second joint of palpi without rough projecting scales above towards apex	262.
258. Second joint of palpi rough-scaled beneath	259.
Second joint of palpi with appressed scales	260.
259. Terminal joint of palpi half second	221. Genus CROSSEBELA, Meyrick.
Terminal joint of palpi as long as second or nearly	35. Genus IULOTA, Meyrick.
260. Terminal joint of palpi 1/5 of second	217. Genus HOLCOPHORA, Staudinger.
Terminal joint of palpi as long as second or not much shorter	261.
261. Forewings with 6 out of 8	34. Genus PYCNODYTIS, Meyrick.
Forewings with 6 separate	218. Genus SYMBOLISTIS, Meyrick.
262. Forewings with 4 absent	263.
Forewings with 4 present	264.
263. Forewings with 7 and 8 out of 6	46. Genus STENOLECHIA, Meyrick.
Forewings with 6 separate	47. Genus PARACHRONISTIS, Meyrick.
264. Hindwings with 6 obsolete	94. Genus THOLEROSTOLA, Meyrick.
Hindwings with 6 present	265.
265. Hindwings with 6 and 7 stalked or connate or closely approximated	266.
Hindwings with 6 and 7 separate	274.
266. Forewings with 8 out of 6	267.
Forewings with 8 out of 7	268.
267. Terminal joint of palpi longer than second	60. Genus NESOLECHIA, Meyrick.
Terminal joint of palpi shorter than second	39. Genus PRAGMATODES, Walsingham.
268. Forewings with 9 out of 7	269.
Forewings with 9 separate	270.
269. Forewings with 6 out of 7	82. Genus SYNCOPACMA, Meyrick.
Forewings with 6 separate	136. Genus CATALEXIS, Walsingham.
270. Second joint of palpi rough beneath	80. Genus PITHANURGA, Meyrick.
Second joint of palpi with appressed scales	271.
271. Terminal joint of palpi longer than second	272.
Terminal joint of palpi not longer than second	273.

272. Forewings with 6 out of 8	32. Genus ACRAEOLOGA, Meyrick.
Forewings with 6 separate	120. Genus INOTICA, Meyrick.
273. Hindwings with termen excised, prominent below	7. Genus NEODACTYLOTA, Busck.
Hindwings with termen sinuate or emarginate	79. Genus LEURONOMA, Meyrick.
274. Forewings with 5 out of 6	36. Genus PSAMATHOCRITA, Meyrick.
Forewings with 5 separate	275.
275. Terminal joint of palpi much shorter than second, thickened with scales	276.
Terminal joint of palpi not much shorter than second	277.
276. Forewings with 6 separate	17. Genus EPIPARASIA, Rebel.
Forewings with 7 and 8 out of 6	18. Genus METZNERIA, Zeller.
277. Forewings with 4 and 5 long-stalked	20. Genus MERIDORMA, Meyrick.
Forewings with 4 and 5 separate	278.
278. Terminal joint of palpi much thickened and roughened with scales	21. Genus PITYOCONA, Meyrick.
Terminal joint of palpi not much thickened and roughened with scales	279.
279. Hindwings in ♂ with subcostal pecten of stiff flat bristles towards base	40. Genus GLAUCE, Chambers.
Hindwings without such pecten	37. Genus ARISTOTELIA, Hübner.
280. Hindwings usually lanceolate, 5 and 6 (or 7) absent	45. Genus HELICE, Chambers.
Hindwings usually trapezoidal, 5 and 6 not both absent	281.
281. Forewings with one or more veins absent	282.
Forewings with all veins present	291.
282. Forewings with 2 and 3 coincident (or 3 or 4 or 5 absent).	283.
Forewings with 2-5 present	290.
283. Second joint of palpi tufted beneath	284.
Second joint of palpi not tufted beneath	285.
284. Hindwings with 6 and 7 stalked	233. Genus RHADINOPHYLLA, Turner.
Hindwings with 6 and 7 parallel	212. Genus THRIOPHORA, Meyrick.
285. Hindwings with 6 obsolete	286.
Hindwings with 6 present	287.
286. Second joint of palpi with expanded scales above	54. Genus HAPALOSARIS, Meyrick.
Second joint of palpi without expanded scales above	49. Genus TOSCA, Heinrich.
287. Hindwings with 6 and 7 approximated or stalked	288.
Hindwings with 6 and 7 nearly parallel	183. Genus PRODOSIARCHA, Meyrick.
288. Hindwings with 6 and 7 stalked	289.
Hindwings with 6 and 7 approximated	177. Genus TRITADELPHA, Meyrick.
289. Terminal joint of palpi longer than second	104. Genus SYMPHANACTIS, Meyrick.
Terminal joint of palpi shorter than second	70. Genus CONIOGYRA, Meyrick.
290. Forewings with 6 absent	201. Genus APOTACTIS, Meyrick.
Forewings with 6 present	172. Genus DECATOPSEUSTIS, Meyr. (♂).
291. Terminal joint of palpi with projecting scales or roughened posteriorly	292.
Terminal joint of palpi not with projecting scales or roughened posteriorly	304.
292. Terminal joint of palpi longer than second	293.
Terminal joint of palpi not longer than second	295.

293. Forewings with 3 and 4 stalked	210. Genus TITUACIA, Walker.	
Forewings with 3 and 4 separate		294.
294. Forewings with scaletufts	197. Genus HAPLOCHELA, Meyrick.	
Forewings without scaletufts	198. Genus SCLEROGRAPTIS, Meyrick.	
295. Forewings with 2 and 3 stalked		296.
Forewings with 2 and 3 separate		299.
296. Terminal joint of palpi as long as second		297.
Terminal joint of palpi shorter than second		298.
297. Hindwings with 5 parallel	117. Genus HYPELICTIS, Meyr. (part).	
Hindwings with 5 rather approximated to 4	163. Genus TEUCHOPHANES, Meyrick.	
298. Hindwings with cubital pecten	256. Genus PACHYSARIS, Meyrick.	
Hindwings without cubital pecten	268. Genus COTYLOSCIA, Meyrick.	
299. Hindwings with cubital pecten		300.
Hindwings without cubital pecten		301.
300. Forewings with tufts of scales	138. Genus SOROTACTA, Meyrick.	
Forewings without tufts of scales	134. Genus COBOSTOMA, Meyrick.	
301. Forewings with tufts of scales	206. Genus PORPODRYAS, Meyrick.	
Forewings without tufts of scales		302.
302. Second joint of palpi with very long rough hairscales beneath	89. Genus LOCHARCHA, Meyrick.	
Second joint of palpi without very long rough hairscales beneath		303.
303. Hindwings with 6 and 7 approximated or stalked	87. Genus MOMETA, Durrant.	
Hindwings with 6 and 7 parallel	92. Genus THYMOSOPHA, Meyrick.	
304. Forewings with 2 and 3 stalked or connate (seldom nearly approximated from angle)		305.
Forewings with 2 and 3 more or less remote		330.
305. Second joint of palpi with apical tuft beneath		306.
Second joint of palpi without apical tuft beneath		309.
306. Terminal joint of palpi shorter than second		307.
Terminal joint of palpi as long as second		308.
307. Second joint of palpi straight, porrected	232. Genus AGELIARCHIS, Meyrick.	
Second joint of palpi curved, ascending	241. Genus BROCHOMETIS, Meyrick.	
308. Hindwings with cubital pecten	235. Genus DICHOMBRIS, Hübner.	
Hindwings without cubital pecten	234. Genus TELEPHILA, Meyrick.	
309. Hindwings with cubital pecten		310.
Hindwings without cubital pecten		316.
310. Basal joint of antennae with slight pecten	225. Genus AROTRIA, Meyrick.	
Basal joint of antennae without pecten		311.
311. Forewings with 9 out of 7	246. Genus CATELAPHRIS, Meyrick.	
Forewings with 9 separate		312.
312. Second joint of palpi with scales roughly expanded above towards apex		313.
Second joint of palpi with scales not roughly expanded above towards apex		315.
313. Terminal joint of palpi half second	251. Genus EPORGASTIS, Meyrick.	
Terminal joint of palpi more than half second		314.

314. Terminal joint of palpi rather thickened with loose scales, somewhat shorter than second	252. Genus CARBATINA, Meyrick.
Terminal joint of palpi slender, as long as second	260. Genus CYMOTRICA, Meyr. (part).
315. Antennae in ♂ moderately ciliated	154. Genus ZALITHIA, Meyrick.
Antennae in ♂ simple	160. Genus CERYCANGELA, Meyrick.
316. Second joint of palpi greatly dilated towards apex with appressed hairs	184. Genus CORYNÆA, Turner.
Second joint of palpi not greatly dilated towards apex with appressed hairs	317.
317. Forewings with 2 and 3 out of 4	176. Genus METEORISTIS, Meyrick.
Forewings with 4 separate	318.
318. Second joint of palpi with appressed scales beneath	319.
Second joint of palpi roughened beneath	326.
319. Second joint of palpi with scales expanded towards apex above	320.
Second joint of palpi with scales not expanded towards apex above	321.
320. Hindwings with 6 and 7 parallel	188. Genus HETEROZANCLA, Turner.
Hindwings with 6 and 7 connate or stalked	273. Genus TRICHOTAPHE, Clemens.
321. Forewings with 6 to costa	155. Genus HYPERECTA, Meyrick.
Forewings with 6 to apex	158. Genus CHARISTICA, Meyr. (part).
Forewings with 6 to termen	322.
322. Antennae in ♀ thickened with rough scales	169. Genus THELYASCETA, Meyrick.
Antennae in ♀ not thickened with rough scales	323.
323. Forewings with 7 and 8 out of 6	182. Genus LEXIARCHA, Meyrick.
Forewings with 6 separate	324.
324. Hindwings with 6 absent	186. Genus SEMOCHARISTA, Meyrick.
Hindwings with 6 present	325.
325. Terminal joint of palpi as long as second	152. Genus HOLOPHYSIS, Walsingham.
Terminal joint of palpi much longer than second	157. Genus EUZONOMACHA, Meyrick.
326. Hindwings with 6 and 7 stalked	327.
Hindwings with 6 and 7 separate	329.
327. Second joint of palpi long, rough-scaled above and beneath, terminal joint very short	215. Genus APONOEA, Walsingham.
Second joint of palpi not so formed	328.
328. Forewings with large subdorsal scale-tuft	191. Genus ORTHOPTILA, Meyrick.
Forewings without large subdorsal scale-tuft	185. Genus HEMIARCHA, Meyrick.
329. Second joint of palpi rough-scaled above	189. Genus EPHELICTIS, Meyrick.
Second joint of palpi not rough-scaled above	187. Genus PROTOLECHIA, Meyrick.
330. Second joint of palpi (♂) with expansible tuft of hairs above	59. Genus EUCORDYLEA, Dietz.
Second joint of palpi without expansible tuft of hairs above	331.
331. Forewings with 3 and 4 stalked	332.
Forewings with 3 and 4 not stalked	335.
332. Forewings with 7 and 8 out of 6	65. Genus TRYPANISMA, Clemens.
Forewings with 6 separate, or 6 and 7 out of 8	333.
333. Second joint of palpi with scales expanded above towards apex	223. Genus BRACHYACMA, Meyrick.
Second joint of palpi with scales not expanded above towards apex	334.
334. Forewings in ♂ with costal fold and hair-pencil beneath	91. Genus STEGASTA, Meyrick.
Forewings in ♂ without costal fold and hair-pencil beneath	205. Genus CRASIMORPHA, Meyrick.

335. Hindwings with 5 absent	336.
Hindwings with 5 present	339.
336. Palpi with second joint tufted beneath	324. Genus LIOCLEPTA, Meyrick.
Palpi with second joint not tufted beneath	337.
337. Second joint of palpi rough anteriorly towards apex, especially in ♂	141. Genus ANTHINORA, Meyrick.
Second joint of palpi not rough	338.
338. Terminal joint of palpi (♂) minute, rudimentary	326. Genus PHTHORACMA, Meyrick.
Terminal joint of palpi moderate	249. Genus DEOCLONA, Busck.
339. Second joint of palpi tufted beneath	340.
Second joint of palpi not tufted beneath	355.
340. Terminal joint of palpi in ♂ concealed or nearly so	341.
Terminal joint of palpi exposed	342.
341. Terminal joint of palpi moderately long	194. Genus DOLEROTRICA, Meyrick.
Terminal joint of palpi in ♂ very short or obsolete	193. Genus ANARSIA, Zeller (part).
342. Forewings with tufts of scales	343.
Forewings without tufts of scales	345.
343. Hindwings with 6 and 7 long-stalked	344.
Hindwings with 6 and 7 closely approximated	139. Genus ALSODRYAS, Meyrick.
344. Terminal joint of palpi as long as second	77. Genus ANTHISTARCHA, Meyrick.
Terminal joint of palpi much longer than second	200. Genus PESSOGRAPTIS, Meyr. (part).
345. Hindwings with 6 and 7 stalked or approximated	346.
Hindwings with 6 and 7 remote, parallel	354.
346. Second joint of palpi with basal tuft (?)	27. Genus ZIZYPHIA, Chrétien.
Second joint of palpi with apical tuft	347.
347. Hindwings with cubital pecten	348.
Hindwings without cubital pecten	349.
348. Second joint of palpi with fringe of long rough hairs beneath	130. Genus ETHIOSTOMA, Meyrick.
Second joint of palpi with compact apical tuft	228. Genus ACIBOLOGA, Meyrick.
349. Terminal joint of palpi longer than second	100. Genus NOTHRIS, Hübner.
Terminal joint of palpi not longer than second	350.
350. Hindwings 1, 5 nearly parallel	351.
Hindwings under 1, 5 closely approximated to 4 at base	226. Genus EMPALACTIS, Meyrick.
351. Apex of forewings prominent or subfalcate	352.
Apex of forewings not prominent or subfalcate	353.
352. Hindwings with 6 and 7 approximated at base	113. Genus ANASPHALTIS, Meyrick.
Hindwings with 6 and 7 stalked	112. Genus SOPHRONIA, Hübner.
353. Forewings with 2 from near angle, palpi with several black rings	213. Genus DACTYLETHRA, Meyrick.
Forewings with 2 remote, palpi without black rings	227. Genus XEROMETRA, Meyrick.
354. Second joint of palpi more or less rough-scaled above	95. Genus TECIA, Strand (part).
Second joint of palpi not rough above	219. Genus STRENIASTIS, Meyrick.
355. Second joint of palpi with expanded scales towards apex above	356.
Second joint of palpi without expanded scales towards apex above	363.
356. Forewings with 4 and 5 connate	133. Genus PROMOLOPICA, Meyrick.
Forewings with 4 and 5 separate	357.
357. Terminal joint of palpi 1/3 of second or less	358.
Terminal joint of palpi as long as second or little shorter	360.

358. Hindwings with 6 and 7 parallel	95. Genus TECIA, Strand (part).
Hindwings with 6 and 7 stalked or approximated	359.
359. Second joint of palpi above with rough hairs longer towards base	222. Genus MESOPHLEPS, Hübner.
Second joint of palpi above with rough scales towards apex only	224. Genus EPIMESOPHLEPS, Rebel.
360. Hindwings with cubital pecten	361.
Hindwings without cubital pecten	362.
361. Forewings with tufts of scales	140. Genus ANACAMPSIS, Curtis.
Forewings without tufts of scales	131. Genus BATTARISTIS, Meyrick.
362. Terminal joint of palpi thickened with scales	73. Genus PARAPSECTRIS, Meyrick.
Terminal joint of palpi slender	83. Genus MACRENCHES, Meyrick.
363. Forewings with 8 out of 6	364.
Forewings with 8 out of 7	370.
364. Hindwings with 6 obsolete	365.
Hindwings with 6 developed	366.
365. Forewings with 4 and 5 connate or stalked	50. Genus EVIPPE, Chambers.
Forewings with 4 and 5 separate	52. Genus SMENODOCA, Meyrick.
366. Forewings with 9 out of 6	63. Genus EPHYSTERIS, Meyrick.
Forewings with 9 separate	367.
367. Hindwings with 6 and 7 stalked	368.
Hindwings with 6 and 7 somewhat approximated or nearly parallel	369.
368. Forewings with tufts of scales	66. Genus ISTRIANIS, Meyrick.
Forewings without tufts of scales	67. Genus EPITHECTIS, Meyrick.
369. Forewings with tufts of scales	58. Genus RECURVARIA, Haworth.
Forewings without tufts of scales	97. Genus PHTHORIMAEA, Meyr. (part).
370. Palpi blade-like, sharp-edged	170. Genus PROSTOMEUS, Busck.
Palpi not so formed	371.
371. Forewings with 9 out of 7	62. Genus BATENIA, Chrétien.
Forewings with 9 separate	372.
372. Hindwings with 4 and 5 out of 3	74. Genus SYMBATICA, Meyrick.
Hindwings with 5 separate	373.
373. Tongue absent	276. Genus APOTISTATUS, Walsingham.
Tongue present	374.
374. Second joint of palpi rough or furrowed beneath	375.
Second joint of palpi smooth	386.
375. Forewings with two large tufts in disc anteriorly	75. Genus DISSOPTILA, Meyrick.
Forewings without two large tufts in disc anteriorly	376.
376. Hindwings with 6 and 7 approximated or stalked	377.
Hindwings with 6 and 7 nearly parallel	380.
377. Basal joint of antennae with pecten	86. Genus PLATYEDRA, Meyrick.
Basal joint of antennae without pecten	378.
378. Antennae thickened with rough scales on basal half	84. Genus APOTHETOCA, Meyrick.
Antennae not thickened with rough scales on basal half	379.
379. Forewings with 2 from 3/4	101. Genus ANISOPLACA, Meyrick.
Forewings with 2 from towards angle	85. Genus GELECHIA, Hübner.
380. Second joint of palpi with projecting lateral bristles	98. Genus POGOCHAETIA, Standinger.
Second joint of palpi without projecting lateral bristles	381.

LEPIDOPTERA HETEROCHERA

381. Forewings with 6 to apex	162. Genus <i>ZELOSYNE</i> , Walsingham. (part).
Forewings with 6 to termen	382.
382. Second joint of palpi brushlike and furrowed beneath	383.
Second joint of palpi not brushlike and furrowed beneath	384.
383. Terminal joint of palpi much shorter than second, anterior edge serrate	96. Genus <i>GNORIMOSCHEMA</i> , Busck.
Terminal joint of palpi about as long as second, with appressed scales	97. Genus <i>PHTHORIMAEA</i> , Meyr. (part).
384. Forewings with tufts of scales.	385.
Forewings without tufts of scales	99. Genus <i>SAROTORNA</i> , Meyrick.
385. Terminal joint of palpi longer than second	57. Genus <i>PHYLOPATRIS</i> , Meyrick.
Terminal joint of palpi shorter than second	61. Genus <i>EXOTELEIA</i> , Wallengren.
386. Forewings with 7 and 8 very long-stalked (or coincident)	110. Genus <i>POLYHYNNO</i> , Chamb. (part).
Forewings with 7 and 8 normal	387.
387. Terminal joint of palpi longer than second	388.
Terminal joint of palpi not longer than second	389.
388. Hindwings lanceolate	279. Genus <i>STRENOPHILA</i> , Meyrick.
Hindwings trapezoidal	389.
389. Hindwings with cubital pecten	135. Genus <i>COMPSOLECHIA</i> , Meyrick.
Hindwings without cubital pecten	390.
390. Hindwings with 6 and 7 stalked	121. Genus <i>STOMOPTERYX</i> , Heinemann.
Hindwings with 6 and 7 approximated	391.
391. Hindwings pointed	142. Genus <i>HARPOGRAPTRIS</i> , Meyrick.
Hindwings obtuse	123. Genus <i>MELITOXESTIS</i> , Meyrick.
392. Hindwings with 6 and 7 subparallel	93. Genus <i>PHLOEOCRCIS</i> , Chrétien.
Hindwings with 6 and 7 approximated or stalked	393.
393. Hindwings with termen sinuate	394.
Hindwings with termen not sinuate	280. Genus <i>OEGOCONIA</i> , Stainton.
394. Hindwings in ♂ greatly modified	88. Genus <i>TABERNILLAEA</i> , Walsingham.
Hindwings in ♂ normal	395.
395. Forewings with 6 out of 8	396.
Forewings with 6 separate	397.
396. Hindwings with 3 and 4 long-stalked	118. Genus <i>SCINDALMOTA</i> , Turner.
Hindwings with 3 and 4 connate	174. Genus <i>LEOBATUS</i> , Walsingham.
397. Second joint of palpi with scales rather rough above, terminal joint short	216. Genus <i>RHYNCHOPACHA</i> , Standinger.
Second joint of palpi with scales not rough above	398.
398. Forewings with 2 from angle	119. Genus <i>STIPHOSTOLA</i> , Meyrick.
Forewings with 2 from well before angle	399.
399. Hindwings over 1	400.
Hindwings not over 1	401.
400. Terminal joint of palpi half second	165. Genus <i>PERIORISTICA</i> , Walsingham.
Terminal joint of palpi more than half second	175. Genus <i>ACOMPSIA</i> , Hübner.
401. Terminal joint of palpi as long as second	402.
Terminal joint of palpi much shorter than second	71. Genus <i>COUDIA</i> , Chrétien.
402. Second joint of palpi laterally compressed	145. Genus <i>DIASTALTICA</i> , Walsingham.
Second joint of palpi not laterally compressed	172. Genus <i>DECATOPSEUSTIS</i> , Meyr. (♀).

Group 1 (Apatetris type)

A specialised development from the *Aristotelia* group, with pronounced tendency to the possession of a basal pecten in the antennae (rare otherwise in this family), and to characteristic excision of hindwings into two lobes; in extreme forms the palpi are much reduced or aborted. The larvae usually feed in galls on stems.

I. GENUS AMBLYPALPIS, RAGONOT

Amblypalpis, Ragonot, Bull. Soc. Ent. Fr. p. 209 (1885). — Type : *A. olivierella*, Ragonot.

Characters. — Head with appressed scales; tongue obsolete. Antennæ 4/5, basal joint dilated, without pecten. Labial palpi forming a minute porrected triangular tuft, hardly passing the face. Maxillary palpi obsolete. Forewings with 7 absent. Hindwings elongate-trapezoidal, termen strongly emarginate beneath very prolonged apex; 2-7 all separate.

Remarks. — I have not seen this genus, but the affinity to the two following is obvious.

Geographical distribution of species. — North African.

Larva feeding in galls on stems.

Foodplant *Tamarix*.

1. *A. olivierella*, Ragonot, Bull. Soc. Ent. Fr. p. 209 (1885). Algeria.

2. GENUS SICERA, CHRÉTIEN

Sicera, Chrétien, Bull. Soc. Ent. Fr. p. 144 (1908). — Type : *S. albidella*, Chrétien.

Characters. — Head smooth; ocelli posterior. Antennæ pubescent, basal joint enlarged, with pecten of scales. Labial palpi very short, with loosely appressed scales, hardly pointed. Maxillary palpi obsolete. Posterior tibiae clothed with long hairs. Forewings with 2 remote from angle, 4 absent, 7 absent, 8 and 9 closely approximated at base. Hindwings very narrow, apex acutely produced, termen rectangularly emarginate beneath apex; 2-4 separate, 5 and 6 approximated, 6 and 7 parallel.

Remarks. — Also unknown to me.

Geographical distribution of species. — North African.

Larva unknown.

1. *S. albidella*, Chrétien, Bull. Soc. Ent. Fr. p. 144 (1908). Algeria.

3. GENUS SCLEROCECIS, CHRÉTIEN

Sclerocecis, Chrétien, Bull. Soc. Ent. Fr. p. 142 (1908). — Type : *S. pulverosella*, Chrétien.

Characters. — Head smooth; ocelli posterior; tongue obsolete. Antennæ 2/3, pubescent, basal joint enlarged, with strong pecten of scales. Labial palpi short, drooping, scaled, hardly pointed. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs. Forewings with 2 from towards angle, 5 almost connate with 8, 6 out of 8, 7 absent, 11 from middle. Hindwings under 1, elongate-

trapezoidal, apex produced, hardly acute, termen oblique, slightly bisinuate; 2-4 separate, 5 approximated to 6, cell open between 4 and 5, 7 separate.

Remarks. — This and the two preceding genera are remarkable for the great reduction of the labial palpi, which is a quite abnormal feature in this family; it appears to have some association with the gall-feeding habit.

Geographical distribution of species. — North African.

Larva feeding in stem-galls of variable form.

Pupa not protruded from gall in emergence.

Foodplant *Limoniastrum* (*Plumbaginaceae*).

1. *S. pulverosella*, Chrétien, Bull. Soc. Ent. Fr. p. 142 (1908). Algeria.

4. GENUS OECOCECIS, GUENÉE

Oecocecis, Guenée, Ann. Soc. Ent. Fr. p. 14 (1870). — Type : *Oe. guyonella*, Guenée.

Characters. — Head with appressed hairs; ocelli small, just posterior; tongue absent. Antennae 3/4, in ♂ slender, serrulate, shortly ciliated, basal joint rather short, stout, with strong dense pecten. Labial palpi moderate, curved, subascending, second joint with dense rough projecting tuft beneath, terminal joint short, pointed, projecting little from hairs of second. Maxillary palpi obsolete. Posterior tibiae clothed with rough projecting hairs above and beneath. Forewings with 2-4 remote, 5 connate with 7 from angle, 6 out of 7, 7 to costa, 8 and 9 out of 7, 11 from 2/3. Hindwings 1, elongate-trapezoidal, apex somewhat produced, tolerably pointed, termen slightly bisinuate, cilia 1; 2-5 remote, nearly parallel, 4 from angle, 6 and 7 stalked.

Remarks. — Probably a development of *Apatebris*.

Geographical distribution of species. — Mediterranean.

Larva very active, feeding in fusiform or subglobose galls on stems.

Foodplant *Limoniastrum* (*Plumbaginaceae*).

1. *Oe. guyonella*, Guenée, Ann. Soc. Ent. Fr. p. 14, pl. VII, f. 1-11 (1870). — S. Europe, N. Africa, Pl. I, Fig. 4. Syria.

5. GENUS APATETRIS, STAUDINGER

Apatebris, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 316 (1880). — Type : *A. mirabella*, Staudinger.

Dactylocta, Snellen, Tijdschr. v. Ent. Vol. 19, p. 23 (1875) (praeocc.). — Type : *A. kinkerella*, Snellen.

Epiphthora, Meyrick, Trans. N. Zeal. Inst. Vol. 20, p. 77 (1888). — Type : *A. melanombra*, Meyrick.

Calyptrotis, Meyrick, Ent. M. Mag. Vol. 27, p. 56 (1891). — Type : *A. alpitodes*, Meyrick.

Didactylocta, Walsingham, Proc. Zool. Soc. Lond. p. 522 (1891). — Type : *A. kinkerella*, Snellen.

Stenopherna, Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 78 (1901). — Type : *A. chionocephala*, Lower.

Proactica, Walsingham, Ent. M. Mag. Vol. 40, p. 268 (1904). — Type : *A. halimilignella*, Walsingham.

Cecidophaga, Walsingham, Ent. M. Mag. Vol. 47, p. 189 (1911). — Type : *A. tamaricicola*, Walsingham.

Characters. — Head smooth ; ocelli posterior ; tongue developed. Antennæ 2/3 — 4/5, in ♂ simple or ciliated, basal joint elongate, with dense strong pecten. Labial palpi rather long, curved, ascending, second joint more or less rough-scaled or expanded with projecting scales beneath towards apex, terminal joint shorter than second, sometimes very short, slightly thickened, roughened anteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above and beneath. Forewings with 1 b furcate, 2 remote, 7 and 8 stalked, 7 to costa, seldom 7 and 8 out of 6, or 6 out of 7 near base, 11 from middle. Hindwings under 1, narrow-trapezoidal, termen abruptly or acutely emarginate beneath acutely produced apex, cilia 2-4 ; 3 and 4 rather approximated, 5 tolerably parallel, 6 and 7 rather approximated towards base.

Remarks. — There is some variability of structure, but the genus forms a natural whole and does not need subdivision. The curious hindwings often afford good specific characters, in the more pronounced forms having the apical projection very long or the subapical excision so deep that the wing appears to form a prominence below it.

Geographical distribution of species. — At present 43 species are known, of which number there are 11 African, 3 European, 1 Asia Minor, 1 Indian, 26 Australian, and 1 New Zealand. This is a singular distribution ; as the species are generally small, inconspicuous, and retired in habit, it may be modified by future discoveries, but is probably explicable as follows ; the species have a distinct predilection for arid and xerophytic regions, and it seems likely that Indian collectors have been disposed to neglect country of this character, on the mistaken assumption that it is unproductive, and that the genus will ultimately prove to be fairly represented in India, which should be its original home.

Larva sometimes apodal, usually feeding in galls on stems, but the New Zealand species is a leaf-miner.

Foodplants *Chenopodiaceæ*, *Tamaricaceæ*; the New Zealand species on *Olearia (Compositæ)*.

- | | |
|---|----------------------|
| 1. <i>A. mirabella</i> , Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 317 (1880). | Asia Minor. |
| 2. <i>A. halimilignella</i> , Walsingham, Ent. M. Mag. Vol. 40, p. 269 (1904). | Algeria. |
| <i>acutella</i> , Chrétien, Bull. Soc. Ent. Fr. p. 165 (1908). | |
| 3. <i>A. echiochilonella</i> , Chrétien, Bull. Soc. Ent. Fr. p. 201 (1908). | Algeria. |
| 4. <i>A. leucoglypta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 117 (1918). | India. |
| 5. <i>A. albiramis</i> , Meyrick, ibidem, Vol. 3, p. 51 (1923). | Egypt. |
| 6. <i>A. alphitodes</i> , Meyrick, Ent. M. Mag. Vol. 27, p. 56 (1891). | Algeria. |
| 7. <i>A. anisaula</i> , Meyrick, Exot. Microlep. Vol. 2, p. 424 (1921). | N. Australia. |
| 8. <i>A. thyellias</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 262 (1904). | Victoria. |
| 9. <i>A. delochorda</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 42, p. 237 (1918). | S. Australia. |
| 10. <i>A. lemurella</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 262 (1904). | New South Wales. |
| 11. <i>A. niphaula</i> , Meyrick, ibidem, Vol. 29, p. 263 (1904). | Tasmania. |
| 12. <i>A. hexagramma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 424 (1921). | Queensland. |
| 13. <i>A. leucogaea</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 65 (1921). | Rhodesia. |
| 14. <i>A. psychrodes</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 263 (1904). | New South Wales. |
| 15. <i>A. leptoconia</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 110 (1919). | New South Wales. |
| 16. <i>A. miarodes</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 263 (1904). | S. E. Australia. |
| <i>psoloistica</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 109 (1919). | Tasmania. |
| 17. <i>A. altithermella</i> , Walsingham, Ent. M. Mag. Vol. 39, p. 265 (1903). | S. France, Spain. |
| 18. <i>A. kinkerella</i> , Snellen, Tijdschr. v. Ent. Vol. 19, p. 23, pl. 1 (1875). | Holland, N. Germany. |
| 19. <i>A. undina</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 64 (1921). | Rhodesia. |
| 20. <i>A. cirrhaea</i> , Meyrick, ibidem, Vol. 4, p. 190 (1914). | Transvaal. |
| 21. <i>A. autoleuca</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 264 (1904). | Victoria. |
| 22. <i>A. tamaricicola</i> , Walsingham, Ent. M. Mag. Vol. 47, p. 190 (1911). | Algeria. |
| 23. <i>A. cryolopha</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 265 (1904). | N. Queensland. |

LEPIDOPTERA HETEROCEERA

24. *A. chionocephala*, Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 79 New South Wales.
(1901).
25. *A. leucomichla*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 265 S. E. Australia,
Tasmania.
26. *A. nea*, Walsingham, Ent. M. Mag. Vol. 56, p. 9 (1920).
27. *A. melanombra*, Meyrick, Trans. N. Zeal. Inst. Vol. 20, p. 77 (1888).
28. *A. phantasta*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 266 (1904).
29. *A. achnias*, Meyrick, ibidem, Vol. 29, p. 266 (1904).
30. *A. isonira*, Meyrick, ibidem, Vol. 29, p. 264 (1904). — Pl. I, Fig. 2.
31. *A. spectrella*, Meyrick, ibidem, Vol. 29, p. 266 (1904).
32. *A. harpastis*, Meyrick, ibidem, Vol. 29, p. 267 (1904).
33. *A. crystallista*, Meyrick, Ann. Transv. Mus. Vol. 2, p. 229 (1911).
34. *A. acropasta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 110 (1919).
35. *A. microtima*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 267 (1904).
36. *A. coniombra*, Meyrick, ibidem, Vol. 29, p. 267 (1904).
37. *A. poliopasta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 110 (1919).
38. *A. megalornis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 261
(1904).
39. *A. belonodes*, Meyrick, ibidem, Vol. 29, p. 261 (1904). W. Australia.
40. *A. drosias*, Meyrick, ibidem, Vol. 29, p. 262 (1904). S. Australia.
41. *A. salias*, Meyrick, Exot. Microlep. Vol. 2, p. 500 (1922). S. Australia.
42. *A. collecta*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 64 (1921). Rhodesia.
43. *A. incola*, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 60 (1912). Cape Colony.

6. GENUS NEALYDA, DIETZ

Nealyda, Dietz, Ent. News, Philad. Vol. 11, p. 350 (1900). — Type: *N. bifidella*, Dietz.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 2/3, in ♂ serrate, simple, basal joint moderate, without pecten. Labial palpi moderate, curved, ascending, somewhat thickened with appressed scales, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with median dorsal scale-tooth; 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 2/3, trapezoidal, apex strongly produced, termen abruptly emarginate, projecting beneath emargination, cilia 2; 2-4 remote, 5 and 6 obsolete, transverse vein absent between 4 and 7.

Remarks. — This and the next genus, which are nearly related together, are the only American forms of this group; at present their immediate origin is uncertain.

Geographical distribution of species. — North and South American.

Larva mining in leaves (3 species known).

Foodplants *Nyctaginaceae*.

1. *N. accincta*, Meyrick, Exot. Microlep. Vol. 3, p. 5 (1923). Brazil.
2. *N. bicolor*, Walsingham, Proc. Zool. Soc. Lond. p. 522 (1891). St Vincent.
3. *N. leucosatra*, Meyrick, Exot. Microlep. Vol. 3, p. 5 (1923). Brazil.
4. *N. pisoniae*, Busck, Proc. U. S. Mus. Vol. 23, p. 229, pl. 1, f. 5 (1900). Florida.
5. *N. kinzelella*, Busck, ibidem, Vol. 23, p. 230 (1900). Florida.
6. *N. bifidella*, Dietz, Ent. News, Philad. Vol. 11, p. 351, pl. 1, fig. 2 (1900). — Colorado.

Pl. I, Fig. 3.

7. GENUS NEODACTYLOTA, BUSCK

Neodactylota, Busck, Proc. U. S. Mus. Vol. 25, p. 835 (1903). — Type : *N. snellenella*, Walsingham.

Eudactylota, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 54 (1911). — Type : *N. barberella*, Busck.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae $\frac{3}{4}$, in ♂ simple, basal joint without pecten. Labial palpi long, recurved, second joint somewhat thickened with appressed scales, terminal joint as long as second or longer, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex produced, termen excised, more or less prominent below excision, cilia over 1; 3 and 4 separate, 5 in ♂ sometimes subobsolete, in ♀ sometimes stalked with 4, 6 and 7 stalked.

Remarks. — Apparently a somewhat divergent form allied to the preceding.

Geographical distribution of species. — North American.

Larva unknown.

1. *N. snellenella*, Walsingham, Insect Life, Vol. 1, p. 83 (1888).

Arizona.

2. *N. barberella*, Busck, Proc. U. S. Mus. Vol. 25, p. 836 (1903).

Arizona, Mexico.

8. GENUS PARANARSIA, RAGONOT

Paranarsia, Ragonot, Bull. Soc. Ent. Fr. p. 195 (1895). — Type : *P. joannisiella*, Ragonot.

Characters. — Head smooth; tongue very short. Antennae in ♂ rather strongly pubescent, basal joint rather long, without pecten. Labial palpi long, subascending, clothed with rough projecting scales longer beneath, terminal joint short, pointed, concealed. Maxillary palpi imperceptible. Posterior tibiae with appressed scales. Forewings with 2 from near angle, 7 and 8 stalked, 7 to costa. Hindwings 1, narrow-trapezoidal, termen strongly emarginate, apex produced, acute; 3 and 4 separate, 5 nearer 6 than 4, 6 and 7 separate.

Remarks. — Probably a development of *Metanarsia*.

Geographical distribution of species. — South-West Europe.

Larva unknown.

1. *P. joannisiella*, Ragonot, Bull. Soc. Ent. Fr. p. 196 (1895).

Spain, France, Italy.

9. GENUS METANARSIA, STAUDINGER

Metanarsia, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 315 (1870). — Type : *M. modesta*, Staudinger.

Characters. — Head with appressed scales, sidetufts raised; tongue rudimentary. Antennae in ♂ simple, basal joint with pecten. Labial palpi moderately long, porrected, second joint compressed, somewhat rough-scaled above and with pointed apical scale-projection beneath, terminal joint extremely short in both sexes, almost concealed in scales of second. Maxillary palpi rudimentary. Forewings with 2

from towards angle, 7 and 8 stalked, 7 to costa. Hindwings 1, trapezoidal, apex somewhat produced, termen sinuate; 3 and 4 remote, 6 and 7 remote, parallel.

Remarks. — Apparently an early form of this group.

Geographical distribution of species. — South-East Europe, West-Central Asia.

Larva unknown.

- | | |
|--|--------------------------|
| 1. <i>M. modesta</i> , Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 315 (1870). | S.E. Russia, Turcomania. |
| 2. <i>M. junctivittella</i> , Christoph, Mém. Lép. Roman. Vol. 2, p. 161, pl. 8, f. 11 (1885). | Turcomania. |
| 3. <i>M. onzella</i> , Christoph, ibidem, Vol. 3, p. 120, pl. 5, f. 13 (1887). | Turcomania. |

10. GENUS COLOPTERYX, HOFMANN

Colopteryx, Hofmann, Iris, Vol. 10, p. 239 (1897). — Type: *C. conchylidella*, Hofmann.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae with basal joint thickened with scales. Labial palpi long, curved, ascending, second joint stout, rough-scaled beneath, terminal joint half second, thick, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs. Forewings with 1b short-furcate, 2 remote, 7 absent, 11 from before middle. Hindwings under 1, trapezoidal, apex long-produced, termen emarginate; 2-7 remote, tolerably parallel.

Remarks. — Probably related to *Epidola*.

Geographical distribution of species. — Asia Minor.

Larva unknown.

- | | |
|---|-------------|
| 1. <i>C. conchylidella</i> , Hofmann, Iris, Vol. 10, p. 239 (1897). | Asia Minor. |
|---|-------------|

11. GENUS EPIDOLA, STAUDINGER

Epidola, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 244 (1859). — Type: *E. stigma*, Staudinger.

Characters. — Head with appressed scales; ocelli small, posterior; tongue short. Antennae 5/6, in ♂ simple, basal joint moderately elongate, with pecten. Labial palpi moderate, curved, subascending, second joint with short tuft of rough projecting scales beneath, terminal joint much shorter than second, somewhat thickened with scales, scarcely pointed. Maxillary palpi obsolete. Posterior tibiae clothed with long fine hairs above. Forewings with 2 from towards angle, 2-6 parallel, 7 absent, 11 from middle. Hindwings nearly 1, elongate-trapezoidal, apex scarcely produced, termen slightly sinuate, cilia 1 1/2; 3 and 4 connate or stalked, 5 rather approximated, 6 and 7 parallel.

Remarks. — A singular form; but Staudinger's remarks on the tibial spurs appear unjustified, since they are quite normal.

Geographical distribution of species. — Western Mediterranean.

Larva case-bearing, feeding on leaves.

Foodplants *Quercus*, *Cistus*, etc., seemingly rather indiscriminate.

- | | |
|---|--|
| 1. <i>E. stigma</i> , Staudinger, Stett. Ent. Zeit. Vol. 20, p. 244 (1859). | N. W. Africa, Corsica,
Spain, Canaries. |
| 2. <i>E. barcinonella</i> , Millière, Icon. Descr. Lép. Vol. 2, p. 313, pl. 83, f. 13-15 (1868) | France, Spain. |

12. GENUS TIRANIMIA, CHRÉTIEN

Tiranimia, Chrétien, Ann. Soc. Ent. Fr. Vol. 84, p. 334 (1915). — Type : *T. epidolella*, Chrétien.

Characters. — Head smooth; tongue developed. Antennae in ♂ simple, basal joint without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short. Forewings with 1b furcate, 2 almost from angle, 7 absent, 11 from middle. Hindwings under 1, elongate-trapezoidal, termen slightly sinuate; 3 and 4 connate, 5 approximated, 6 and 7 approximated towards base.

Remarks. — I have not seen this genus, which is of dubious relationship, but considered by its author to be near *Epidola*.

Geographical distribution of species. — North African.

Larva unknown.

1. *T. epidolella*, Chrétien, Ann. Soc. Ent. Fr. Vol. 84, p. 334 (1915). Algeria.

Group 2 (Aristotelia type)

Originating from a form of the *Gelechia* group approaching *Recurvaria*, by basal drawing apart of veins 3 and 4 of hindwings, with the result that veins 2-7 are all remote and more or less parallel, the hindwings having typically the apex acutely produced and an abrupt emargination beneath it; this combination of features is very characteristic and generally easily recognisable.

13. GENUS ENCENTROTIS, MEYRICK

Encentrotis, Meyrick, Ann. Transv. Mus. Vol. 8, p. 65 (1921). — Type : *E. catagrapha*, Meyrick.

Characters. — Head smooth; ocelli very small, posterior; tongue developed. Antennae 4/5, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from angle, 7 and 8 out of 6, 7 to costa, 9 absent, 11 from middle. Hindwings 2/3, elongate-trapezoidal, termen faintly sinuate, cilia 1/2; 2 remote, 3-5 somewhat approximated, 4 from angle, 6 and 7 short-stalked.

Remarks. — Characterised by the peculiar neuration.

Geographical distribution of species. — South African.

Larva unknown.

1. *E. catagrapha*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 65 (1921) Natal.

14. GENUS NEVADIA, CARADJA

Nevadia, Caradja, Iris, Vol. 34, p. 117 (1920). — Type : *N. ribbeella*, Caradja.

Characters. — Head with appressed scales; tongue developed. Antennae under 1, filiform. Labial palpi with second joint very long, prorected, beneath with rough projecting tuft of scales, above

with projecting hairs on apical third, terminal joint concealed. Forewings with 2 separate, 7 and 8 stalked, connate with 6 from apex of cell, 7 to apex, 10 from 1/2, 11 from 2/5. Hindwings over 1 (?), trapezoidal, apex pointed, termen sinuate, cilia very long (assumed 3-5 separate, 6 and 7 stalked).

Remarks. — Not known to me, and the characters are incompletely given by the author, who apparently considers the genus allied to *Chilopselaphus*, which is doubtful; perhaps belonging to preceding group.

Geographical distribution of species. — S. W. Europe.

Larva unknown.

1. *N. ribbeella*, Caradja, Iris, Vol. 34, p. 118 (1920). Spain.

15. GENUS CHILOPSELAPHUS, MANN

Chilopselaphus, Mann, Verh. Zool.-bot. Ges. Wien, Vol. 17, p. 849 (1867). — Type : *C. fallax*, Mann.

Characters. — Head with appressed scales; ocelli posterior; tongue short. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, straight, porrected, second joint very long, compressed, clothed with rough projecting scales above and beneath, terminal joint short, pointed, sometimes concealed in scales of second. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely haired above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex produced, termen emarginate, cilia 1; 3-5 remote, parallel, 6 and 7 stalked.

Remarks. — A derivative of *Trichembola*.

Geographical distribution of species. — South Europe, West-Central Asia, North and East Africa.

Larva unknown.

- | | |
|---|---------------------|
| 1. <i>C. fallax</i> , Mann, Verh. Zool.-bot. Ges. Wien, Vol. 17, p. 850 (1857). | Hungary, S. Russia. |
| 2. <i>C. balneariella</i> , Chrétien, Le Naturaliste, Vol. 29, p. 179 (1907). | S. France. |
| 3. <i>C. numidella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 333 (1915). | Algeria. |
| 4. <i>C. ethicodes</i> , Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 68 (1920). | Kenya Colony. |

16. GENUS TRICHEMBOLA, MEYRICK

Trichembola, Meyrick, Exot. Microlep. Vol. 2, p. 115 (1918). — Type : *T. segnis*, Meyrick.

Characters. — Head smooth-scaled; ocelli very small, posterior; tongue short. Antennae 5/6, in ♂ shortly ciliated, basal joint elongate, without pecten. Labial palpi with second joint extremely long, straight, porrected, loosely scaled beneath, above with long rough projecting hairscales diminishing to apex, terminal joint short, obliquely projecting, slender, pointed. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2-6 remote, 7 and 8 stalked, 7 to costa, 9 sometimes absent, 11 from middle. Hindwings 1, narrow-trapezoidal, apex pointed, termen more or less emarginate, cilia nearly 2; 2-5 remote, parallel, 6 and 7 rather approximated at base.

Remarks. — Probably related to *Megacraspedus*. The absence of vein 9 is a specific character only.

Geographical distribution of species. — India and Ceylon.

Larva unknown.

- | | |
|---|---------------|
| 1. <i>T. opisthopa</i> , Meyrick, Exot. Microlep. Vol. 2, p. 115 (1918). | Coorg. |
| 2. <i>T. segnis</i> , Meyrick, ibidem, Vol. 2, p. 116 (1918). — Pl. I, Fig. 6; | Assam, Coorg. |
| Pl. 5, Fig. 127. | |
| 3. <i>T. epichorda</i> , Meyrick, ibidem, Vol. 2, p. 116 (1918). | Assam. |
| 4. <i>T. fuscata</i> , Meyrick, ibidem, Vol. 2, p. 116 (1918). | Ceylon. |

17. GENUS EPIPARASIA, REBEL

Epiparasia, Rebel, Iris, Vol. 28, p. 276 (1914). — Type : *E. incertella*, Herrich-Schäffer.

Characters. — Head with appressed scales; tongue developed. Antennae 3/4, in ♂ rather thick. Labial palpi very long, curved, ascending, second joint much thickened with appressed scales, terminal joint very short (in ♂ 1/4, in ♀ 1/5 of second), thickened, pointed. Forewings with 2 remote, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, elongate-trapezoidal, apex pointed, termen sinuate; 3 and 4 remote, 5 nearer 6, 6 and 7 somewhat approximated towards base.

Remarks. — I have not seen this genus, probably correlated with *Metzneria*.

Geographical distribution of species. — South-East Europe, West-Central Asia.

Larva unknown.

- | | |
|---|--------------------------|
| 1. <i>E. incertella</i> , Herrich-Schäffer, Neue Schmett. p. 31, f. 156 (1861). | S. Russia, E. Turkestan. |
| <i>longivitella</i> , Rebel, Iris, Vol. 28, p. 276, pl. 4, f. 12 (1914). | |

18. GENUS METZNERIA, ZELLER

Metzneria, Zeller, Isis, p. 197 (1839). — Type : *M. paucipunctella*, Zeller.

Cleodora, Stephens, Ill. Brit. Ent. Haust. Vol. 4, p. 220 (1834) (praeocc.). — Type : *M. lappella*, Linnaeus.

Parasia, Duponchel, Cat. Lép. Eur. p. 350 (1846). — Type : *M. neuropterella*, Zeller.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ serrate, simple, basal joint moderately elongate, without pecten. Labial palpi very long, curved, ascending, more or less thickened with somewhat loose scales, sometimes rough above, terminal joint much shorter than second, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 widely remote from angle, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex acute, produced, termen sinuate, cilia 1-2; 3 and 4 remote, 5 nearer 6, 6 and 7 somewhat approximated.

Remarks. — Generally related to *Pycnostola*. Imago often retired in habit.

Geographical distribution of species. — Palaearctic, with one species in South Africa, and one spreading into North America.

Larva (9 known) always feeding in seedheads.

Foodplants almost always *Compositae*, but for one species *Plantaginaceae*.

- | | |
|---|----------|
| 1. <i>M. hastella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 315 (1915). | Algeria. |
|---|----------|

2. *M. insignicans*, Walsingham, Proc. Zool. Soc. Lond. p. 926 (1907) Canaries.
 3. *M. littorella*, Douglas, Trans. Ent. Soc. Lond. (2) Vol. 1, p. 67 (1850). C. & S. Europe, N. Africa.
 quinquepunctella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 172, f. 573 (1855).
 4. *M. acrena*, Meyrick, Proc. Zool. Soc. Lond. p. 724 (1908). Transvaal, Cape Colony.
 heptacentra, Meyrick, Ann. Transv. Mus. Vol. 3, p. 64 (1912).
 5. *M. paucipunctella*, Zeller, Isis, p. 202 (1839). Europe, W. C. Asia.
 intestinella, Mann, Wien. Ent. Monatsschr. Vol. 8, p. 187, pl. 4, f. 15 (1864).
 6. *M. infelix*, Walsingham, Proc. Zool. Soc. Lond. p. 926 (1907). Canaries.
 7. *M. hilarella*, Caradja, Iris, Vol. 34, p. 95 (1920). Spain.
 8. *M. tristella*, Rebel, Iris, Vol. 13, p. 164 (1900). Spain, Algeria.
 9. *M. dichroa*, Walsingham, Proc. Zool. Soc. Lond. p. 927, pl. 51, f. 4 (1907). Canaries.
 10. *M. castiliella*, Möschler, Berl. Ent. Zeitsch. Vol. 10, p. 142 (1866). Spain, Portugal.
 11. *M. monochroa*, Walsingham, Proc. Zool. Soc. Lond. p. 927, pl. 51, f. 5 (1907). Canaries.
 [Asia Minor.
 12. *M. torridella*, Mann, Wien. Ent. Monatsschr. Vol. 3, p. 174 (1859). Corsica, Sicily, Dalmatia,
 13. *M. lappella*, Linnaeus, Syst. Nat. (10) Vol. 1, p. 537 (1758). Europe, W. Asia, Canada,
 silacea, Haworth, Lep. Brit. p. 555 (1828). E. United States.
 aestivella, Herrich-Schäffer, Schmett. Eur. Vol. 5, f. 559 (1855).
 14. *M. obsoleta*, Christoph, Hor. Soc. Ent. Ross. Vol. 22, p. 313 (1888). Turcomania.
 15. *M. torosulella*, Rebel, Stett. Ent. Zeit. Vol. 54, p. 49 (1893). Spain.
 16. *M. aspretella*, Lederer, Hor. Soc. Ent. Ross. Vol. 6, p. 92, pl. 5, f. 13 (1869). N. Persia.
 17. *M. litigiosella*, Millière, Le Naturaliste, Vol. 1, p. 139 (1879). S. France.
 18. *M. carlinella*, Stanton, Cat. Brit. Tin. Suppl. p. 5 (1851). C. & S. E. Europe,
 aestivella, Herrich-Schäffer, Schmett. Eur. Vol. 5, f. 558 (1855). Algeria.
 19. *M. metzneriella*, Stanton, Cat. Brit. Tin. Suppl. p. 5 (1851). C. Europe.
 paucipunctella, Douglas, Trans. Ent. Soc. Lond. (2) Vol. 1, p. 14 (1850).
 20. *M. aprilella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 207, f. 963 (1856). S.E. Europe, Asia Minor.
 21. *M. inflammatella*, Christoph, Bull. Soc. Nat. Mosc. p. 26 (1882). — Pl. I, E. Siberia, China.
Fig. 5.
 22. *M. selaginella*, Mann, Verh. Zool.-bot. Ges. Wien, Vol. 5, p. 565 (1855). Corsica, Dalmatia.
 23. *M. Eatoni*, Walsingham, Ent. M. Mag. Vol. 35, p. 183 (1899). Spain, Algeria.
 citella, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 53, p. 411 (1903).
 24. *M. incognita*, Walsingham, Ent. M. Mag. Vol. 40, p. 220 (1904). Algeria.
 25. *M. agraphella*, Ragonot, Bull. Soc. Ent. Fr. p. 106 (1895). Asia Minor.
 26. *M. neuropterella*, Zeller, Isis, p. 202 (1839). C. & S. E. Europe.
 27. *M. sanguinolentella*, Joannis, Bull. Soc. Ent. Fr. p. 295 (1910). France.
 28. *M. igneella*, Tengström, Not. Sällsk. Faun. Fenn. Förh. p. 183 (1859). Russia.
 29. *M. albiramosella*, Christoph, Mém. Lép. Roman. Vol. 2, p. 159, pl. 8, Turcomania.
 f. 9 (1885).

19. GENUS PROSELOTIS, MEYRICK

Proselotis, Meyrick, Exot. Microlep. Vol. 1, p. 276 (1914). — Type : *P. sceletodes*, Meyrick.

Idiobela, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 111 (1919). — Type : *P. ischnoptila*, Turner.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with dense appressed scales, dilated towards apex, terminal joint as long as second, thickened with dense scales projecting roughly posteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, very narrowly elongate-trapezoidal, apex long-produced,

termen obliquely bisinuate, cilia 3; 2-4 remote, parallel, 5 curved, approximated, 6 and 7 parallel or rather approximated.

Remarks. — Probably correlated with the preceding.

Geographical distribution of species. — Africa, India and Australia.

Larva unknown.

1. *P. ischnoptila*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 111 (1919). Queensland.
amphiptila, Meyrick, Exot. Microlep. Vol. 2, p. 425 (1921).
2. *P. apicipunctella*, Stainton, Trans. Ent. Soc. Lond. (2), Vol. 5, p. 119 (1861). Bengal.
3. *P. sceletodes*, Meyrick, Exot. Microlep. Vol. 1, p. 276 (1914). Nyassaland.

20. GENUS MERIDORMA, NOV. GEN.

Type : *M. thrombodes*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, somewhat thickened with scales throughout, second joint with scales rough towards apex beneath, terminal joint as long as second, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 4 and 5 long-stalked, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings 1/2, narrow-trapezoidal, apex long-produced, acute, termen emarginate, cilia 4; 3 and 4 separate, 5 parallel, 6 and 7 separate, diverging.

Remarks. — Probably related to *Pityocona*.

Geographical distribution of species. — South American.

Larva unknown.

1. *M. thrombodes*, Meyrick, Trans. Ent. Soc. Lond. p. 232 (1914). Brit. Guiana.

21. GENUS PITYOCONA, MEYRICK

Pityocona, Meyrick, Exot. Microlep. Vol. 2, p. 116 (1918). — Type : *P. xeropis*, Meyrick.

Characters. — Head smooth-scaled; ocelli small, posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, much thickened throughout with scales, rough posteriorly, especially on terminal joint, second joint thickest towards apex, terminal as long as second, thickest in middle, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with 1b furcate, 2 from 3/4, 3 from before angle, 4 from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, narrowly elongate-trapezoidal, apex pointed, produced, termen sinuate-emarginate, cilia 4; 2-4 remote, 5 and 6 somewhat approximated at base, 6 and 7 parallel.

Remarks. — A peculiar type, also allied to the foregoing.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

1. *P. xeropis*, Meyrick, Exot. Microlep. Vol. 2, p. 117 (1918). India, Ceylon, Java.

22. GENUS CATAMECES, TURNER

Catameces, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 122 (1919). — Type : *C. thiophara*, Turner.

Characters. — Head with appressed scales; tongue developed. Antennae 3/4, basal joint without pecten. Labial palpi long, recurved, second joint very long, thickened anteriorly with long rough scales forming a small apical tuft, terminal joint about half second, rather stout, acute. Maxillary palpi very short, appressed to tongue. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to costa. Hindwings 1, trapezoidal, apex pointed, strongly produced, termen sinuate; 3 and 4 remote, 5 nearer 6 than 4, 6 and 7 nearly parallel.

Remarks. — I have not seen this genus, which Dr. Turner considers to be of uncertain affinity. It must however by the hindwings undoubtedly belong to this group, and I have therefore ventured to assume that his description of the termination of vein 7 of the forewings as apical is an inaccuracy, since in this group it is invariably costal.

Geographical distribution of species. — Australian.

Larva unknown.

1. *C. thiophara*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 122 (1919). Queensland.

23. GENUS PYCNOTOLA, MEYRICK

Pycnotola, Meyrick, Ent. M. Mag. Vol. 53, p. 113 (1917). — Type : *P. operosa*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint beneath with compact apical projecting tuft of dense scales, terminal joint as long as second (in *auturga* half second), slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings 1 or under 1, narrow-trapezoidal, apex produced, termen emarginate, cilia 1 1/2-2 1/2; 3 and 4 remote, 5 nearer 6, 6 and 7 remote, diverging.

Remarks. — This and the next three genera are perhaps correlated developments from *Aristotelia*.

Geographical distribution of species. — Characteristically South African and rather numerous in that region, whence the single South European and group of three Australian species are probably stragglers.

Larva (*helicaula*) living in a case formed of a silken refuse-covered tube coiled in a spiral cone of three whorls, resembling a snail-shell (*Helix*).

Foodplant unrecorded.

- | | |
|---|------------------------------|
| 1. <i>P. actias</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 272 (1904). | S. & W. Australia, Tasmania. |
| 2. <i>P. sciopola</i> , Meyrick, ibidem, Vol. 29, p. 272 (1904). | New S. Wales. [mania. |
| 3. <i>P. stalactis</i> , Meyrick, ibidem, Vol. 29, p. 271 (1904). | W. Australia, Tasmania. |
| 4. <i>P. bohemica</i> , Nickerl, Wien. Ent. Monatsschr. Vol. 8, p. 5, pl. 5, f. 9 (1864). | Bohemia, S. Russia. |
| 5. <i>P. merista</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 14 (1918). | Transvaal. |

6. *P. psacasta*, Meyrick, Proc. Zool. Soc. Lond. p. 723 (1908).
celyphodes, Meyrick, Ann. Transv. Mus. Vol. 2, p. 10, pl. 4, f. 3 (1910).
7. *P. melanotracta*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 413 (1910).
8. *P. helicaula*, Meyrick, ibidem, Vol. 10, p. 59 (1912).
9. *P. hiberna*, Meyrick, ibidem, Vol. 10, p. 60 (1912).
10. *P. perlustrata*, Meyrick, ibidem, Vol. 17, p. 280 (1920).
11. *P. suffusella*, Walsingham, Trans. Ent. Soc. Lond. p. 109, pl. 5, f. 47 (1891).
12. *P. auturga*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 65 (1921).
13. *P. operosa*, Meyrick, ibidem, Vol. 2, p. 10, pl. 4, f. 1 (1910). — PI. I, Fig. 8.
14. *P. ochraula*, Meyrick, ibidem, Vol. 6, p. 14 (1918).
15. *P. illuminata*, Meyrick, ibidem, Vol. 3, p. 281 (1913).
16. *P. pentacentra*, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 59 (1912).
17. *P. iospila*, Meyrick, Ann. Transv. Mus. Vol. 2, p. 10, pl. 4, f. 2 (1910).
18. *P. celeris*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 280 (1920).
19. *P. magnanima*, Meyrick, ibidem, Vol. 10, p. 59 (1912).
20. *P. oeconomica*, Meyrick, ibidem, Vol. 17, p. 280 (1920).
21. *P. invida*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 64 (1912).
22. *P. crateraula*, Meyrick, ibidem, Vol. 6, p. 14 (1918).
23. *P. pammacha*, Meyrick, ibidem, Vol. 3, p. 281 (1913).
24. *P. semnochroa*, Meyrick, ibidem, Vol. 3, p. 281 (1913).
- Transvaal, Natal.
[R. Colony.
Transvaal, Orange,
Cape Colony.
Natal, Zululand.
Cape Colony.
Transvaal, Natal,
Zululand.
Natal.
Transvaal, Natal,
Zululand.
Transvaal.
Transvaal.
Zululand. [Protectorate.
Transv., Orange R., S.W.
Cape Colony.
Orange R. Colony.
Cape Colony.
W. Pondoland.
Natal.
Transvaal.
Transvaal.

24. GENUS MEGACRASPEDUS, ZELLER

Megacraspedus, Zeller, Isis, p. 189 (1839). — Type : *P. dolosella*, Zeller.

Neda, Chambers, Canad. Ent. Vol. 6, p. 243 (1874) (praeocc.). — Type : *P. plutella*, Chambers.

Pycnobathra, Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 80 (1901). — Type : *P. achroa*, Lower.

Autoneda, Busck, Dyar's Cat. N. Amer. Lep. p. 496 (1902). — Type : *P. plutella*, Chambers.

Toxoceras, Chrétien, Ann. Soc. Ent. Fr. Vol. 84, p. 329 (1915). — Type : *P. violacella*, Chrétien.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrate, ciliated, basal joint elongate, slender, without pecten. Labial palpi long, recurved, second joint beneath with dense projecting apical tuft, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from 1/2-3/4 of cell, 6 sometimes out of 7 near base, 7 and 8 stalked, 7 to costa, 11 from or before middle. Hindwings 1, trapezoidal, apex produced, acute, termen emarginate, cilia 2 1/2-3; 3 and 4 remote, 5 nearer 6, 6 and 7 remote; in ♀ sometimes semiaborted or rudimentary.

Remarks. — The many curious analogies between this and the quite alien Oecophorid genus *Pleurota* (probably connected with similarity of habits) are worthy of study.

Geographical distribution of species. — Europe (southern half), West-Central Asia, and North Africa, 16; South Africa, 3; North America, 2; Australia, 24; New Zealand, 1. This curious distribution leaves the Australian and New Zealand section (which are similar in character to the European) disconnected from the rest; I have not received a single individual from India. I can only infer (as in *Pleurota*, of which the distribution is exactly similar) that the genus is really Australian in origin, and that at an early period (perhaps coeval with *Pleurota*) wind-borne immigrants were carried

to Europe, whence subsequently America received a straggler; the South African Colony probably indicates a separate southern immigration from Australia.

Larva (*calamogona*) feeding on seeds of grasses; it is quite likely this habit may be general.

Foodplants *Gramineae*.

1. *M. exilis*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 21, pl. 1, Mexico.
f. 20 (1909).
2. *M. plutella*, Chambers, Canad. Ent. Vol. 6, p. 244 (1874). Kentucky.
3. *M. photinopa*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 281 (1920). Cape Colony.
4. *M. serica*, Meyrick, ibidem, Vol. 5, p. 369 (1909). Cape Colony.
5. *M. peracuta*, Meyrick, ibidem, Vol. 17, p. 281 (1920). Cape Colony.
6. *M. argyroneurella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 316 (1870). S. Russia, W. C. Asia.
7. *M. exoletella*, Erschoff, Fedtsch. Lep. Turk. p. 103, pl. 6, f. 117 (1874). Turkestan.
8. *M. lagopella*, Herrich-Schäffer, Neue Lep. p. 13, f. 81 (1860). Hungary, S. Russia.
9. *M. attritella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 316 (1870). S. Russia.
10. *M. pentheres*, Walsingham, Ent. M. Mag. Vol. 56, p. 10 (1920). France.
11. *M. consortella*, Caradja, Iris, Vol. 34, p. 117 (1920). W. Turkestan.
12. *M. imparella*, Fischer von Röslerstamm, Abbild. Schmett. p. 303, pl. 100, f. 2 (1844). S.E. Europe, Asia Minor.
13. *M. tristicta*, Walsingham, Ent. M. Mag. Vol. 46, p. 231 (1910). France.
14. *M. binotella*, Fischer von Röslerstamm, Abbild. Schmett. p. 301, pl. 99, f. 2 (1844). C. & S. E. Europe.
15. *M. monolorella*, Rebel, Ann. Hofmus. Wien, Vol. 20, p. 25 (1905). Asia Minor.
16. *M. lanceolella*, Zeller, Stett. Ent. Zeit. Vol. 11, p. 143 (1850). C. & S. Europe.
- hessleriellus*, Rössler, Jahrb. Nass. Ver. Naturk. Vol. 20, p. 247 (1866).
17. *M. cuencella*, Caradja, Iris, Vol. 34, p. 117 (1920). Spain.
18. *M. subdolella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 243 (1859). S. Europe.
19. *M. separatella*, Fischer von Röslerstamm, Abbild. Schmett. p. 302, pl. 100, f. 1 (1844). E. C. & S. Europe.
20. *M. pusilla*, Walsingham, Ent. M. Mag. Vol. 39, p. 266 (1903). Spain.
21. *M. Tutti*, Walsingham, Entom. Rec. Vol. 9, p. 140 (1897). France.
22. *M. dolosella*, Zeller, Isis, p. 190 (1839). E. C. & S. Europe.
23. *M. violacella*, Chrétien, Ann. Soc. Ent. Fr. p. 330 (1915). Algeria. [Tasmania.
24. *M. platyleuca*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 274 (1904). S. E. & W. Australia,
25. *M. astemprella*, Meyrick, ibidem, Vol. 29, p. 275 (1904). S. & W. Australia.
26. *M. centrosema*, Meyrick, ibidem, Vol. 29, p. 275 (1904). S. E. Australia, Tasmania.
27. *M. oxyphanes*, Meyrick, ibidem, Vol. 29, p. 276 (1904). W. Australia. [nia.
28. *M. chalcoscia*, Meyrick, ibidem, Vol. 29, p. 276 (1904). S. & W. Australia.
29. *M. melitopis*, Meyrick, ibidem, Vol. 29, p. 276 (1904). W. Australia.
30. *M. euxena*, Meyrick, ibidem, Vol. 29, p. 277 (1904). W. Australia.
31. *M. hoplitis*, Meyrick, ibidem, Vol. 29, p. 277 (1904). W. Australia.
32. *M. inficeta*, Meyrick, ibidem, Vol. 29, p. 277 (1904). S. E. Australia, Tasmania.
33. *M. stratimera*, Lower, Trans. Roy. Soc. S. Australia, Vol. 39, p. 58 (1897). S. Australia. [nia.
34. *M. conogramma*, Meyrick, Exot. Microlep. Vol. 2, p. 424 (1921). N. Queensland.
35. *M. calamogona*, Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 163 (1886). New Zealand.
36. *M. sematacma*, Meyrick, Exot. Microlep. Vol. 2, p. 424 (1921). Queensland.
37. *M. niphodes*, Lower, Trans. Roy. Soc. S. Australia, Vol. 39, p. 58 (1897). Victoria, Tasmania.
38. *M. pityritis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 278 (1904). S.E. Australia, Tasmania.
39. *M. argonota*, Lower, Trans. Roy. Soc. S. Australia, Vol. 43, p. 81 (1901). New South Wales.
40. *M. aenictodes*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 112 (1919). Queensland.
41. *M. sclerotricha*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 279 (1904). New South Wales.
42. *M. aphileta*, Meyrick, ibidem, Vol. 29, p. 280 (1904). S. & W. Australia.
43. *M. isotis*, Meyrick, ibidem, Vol. 29, p. 280 (1904). W. Australia.
44. *M. sagittifera*, Lower, ibidem, Vol. 25, p. 416 (1900). New South Wales.

45. *M. coniodes*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 281 (1904). S. Australia.
 46. *M. ischnota*, Meyrick, ibidem. Vol. 29, p. 282 (1904). W. Australia.
 47. *M. achroa*, Lower, Trans. Roy. Soc. S. Australia, Vol. 43, p. 80 (1901). New South Wales.
 48. *M. popularis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 282 (1904). S.E. Australia, Tasmania.

25. GENUS ISOPHRICHTIS, MEYRICK

Isophrictis, Meyrick, Ent. M. Mag. Vol. 53, p. 113 (1917). — Type : *I. tanacetella*, Schrank.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint beneath throughout with long rough spreading hairs, terminal joint as long as second or rather shorter, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings 1 or under 1, narrow-trapezoidal, apex produced, termen emarginate, cilia 2-3; 3 and 4 remote, 5 nearer 6, 6 and 7 separate or connate, diverging.

Remarks. — The species have a characteristic type of marking, with inwards-oblique costal strigulae before apex.

Geographical distribution of species. — European region, 8; North America, 13; and also 1 common to both regions. Probably North America may be the place of origin, and the genus is a comparatively recent one.

Larva feeding in flowers, seed-heads, and stems (4 known).

Foodplants always *Compositae*.

- | | |
|--|------------------------------------|
| 1. <i>I. robinella</i> , Chrétien, Le Naturaliste, Vol. 29, p. 178 (1907). | France. |
| 2. <i>I. meridionella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 204, f. 595 (1855). | France, Spain. |
| 3. <i>I. lineatella</i> , Zeller, Stett. Ent. Zeit. Vol. 11, p. 142 (1850). | S. Europe, Asia Minor. |
| 4. <i>I. kefersteiniella</i> , Zeller, ibidem, Vol. 11, p. 142 (1850). | S. Europe, Asia Minor, |
| 5. <i>I. invisella</i> , Constant, Ann. Soc. Ent. Fr. p. 257, pl. 10, f. 18 (1884). | Corsica. [Palestine. |
| 6. <i>I. senicula</i> , Meyrick, Ent. Mitth. Berlin, Vol. 2, p. 298 (1913). | Tunis. |
| 7. <i>I. constantina</i> , Baker, Ent. M. Mag. Vol. 24, p. 255 (1888). | Algeria. |
| 8. <i>I. anthemidella</i> , Wocke, Cat. Lép. Eur. p. 301 (1871).
<i>striatella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 204, f. 563 (1855). | C. & S. Europe, Asia
Minor. |
| 9. <i>I. tanacetella</i> , Schranck, Fauna Boic. Vol 2, p. 122 (1802).
<i>striatella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 288 (1805). | Europe, Asia Minor,
California. |
| 10. <i>I. pallidistrigella</i> , Chambers, Canad. Ent. Vol. 6, p. 244 (1874) (-dast-). | Texas. |
| 11. <i>I. pennella</i> , Busck, Proc. Ent. Soc. Wash. p. 88 (1907). | Arizona. |
| 12. <i>I. magnella</i> , Busck, Proc. U. S. Mus. Vol. 25, p. 776 (1903) — Pl. I, Fig. 7. | Colorado. |
| 13. <i>I. ciliolineella</i> , Chambers, Canad. Ent. Vol. 6, p. 242 (1874) (cilia-). | Texas. |
| 14. <i>I. dietziella</i> , Busck, Proc. U. S. Mus. Vol. 25, p. 777 (1903). | Colorado. |
| 15. <i>I. canicostella</i> , Walsingham, Insect Life, Vol. 1, p. 82 (1888). | California, Colorado. |
| 16. <i>I. actiella</i> , Barnes, Contr. Lep. N. Amer. Vol. 4, p. 224 (1920). | California. |
| 17. <i>I. anteliella</i> , Busck. Proc. U. S. Mus. Vol. 25, p. 778 (1903). | New Jersey. |
| 18. <i>I. sabulella</i> , Walsingham, Insect Life, Vol. 1, p. 83 (1888). | California. |
| 19. <i>I. similella</i> , Chambers, Canad. Ent. Vol. 4, p. 193 (1872) (-liella).
<i>piscipellis</i> , Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 277 (1873). | Kentucky, Texas. |
| 20. <i>I. tophella</i> , Walsingham, Insect Life, Vol. 1, p. 83 (1888). | California, New Mexico. |
| 21. <i>I. pallidella</i> , Chambers, Canad. Ent. Vol. 6, p. 245 (1874). | Colorado. |
| 22. <i>I. modesta</i> , Walsingham, Insect Life, Vol. 1, p. 82 (1888). | California. |

26. GENUS PALTODORA, MEYRICK

Paltodora, Meyrick, Ent. M. Mag. Vol. 30, p. 230 (1894). — Type : *P. cytisella*, Curtis.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint with rough spreading hairs beneath throughout, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex pointed, produced, termen emarginate, cilia 2 1/2-3; 3 and 4 remote, parallel, 5 nearer 6, 6 and 7 remote, diverging.

Remarks. — An interesting monotypic genus related to the preceding.

Geographical distribution of species. — Europe, Assam; probably also in the intervening Himalayan and West-Central Asiatic regions and perhaps elsewhere, as it is easily overlooked and its foodplant is nearly cosmopolitan.

Larva feeding on fronds of fern.

Foodplant *Pteris aquilina* (*Filices*).

1. *P. cytisella*, Curtis, Brit. Ent. Vol. 14, p. 671 (1837). Europe, Assam.

walkeriella, Douglas, Trans. Ent. Soc. Lond. (2) Vol. 1, p. 21 (1850).

fuscipennis, Westwood, Brit. Moths, Vol. 2, p. 192 (1851).

coeruleostella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 204, f. 562 (1855).

27. GENUS ZIZYPHIA, CHRÉTIEN

Zizyphia, Chrétien, Bull. Soc. Ent. Fr. p. 166 (1908). — Type : *Z. cleodorella*, Chrétien.

Characters. — Head with dense appressed scales. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, acute, with tuft of hairs of equal length (?) sexual rising from its base internally. Forewings with 2 widely remote, 3-5 approximated, 7 and 8 stalked, 7 to costa. Hindwings under 1, elongate-trapezoidal, apex obtuse; 5 curved near origin, 6 and 7 separate, diverging.

Remarks. — I have not seen this genus, of which the published characters are unfortunately incomplete; it is stated by its author to be near *Sophronia*, but this does not seem likely.

Geographical distribution of species. — North African.

Larva feeding on leaves (?) (particulars unrecorded).

Foodplant *Zizyphus* (*Rhamnaceae*).

1. *Z. cleodorella*, Chrétien, Bull. Soc. Ent. Fr. p. 167 (1908). Algeria.

28. GENUS DORYCNOPA, LOWER

Dorycnopa, Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 77 (1901). — Type : *D. heliochares*, Lower.

Bactrolopha, Lower, ibidem, Vol. 25, p. 79 (1901). — Type : *D. orthodesma*, Lower.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple or ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with short rough apical tuft beneath, terminal joint as long as second, with rough scales anteriorly throughout. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 parallel, 7 and 8 stalked, 6 sometimes out of 7 near base, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, produced, termen emarginate, cilia 2 1/2-3; 3 and 4 remote, parallel, 5 nearer 6, 6 and 7 rather approximated towards base.

Remarks. — A derivative of *Aristotelia*.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|---|------------------|
| 1. <i>D. orthodesma</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 79 (1901). | New South Wales. |
| 2. <i>D. marmorea</i> , Lower, Proc. Linn. Soc. N. S. Wales, Vol. 24, p. 96 (1899). | New South Wales. |
| 3. <i>D. heliochares</i> , Lower, ibidem, Vol. 25, p. 417 (1900).
<i>acroxantha</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 78 (1901). | South Australia. |
| 4. <i>D. triphora</i> , Lower, ibidem, Vol. 44, p. 67 (1920) (<i>triphera</i>). | South Australia. |

29. GENUS STEREOMITA, BRAUN

Stereomita, Braun, Ent. News, Philad. Vol. 33, p. 43 (1922). — Type : *S. andropogonis*, Braun.

Characters. — Head smooth. Antennae nearly 1, subserrate, basal joint long, slender. Labial palpi long, recurved, second joint thickened with scales beneath and slightly tufted, terminal joint as long as second, thickened with scales towards middle, acute. Maxillary palpi short, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 1b furcate, 2 from angle, 3 absent, 4 and 5 connate, nearer 6, 7 and 8 out of 6, 7 to costa, 9 remote, 11 from beyond middle. Hindwings 1/2, apex produced, termen emarginate; 2-5 remote, 5 nearer 6, 6 and 7 very short-stalked.

Remarks. — Perhaps related to *Megacraspedus*; there is marked superficial resemblance to *Batrachedra*.

Geographical distribution of species. — North American.

Larva feeding in inflorescence of grass.

Foodplant *Andropogon* (*Gramineæ*).

- | | |
|--|-------|
| 1. <i>S. andropogonis</i> , Braun, Ent. News, Philad. Vol. 33, p. 44 (1922). | Ohio. |
|--|-------|

30. GENUS PTYCERATA, ELY

Ptycerata, Ely, Proc. Ent. Soc. Wash. Vol. 12, p. 69 (1910). — Type : *P. buschella*, Ely.

Characters. — Labial palpi long, recurved, second joint with projecting tuft of scales beneath, terminal joint slender, acute. Forewings with 2 from towards angle, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, termen emarginate; 3 and 4 separate, 6 and 7 connate.

Remarks. — I have not seen this genus, which seems allied to the preceding.

Geographical distribution of species. — North American.

Larva unknown.

1. *P. buschella*, Ely, Proc. Ent. Soc. Wash., Vol. 12, p. 69 (1910).

N. E. United States.

31. GENUS SITOTROGA, HEINEMANN

Sitotroga, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 287 (1870). — Type : *S. cerealella*, Olivier.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint long, with pecten. Labial palpi long, recurved, second joint rough-scaled beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 parallel, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex pointed, produced, termen emarginate, cilia 2; 3-5 remote, parallel, 6 and 7 stalked.

Remarks. — A monotypic genus, specially interesting through the presence of the antennal pecten, which does not occur in any other genus of this group known to me; otherwise apparently allied to the two preceding.

Geographical distribution of species. — This well-known and very destructive insect is probably North American in origin, but is now practically cosmopolitan except in colder climates, being artificially imported with its food.

Larva feeding in stored grain (maize, rice, wheat, etc).

Foodplants Gramineae.

1. *S. cerealella*, Olivier, Encycl. Méth. Ent. Vol. 1, p. 121 (1789).

Europe, Asia, Africa,

hordei, Kirby, Introd. Ent. Vol. 1, p. 172 (1815).

America, Australia,

arettella, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 635 (1864).

New Zealand, Fiji

? *coarctatella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 353 (1877).

melanarthra, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 416 (1900).

palearis, Meyrick, Exot. Microlep. Vol. 1, p. 65 (1913).

32. GENUS ACRAEOLOGA, MEYRICK

Acraeologa, Meyrick, Ann. Transv. Mus. Vol. 8, p. 66 (1921). — Type : *A. xerochroa*, Meyrick.

Characters. — Head smooth; ocelli small, posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint very elongate, without pecten. Labial palpi very long, recurved, second joint somewhat thickened with scales, somewhat rough towards apex beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae shortly rough-scaled above. Forewings with 1b furcate, 2 from towards angle, 6 and 7 out of 8, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex long-produced, acute, termen rectangularly emarginate, cilia 2; 3-5 somewhat approximated, 6 and 7 long-stalked.

Remarks. — Perhaps allied to the preceding.

Geographical distribution of species. — South African.

Larva unknown.

1. *A. xerochroa*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 66 (1921).

Transvaal.

33. GENUS ANOMOXENA, MEYRICK

Anomoxena, Meyrick, Trans. Ent. Soc. Lond. p. 28 (1917). — Type: *A. spinigera*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with whorls of scales forming a series of separate acute projecting teeth beneath, terminal joint as long as second, slightly thickened with scales, acute. Maxillary palpi minute, filiform, appressed to tongue. Posterior tibiae clothed with rough projecting hairs above and beneath. Forewings with 1b furcate, 2 from angle, 2-5 parallel, transverse vein very oblique outwards from 2 to 5, faint between 5 and 6, 6 and 7 stalked, 7 to costa, 8 separate, approximated to 6 at base, 11 from near middle. Hindwings 3/5, narrow-trapezoidal, apex acute, strongly produced, termen emarginate, cilia 4; 2 remote, 3 and 4 nearly parallel, 4 from angle, 4 and 5 somewhat approximated, 6 and 7 approximated at base.

Remarks. — This curious genus differs from the whole of the family in having vein 8 of the forewings separate from 7; in all the rest these veins are stalked or coincident. There can be no doubt as to homology, since all veins are present. It is equally certain that the genus belongs to this family and to the most specialised form of it as represented by the first two groups, since the highly characteristic hindwings are found in no other. Hence it follows that the genus must be supposed to have descended from a long line of ancestors in which the stalking of 7 and 8 was an invariable feature, and to have lost that structure, probably by the gradual reduction and absorption of the stalk, thus affording a proved instance of a modification contrary to expectation. The genus is probably derived from *Aristotelia*.

Geographical distribution of species. — South American.

Larva unknown.

1. *A. spinigera*, Meyrick, Trans. Ent. Soc. Lond. p. 29 (1917). — PI. I, Colombia, Ecuador,

Fig. I; PI. 5, Fig. II6.

Peru, Brazil.

tetraxoa, Meyrick, ibidem, p. 30 (1917).

34. GENUS PYCNOODYTIS, MEYRICK

Pycnodynitis, Meyrick, Ann. Transv. Mus. Vol. 6, p. 15 (1918). — Type: *P. erebaula*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with dense scales rather loose beneath and triangularly expanded with hairs on apical half above, terminal joint shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with 2-4 parallel, 6 and 7 out of 8, 7 to costa, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex strongly produced, acute, termen emarginate, cilia 2; 2-5 remote, parallel, 6 and 7 approximated towards base.

Remarks. — Allied to *Aristotelia*.

Geographical distribution of species. — South Africa, Madagascar.

Larva unknown.

1. *P. erebaula*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 15 (1918).

Zululand.

2. *P. irrigata*, Meyrick, Exot. Microlep. Vol. 2, p. 120 (1918).

Madagascar.

35. GENUS IULOTA, MEYRICK

Iulota, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 283 (1904). — Type : *I. ithyxyla*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4. in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi long, curved, ascending, second joint clothed with dense scales, rough beneath, and roughly projecting above towards apex, terminal joint as long as second or shorter, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 nearly parallel, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, produced, termen sinuate, cilia 1 1/2-2; 3 and 4 separate, 5 somewhat approximated to 4, 6 and 7 approximated at base.

Remarks. — Apparently related to *Aristotelia*.

Geographical distribution of species. — Australian.

Larva unknown.

1. <i>I. phaenoptila</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 112 (1919).	New South Wales.
2. <i>I. ischnora</i> , Turner, ibidem, Vol. 31, p. 112 (1919).	Queensland.
3. <i>I. ithyxyla</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 283 (1904).	W. Australia.
4. <i>I. triglossa</i> , Meyrick, ibidem, Vol. 29, p. 284 (1904).	Tasmania.
5. <i>I. episila</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 58 (1897).	S. E. & W. Australia, Tasmania.

36. GENUS PSAMATHOCRITA, NOV. GEN.

Type : *P. osseella*, Stainton.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi moderately long, curved, ascending, slender, smooth-scaled, terminal joint shorter than second, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 2 from angle, 3 and 4 remote, 5 out of 6 below 8, 7 and 8 out of 6, 7 to costa, 9 from near 6, 10 remote, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex long-produced, acute, termen emarginate, cilia 3; 3 and 4 separate, 5 parallel, 6 and 7 separate, diverging.

Remarks. — A derivative of advanced forms of *Aristotelia*.

Geographical distribution of species. — Mediterranean.

Larva unknown.

1. <i>P. innotatella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 327 (1915).	Algeria.
2. <i>P. albidella</i> , Rebel, Verh. Zool.-bot. Ges. Wien. Vol. 53, p. 412 (1903).	Algeria.
3. <i>P. dejectella</i> , Staudinger, Stett. Ent. Zeit. Vol. 20, p. 242 (1859).	Spain.
4. <i>P. osseella</i> , Stainton, Ent. Annual, p. 87 (1861).	C. & S. Europe, Algeria.

37. GENUS ARISTOTELIA, HÜBNER

Aristotelia, Hübner, Verz. bek. Schmett. p. 424 (1826). — Type : *A. decurtella*, Hübner.

Chrysoesthia, Hübner, ibidem, p. 422 (1826). — Type : *A. hermannella*, Fabricius.

Nomia, Clemens, Proc. Acad. Nat. Sc. Philad. p. 167 (1860) (praeocc.). — Type : *A. lingulacella*, Clemens.

Chrysopora, Clemens, ibidem, p. 362 (1860). — Type : *A. lingulacella*, Clemens.

Nannodia, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 280 (1870). — Type : *A. stipella*, Hübner.

Apodia, Heinemann, ibidem, p. 286 (1870). — Type : *A. bifractella*, Douglas.

Ptocheuusa, Heinemann, ibidem, p. 288 (1870). — Type : *A. inopella*, Zeller.

Ergatis, Heinemann, ibidem, p. 295 (1870). — Type : *A. brizella*, Treitschke.

Doryphora, Heinemann, ibidem, p. 298 (1870) (praeocc.). — Type : *A. pulveratella*, Herrich-Schäffer.

Monochroa, Heinemann, ibidem, p. 308 (1870). — Type : *A. tenebrella*, Hübner.

Lamprotes, Heinemann, ibidem, p. 309 (1870). — Type : *A. atrella*, Haworth.

Enchrysa, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 282 (1873). — Type : *A. dissectella*, Zeller.

Xystophora, Heinemann, Schmett. Deutschl. (2), Vol. 2, Tab. p. 6 (1876). — Type : *A. pulveratella*, Herrich-Schäffer.

Isochasta, Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 163 (1886). — Type : *A. paradesma*, Meyrick.

Eucatoptus, Walsingham, Proc. Zool. Soc. Lond. p. 69 (1897). — Type : *A. penicillata*, Walsingham.

Anaphaula, Walsingham, Ent. M. Mag. Vol. 40, p. 268 (1904). — Type : *A. gaditella*, Staudinger.

Parapodia, Joannis, Bull. Soc. Ent. Fr. p. 305 (1912). — Type : *A. sinica*, Frauenfelder.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennæ 4/5, in ♂ simple or ciliated, basal joint elongate, without pecten. Labial palpi from moderate to very long, second joint variably thickened with appressed scales or sometimes rough beneath, terminal joint nearly as long as second or longer, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, sometimes 6 out of 7 near base, or 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings 1 or less than 1, elongate-trapezoidal, apex pointed, produced, termen sinuate or emarginate, cilia 1 1/4-3; 3-5 remote, nearly parallel, 6 and 7 remote or somewhat approximated.

Remarks. — This extensive genus, which must be derived from a form approaching *Recurvaria*, is doubtless of considerable age. It presents many difficulties, since there are numerous species so closely allied together that their relations need very minute study, whilst others seem abruptly disconnected. There is much variation in the structure of the palpi, yet these differences are only specific, and sometimes occur in species otherwise very closely related; and similarly the variation in the structure of veins 6-8 of the forewings occurs to some extent erratically, and would not define natural groups or permit generic separation. The earlier forms of the genus are often elegantly and even brilliantly coloured, but the later developments are eminently sombre and obscure. As they are also frequently small and very retired in habit, it is likely that many species remain to be discovered in regions imperfectly explored.

Geographical distribution of species. — Summarised as follows : 102 Palaearctic, 9 South African, 19 Indian, 20 Australian, 1 New Zealand, 10 Hawaiian, 74 American, and 1 artificially widespread (*peltosema*) of uncertain origin. This distribution and the characters noted above seem to indicate that the home of the genus might be looked for in regions (unexplored for Micro-Lepidoptera) of South-Central Asia. The Hawaiian species indicate a single original immigrant perhaps from South America, whence also the New Zealand species.

LEPIDOPTERA HETEROCEA

Larva more usually feeding internally (then sometimes apodal) in seed-heads, stems, roots, galls (not infrequently), or mining blotches in leaves, but also commonly amongst spun inflorescence or leaves.

Foodplants (known for 61 species): *Compositae* (7), *Leguminosae* (7), *Rosaceae* (5), *Chenopodiaceae* (5), *Polygonaceae* (5), *Tamaricaceae* (3), *Primulaceae* (3), *Rubiaceae* (3), *Cyperaceae* (3), and 12 other Natural Orders, showing great diversity and adaptability.

- | | |
|---|---------------------------|
| 1. <i>A. bifractella</i> , Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 66 (1850). | C. & S. Europe, Asia |
| 2. <i>A. Martini</i> , Petry, Iris, Vol. 25, p. 101 (1911). | Thuringia. [Minor.] |
| 3. <i>A. seminivora</i> , Walsingham, Ent. M. Mag. Vol. 39, p. 263 (1903). | Spain, Morocco. |
| 4. <i>A. asterisci</i> , Walsingham, ibidem, Vol. 39, p. 263 (1903). | Spain. |
| 5. <i>A. scholastica</i> , Walsingham, ibidem, Vol. 39, p. 264 (1903). | Spain. |
| 6. <i>A. guimarensis</i> , Walsingham, Proc. Zool. Soc. Lond. p. 930, pl. 51, f. 6 (1907). | Canaries. |
| 7. <i>A. cuprimarginella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 326 (1915). | Algeria. |
| 8. <i>A. multistrigella</i> , Ragonot, Bull. Soc. Ent. Fr. p. 83 (1892). | Algeria. |
| 9. <i>A. sublutella</i> , Christoph, Hor. Soc. Ent. Ross. Vol. 9, p. 23, pl. 2, f. 20 (1872). | S. Russia. |
| 10. <i>A. abnormella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 199, f. 549 (1855). | Germany, Austria, Sicily. |
| 11. <i>A. inopella</i> , Zeller, Isis, p. 201 (1839). | C. & S. Europe, Algeria. |
| <i>papella</i> , Zeller, Iris, p. 858 (1847). | |
| <i>inulella</i> , Curtis, Ann. Mag. Nat. Hist. (2), Vol. 5, p. 117 (1850). | |
| <i>amesella</i> , Chrétien, Bull. Soc. Ent. Fr. p. 339 (1907). | |
| 12. <i>A. atrella</i> , Haworth, Lep. Brit. p. 587 (1828). | C. Europe. |
| <i>umbriterella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 195, f. 524 (1855). | |
| <i>aurimaculella</i> , Höfner, Jahrb. Mus. Kärnth. Klagenfurt, Vol. 24, p. 170 (1896). | |
| 13. <i>A. unicolorella</i> , Duponchel, Hist. Nat. Lép. Fr. Suppl. Vol. 4, p. 458, pl. 85, f. 8 (1842). | C. Europe, Spain. |
| <i>immaculatella</i> , Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 67 (1850). | |
| 14. <i>A. tenebrella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 434 (1818). | Europe. |
| <i>tenebrosella</i> , Zeller, Isis, p. 201 (1839). | |
| <i>buffonella</i> , Millière, Cat. Lép. Alp. Mar. p. 361 (1875). | |
| 15. <i>A. plumbella</i> , Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 310 (1870). | Germany. |
| 16. <i>A. rhenanella</i> , Heyden, Stett. Ent. Zeit. Vol. 24, p. 343 (1863). | Germany. |
| 17. <i>A. arundinetella</i> , Stainton, Ent. Ann. p. 91 (1858). | England, Holland, |
| 18. <i>A. Hornigi</i> , Staudinger, Stett. Ent. Zeit. Vol. 44, p. 184 (1882). | Austria. [N. Germany.] |
| 19. <i>A. rumicetella</i> , Hofmann, ibidem, Vol. 29, p. 28 (1868). | C. Europe. |
| 20. <i>A. acutangulella</i> , Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 305 (1870). | Germany. |
| 21. <i>A. sepicolella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 192 (1855). | Germany, Austria. |
| <i>rectifasciella</i> , Fuchs, Stett. Ent. Zeit. Vol. 63, p. 326 (1902). | |
| 22. <i>A. cacomictra</i> , Walsingham, Proc. Zool. Soc. Lond. p. 931 (1907). | Canaries. |
| 23. <i>A. bicolorella</i> , Rebel, Ann. Hofmus. Wien, Vol. 18, p. 329, pl. 3, f. 14 (1903). | Bulgaria. |
| 24. <i>A. morosa</i> , Mühlig, Stett. Ent. Zeit. Vol. 25, p. 101 (1864). | C. Europe. |
| 25. <i>A. farinosae</i> , Stainton, Nat. Hist. Tin. Vol. 10, p. 164, pl. 12, f. 2 (1867). | C. Europe. |
| 26. <i>A. quaestionella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 193, f. 587 (1855). | C. Europe. |
| 27. <i>A. orthogonella</i> , Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 307 (1870). | S. Russia. |
| 28. <i>A. conspersella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 177, f. 591 (1855). | C. Europe. |
| 29. <i>A. antipala</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 290 (1904). | New South Wales. |
| 30. <i>A. pamphaea</i> , Meyrick, ibidem, Vol. 29, p. 290 (1904). | New South Wales. |
| 31. <i>A. elachistella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 356 (1877). | Colombia. |
| 32. <i>A. repudiata</i> , Meyrick, Exot. Microlep. Vol. 3, p. 6 (1923). | Assam. |
| 33. <i>A. Brunicki</i> , Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 63, p. (43) (1913). | Galicia. |
| 34. <i>A. Benedeni</i> , Weyenbergh, Ent. M. Mag. Vol. 10, p. 122 (1873). | Cape Verde Islands. |

35. *A. helotella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 240 (1859).
damonella, Millière, Cat. Lép. Alp. Mar. p. 328 (1875).
algeriella, Baker, Ent. M. Mag. Vol. 24, p. 255 (1888).
doliodes, Meyrick, ibidem, Vol. 27, p. 55 (1891).
36. *A. striatopunctella*, Rebel, Verh. Zool.-bot. Ges. Wien. Vol. 41, p. 630, S. Europe, Asia Minor. (1891).
37. *A. retusella*, Rebel, ibidem, Vol. 41, p. 632 (1891).
38. *A. decolorella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 199, f. 550 (1855).
luteella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 300 (1870).
39. *A. perterrita*, Meyrick, Exot. Microlep. Vol. 3, p. 11 (1923).
40. *A. servella*, Zeller, Isis, p. 201 (1839).
ferrea, Frey. Mith. Schweiz. Ent. Ges. Vol. 3, p. 253 (1870).
41. *A. robustella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 312 (1870).
42. *A. tenuella*, Mann, Wien. Ent. Monatsschr. Vol. 8, p. 186, pl. 4, f. 16 (1864).
43. *A. griseella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 301 (1870).
44. *A. latiuscula*, Heinemann, ibidem (2), Vol. 2, p. 300 (1870).
45. *A. flavicapitella*, Chrétien, Ann. Soc. Ent. Fr. p. 323 (1915).
46. *A. carchariella*, Zeller, Isis, p. 201 (1839).
47. *A. pulveratella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 199, f. 552 (1855).
intaminatella, Stainton, Ent. Weekly Intell. Vol. 7, p. 140 (1861).
48. *A. asthenodes*, Meyrick, Exot. Microlep. Vol. 3, p. 6 (1923).
49. *A. parvula*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 321 (1880).
50. *A. gaditella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 243 (1859).
51. *A. leptocentra*, Meyrick, Exot. Microlep. Vol. 1, p. 64 (1912).
52. *A. artificella*, Herrich-Schäffer, Neue Schmett. p. 31, f. 158 (1861).
53. *A. rufulella*, Snellen, Tijdschr. v. Ent. Vol. 27, p. 175, pl. 9, f. 9 (1884).
54. *A. sirota*, Meyrick, Proc. Zool. Soc. Lond. p. 724 (1908).
55. *A. chloroneura*, Meyrick, Exot. Microlep. Vol. 3, p. 6 (1923).
56. *A. mesoxysta*, Meyrick, Ent. Mith. Berlin, Vol. 2, p. 299 (1913).
57. *A. tripunctella*, Snellen, Tijdschr. v. Ent. Vol. 27, p. 172, pl. 9, f. 7 (1884).
58. *A. palustrella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 14 (1850).
59. *A. divisella*, Douglas, ibidem (2), Vol. 1, p. 60 (1850).
60. *A. centrosema*, Lower, Trans. Roy. Soc. S. Australia, Vol. 35, p. 171 (1893). — PI. I, Fig. 13.
61. *A. monactis*, Meyrick, Exot. Microlep. Vol. 3, p. 10 (1923).
62. *A. peribapta*, Lower, Trans. Roy. Soc. S. Australia, Vol. 42, p. 237 (1918).
63. *A. placidella*, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 24, p. 441, pl. 12, f. 11 (1874).
natalella, Busck, Proc. U. S. Mus. Vol. 27, p. 756 (1904).
64. *A. disconotella*, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 86 (1878).
65. *A. agatha*, Meyrick, Exot. Microlep. Vol. 2, p. 119 (1918).
66. *A. pantalaena*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 78, pl. 2, f. 33 (1911).
67. *A. angustipennella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 119 (1863).
kearfottella, Busck, Proc. U. S. Mus. Vol. 25, p. 803 (1903).
68. *A. gilvolinella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 119 (1863).
69. *A. harrisonella*, Busck, Proc. U. S. Mus. Vol. 27, p. 756 (1904).
70. *A. discriminata*, Meyrick, Exot. Microlep. Vol. 3, p. 10 (1923).
71. *A. robusta*, Braun, Ent. News, Philad. Vol. 32, p. 8 (1921).
72. *A. pullella*, Chambers, Canad. Ent. Vol. 6, p. 237 (1874) (*pullusella*).
minimella, Chambers, ibidem, Vol. 6, p. 243 (1874).
73. *A. melanaphra*, Meyrick, Exot. Microlep. Vol. 3, p. 10 (1923).
74. *A. fragariae*, Busck, Proc. Ent. Soc. Wash. Vol. 21, p. 52 (1919).
- S. W. Europe,
N. W. Africa.
- Asia Minor.
- Austria.
- Ontario.
- Europe, Asia Minor.
- S. Russia.
- Dalmatia, Asia Minor.
- Germany.
- Silesia, Latvia.
- Algeria. [Minor.
- C. & S. E. Europe, Asia
- C. & S. E. Europe.
- Himalayas.
- Asia Minor.
- Spain, Algeria.
- Bengal.
- S. Russia.
- E. Siberia.
- Transvaal.
- Brazil.
- Tunis.
- E. Siberia.
- England, Holland.
- England.
- S. E. Australia,
- Tasmania.
- Ontario, N. Carolina.
- S. Australia.
- Br. Columbia, California.
- Kentucky.
- Assam.
- Mexico.
- Pennsylvania,
- New Jersey.
- Pennsylvania.
- Brit. Columbia, Cali-
- Ontario. [fornia.
- Ontario, Ohio.
- Texas.
- California.
- Brit. Columbia, Oregon.

75. *A. absconditella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 595 (1864). N. E. United States.
palpiannulella. Chambers, Canad. Ent. Vol. 4, p. 68 (1872).
76. *A. physaliella*, Chambers, ibidem, Vol. 4, p. 173 (1872).
77. *A. palpiabellula*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 253 (1875).
78. *A. suffusella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 64 (1850).
79. *A. elongella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 307 (1870).
80. *A. lucidella*, Stephens, Ill. Brit. Ent. Haust. Vol. 4, p. 221 (1835).
81. *A. nomadella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 18, p. 616 (1868).
82. *A. scordiscella*, Rebel, Ann. Hofmus. Wien, Vol. 19, p. 352 (1904).
83. *A. lutulentella*, Zeller, Iris, p. 201 (1839).
84. *A. remissella*, Zeller, ibidem, p. 854 (1847).
85. *A. vetustella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 191, f. 526 (1855).
86. *A. surda*, Meyrick, Exot. Microlep. Vol. 3, p. 6 (1923).
87. *A. aulacopis*, Meyrick, ibidem, Vol. 3, p. 6 (1923).
88. *A. fluidescens*, Meyrick, Ann. Transv. Mus. Vol. 4, p. 190 (1914).
89. *A. balanocentra*, Meyrick, ibidem, Vol. 4, p. 190 (1914).
90. *A. trematias*, Meyrick, ibidem, Vol. 3, p. 282 (1913).
91. *A. crassicornis*, Walsingham, Proc. Zool. Soc. Lond. p. 68 (1897).
92. *A. pulicella*, Walsingham, ibidem, p. 67 (1897).
93. *A. campicolella*, Mann, Wien. Ent. Monatsschr. Vol. 1, p. 181 (1857).
94. *A. vallicola*, n. sp.
onella, Chrétien, Bull. Soc. Ent. Fr. p. 141 (1908) (*var.*).
95. *A. gracilella*, Chrétien, ibidem, p. 140 (1908).
96. *A. ainella*, Chrétien, ibidem, p. 91 (1908).
97. *A. psamathias*, Meyrick, Ent. M. Mag. Vol. 27, p. 56 (1891).
98. *A. tetragonella*, Stainton, ibidem, Vol. 22, p. 99 (1885).
99. *A. morphochroma*, Walsingham, ibidem, Vol. 36, p. 80 (1900).
100. *A. punctatella*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 321 (1880).
101. *A. quinquepunctella*, Busck, Proc. U. S. Mus. Vol. 25, p. 804 (1903).
102. *A. maculaticornis*, Walsingham, Faun. Hawaiensis, Vol. 1, p. 478, pl. 13, f. 17 (1907).
103. *A. nigricitella*, Walsingham, ibidem, Vol. 1, p. 479, pl. 13, f. 18 (1907).
104. *A. gigantea*, Swezey, Proc. Hawa. Ent. Soc. Vol. 2, p. 274 (1913).
105. *A. ichthyochora*, Walsingham, Faun. Hawaiensis, Vol. 1, p. 479, pl. 13, f. 19 (1907).
106. *A. epermeniella*, Walsingham, ibidem, Vol. 1, p. 480, pl. 13, f. 20 (1907).
107. *A. notata*, Walsingham, ibidem, Vol. 1, p. 480, pl. 13, f. 21 (1907).
108. *A. lanaiensis*, Walsingham, ibidem, Vol. 1, p. 481, pl. 13, f. 22 (1907).
109. *A. elegantior*, Walsingham, ibidem, Vol. 1, p. 481, pl. 13, f. 23 (1907).
110. *A. mendax*, Walsingham, ibidem, Vol. 1, p. 481, pl. 13, f. 24 (1907).
111. *A. arcuata*, Walsingham, ibidem, Vol. 1, p. 482, pl. 13, f. 25 (1907).
112. *A. paradesma*, Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 163 (1886).
113. *A. tetracosma*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 289 (1904).
114. *A. furtiva*, Meyrick, ibidem, Vol. 29, p. 288 (1904).
115. *A. sticheris*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 113 (1919).
116. *A. thetica*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 289 (1904).
117. *A. themerastis*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 113 (1919).
118. *A. cypriantha*, Turner, ibidem, Vol. 31, p. 114 (1919).
119. *A. sinaica*, Frauenfelder, Verh. Zool.-bot. Ges. Wien, Vol. 9 p. 324, pl. 7, f. 4 (1859).
tamaricicola, Joannis, Bull. Soc. Ent. Fr. p. 305 (1912).
120. *A. brochodesma*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 438 (1908).
- Kentucky.
- Texas.
- C. Europe.
- England, Germany.
- N. & C. Europe.
- S. Europe.
- Herzegovina.
- N., C. & S. E. Europe,
Asia Minor.
- C. & S. Europe.
- C. & S. E. Europe.
- Burma.
- Assam, Himalayas.
- Natal.
- Natal.
- Transvaal, Port. E. Afr.
- Virgin Is.
- Virgin Is.
- Corsica, Dalmatia, Asia
Algeria. [Minor.]
- Algeria.
- Algeria.
- Algeria.
- England, Holland.
- S. France, Italy.
- Asia Minor.
- Pennsylvania, Ontario.
- Hawaiian Is.
- E. Australia, Tasmania.
- Queensland.
- S.E. Australia, Tasmania.
- New South Wales.
- New South Wales.
- S. Europe, S. W. Asia,
- N. Africa.
- Assam.

121. *A. argodecta*, Meyrick, Exot. Microlep. Vol. 2, p. 118 (1918). Ceylon.
122. *A. leucophanta*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 438 (1908). Assam.
123. *A. epicharla*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 113 (1919). New South Wales.
124. *A. iomarmara*, Meyrick, Exot. Microlep. Vol. 2, p. 425 (1921). Queensland.
125. *A. galeotis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 438 (1908). Ceylon, Transvaal.
126. *A. prominula*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 282 (1913). Transvaal.
127. *A. trinotella*, Herrich-Schäffer, Neue Schmett. p. 6, f. 46 (1856). S. Europe.
- aurantiella*, Rebel, Rov. Lap. Vol. 22, p. 188.
128. *A. incilata*, Meyrick, Exot. Microlep. Vol. 2, p. 117 (1918). Assam.
129. *A. thalamitis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 437 (1908). Assam.
130. *A. resinosa*, Meyrick, Exot. Microlep. Vol. 2, p. 118 (1918). Assam.
131. *A. citrocosma*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 437 (1908). Ceylon, Assam.
- (1908). — **Pl. I, Fig. 14.**
132. *A. sinistra*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 287 (1904). New South Wales.
133. *A. macrothecta*, Meyrick, ibidem, Vol. 29, p. 288 (1904). New South Wales.
134. *A. eurypsola*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 115 (1919). New South Wales.
135. *A. turbida*, Turner, ibidem, Vol. 31, p. 115 (1919). Queensland.
136. *A. ferritincta*, Turner, ibidem, Vol. 31, p. 115 (1919). Queensland.
137. *A. ancillula*, Walsingham, Proc. Zool. Soc. Lond. p. 930 (1907). Canaries.
138. *A. ingravata*, Meyrick, Exot. Microlep. Vol. 2, p. 118 (1918). Bengal.
139. *A. aulonota*, Meyrick, Trans. Ent. Soc. Lond. p. 30 (1917). Ecuador.
140. *A. corallina*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 23 (1909). Mexico.
141. *A. plumata*, Meyrick, Trans. Ent. Soc. Lond. p. 30 (1917). Brit. Guiana.
142. *A. veteranella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 361, pl. 5, f. 121 (1877). ?C. America.
143. *A. cytheraea*, Meyrick, Trans. Ent. Soc. Lond. p. 32 (1917). Colombia.
144. *A. erycina*, Meyrick, ibidem, p. 31 (1917). Ecuador, Peru.
145. *A. chalybochroa*, Walsingham, Proc. Zool. Soc. Lond. p. 69 (1897) (-*beichroa*). Virgin Is.
146. *A. penicillata*, Walsingham, ibidem, p. 70 (1897). Hayti, Virgin Is.
147. *A. rubidella*, Clemens, Proc. Acad. Nat. Sc. Philad. Vol. 12, p. 163 (1860). E. United States, Virgin Is.
- cassella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 594 (1864).
- rubensella*, Chambers, Canad. Ent. Vol. 4, p. 193 (1872).
- pudibundella*, Chambers, ibidem, Vol. 9, p. 23 (1877).
148. *A. lycopersicella*, Walsingham, Proc. Zool. Soc. Lond. p. 71 (1897). Virgin Is.
149. *A. calens*, Meyrick, Exot. Microlep. Vol. 3, p. 9 (1923). California.
150. *A. radicata*, Meyrick, Trans. Ent. Soc. Lond. p. 35 (1917). Colombia.
151. *A. aphiltra*, Meyrick, ibidem, p. 34 (1917). Colombia, Ecuador,
152. *A. hieroglyphica*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 24 (1909). Mexico. [Peru.]
153. *A. dasypoda*, Walsingham, ibidem, Vol. 4, p. 25, pl. 1, f. 22 (1909). Mexico, Jamaica.
154. *A. pyrodercia*, Walsingham, ibidem, Vol. 4, p. 26 (1909). Mexico.
155. *A. calculatrix*, Meyrick, Exot. Microlep. Vol. 3, p. 7 (1923). Brazil.
156. *A. perfossa*, Meyrick, Trans. Ent. Soc. Lond. p. 33 (1917). Ecuador, Peru.
157. *A. saturnina*, Meyrick, ibidem, p. 33 (1917). Peru.
158. *A. cynthia*, Meyrick, ibidem, p. 32 (1917). Peru.
159. *A. subrosea*, Meyrick, ibidem, p. 230 (1914). Guiana, Brazil.
160. *A. oribatis*, Meyrick, ibidem, p. 35 (1917). Peru.
161. *A. paphia*, Meyrick, ibidem, p. 34 (1917). Peru.
162. *A. vicana*, Meyrick, ibidem, p. 36 (1917). Peru.
163. *A. ephoria*, Meyrick, ibidem, p. 36 (1917). Peru.
164. *A. callirrhoda*, Meyrick, Exot. Microlep. Vol. 3, p. 9 (1923). N. Carolina.
165. *A. probolopis*, Meyrick, ibidem, Vol. 3, p. 8 (1923). Brazil, Peru.
166. *A. paterata*, Meyrick, Trans. Ent. Soc. Lond. p. 229 (1914). Guiana, Peru.
167. *A. salicifungiella*, Clemens, Proc. Ent. Soc. Philad. Vol. 3, p. 508 (1864). Illinois.

168. *A. fungivorella*, Clemens, Proc. Ent. Soc. Philad. Vol. 3, p. 507 (1864). E. United States.
 169. *A. ivae*, Busck, Proc. U. S. Mus. Vol. 23, p. 225, pl. 1, f. 1 (1900). Florida.
 170. *A. trossulella*, Walsingham, Proc. Zool. Soc. Lond. p. 67 (1897). Jamaica, Hayti.
 171. *A. Howardi*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 23, California, Mexico.
 pl. 1, f. 21 (1909).
 172. *A. squamigera*, Walsingham, ibidem, Vol. 4, p. 24 (1909). Mexico.
 173. *A. molestella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 274 (1873). Texas.
 174. *A. pudibundella*, Zeller, ibidem, Vol. 23, p. 273 (1873). E. United States, Hayti,
intermediella, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 89 (1878). Virgin Is.
 175. *A. pulvrea*, Braun, Proc. Calif. Acad. Sc. (4), Vol. 12, p. 117 (1923) California.
 (*pulvera*).
 176. *A. eumeris*, Meyrick, Exot. Microlep. Vol. 3, p. 9 (1923). Arizona.
 177. *A. roseosuffusella*, Clemens, Proc. Acad. Nat. Sc. Philad. Vol. 12, p. 162 E. United States,
 (1860). — Pl. I, Fig. 12. Canada.
bellella, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 595 (1864).
 178. *A. bifasciella*, Busck, Proc. U. S. Mus. Vol. 25, p. 799 (1903). Arizona.
 179. *A. lindanella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 226 (1920). California.
 180. *A. articulata*, Meyrick, Exot. Microlep. Vol. 2, p. 119 (1918). S. India.
 181. *A. aquosa*, n. sp. Kentucky.
suffusella, Chambers, Canad. Ent. Vol. 4, p. 171 (1872) (praeocc.).
 182. *A. aphromorpha*, Meyrick, Exot. Microlep. Vol. 3, p. 7 (1923). Burma.
 183. *A. interstratella*, Christoph, Hor. Soc. Ent. Ross. Vol. 9, p. 21, pl. 1, S. Russia.
 f. 18 (1872).
 184. *A. hemisarca*, Lower, Trans. Roy. Soc. S. Australia, Vol. 40, p. 542 New South Wales.
 (1916).
 185. *A. semiophanes*, Meyrick, Exot. Microlep. Vol. 2, p. 119 (1918). India, Ceylon, Java.
 186. *A. cosmographa*, Meyrick, Trans. Ent. Soc. Lond. p. 36 (1917). Peru.
 187. *A. chlorographa*, Meyrick, Ann. Transv. Mus. Vol. 4, p. 190 (1914). Transvaal.
 188. *A. callyntrophora*, Rebel, Anz. Akad. Wiss. Wien, Vol. 27, p. 3 (1899). Arabia.
 189. *A. subericinella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 197, f. 541 S. Europe, Asia Minor,
 (1855). N. Africa.
prohaskaella, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 57, p. 213 (1907). Uralsk.
 190. *A. bolschewichiella*, Caradja, Iris, Vol. 34, p. 109 (1920). England, Germany.
 191. *A. subdecurtella*, Stainton, Ent. Ann. p. 152 (1859). C. & S. Europe, Asia
 192. *A. decurtella*, Hübner, Samml. Eur. Schmett. Tin. f. 311 (1817). Minor, W. C. Asia.
turbatella, Treitschke, Schmett. Eur. Vol. 10 (3), p. 195 (1835).
amoenella, Joannis, Bull. Soc. Ent. Fr. p. 83 (1891).
 193. *A. decoratella*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 320 (1880). Sicily, S. E. Europe,
 194. *A. ericinella*, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, p. 497, pl. 306, C. Europe. [Asia Minor.
 f. 12 (1838).
micella, Hübner, Samml. Eur. Schmett. Tin. f. 210 (1796) (praeocc.).
 195. *A. Leonhardi*, Krone, Jahresber. Wien. Ent. Ver. Vol. 17, p. 25 (1907). Austria.
 196. *A. panchaliella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 312 (1870). S. Russia, Asia Minor.
 197. *A. mirandella*, Chrétien, Bull. Soc. Ent. Fr. p. 166 (1908). Algeria.
 198. *A. argyractis*, Meyrick, Exot. Microlep. Vol. 3, f. 8 (1923). Brazil.
 199. *A. heliacella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 198, f. 544 (1855). C. Europe.
Rogenhoferi, Staudinger, Verh. Zool.-bot. Ges. Wien, Vol. 22, p. 734 (1872).
 200. *A. mirabilis*, Christoph, Hor. Soc. Ent. Ross. Vol. 22, p. 314 (1888). S. Russia.
 201. *A. calastomella*, Christoph, ibidem, Vol. 9, p. 24, pl. 2, f. 21 (1872). S. Russia. [tine.
 202. *A. brisella*, Treitschke, Schmett. Eur. Vol. 9 (2), p. 173 (1833). N. & C. Europe, Pales-
 203. *A. staticella*, Millière, Bull. Soc. Ent. Fr. p. 167 (1875). S. France, Algeria.
 204. *A. frankeniae*, Walsingham, Ent. M. Mag. Vol. 34, p. 132 (1898). Corsica, Algeria.
 205. *A. micella*, Schiffersmüller, Syst. Verz. Schmett. Wien, p. 140 (1776). N. & C. Europe.
asterella, Treitschke, Schmett. Eur. Vol. 9 (2), p. 172 (1833).
 206. *A. coeruleopictella*, Caradja, Iris, Vol. 34, p. 106 (1920). E. Siberia.

207. *A. rutilella*, Snellen, Tijdschr. v. Ent. Vol. 27, p. 174, pl. 9, f. 8 (1884). E. Siberia.
 208. *A. epimetalla*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 290 (1904). New South Wales.
209. *A. ochricapilla*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 53, p. 96 (1903). Austria.
 210. *A. libertinella*, Zeller, Stett. Ent. Zeit. Vol. 33, p. 112 (1872).
 ? *germarella*, Hübner, Samml. Eur. Schmett. Tin. f. 467-8 (1832). Alps, Spain.
211. *A. tarquinella*, Stainton, Ent. Ann. p. 112 (1862). Ireland.
 212. *A. pictella*, Zeller, Isis, p. 202 (1839). Europe, Asia Minor.
- germarella, Nolcken, Lep. Faun. Livl. Vol. 2, p. 585 (1870).
 ? wilkella, Linnaeus, Faun. Suec. p. 361 (1761).
213. *A. superbella*, Zeller, Isis, p. 202 (1839). [Minor, Algeria.
 214. *A. arnoldiella*, Rebel, Ann. Hofmus. Wien, Vol. 20, p. 24 (1905). N. & C. Europe, Asia
 215. *A. cervinella*, Eversmann, Faun. Lep. Volg.-Ural. p. 585 (1844). Asia Minor.
 216. *A. nitidula*, Stainton, Tin. Syr. p. 44 (1867). S.E. Europe, Asia Minor.
 217. *A. cockerella*, Busck, Proc. U. S. Mus. Vol. 25, p. 800 (1903). Palestine.
 218. *A. elegantella*, Chambers, Canad. Ent. Vol. 4, p. 239 (1872). New Mexico.
 superbella, Chambers, ibidem, Vol. 7, p. 32 (1875). Missouri, Texas, Arizona.
219. *A. monilella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 225 (1920). N. Carolina.
 220. *A. sarcodes*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 26, pl. 1, f. 23 (1909). Panama.
221. *A. argentifera*, Busck, Proc. U. S. Mus. Vol. 25, p. 800 (1903). California.
 222. *A. primipilana*, Meyrick, Exot. Microlep. Vol. 3, p. 8 (1923). Ontario.
 223. *A. fervidella*, Mann, Wien. Ent. Monatsschr. Vol. 8, p. 187, pl. 5, f. 4 (1864). Asia Minor.
224. *A. hermannella*, Fabricius, Spec. Insect. Vol. 2, p. 509 (1781). Europe, Asia Minor,
 zinckenella, Hübner, Samml. Eur. Schmett. Tin. f. 401-2 (1817). N. Africa, Canada.
 225. *A. lingulacella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 167 (1860). E. United States.
 arminella, Frey, Stett. Ent. Zeit. Vol. 39, p. 249 (1878).
226. *A. versicolorella*, Kearfott, Journ. N. York Ent. Soc. Vol. 16, p. 183 (1908). California.
 227. *A. Eppelsheimi*, Staudinger, Stett. Ent. Zeit. Vol. 46, p. 351 (1885). Germany.
 228. *A. aletris*, Walsingham, Ent. Record, Vol. 31, p. 10 (1919). Sicily.
 229. *A. boseae*, Walsingham, Proc. Zool. Soc. Lond. p. 931, pl. 51, f. 7 (1907). Canaries.
 230. *A. stipella*, Hübner, Samml. Eur. Schmett. Tin. f. 138 (1796). C. & S. Europe, Asia
 naeviferella, Duponchel, Hist. Nat. Lép. Fr. Suppl. Vol. 4, pl. 85, f. 4 (1842). Minor, Natal.
 231. *A. dissectella*, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 283, pl. 4, f. 29 (1873). Ohio, Ontario.
- youngella, Kearfott, Canad. Ent. Vol. 37, p. 15 (1905).
232. *A. maculata*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 314 (1880). Asia Minor, Palestine.
 ? Korbi, Caradjia, Iris, Vol. 34, p. 105 (1920).
233. *A. clavata*, Meyrick, Exot. Microlep. Vol. 1, p. 197 (1914). Victoria.
 234. *A. schematas*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 271 (1911). Seychelles.
 235. *A. peltosema*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 50 (1900). E. & W. Australia, Ceylon,
 pyramidophora, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 123 (1919). S. Africa, S. America.
 236. *A. brucinella*, Mann, Verh. Zool.-bot. Ges. Wien. Vol. 22, p. 37 (1872). S. Europe, Asia Minor,
 237. *A. Wachtli*, Rogenhofer, ibidem, Vol. 30, p. (48) (1880). Egypt. [Egypt.
 238. *A. comis*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 282 (1913). Transvaal.

38. GENUS EMPEDAULA, MEYRICK

Empedaula, Meyrick, Exot. Microlep. Vol. 2, p. 148 (1918). — Type : *E. insipiens*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennæ 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, considerably thickened

throughout with appressed scales, laterally compressed, terminal joint as long as second, with scales loosely projecting posteriorly, apex shortly projecting or concealed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely haired above. Forewings with 1b furcate, 2 from towards angle, 3 from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings somewhat under 1, elongate-trapezoidal, apex acute, produced, termen obliquely bisinuate beneath apex, cilia 1 1/4; 3 and 4 rather approximated towards base, 5 remote, parallel, 6 and 7 parallel.

Remarks. — The species are superficially similar to the *fungivorella* group of *Aristotelia*, from which they seem to be derived.

Geographical distribution of species. — Indian and South American; the species from these two regions seem to be really allied, and doubtless others will be discovered.

Larva (only the Indian species known) feeding in rolled leaves.

Foodplant *Breynia* (*Euphorbiaceae*).

- | | |
|--|-----------------|
| 1. <i>E. insipiens</i> , Meyrick, Exot. Microlep. Vol. 2, p. 149 (1918), Vol. 3, p. 11 | Bengal. |
| (1923). | |
| 2. <i>E. rhodocosma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 229 (1914). | Guiana, Brazil. |
| 3. <i>E. phanerosoma</i> , Meyrick, ibidem, p. 65 (1922). — Pl. I, Fig. III. | Brazil. |

39. GENUS PRAGMATODES, WALSINGHAM

Pragmatodes, Walsingham, Proc. Zool. Soc. Lond. p. 928 (1907). — Type : *P. fruticosella*, Walsingham.

Characters. — Head smooth; tongue developed. Antennae 4/5, in ♂ serrulate, basal joint without pecten. Labial palpi moderately long, recurved, second joint with appressed scales, terminal joint shorter, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely haired. Forewings 1b furcate, 2 from towards angle, 7 and 8 out of 6, 7 to costa. Hindwings under 1, elongate trapezoidal, apex strongly produced, cilia 3; 2-5 remote, 6 and 7 stalked, 6 weak.

Remarks. — Not known to me; perhaps allied to early forms of *Aristotelia*.

Geographical distribution of species. — Canary Islands.

Larva mining in leaves.

Foodplant *Rubia* (*Rubiaceae*).

1. *P. fruticosella*, Walsingham, Proc. Zool. Soc. Lond. p. 927, pl. 51, f. 10 (1907). — Canaries.

40. GENUS GLAUCE, CHAMBERS

Glauce, Chambers, Canad. Ent. Vol. 7, p. 11 (1875). — Type : *G. pectinalella*, Chambers.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, simple, basal joint elongate, without pecten. Labial palpi long, curved, ascending, second joint with appressed scales, slightly roughened beneath, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 6 and 7 out of 8, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex pointed, produced, termen sinuate-oblique beneath apex, cilia 2;

3 and 4 separate, 5 rather approximated, 6 and 7 approximated; in ♂ an expansible fringe or fan of strong flattened black bristles (or spines) from beneath costa towards base.

Remarks. — Probably related to transitional forms between *Aristotelia* and *Recurvaria*, but the curious fringe of bristles is unique. Busck's correction of Chambers with regard to the structure of vein 6 of forewings (Proc. U. S. Mus. Vol. 25, p. 783) is itself mistaken, and Chambers' original statement is correct.

Geographical distribution of species. — North American.

Larva unknown.

1. *G. pectinella*, Chambers, Canad. Ent. Vol. 7, p. 12 (1875) (*pectenalaella*). E. United States, Canada.

41. GENUS LEUCE, CHAMBERS

Leuce, Chambers, Canad. Ent. Vol. 7, p. 51 (1875). — Type : *L. fuscocristatella*, Chambers.

Naera, Chambers, ibidem, Vol. 7, p. 9 (1875) (praeocc.). — Type : *L. fuscocristatella*, Chambers.

Characters. — Head with appressed scales; tongue developed. Labial palpi moderately long, curved, ascending, second joint with large rough tuft of scales beneath, terminal joint shorter than second, thickened with scales, hardly pointed. Forewings with tufts of scales on surface; 1b furcate, 2 from 4/5, 3 and 4 connate from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, termen sinuate-emarginate; 3 and 4 approximated, 5 rather approximated to 4, 6 and 7 remote, nearly parallel.

Remarks. — Not known to me; seemingly allied to the two following.

Geographical distribution of species. — North American.

Larva unknown.

1. *L. fuscocristatella*, Chambers, Canad. Ent. Vol. 7, p. 9 (1875). Texas.
belfragesella, Chambers, Journ. Cinc. Soc. Nat. Hist. Vol. 2, p. 183 (1880).

42. GENUS PACHYGENEIA, MEYRICK

Pachygeneia, Meyrick, Exot. Microlep. Vol. 3, p. 11 (1923). — Type : *P. clitellaria*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, simple, basal joint moderate, without pecten. Labial palpi moderately long, curved, ascending, second joint much thickened with dense scales roughly projecting towards apex beneath, terminal joint as long as second, stout, roughened anteriorly, pointed. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above. Forewings with tufts of scales; 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex produced, acute, termen emarginate, cilia 1 2/3; 3-5 rather approximated towards base, 6 and 7 somewhat approximated.

Remarks. — Allied to early forms of *Aristotelia*.

Geographical distribution of species. — South American.

Larva unknown.

1. *P. clitellaria*, Meyrick, Exot. Microlep. Vol. 3, p. 11 (1923). Brazil, Peru.

43. GENUS LEPTOGENEIA, MEYRICK

Leptogeneia, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 412 (1904). — Type : *L. bicristata*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ shortly ciliated, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint with large rough projecting tuft of scales beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with tufts of scales on surface; 1b furcate, 2-4 approximated at base, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex produced, round-pointed, termen sinuate-emarginate, cilia 1 1/2-2; 3 and 4 approximated at base, 5 rather approximated to 4, 6 and 7 remote, nearly parallel.

Remarks. — Undoubtedly nearly allied to *Pachygeneia*, and therefore geographically interesting.

Geographical distribution of species. — Australian.

Larva unknown.

1. *L. bicristata*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 413 (1904). New South Wales,
W. Australia.

44. GENUS PHOTODOTIS, MEYRICK

Photodotis, Meyrick, Ann. Transv. Mus. Vol. 2, p. 229 (1911). — Type : *P. prochalina*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with dense projecting scales forming a short tuft beneath, terminal joint as long as second, thickened with appressed scales slightly rough anteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from angle, 3 sometimes absent, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, termen obliquely bisinuate beneath apex, cilia 1/2; 3 and 4 somewhat approximated, 5 rather bent, 6 and 7 parallel.

Remarks. — Although the forewings have no apparent scale-tufts, this genus is probably allied to the two preceding.

Geographical distribution of species. — South African.

Larva unknown.

1. *P. spilodoma*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 15 (1918). Zululand.
2. *P. prochalina*, Meyrick, ibidem, Vol. 2, p. 229 (1911). — Pl. I, Fig. 10. Transvaal, Natal, Kenya.

Group 3 (Gelechia type)

This and the two next groups may be regarded as parallel lines of development, referable to a common origin in some earlier form of the *Brachmia* group. The present group has vein 7 of the forewings always terminating in the costa, vein 2 rising separately and usually from before the angle, and commonly a characteristic prevalent type of rough brush-like scaling on the second joint of the labial palpi

beneath, which however is lost again in the higher forms. The terminal joint of palpi tends to be marked with two or sometimes three dark rings. Though very numerous the group is almost absent from India, and little represented in Australia, its greatest development being in North America and Europe.

45. GENUS HELICE, CHAMBERS

Helice, Chambers, Canad. Ent. Vol. 5, p. 187 (1873). — Type : *H. pallidochrella*, Chambers.

Theisoa, Chambers, ibidem, Vol. 6, p. 75 (1874). — Type : *H. constrictella*, Zeller.

Cacelice, Busck, Journ. N. York Ent. Soc. Vol. 10, p. 93 (1902). — Type : *H. pallidochrella*, Chambers.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennæ 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, smooth-scaled, second joint somewhat thickened towards apex, terminal joint longer than second, slender, acute. Maxillary palpi very small, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with tufts of scales on surface; 1b furcate, 2 from angle, 3 and 4 stalked, 5 absent, 7 and 8 out of 6 or in ♂ 8 absent, 7 to costa, 9 sometimes out of 6, 11 from middle. Hindwings 2/3, in both sexes narrow-lanceolate, or in ♀ narrow-trapezoidal, apex produced, acute, termen sinuate, cilia 3-4; 3 and 4 stalked, 5 absent, in ♀ 6 and transverse vein absent, in ♂ 7 absent (or rarely abnormally present).

Remarks. — This extraordinary genus is an aberrant member of this group. The forms with narrow-lanceolate hindwings and those with Gelechiad hindwings were originally regarded as belonging to distinct genera and families, the former being referred to the *Cosmopterygidae* (in which however their neuration would be quite abnormal), and the similarity in the characteristic markings being assumed to be due to analogy or mimicry; but Miss Braun has now shown (Canad. Ent. Vol. 51, p. 201, 1919) that the two different forms denote the sexes of *pallidochrella*; it therefore becomes evident that *pallidochrella* is a Gelechiad of which the ♂ has developed a hindwing of the Cosmopterygid form, and that the nearly allied *constrictella* is a more advanced type in which a similar form of wing has been transferred to the ♀ also. The explanation of these curious changes (which would have been deemed *prima facie* improbable in the highest degree) still remains to be investigated. I am indebted to Miss Braun for examples of both these species.

Geographical distribution of species. — North American.

Larva (*constrictella*) feeding beneath a web on under-side of leaves.

Foodplant *Ulmus*.

1. *H. multifasciella*, Chambers, Canad. Ent. Vol. 7, p. 93 (1875). Texas.
2. *H. constrictella*, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 291, pl. 4, f. 32 (1873). — **Pl. 1, Fig. 16; Pl. 5, Fig. 122a, b.** Ohio, Texas.

bifasciella, Chambers, Canad. Ent. Vol. 6, p. 75 (1874).

3. *H. pallidochrella*, Chambers, ibidem, Vol. 5, p. 188 (1873). — **Pl. 5**, Kentucky, Ohio.

Fig. 123a, b.

gledtschaeella, Chambers, ibidem, Vol. 9, p. 232 (1877).

permolestella, Busck, Journ. N. York Ent. Soc. Vol. 10, p. 93, pl. 12, f. 2 (1902).

46. GENUS STENOLECHIA, MEYRICK

Stenolechia, Meyrick, Ent. M. Mag. Vol. 30, p. 230 (1894). — Type : *S. gemmella*, Linnaeus.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennæ 4/5, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with appressed scales,

hardly rough beneath, terminal joint shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 absent, 3 from angle, 4 absent, 5 rather approximated, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex acute, produced, termen emarginate, cilia 3; 2-4 remote, parallel, 5 curved, rather approximated, 6 and 7 nearly parallel.

Remarks. — Perhaps a development of the following, and hitherto included with it.

Geographical distribution of species. — European and Indian.

Larva (only *gemmella* known) feeding in buds and shoots.

Foodplant *Quercus*.

1. <i>S. gemmella</i> , Linnaeus, Syst. Nat. Vol. 1, p. 539 (1758).	C. & S. E. Europe.
<i>nivea</i> , Haworth, Lep. Brit. p. 554 (1828).	
<i>nigrovittella</i> , Duponchel, Hist. Nat. Lép. Fr. Vol. 11, pl. 298, f. 5 (1838).	
<i>lepidella</i> , Zeller, Isis, p. 202 (1839).	
2. <i>S. sagittella</i> , Caradja, Iris, Vol. 34, p. 110 (1920).	Asia Minor.
3. <i>S. nigrinotella</i> , Zeller, Isis, p. 856 (1847).	S.E. Europe, Asia Minor.
<i>nigralbella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, f. 565 (1855).	
4. <i>S. orsicoma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 130 (1918) — Pl. I, Fig. 15.	Ceylon.
5. <i>S. trichaspis</i> , Meyrick, ibidem, Vol. 2, p. 131 (1918).	Ceylon.
6. <i>S. frustulenta</i> , Meyrick, ibidem, Vol. 3, p. 13 (1923).	Assam.
7. <i>S. marginipunctella</i> , Stainton, Trans. Ent. Soc. Lond. (2) Vol. 5, p. 118 (1859).	Bengal.
8. <i>S. zelosaris</i> , Meyrick, Exot. Microlep. Vol. 3, p. 12 (1923).	Assam.

47. GENUS PARACHRONISTIS, NOV. GEN.

Poecilia, Heinemann, Schmett. Deutschl. (2) Vol. 2, p. 281 (1870) (praeocc.). — Type: *P. albiceps*, Zeller.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint somewhat thickened with appressed scales, terminal joint shorter than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from angle, 4 absent, 3 and 5 approximated, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, pointed, termen emarginate, cilia 3; 2 remote, 3-5 rather approximated towards base, 6 and 7 nearly connate.

Remarks. — Nearly related to *Recurvaria*.

Geographical distribution of species. — European.

Larva unknown.

1. <i>P. albiceps</i> , Zeller, Isis, p. 202 (1839).	C. & S. E. Europe.
<i>alella</i> , Wood, Ind. Ent. f. 1225 (1839).	

48. GENUS OXYLECHIA, MEYRICK

Oxylechia, Meyrick, Trans. Ent. Soc. Lond. p. 39 (1917). — Type: *O. confirmata*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint with short rough

tuft beneath and hairs roughly expanded towards apex above, terminal joint longer than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with 1_b furcate, 2 from near angle, 4 absent, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 3/4, elongate-trapezoidal, apex somewhat produced, acute, termen obliquely bisinuate beneath apex, cilia 4; 2-4 remote, 5 from angle, 5 approximated to 4 at base, 6 and 7 connate.

Remarks. — Nearly allied to the preceding.

Geographical distribution of species. — South American.

Larva unknown.

1. *O. confirmata*, Meyrick, Trans. Ent. Soc. Lond. p. 39 (1917). — **Pl. I, Fig. 20.** Colombia.

49. GENUS TOSCA, HEINRICH

Tosca, Heinrich, Proc. U. S. Mus. Vol. 57, p. 65 (1920). — Type : *T. platonella*, Heinrich.

Characters. — Head smooth; tongue developed. Antennae 4/5, in ♂ simple, basal joint without pecten. Labial palpi long, slender, ascending, second joint slightly thickened and roughened beneath, terminal joint as long as second, acute. Maxillary palpi rudimentary. Forewings with 1_b furcate, 2 from angle, 4 absent, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, termen sinuate; 3 and 4 closely approximated at base, 5 and 6 nearly obsolete, 7 weak towards base.

Remarks. — A derivative of *Recurvaria*.

Geographical distribution of species. — North American.

Larva (*plutonella*) mining blotches in leaves.

Foodplant *Prunus*.

- | | |
|---|-------------|
| 1. <i>T. platonella</i> , Heinrich, Proc. U. S. Mus. Vol. 57, p. 68 (1920). | New Mexico. |
| 2. <i>T. pollostella</i> , Busck, ibidem, Vol. 30, p. 725 (1906). | Texas. |
| 3. <i>T. elachistella</i> , Busck, ibidem, Vol. 30, p. 725 (1906). | Texas. |

50. GENUS EVIPPE, CHAMBERS

Evippe, Chambers, Canad. Ent. Vol. 5, p. 185 (1873). — Type : *E. prunifoliella*, Chambers.

Phaetusa, Chambers, ibidem, Vol. 7, p. 105 (1875) (praeocc.). — Type : *E. leuconota*, Zeller.

Characters. — Head smooth-scaled; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint without pecten. Labial palpi long, recurved, second joint somewhat thickened beneath, hardly roughened, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1_b furcate, 2 from angle, 4 and 5 connate or short-stalked, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, pointed, termen sinuate, cilia 2 1/2, 3 and 4 connate, 5 approximated at base, 6 obsolete.

Remarks. — A derivative of *Recurvaria*.

Geographical distribution of species. — North American.

Larva (*prunifoliella*) feeding under turned-down tip of leaf.

Foodplant *Prunus*.

- | | |
|--|--------------------------|
| 1. <i>E. prunifoliella</i> , Chambers, Canad. Ent. Vol. 5, p. 186 (1873). | N. E. United States, |
| 2. <i>E. leuconota</i> , Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 268, pl. 3, f. 21 (1873). | Texas, Canada. [Canada.] |
| <i>plutella</i> , Chambers, Canad. Ent. Vol. 7, p. 106 (1875). | |

51. GENUS AGNIPPE, CHAMBERS

Agnippe, Chambers, Canad. Ent. Vol. 4, p. 194 (1872). — Type : *A. bicolorella*, Chambers.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint slightly thickened, rather rough-scaled beneath towards apex, terminal joint nearly as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with subdorsal scaletuft; 1b furcate, 2 from towards angle, 4 and 5 stalked or coincident, 7 and 8 out of 6, 7 to costa, 11 absent. Hindwings nearly 1, elongate-trapezoidal, apex produced, pointed, termen sinuate; 3-5 rather approximated at base, 6 obsolete, cell open between 5 and 7.

Remarks. — Apparently related to the preceding.

Geographical distribution of species. North American.

Larva unknown.

- | | |
|---|-----------|
| 1. <i>A. fuscopulvella</i> , Chambers, Canad. Ent. Vol. 4, p. 195 (1872). | Kentucky. |
| 2. <i>A. bicolorella</i> , Chambers, ibidem, Vol. 4, p. 195 (1872) (<i>bisc-</i>). | Kentucky. |
| 3. <i>A. evippella</i> , Busck, Proc. U. S. Mus. Vol. 30, p. 723 (1906) (- <i>peella</i>). | Texas. |

52. GENUS SMENODOCA, MEYRICK

Smenodoca, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 302 (1904). — Type : *S. erebenna*, Meyrick.

Characters. — Head smooth; ocelli present; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, slightly rough beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex produced, pointed, termen sinuate, cilia 2; 3 and 4 stalked, 5 becoming obsolete basally, 6 and transverse vein apparently obsolete, 8 anastomosing strongly with margin of cell.

Remarks. — Allied to *Recurvaria*.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|--|-----------------------|
| 1. <i>S. erebenna</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 303 (1904). | S. E. & W. Australia. |
|--|-----------------------|

53. GENUS ELASIPRORA, MEYRICK

Elasiprora, Meyrick, Trans. Ent. Soc. Lond. p. 230 (1914). — Type: *E. rostrifera*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae $\frac{3}{4}$, in ♂ serrulate, minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint somewhat thickened, slightly roughened beneath towards apex, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long fine hairs above. Forewings with 1 *b* simple, 2 from angle, 4 absent, 7 absent, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex somewhat produced, pointed, termen somewhat emarginate, oblique. Cilia 2; cell rather wide, transverse vein absent, 3 and 4 connate, 5 nearly approximated to 4 at base, 6 and 7 stalked, 6 to apex.

Remarks. — A rather peculiar form of the *Recurvaria* group.

Geographical distribution of species. — South American.

Larva unknown.

- E. rostrifera*, Meyrick, Trans. Ent. Soc. Lond. p. 231 (1914). Guiana, Brazil.

54. GENUS HAPALOSARIS, MEYRICK

Haplosarxis, Meyrick, Trans. Ent. Soc. Lond. p. 37 (1917). — Type: *H. petulans*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae $\frac{3}{4}$, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, curved, ascending, second joint above in ♂ clothed with long fine expandible hairs, in ♀ with triangularly expanded hair-scales towards apex, beneath smooth-scaled, terminal joint somewhat shorter than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long rough hairs above. Forewings with 1 *b* furcate, 2 from very near angle, 3 from angle, 4 absent, 5 approximated, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings $\frac{3}{4}$, elongate-trapezoidal, apex acute, termen obliquely bisinuate beneath apex, cilia 3; 2 remote, 3 and 4 almost connate from angle, 5 approximated, 6 obsolete.

Remarks. — Nearly related to *Recurvaria*.

Geographical distribution of species. — South American.

Larva unknown.

- I. H. petulans*, Meyrick, Trans. Ent. Soc. Lond. p. 37 (1917). Colombia, Ecuador, Peru.

55. GENUS COLONANTHES, MEYRICK

Colonanthes, Meyrick, Exot. Microlep. Vol. 3, p. 12 (1923). — Type : *C. plectanopa*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae $\frac{3}{4}$, in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint beneath with triangular median and apical tufts of scales, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Anterior tibiae and basal joint of tarsi thickened with rough scales above; posterior tibiae clothed with hairs above. Forewings with tufts of scales; 1^b furcate, 2 from

towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, termen emarginate beneath produced apex, cilia 2; 3-5 somewhat approximated at base, 6 and 7 parallel.

Remarks. — This interesting form probably approximates to the line of descent passing from *Recurvaria* to *Aristotelia*.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. plectanopa*, Meyrick, Exot. Microlep. Vol. 3, p. 12 (1923).

Brazil, Peru.

56. GENUS COMPOSARIS, MEYRICK

Composaris, Meyrick, Trans. Ent. Soc. Lond. p. 233 (1914). — Type : *C. testacea*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint long, without pecten. Labial palpi very long, curved, ascending, second joint above with scales expanded at apex, beneath with long rough projecting hair-scales throughout and denser projecting apical tuft, terminal joint shorter, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with tufts of scales; 1b furcate, 2 from towards angle, 3-5 approximated, 6 near 7, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 3/4, narrow-trapezoidal, apex produced, termen obliquely bisinuate, cilia 3; 3 and 4 rather remote, 5 approximated to 4 at base, 6 and 7 nearly parallel, transverse vein very oblique inwards from 6 to 7.

Remarks. — Closely related to *Recurvaria*.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. flavidella*, Busck, Proc. U. S. Mus. Vol. 47, p. 12 (1914).

Panama.

2. *C. testacea*, Meyrick, Trans. Ent. Soc. Lond. p. 234 (1914).

Guiana, Brazil.

57. GENUS PHYLOPATRIS, MEYRICK

Phylopatria, Meyrick, Exot. Microlep. Vol. 3, p. 14 (1923). — Type : *P. terpnodes*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ rather stout, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with rough projecting scales beneath, terminal joint longer than second, rather stout, pointed. Maxillary palpi rudimentary. Posterior tibiae clothed with long fine hairs above. Forewings with small tufts of scales; 2 from near angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, termen faintly bisinuate, angulated on 4. cilia 1 1/2; 3 and 4 connate, 5 approximated at base, 6 and 7 rather approximated towards base.

Remarks. — Also closely allied to *Recurvaria*, of which it must approach the ancestral form.

Geographical distribution of species. — South American.

Larva unknown.

1. *P. terpnodes*, Meyrick, Exot. Microlep. Vol. 3, p. 15 (1923).

Brazil, Peru.

58. GENUS RECURVARIA, HAWORTH

Recurvaria, Haworth, Lep. Brit. p. 547 (1828). — Type : *R. nanella*, Hübner.

Evagora, Clemens, Proc. Acad. Nat. Sc. Philad. p. 165 (1860). — Type : *R. apicitripunctella*, Clemens.

Eidothea, Chambers, Canad. Ent. Vol. 5, p. 186, 229 (1873). — Type : *R. dorsivittella*, Zeller.

Sinoe, Chambers, ibidem, Vol. 5, p. 229 (1873). — Type : *R. robiniella*, Fitch.

Aphanaula, Meyrick, Handb. Brit. Lep. p. 579 (1895). — Type : *R. leucatella*, Clerck.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with appressed scales or more or less rough beneath, terminal joint shorter than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with small tufts of scales; 1b furcate, 2 from angle, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex acute, termen sinuate, cilia 2-3; 3 and 4 connate, 5 approximated, 6 and 7 somewhat approximated.

Remarks. — Probably a development from *Telphusa*.

Geographical distribution of species. — Characteristically American, with only isolated stragglers elsewhere.

Larva (20 species known) feeding between spun leaves, or mining in leaves or inflorescence or (*annulicornis*) stated bred from nests of wasp (*Polistes*).

Foodplants especially *Coniferae* (10), but also *Rhamnaceae*, *Cupuliferae*, and several other Orders, usually trees or shrubs.

1. <i>R. thiodes</i> , Meyrick, Trans. Ent. Soc. Lond. p. 38 (1917).	Colombia.
2. <i>R. annulicornis</i> , Walsingham, Proc. Zool. Soc. Lond. p. 63 (1897)	Virgin Is.
3. <i>R. thysanota</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 45, pl. 2, f. 5 (1910).	Mexico.
4. <i>R. obliquistrigella</i> , Chambers, Canad. Ent. Vol. 4, p. 65 (1872).	Kentucky.
5. <i>R. juniperella</i> , Kearfott, Journ. N. York Ent. Soc. Vol. 11, p. 157, pl. 9, f. 3, 17 (1903).	New Jersey.
6. <i>R. penetrans</i> , Meyrick, Exot. Microlep. Vol. 3, p. 13 (1923).	Brazil.
7. <i>R. saxea</i> , Meyrick, ibidem, Vol. 3, p. 14 (1923).	Brazil.
8. <i>R. rhinota</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 44, pl. 2, f. 4 (1910).	Guatemala.
9. <i>R. robiniella</i> , Fitch, Rep. Ins. N. York. Vol. 5, p. 334 (1859). <i>fuscopallidella</i> , Chambers, Canad. Ent. Vol. 5, p. 231 (1873). <i>robiniæfoliella</i> , Chambers, Rep. U. S. Dep. Agric. p. 224 (1879).	E. United States.
10. <i>R. putella</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 11 (1914).	Panama.
11. <i>R. sticta</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 46 (1910).	Mexico.
12. <i>R. trigonophorella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 343, pl. 4, f. 112 (1877).	Colombia.
13. <i>R. rhombophorella</i> , Zeller, ibidem, Vol. 13, p. 344, pl. 4, f. 113 (1877).	Colombia.
14. <i>R. intermissella</i> , Zeller, ibidem, Vol. 13, p. 347, pl. 4, f. 115 (1877).	Colombia.
15. <i>R. senariella</i> , Zeller, ibidem, Vol. 13, p. 346, pl. 4, f. 114 (1877).	Colombia.
16. <i>R. ostariella</i> , Walsingham, Proc. Zool. Soc. Lond. p. 65 (1897).	Virgin Is.
17. <i>R. merismatella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 352, pl. 5, f. 119, (1877).	Colombia.
18. <i>R. pleurosaris</i> , Meyrick, Exot. Microlep. Vol. 3, p. 14 (1923).	Brazil.

19. *R. filicornis*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 350, pl. 5, f. 117 Colombia.
(1877).
20. *R. febriculella*, Zeller, ibidem. Vol. 13, p. 341, pl. 4, f. 111 (1877). Colombia.
21. *R. melanostictella*, Zeller, ibidem, Vol. 13, p. 351, pl. 5, f. 118 (1877). Colombia.
22. *R. argentialbella*, Chambers, Canad. Ent. Vol. 6, p. 24 (1874). Texas.
23. *R. variella*, Chambers, ibidem, Vol. 4, p. 174 (1872). Kentucky.
24. *R. apicitrispunctella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 165 (1860). E. United States, Canada.
altritella, Walker, List. Lep. Het. Brit. Mus. Vol. 29, p. 592 (1864).
abietisella, Packard, Rep. U. S. Dep. Agr. p. 150 (1883).
25. *R. nothostigma*, Meyrick, Trans. Ent. Soc. Lond. p. 233 (1914). Brit. Guiana.
26. *R. colubrinae*, Busck, Proc. U. S. Mus. Vol. 25, p. 810 (1903). Texas.
27. *R. cristatella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 241 (1875). Kentucky.
28. *R. moreonella*, Heinrich, Proc. U. S. Mus. Vol. 57, p. 65 (1920). Colorado.
29. *R. Milleri*, Busck, Proc. Ent. Soc. Wash. Vol. 16, p. 144 (1914). California.
30. *R. thujella*, Kearfott, Journ. N. York Ent. Soc. Vol. 11, p. 154, pl. 9, f. 8, 21 (1903) (-jacella). — Pl. I. Fig. 19. New Jersey.
31. *R. conifarella*, Kearfott, Canad. Ent. Vol. 39, p. 3 (1907). Ontario.
32. *R. pinella*, Busck, ibidem, Vol. 38, p. 212 (1906). Colorado.
33. *R. piceella*, Kearfott, Journ. N. York Ent. Soc. Vol. 11, p. 155, pl. 9, f. 10, 19 (1903) (-cearella). New Jersey.
34. *R. gibsonella*, Kearfott, Canad. Ent. Vol. 39, f. 4 (1907). Ontario.
35. *R. eromene*, Walsingham, Proc. Zool. Soc. Lond. p. 64 (1897). Virgin Is.
36. *R. ornatipalpella*, Walsingham, ibidem, p. 64 (1897). Grenada.
37. *R. aequorea*, Meyrick, Trans. Ent. Soc. Lond. p. 39 (1917). Peru.
38. *R. alnifructella*, Busck, Proc. Ent. Soc. Wash. Vol. 17, p. 82 (1915). Virginia.
39. *R. dorsivittella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 267, pl. 3, f. 20 (1873). E. United States, Virgin Is.
vagatioella, Chambers, Canad. Ent. Vol. 5, p. 186 (1873).
40. *R. quercivorella*, Chambers, ibidem, Vol. 4, p. 173 (1872). E. United States.
gilviscopella, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 266 (1873).
41. *R. flagellifera*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 45 (1910). Mexico.
42. *R. sartor*, Walsingham, ibidem, Vol. 4, p. 44, pl. 2, f. 7 (1910). Mexico.
43. *R. dryozona*, Meyrick, Exot. Microlep. Vol. 1, p. 568 (1916). Bengal, Ceylon.
44. *R. invictella*, Busck, Ent. News, Philad. p. 316 (1908). California.
45. *R. ptilastis*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 369 (1909). Cape Colony.
ptychophora, Meyrick, ibidem, Vol. 10, p. 244 (1914).
46. *R. cinerella*, Chrétien, Bull. Mus. Hist. Nat. Paris, Vol. 14, p. 361 (1908). Canaries.
47. *R. xanthotricha*, Meyrick, Trans. Ent. Soc. Lond. p. 38 (1917). Peru.
48. *R. nanella*, Hübner, Samml. Eur. Schmett. Tin. f. 267 (1805). Europe, Canada,
pruniella, Hübner, ibidem, f. 268 (1805). Pennsylvania.
crataegella, Busck, Proc. U. S. Mus. Vol. 25, p. 812 (1903).
49. *R. picula*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 46, pl. 2, f. 6 (1910). Mexico.
50. *R. ceanothiella*, Braun, Ent. News, Philad. Vol. 32, p. 10 (1921). California.
51. *R. nigra*, Busck, Proc. U. S. Mus. Vol. 25, p. 814 (1903). District of Columbia.
52. *R. kittella*, Walsingham, Proc. Zool. Soc. Lond. p. 65 (1897). Hayti.
53. *R. graphicella*, Busck, Proc. Ent. Soc. Wash. Vol. 9, p. 86 (1908). California.
54. *R. albidorsella*, Snellen, Tijdschr. v. Ent. Vol. 27, p. 169, pl. 9, f. 4 (1884). E. Siberia.
55. *R. leucatella*, Clerck, Icon. Ins. p. 2, f. 3 (1759). Europe, Asia Minor.
albocinetella, Duponchel, Hist. Nat. Lep. Fr. Vol. 11, pl. 298, f. 13 (1838).

59. GENUS EUCORDYLEA, DIETZ

Eucordylea, Dietz, Ent. News, Philad. Vol. 11, p. 349 (1900). — Type : *E. atropictella*, Dietz.

Characters. — Head smooth; ocelli posterior; tongue absent. Antennae 4/5, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint in ♂ clothed above with dense expandible tuft of long hairs, in ♀ with appressed scales, terminal joint much shorter than second, moderate, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long rough hair-scales above. Forewings with small tufts of scales; 1b furcate, 2 from towards angle, 3-5 closely approximated at base, 7 and 8 out of 6, 7 to costa, 11 from beyond middle. Hindwings nearly 1, trapezoidal, apex obtuse, termen bisinuate, cilia 2; 3-5 closely approximated from angle, 6 and 7 connate.

Remarks. — Correlated with *Recurvaria*.

Geographical distribution of species. — North American.

Larva (*gallicola*) feeding in galls caused by a sawfly (*Euura*).

Foodplant *Salix*.

1. <i>E. gallicola</i> , Busck, Proc. Ent. Soc. Wash. Vol. 17, p. 81 (1915).	Colorado.
2. <i>E. elucidella</i> , Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 227 (1920).	California.
3. <i>E. atropictella</i> , Dietz, Ent. News, Philad. Vol. 11, p. 350 (1900).	Pennsylvania, Ontario.

60. GENUS NESOLECHIA, MEYRICK

Nesolechia, Meyrick, Exot. Microlep. Vol. 2, p. 425 (1921). — Type : *N. horogramma*, Meyrick.

Characters. — Head with appressed scales, sidetufts slightly raised; ocelli small, posterior; tongue developed. Antennae 2/3, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, slightly roughened anteriorly, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2-5 parallel, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings somewhat under 1, elongate-trapezoidal, apex somewhat produced, termen sinuate, cilia 1; 3-5 rather approximated at base, 6 and 7 stalked.

Remarks. — Whilst belonging to this neighbourhood generally, the precise affinity of this genus is dubious.

Geographical distribution of species. — Polynesian.

Larva unknown.

1. <i>N. horogramma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 425 (1921).	Fiji.
---	-------

61. GENUS EXOTELEIA, WALLENGREN

Exoteleia, Wallengren, Ent. Tidsskr. Vol. 2, p. 94 (1881). — Type : *E. dodecella*, Linnaeus.

Heringia, Hedemann (1894). — Type : *E. dodecella*, Linnaeus.

Paralechia, Busck, Proc. U. S. Mus. Vol. 25, p. 820 (1903). — Type : *E. pinifoliella*, Chambers.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint moderate, without pecten. Labial palpi moderately long, recurved,

second joint somewhat thickened beneath with rough scales, terminal joint shorter than second, moderate, pointed. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with small tufts of scales; 1b furcate, 2 nearly from angle, 7 and stalked, 7 to costa, 11 from middle. Hindwings nearly 1, elongate-trapezoidal, termen sinuate beneath apex, cilia nearly 2; 3 and 4 connate or short-stalked, 5 rather approximated, 6 and 7 parallel.

Remarks. — Nearly related to *Recurvaria*.

Geographical distribution of species. — Palaearctic and North American.

Larva feeding in shoots and between spun leaves.

- | | |
|---|-------------------|
| 1. <i>E. californica</i> , Busck, Proc. Ent. Soc. Wash. Vol. 9, p. 92 (1907). | California. |
| 2. <i>E. pinifoliella</i> , Chambers, Journ. Cinc. Soc. Nat. Hist. Vol. 2, p. 181 (1880). | E. United States. |
| 3. <i>E. trijugella</i> , Erschoff, Hor. Soc. Ent. Ross. Vol. 12, p. 345 (1877). | E. Siberia. |
| 4. <i>E. dodocella</i> , Linnaeus, Syst. Nat. Vol. 1, p. 539 (1758) | C. & S. Europe. |
| <i>annulicornis</i> , Stephens, Ill. Brit. Ent. Haust. Vol. 4, p. 208 (1835). | |
| <i>favillaticella</i> , Zeller, Isis, p. 201 (1839). | |
| <i>reussiella</i> , Ratzeburg, Forst Ins. Vol. 2, p. 240, pl. 15, f. 5 (1840). | |
| 5. <i>E. succinctella</i> , Zeller, Stett. Ent. Zeit. Vol. 33, p. 108 (1872). | Switzerland. |

62. GENUS BATENIA, CHRÉTIEN

Batenia, Chrétien, Bull. Soc. Ent. Fr. p. 57 (1908). — Type : *B. fasciella*, Chrétien.

Characters. — Head smooth, crown loosely scaled; ocelli posterior; tongue developed. Antennae 3/4, basal joint somewhat enlarged, without pecten. Labial palpi moderately long, slightly curved, slender, second joint with appressed scales, terminal joint shorter than second, pointed. Maxillary palpi very short. Forewings with 2-5 nearly equidistant, 6 from near 7, 8 and 9 out of 7, 7 to costa. Hindwings 1/2, narrow-trapezoidal, apex long-produced, termen emarginate; 3 and 4 connate, 5 remote, 6 and 7 stalked.

Remarks. — I have not seen this genus, which is of uncertain affinity.

Geographical distribution of species. — North African.

Larva unknown.

- | | |
|--|----------|
| 1. <i>B. fasciella</i> , Chrétien, Bull. Soc. Ent. Fr. p. 58 (1908). | Algeria. |
|--|----------|

63. GENUS EPHYSTERIS, MEYRICK

Ephysteris, Meyrick, Proc. Zool. Soc. Lond. p. 724 (1908). — Type : *E. chersaea*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately long, without pecten. Labial palpi moderately long, recurved, second joint beneath with rough projecting scales, terminal joint shorter than second, loosely scaled, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 2-5 parallel, 7, 8, 9 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex strongly produced, termen emarginate, cilia 3; 3 and 4 connate or stalked, 5 rather approximated, 6 and 7 tolerably parallel.

Remarks. — Probably a derivative of *Epithetis*.

Geographical distribution of species. — African, Indo-Malayan, and Australian, but without doubt artificially spread, and probably Indian by origin.

Larva feeding in stems of cultivated cereals and grasses.

Foodplant *Gramineae*.

- | | |
|---|--|
| 1. <i>E. chersaea</i> , Meyrick, Proc. Zool. Soc. Lond. p. 725 (1908). | Egypt, S. Africa, India,
Ceylon, New Guinea,
Queensland. |
| <i>oschophora</i> , Meyrick, Rec Ind. Mus. Vol. 5, p. 219 (1910). | |
| ? <i>despectella</i> , Walker, List. Lep. Het. Brit. Mus. Vol. 28, p. 477 (1863). | |

64. GENUS CLISTOTHYRIS, ZELLER

Clistothyris, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 330 (1877). — Type : *C. villosula*, Zeller.

Characters. — Head smooth; tongue short. Antennæ in ♂ simple, basal joint elongate. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short. Posterior tibiae with long hairs above. Forewings above with a naked oval glandular spot towards costa rather near base (beneath scaled). Hindwings 1, elongate-trapezoidal, apex pointed, termen slightly sinuate, oblique, cilia over 1; hyaline median and subdorsal streaks, dorsal area clothed with short hairs.

Remarks. — I have not seen this genus, and Zeller does not attempt to give the neuration, therefore the standing of the genus is problematical; according to the figure and characters given it appears to be related to *Recurvaria* or *Epithecis*; the glandular spot of forewings, and hyaline streaks and dorsal hairs of hindwings should enable the genus and species to be recognised, but I probably should not myself regard these as being of generic value.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. villosula*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 331, pl. 4, f. 104 (1877). Colombia.

65. GENUS TRYPANISMA, CLEMENS

Trypanisma, Clemens, Proc. Acad. Nat. Sc. Philad. p. 168 (1860). — Type : *T. prudens*, Clemens.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennæ 2/3, in ♂ simple, basal joint elongate, without pecten. Labial palpi moderately long, recurved, second joint slightly thickened and roughened beneath, terminal joint as long as second, moderate, pointed. Maxillary palpi rudimentary. Forewings with 1b furcate, 2 from angle, 3 and 4 stalked, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, pointed, termen emarginate; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — Probably related to *Epithecis*.

Geographical distribution of species. — North American.

Larva (both species) feeding beneath a silken web on underside of leaves, also gaining access to upperside through small round holes.

Foodplants *Quercus*, *Fagus*.

- | | |
|---|-----------------|
| 1. <i>T. fagella</i> , Busck, Proc. U. S. Mus. Vol. 25, p. 816 (1903). | Dist. Columbia. |
| 2. <i>T. prudens</i> , Clemens, Proc. Acad. Nat. Sc. Philad. p. 168 (1860). | Dist. Columbia, |
| <i>quinqueannulella</i> , Chambers, Canad. Ent. Vol. 4, p. 191 (1872). | [Pennsylvania.] |

66. GENUS ISTRIANIS, MEYRICK

Istrianis, Meyrick, Exot. Microlep. Vol. 2, p. 130 (1910). — Type : *I. crauropa*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint rather thickened, rough-scaled beneath, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with tufts of scales on surface; 2 from towards angle, 7 and 8 out of 6, 7 to costa, 9 connate, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex produced, acute, termen sinuate, cilia 3; 3 and 4 connate, 5 somewhat approximated, 6 and 7 long-stalked.

Remarks. — Somewhat intermediate in structure between *Recurvaria* and *Epithestis*.

Geographical distribution of species. — Indian.

Larva feeding on underside of leaves.

Pupa in spindle-shaped cocoon on leaf.

Foodplant *Butea* (*Leguminosae*).

1. *I. crauropa*, Meyrick, Exot. Microlep. Vol. 2, p. 130 (1910).

Kanara.

67. GENUS EPITHECTIS, MEYRICK

Epithestis, Meyrick, Handb. Brit. Lep. p. 580 (1895). — Type : *E. lathyri*, Stainton.

Taygete, Chambers, Canad. Ent. Vol. 5, p. 229 (1873) (praeocc.). — Type : *E. attributella*, Walker.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed or rather rough scales beneath, terminal joint as long as second or shorter, moderate, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from near angle, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex pointed, termen more or less sinuate, cilia 1-2; 3 and 4 connate, 5 rather approximated, 6 and 7 closely approximated or connate or stalked.

Remarks. — Probably derivable from *Telphusa*.

Geographical distribution of species. — Represented in all regions, yet not very numerous anywhere; chiefly found in Europe, America and Africa.

Larva (9 known) usually in spun shoots or leaves, also one in *Cynips*-galls and one on dried plants.

Foodplants. — *Leguminosae* (3), *Rosaceae*, *Ericaceae*, *Zygophyllaceae*, etc.

- | | |
|--|---------------|
| 1. <i>E. consociata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 232 (1914). | Brit. Guiana. |
| 2. <i>E. lasciva</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 47, pl. 2, f. 8 (1910). | Panama. |
| 3. <i>E. barydelta</i> , Meyrick, Exot. Microlep. Vol. 3, p. 15 (1923). | Brazil. |
| 4. <i>E. citranthes</i> , Meyrick, ibidem, Vol. 3, p. 15 (1923). | Brazil. |
| 5. <i>E. balsamopa</i> , Meyrick, ibidem, Vol. 3, p. 16 (1923). | Brazil, Peru. |
| 6. <i>E. notospila</i> , Meyrick, ibidem, Vol. 3, p. 15 (1923). | Brazil. |
| 7. <i>E. citrinella</i> , Barnes, Contr. N. Amer. Lep. Vol. 4, p. 224 (1920). | Arizona. |

8. *E. saundersella*, Chambers, Canad. Ent. Vol. 8, p. 173 (1876). Kentucky.
 9. *E. galligenitella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 420 (1864) N. E. United States.
(gallae-).
geminella, Riley, Canad. Ent. Vol. 3, p. 195 (1871).
 10. *E. subsimella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 173 (1860). Pennsylvania.
 11. *E. attributella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 593 (1864). — N. E. United States,
Pl. I, Fig. 21. Canada.
difficilisella, Chambers, Canad. Ent. Vol. 4, p. 65 (1872).
 12. *E. sylvicolella*, Busck, Proc. U. S. Mus. Vol. 25, p. 818 (1903). New York.
 13. *E. platysoma*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 47 (1910). Mexico.
 14. *E. bicostimaculella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 127 Colorado, Arizona.
(1877) (bicosto-).
thoracella, Walsingham, Insect Life, Vol. 1, p. 147 (1888).
 15. *E. critica*, Walsingham, Biol. Centr.-Amer. Lep. Het Vol. 4, p. 48, pl. 2, f. 9 (1910). Mexico.
 16. *E. nitrariella*, Chrétien, Bull. Soc. Ent. Fr. p. 56 (1908). Algeria.
 17. *E. petiginella*, Mann, Verh. Zool.-bot. Ges. Wien, Vol. 17, p. 843 (1867). S. Tyrol.
 18. *E. delminiella*, Rebel, Ann. Hofmus. Wien, Vol. 19, p. 353 (1904). Herzegovina.
 19. *E. amoenella*, Frey, Mitth. Schweiz. Ent. Ges. Vol. 6, p. 366 (1882). S. Tyrol.
 20. *E. lathyri*, Stainton, Ent. Annual. p. 130 (1865). England, Latvia.
nigricostella, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 244 (1850).
 21. *E. nigricostella*, Duponchel, Hist. Nat. Lép. Fr. Suppl. Vol. 4, p. 288, pl. 74, f. 9 (1842). E. C. & S. E. Europe, Asia Minor.
 22. *E. ulicinella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 240 (1859). Spain, S. France.
 23. *E. lacrimosa*, Meyrick, Ent. Mitth. Berlin, Vol. 2, p. 299 (1913). Tunis.
 24. *E. ignavella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 368, pl. 5, f. 125 (1877). Colombia.
 25. *E. telifera*, Meyrick, Rec. Ind. Mus. Vol. 5, p. 220 (1910). Himalayas.
 26. *E. sordidula*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 285 (1913). Transvaal.
 27. *E. farinata*, Meyrick, ibidem, Vol. 3, p. 285 (1913). Transvaal.
 28. *E. carinata*, Meyrick, ibidem, Vol. 3, p. 64 (1912). Transvaal.
 29. *E. emerita*, Meyrick, ibidem, Vol. 8, p. 67 (1921). Transvaal.
 30. *E. lacunosa*, Meyrick, ibidem, Vol. 6, p. 16 (1918). Natal.
 31. *E. profusa*, Meyrick, ibidem, Vol. 8, p. 67 (1921). Rhodesia.
 32. *E. ochrocosma*, Meyrick, ibidem, Vol. 2, p. 230 (1911). Transvaal, Natal.
 33. *E. mesoleuca*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 48 (1900). Victoria.
 34. *E. zophochalca*, Meyrick, Trans. N. Zeal. Inst. Vol. 50, p. 133 (1918). New Zealand.
 35. *E. mouffetella*, Schiffermüller, Syst. Verz. Schmett. Wien, p. 140 (1776). Europe.
pedisequella, Hübner, Samml. Eur. Schmett. Tin. f. 95 (1796).
 36. *E. medjella*, Chrétien, Bull. Soc. Ent. Fr. p. 191 (1900). France.
 37. *E. pruinosa*, Zeller, Isis, p. 288 (1846). Finland, Latvia, Ger-
 38. *E. adumbratella*, Snellen, Tijdschr. v. Ent. Vol. 27, p. 170, pl. 9, f. 5 (1884). E. Siberia. [many.
 39. *E. extincta*, Meyrick, Ann. Transv. Mus. Vol. 2, p. 231 (1911). Transvaal.
 40. *E. triatomea*, Mühlig, Stett. Ent. Zeit. Vol. 25, p. 101 (1864). France, Germany.
 41. *E. studiosa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 591 (1905). Ceylon.

68. GENUS LATROLOGA, MEYRICK

Latrologa, Meyrick, Exot. Microlep. Vol. 2, p. 132 (1918). — Type : *L. aoropis*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 5/6, in ♂ serrulate, pubescent, basal joint moderate, without pecten. Labial palpi long,

recurved, smooth-scaled, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs. Forewings with 1b furcate, 2 from near angle, 6 and 8 stalked, 7 absent, 11 from middle. Hindwings slightly under 1, elongate-trapezoidal, apex obtuse-pointed, termen faintly sinuate, cilia 1 2/3; 3 and 4 stalked, 5 rather approximated, 6 and 7 long-stalked.

Remarks. — Apparently related to the preceding.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *L. aoropis*, Meyrick, Exot. Microlep. Vol. 2, p. 132 (1918). Ceylon.

69. GENUS APOCRITICA, NOV. GEN.

Type : *A. chromatica*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 from near angle, 7 absent, 11 from middle. Hindwings 1, elongate-trapezoidal, apex prominent, termen sinuate, cilia nearly 1; 3 and 4 connate, 5 nearly parallel, 6 and 7 tolerably remote, rather diverging.

Remarks. — Probably related to *Latrologa*.

Geographical distribution of species. — Seychelles Islands.

Larva unknown.

1. *A. chromatica*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 272 (1911). Seychelles.

70. GENUS CONIOGYRA, MEYRICK

Coniogyra, Meyrick, Ann. Transv. Mus. Vol. 8, p. 66 (1921). — Type : *C. dilucescens*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, basal joint moderate, without pecten. Labial palpi moderate, slightly curved, subascending, slender, with appressed scales, terminal joint somewhat shorter than second, pointed. Maxillary palpi obsolete. Posterior tibiae clothed with hairs above. Forewings with 2 from angle, 4 absent, 7 and 8 stalked, 7 to costa, 9 connate, 11 from middle. Hindwings 2/3, elongate-trapezoidal, apex pointed, produced, termen sinuate, cilia 3; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Seemingly also allied to *Epithecis*.

Geographical distribution of species. — South African.

Larva unknown.

1. *C. dilucescens*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 66 (1921). Rhodesia.

71. GENUS COUDIA, CHRÉTIEN

Coudia, Chrétien, Ann. Soc. Ent. Fr. p. 326 (1915). — Type : *C. strictella*, Chrétien.

Characters. — Head smooth; tongue developed. Antennae 4/5, basal joint without pecten. Labial palpi moderately long, curved, ascending, not rising above forehead, second joint with appressed scales, terminal joint much shorter than second, acute. Maxillary palpi rudimentary. Forewings with 1_b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, trapezoidal, apex pointed, termen sinuate; 3 and 4 connate, 5 approximated, 6 and 7 connate.

Remarks. — Not known to me, but probably referable here.

Geographical distribution of species. — North African.

Larva unknown.

1. *C. strictella*, Chrétien, Ann. Soc. Ent. Fr. p. 326 (1915). Algeria.

72. GENUS GUEBLA, CHRÉTIEN

Guebla, Chrétien, Ann. Soc. Ent. Fr. p. 324 (1915). — Type : *G. compositella*, Chrétien.

Characters. — Head smooth; tongue weak. Antennae 4/5, basal joint without pecten. Labial palpi moderate, curved, ascending, not rising above forehead, second joint with appressed scales, terminal joint shorter than second, acute. Maxillary palpi very short. Forewings with 1_b furcate, 2 from near angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, apex produced, pointed, termen emarginate; 3 and 4 connate or closely approximated, 5 approximated, 6 and 7 remote, slightly approximated towards base.

Remarks. — Also unknown to me.

Geographical distribution of species. — North African.

Larva unknown.

1. *G. compositella*, Chrétien, Ann. Soc. Ent. Fr. p. 325 (1915). Algeria.

2. *G. candidella*, Chrétien, ibidem, p. 325 (1915). Algeria.

73. GENUS PARAPSECTRIS, MEYRICK

Parapsectris, Meyrick, Ann. Transv. Mus. Vol. 2, p. 230 (1911). — Type : *P. tholaea*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, laterally compressed, somewhat rough and furrowed beneath, above with rather rough scales towards apex, terminal joint shorter than second, considerably thickened with scales, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 1_b furcate, 2 from angle, 6 sometimes out of 7 near base, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1 or nearly, trapezoidal, apex pointed, termen obliquely sinuate beneath apex, cilia 1 or over 1; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — A derivative of *Gelechia*.

Geographical distribution of species. — African.

Larva unknown.

- | | |
|--|-------------------|
| 1. <i>P. ferulata</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 17 (1918). | Natal. |
| 2. <i>P. anxia</i> , Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 4 (1917). | Cape Colony. |
| 3. <i>P. fastidiosa</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 66 (1912). | Transvaal. |
| 4. <i>P. neograpta</i> , Meyrick, ibidem, Vol. 4, p. 192 (1914). — Pl. I, Fig. 23. | Transvaal. |
| 5. <i>P. mappigera</i> , Meyrick, ibidem, Vol. 4, p. 191 (1914). | Transvaal, Natal. |
| 6. <i>P. tholaea</i> , Meyrick, ibidem, Vol. 2, p. 231 (1911). | Transvaal. |
| 7. <i>P. infrita</i> , Meyrick, Exot. Microlep. Vol. 1, p. 579 (1916). | Nyassaland. |

74. GENUS SYMBATICA, MEYRICK

Symbatica, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 413 (1910). — Type : *S. cryphias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderately elongate, without pecten. Labial palpi long, curved, ascending, second joint with appressed scales, terminal joint shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 almost from angle, 7 and 8 stalked, 7 to costa, 11 from before middle. Hindwings 1, elongate-trapezoidal, termen sinuate, oblique, cilia almost 1; 4 and 5 out of 3, 6 and 7 long-stalked.

Remarks. — Probably allied to *Gelechia*, but with singular neuration of hindwings.

Geographical distribution of species. — South African.

Larva unknown.

- | | |
|---|--------------|
| 1. <i>S. cryphias</i> , Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 413 (1910). | Cape Colony. |
|---|--------------|

75. GENUS DISSOPTILA, MEYRICK

Dissoptila, Meyrick, Trans. Ent. Soc. Lond. p. 234 (1914). — Type : *D. mutabilis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, slender, second joint with scales roughened towards apex beneath, terminal joint much longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae densely clothed with rough hairs. Forewings with two large scale-tufts in disc anteriorly; 1b furcate, 2 from 2/3, 4 and 5 connate from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex rather produced, pointed, termen obliquely emarginate, cilia 2; 3 and 4 connate, 5 approximated, 6 and 7 rather approximated towards base, transverse vein inwardly oblique from 6 to 7.

Remarks. — A well-marked and distinct genus, of which the immediate affinity is not obvious.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|---|-----------------|
| 1. <i>D. asphaltitis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 234 (1914). | Brit. Guiana. |
| 2. <i>D. prosona</i> , Meyrick, ibidem, p. 235 (1914). | Brit. Guiana. |
| 3. <i>D. crocodora</i> , Meyrick, ibidem, p. 65 (1922). — Pl. I, Fig. 24. | Brazil, Peru. |
| 4. <i>D. mutabilis</i> , Meyrick, ibidem, p. 235 (1914). | Guiana, Brazil. |
| 5. <i>D. disrupta</i> , Meyrick, ibidem, p. 235 (1914). | Guiana, Brazil. |

76. GENUS AROGALEA, WALSINGHAM

Aroalea, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 48 (1910). — Type : *A. cristifasciella*, Chambers.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with rough scales beneath, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with tufts of scales on surface; 1b furcate, 2 remote from 3, 7 and 8 stalked; 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen faintly bisinuate, cilia 1 2/3; 3 and 4 approximated, 5 rather approximated, 6 and 7 somewhat approximated towards base.

Remarks. — Probably a derivative of *Telphusa*.

Geographical distribution of species. — North and South American.

Larva (*cristifasciella*) feeding between spun leaves.

Foodplant *Quercus*.

1. <i>A. soronella</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 17 (1914).	Panama, Guiana, Brazil.
<i>pentadora</i> , Meyrick, Trans Ent. Soc. Lond. p. 51 (1917).	
<i>melitoptila</i> , Meyrick, Exot. Microlep. Vol. 3, p. 17 (1923).	
2. <i>A. crocipunctella</i> , Walsingham, Proc. Zool. Soc. Lond. p. 520 (1891).	Windward Islands.
3. <i>A. archaea</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 49, pl. 2, f. 10 (1911).	Mexico.
4. <i>A. senecta</i> , Walsingham, ibidem, Vol. 4, p. 49 (1911).	Mexico.
5. <i>A. albilingua</i> , Walsingham, ibidem; Vol. 4, p. 50 (1911).	Mexico.
6. <i>A. cristifasciella</i> , Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 87 (1878). <i>inscripta</i> , Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 180 (1882).	E. United States.

77. GENUS ANTHISTARCHA, NOV. GEN.

Type : *A. geniatella*, Busck.

Characters. — Head smooth; tongue developed. Antennae 4/5, basal joint without pecten. Labial palpi very long, recurved, second joint beneath with triangular projecting tuft of scales, terminal joint as long as second, slender, acute. Maxillary palpi very short. Forewings with large tufts of scales in disc; 2 from towards angle, 7 and 8 stalked, 7 to costa. Hindwings trapezoidal, in ♂ with long costal hair-pencil from base; 3 and 4 short stalked, 6 and 7 long-stalked.

Remarks. — I have not seen this species, but am unable on the characters given by Busck to include it in *Gelechia*, as he does.

Geographical distribution of species. — Central American.

Larva unknown.

1. <i>A. geniatella</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 13 (1914).	Panama.
--	---------

78. GENUS SCHISTOPHILA, CHRETIEN

Schistophila, Chrétien, Bull. Soc. Ent. Fr. p. 112 (1899). — Type : *S. laurocistella*, Chrétien.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, basal joint moderately elongate, without pecten. Labial palpi moderately long, recurved, second joint thickened with dense scales rough beneath towards apex, terminal joint shorter than second, rather thick, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with small scaletufts; 1b furcate, 2 from towards angle, 2-5 parallel, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings nearly 1, trapezoidal, apex produced, pointed, termen emarginate, cilia 1 1/2; 3 and 4 remote, 5 nearly parallel, 6 and 7 long-stalked.

Remarks. — A development of *Telphusa*.

Geographical distribution of species. — South-West Europe.

Larva feeding in spun shoots and leaves.

Foodplant *Cistus*.

1. *S. laurocistella*, Chrétien, Bull. Soc. Ent. Fr. p. 112 (1899).

S. France, Spain.

79. GENUS LEURONOMA, MEYRICK

Leuronoma, Meyrick, Ann. Transv. Mus. Vol. 6, p. 16 (1918). Type : *L. chlorotoma*, Meyrick.

Characters. — Head smooth; ocelli moderate, posterior; tongue developed. Antennae 3/4-4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint more or less thickened with scales, slightly rough beneath, terminal joint as long as second or nearly, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long rough hairs beneath. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 8 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex more or less produced, termen sinuate or emarginate, cilia 1-2; 3-5 separate, approximated at base, 6 and 7 nearly approximated or stalked.

Remarks. — Probably related to *Telphusa*.

Geographical distribution of species. — South African.

Larva unknown.

- | | |
|---|------------|
| 1. <i>L. nigridorsis</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 68 (1921). | Rhodesia. |
| 2. <i>L. chlorotoma</i> , Meyrick, ibidem, Vol. 6, p. 16 (1918). | Transvaal. |
| 3. <i>L. veterascens</i> , Meyrick, ibidem, Vol. 6, p. 16 (1918). | Natal. |
| 4. <i>L. symotiss</i> , Meyrick, ibidem, Vol. 2, p. 2, pl. 4, f. 7 (1910). | Transvaal. |
| 5. <i>L. eremopis</i> , Meyrick, ibidem, Vol. 8, p. 67 (1921). | Transvaal. |

80. GENUS PITHANURGA, MEYRICK

Pithanurga, Meyrick, Ann. Transv. Mus. Vol. 8, p. 68 (1921). — Type : *P. chariphila*, Meyrick.

Characters. — Head smooth; ocelli small, posterior; tongue developed. Antennae 3/4 (?), basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with rough projecting scales beneath, terminal joint shorter than second, moderate, acute. Maxillary palpi

very short, filiform, appressed to tongue. Posterior tibiae loosely scaled. Forewings with 2 from 3/4, 3 and 4 closely approximated towards base, from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex pointed, termen sinuate; cilia 2; frenulum in ♀ strong, simple; 3 and 4 approximated towards base, 5 nearly parallel, 6 and 7 connate.

Remarks. — Apparently derived from *Telphusa*; characterised by the closely approximated veins 2 and 4 of forewings, and the unusual feature of the simple frenulum of ♀.

Geographical distribution of species. — South African.

Larva unknown.

1. *P. chariphila*, Meyrick, Ann. Transv. Mus. Vol. 8. p. 68 (1921). Transvaal.

81. GENUS TELPHUSA, CHAMBERS

Telphusa, Chambers, Canad. Ent. Vol. 4, p. 132 (1872). — Type : *T. longifasciella*, Clemens.

Teleia, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 272 (1870) (praeocc.). — Type : *T. vulgella*, Hübner.

Adrasteia, Chambers, Canad. Ent. Vol. 4, p. 149 (1872). — Type : *T. alexandriella*, Chambers.

Xenolechia, Meyrick, Handb. Brit. Lep. p. 583 (1895). — Type : *T. aethiops*, Westwood.

Geniadophora, Walsingham, Proc. Zool. Soc. Lond. p. 71 (1897). — Type : *T. extranea*, Walsingham.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennæ 4/5, in ♂ simple or minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with rough projecting often brush-like scales beneath, terminal joint as long as second or nearly, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with raised scales or tufts on surface; 1 b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 6 sometimes out of 7 near base, 11 from middle. Hindwings 1 or nearly, trapezoidal, apex pointed, sometimes produced, termen more or less sinuate, cilia over 1; 3-5 separate, more or less approximated, 6 and 7 connate or stalked.

Remarks. — A derivative of *Gelechia*, from which it is separated by the rough scales of forewings and separation of veins 3 and 4 of hindwings; these characters are variable in development, but both are present in all species known to me, though they have frequently been overlooked.

Geographical distribution of species. — Well developed in Europe, Africa, and North America, probably also in Northern Asia, and extending into India and South America, but not known from the Malayan region, Australia, or New Zealand.

Larva (26 known) feeding between spun leaves, rarely mining in leaves or in stem-galls.

Foodplants (almost always trees or shrubs) : *Cupuliferae* (9), *Betulaceae* (4), *Rosaceae* (2), *Cistaceae* (2), and 9 other Orders.

- | | |
|---|---|
| 1. <i>T. scabra</i> , Staudinger, Hor. Soc. Ent. Ross. Vol. 7, p. 254 (1870). | Greece. |
| 2. <i>T. aethiops</i> , Westwood, Brit. Moths, Vol. 2, p. 192, pl. 107, f. 13 (1851).
<i>quinquecristatella</i> , Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 88 (1878). | C. Europe, N. E. United States. |
| 3. <i>T. phaulosema</i> , Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 70 (1920). | Kenya Colony. |
| 4. <i>T. confixa</i> , Meyrick, Ann. Trans. Mus. Vol. 6, p. 51 (1918). | Transvaal. |
| 5. <i>T. Erschoffi</i> , Frey, Lep. Schweiz. p. 361 (1880). | Switzerland. |
| 6. <i>T. Killiasi</i> , Frey, ibidem, p. 362 (1880). | Switzerland. |
| 7. <i>T. humeralis</i> , Zeller, Isis, p. 200 (1839).
<i>lyellella</i> , Westwood, Brit. Moths, Vol. 2, p. 190, pl. 106, f. 10 (1851).
<i>?decorella</i> , Haworth, Trans. Ent. Soc. Lond. p. 338 (1812). | C. & S. Europe, Asia
Minor, Algeria. |

8. *T. commaculata*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 69 (1921). Portuguese E. Africa.
 9. *T. ochrifoliata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 56, Mexico.
 pl. 2, f. 15 (1911).
 10. *T. perspicua*, Walsingham, Proc. Zool. Soc. Lond. p. 72 (1897). Hayti.
 11. *T. translucida*, Walsingham, ibidem, p. 520 (1891). Mexico, Guiana, Antilles.
 12. *T. callitechna*, Meyrick, Trans. Ent. Soc. Lond. p. 236 (1914). — Pl. I, Guiana, Brazil.
Fig. 25.
13. *T. agrifolia*, Braun, Ent. News, Philad. Vol. 32, p. 9 (1921). California, Brit.
 14. *T. melanozona*, Meyrick, Exot. Microlep. Vol. 1, p. 65 (1913). Bengal. [Columbia.
 15. *T. syndelta*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 69 (1921). Rhodesia.
 16. *T. euryzeucta*, Meyrick, Exot. Microlep. Vol. 2, p. 501 (1922). China.
 17. *T. destillans*, Meyrick, ibidem, Vol. 2, p. 133 (1918). Assam.
 18. *T. caelata*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 287 (1913). Transvaal.
 19. *T. emphanista*, Meyrick, ibidem, Vol. 8, p. 70 (1921). Natal.
 20. *T. delatrix*, Meyrick, Exot. Microlep. Vol. 3, p. 17 (1923). Peru.
 21. *T. scalella*, Scopoli, Ent. Carn. no 654 (1763). Europe, Asia Minor.
 aleella, Fabricius, Ent. Syst. Vol. 3 (2), p. 317 (1794).
 alternella, Hübner, Samml. Eur. Schmett. Tin. f. 151 (1796).
 bicolorella, Treitschke, Schmett. Eur. Vol. 9 (1), p. 233 (1832).
 ? incognitella, Caradja, Iris, Vol. 34, p. 100 (1920).
 22. *T. basifasciella*, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 269, pl. 3, New Jersey, Texas.
 f. 22 (1873).
 23. *T. velatella*, Busck, Proc. Ent. Soc. Wash. p. 90 (1907). Arizona.
 24. *T. glandiferella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 275, pl. 4, Texas, Grenada.
 f. 25 (1873).
 sella, Chambers, Canad. Ent. Vol. 6, p. 238 (1874).
 25. *T. betulella*, Busck, Proc. U. S. Mus. Vol. 25, p. 787 (1903). Dist. Columbia, Virginia.
 26. *T. belangerella*, Chambers, Canad. Ent. Vol. 7, p. 210 (1875). E. United States.
 oronella, Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 179 (1882).
 27. *T. palliderosacella*, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 90 (1878). E. United States.
 28. *T. quinquedentata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, Mexico.
 p. 63, pl. 2, f. 19 (1911).
 29. *T. disclusa*, Meyrick, Exot. Microlep. Vol. 3, p. 16 (1923). Palestine.
 30. *T. paraula*, Meyrick, ibidem, Vol. 1, p. 568 (1916). S. India, Ceylon.
 31. *T. semiusta*, Meyrick, ibidem, Vol. 2, p. 500 (1922). China.
 32. *T. epomidella*, Tengström, Act. Soc. Faun. Flor. Fenn. Vol. 10, p. 365 Finland.
 (1869).
 33. *T. femoralis*, Staudinger, Stett. Ent. Zeit. Vol. 37, p. 146 (1876). Sicily.
 34. *T. myricariella*, Frey, Mitt. Schweiz. Ent. Ges. Vol. 3, p. 251 (1870). C. Europe, Algeria.
 35. *T. wagae*, Nowicki, Enum. Lep. Hal. p. 189 (1860). C. & S. W. Europe.
 36. *T. caecigena*, Meyrick, Exot. Microlep. Vol. 2, p. 134 (1918). N. W. India.
 37. *T. notatella*, Hübner, Samml. Eur. Schmett. Tin. f. 344 (1817). N. & C. Europe.
 auratella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 169, f. 493 (1855).
 38. *T. saltuum*, Zeller, Stett. Ent. Zeit. Vol. 39, p. 140 (1878). Switzerland, Germany,
 39. *T. proximella*, Hübner, Samml. Eur. Schmett. Tin. f. 228 (1796). Europe. [Tyrol.]
 ? mouffetella, Linnaeus, Faun. Suec. p. 1397 (1761).
 40. *T. peritella*, Constant, Ann. Soc. Ent. Fr. p. 254, pl. 10, f. 15 (1884). Corsica.
 41. *T. canariensis*, Walsingham, Proc. Zool. Soc. Lond. p. 936, pl. 51, f. 15 Canaries.
 (1907).
 42. *T. decuriella*, Mann, Verh. Zool.-bot. Ges. Wien, Vol. 22, p. 38 (1872). Carinthia.
 43. *T. anguinella*, Herrich-Schäffer, Neue Schmett. p. 31, f. 159 (1861). Hungary, S. Russia.
 44. *T. tigrina*, Christoph, Hor. Soc. Ent. Ross. Vol. 12, p. 296, pl. 6, f. 68 Turkestan.
 (1877).
 45. *T. partitella*, Christoph, Stett. Ent. Zeit. Vol. 48, p. 167 (1887). Turkestan.

46. *T. albunella*, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, pl. 298, f. 2 (1838). C. Europe, Mongolia.
radiella, Krulikowsky, Rev. Russe Ent. Vol. 3, p. 181 (1903).
47. *T. mersinella*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 313 (1880). Asia Minor.
48. *T. praedicata*, Meyrick, Exot. Microlep. Vol. 3, p. 19 (1923).
melanostictella, Ragonot, Bull. Soc. Ent. Fr. p. 106 (1895) (praeocc.). Asia Minor.
49. *T. fugitivella*, Zeller, Isis, p. 200 (1839). Europe. [W. Russia.
50. *T. fugacella*, Zeller, Isis, p. 200 (1839). Germany, Galicia,
51. *T. squamulella*, Peyerimhoff, Mitth. Schweiz. Ent. Ges. Vol. 3, p. 412 (1871). S. France.
52. *T. vulgella*, Hübner, Samml. Eur. Schmett. Tin. f. 346 (1817). Europe.
- aspera*, Haworth, Lep. Brit. p. 550 (1828)
53. *T. trifasciella*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 66, p. (44) (1916). Bulgaria.
54. *T. triparella*, Zeller, Isis, p. 200 (1839). Europe, Asia Minor.
- dodecella*, Wood, Ind. Ent. f. 1200 (1839).
? paripunctella, Thunberg, Diss. Ent. Vol. 7, p. 96 (1794).
55. *T. inscriptella*, Christoph, Bull. Soc. Nat. Mosc. p. 25 (1882). E. Siberia.
56. *T. scriptella*, Hübner, Samml. Eur. Schmett. Tin. f. 152 (1796). Europe, Asia Minor.
- tremella*, Wood, Ind. Ent. f. 1223 (1839).
57. *T. prasinoleuca*, Meyrick, Zool. Med. Leid. Vol. 6, p. 161 (1921). Java.
58. *T. thomeriella*, Chrétien, Bull. Soc. Ent. Fr. p. 11 (1901). France.
59. *T. violenta*, Meyrick, Exot. Microlep. Vol. 2, p. 235 (1919). Bombay.
60. *T. eodryas*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 20 (1918). Rhodesia, Transvaal,
61. *T. luculella*, Hübner, Samml. Eur. Schmett. Tin. f. 397 (1817). Europe. [Natal.
luctuella, Wood, Ind. Ent. f. 1215 (1839).
subrosea, Wood, ibidem, f. 1216 (1839).
62. *T. semicostella*, Staudinger, Berl. Ent. Zeitschr. p. 311 (1870). S. Russia.
63. *T. craterota*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 287 (1913). Transvaal.
64. *T. cycota*, Meyrick, ibidem, Vol. 3, p. 65 (1912). Transvaal.
65. *T. resecta*, Meyrick, ibidem, Vol. 3, p. 288 (1913). Transvaal.
66. *T. revoluta*, Meyrick, ibidem, Vol. 6, p. 17 (1918). Zululand.
67. *T. objecta*, Meyrick, ibidem, Vol. 8, p. 70 (1921). Rhodesia.
68. *T. amphichroma*, Meyrick, ibidem, Vol. 3, p. 286 (1913). Transvaal.
69. *T. microsperma*, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 69 (1920). Kenya Colony.
70. *T. fecunda*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 17 (1918). Natal.
71. *T. sematica*, Meyrick, ibidem, Vol. 3, p. 286 (1913). Transvaal.
72. *T. retecta*, Meyrick, ibidem, Vol. 8, p. 70 (1921). Natal.
73. *T. lathridia*, Meyrick, ibidem, Vol. 2, p. 11, pl. 4, f. 5, 6 (1909). Transvaal.
74. *T. accensa*, Meyrick, ibidem, Vol. 8, p. 68 (1921). Rhodesia.
75. *T. calathaea*, Meyrick, ibidem, Vol. 3, p. 286 (1913). Transvaal.
76. *T. castrigera*, Meyrick, ibidem, Vol. 3, p. 287 (1913). Transvaal. [rate.
77. *T. iriditis*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 282 (1920). S. W. African Protectorate.
78. *T. basistrigella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 270, pl. 4, f. 23 (1873). Texas.
79. *T. caryivorella*, Packard, Rep. U. S. Dep. Agric. 1885, p. 331 (1886) Rhode Island.
(-yaevorella).
80. *T. quercinigrella*, Chambers, Canad. Ent. Vol. 4, p. 170 (1872) (-gracella). E. United States.
fragmentella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 271 (1873).
81. *T. querciella*, Chambers, Canad. Ent. Vol. 4, p. 127 (1872). E. United States, Canada.
82. *T. fasciella*, Chambers, ibidem, Vol. 4, p. 149 (1872). Kentucky.
83. *T. lophella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 182 (1909). California.
84. *T. nigrimaculata*, Braun, Proc. Calif. Acad. Sc. (4) Vol. 12, p. 118 (1923). California.
85. *T. bicristatella*, Chambers, Canad. Ent. Vol. 7, p. 210 (1875). Kentucky.
86. *T. sedulitella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 181 (1909). California.
87. *T. schizogyna*, Walsingham, Proc. Zool. Soc. Lond. p. 936, pl. 51, f. 12, (1907). Canaries.

88. *T. obligata*, Busck, Proc. U. S. Mus. Vol. 47, p. 15 (1914).
 89. *T. medullella*, Busck, ibidem, Vol. 47, p. 15 (1914) (-*dulella*).
 90. *T. melicentra*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 71 (1921).
 91. *T. orgilopis*, Meyrick, Exot. Microlep. Vol. 3, p. 16 (1923).
 92. *T. ochrotona*, Meyrick, ibidem, Vol. 3, p. 17 (1923).
 93. *T. cisti*, Stainton, Tin. S. Eur. p. 211 (1869).
 94. *T. sequax*, Haworth, Lep. Brit. p. 552 (1828).
 apicistrigella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 176, f. 483 (1855).
 95. *T. comedonella*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 312 (1880).
 96. *T. baldiana*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 225 (1920).
 97. *T. extranea*, Walsingham, Proc. Zool. Soc. Lond. p. 521 (1891).
 98. *T. fuscopunctella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 12 (1863).
 99. *T. praefixa*, Braun, Proc. Acad. Nat. Sc. Philad. p. 6 (1921).
 100. *T. acariella*, Busck, Proc. U. S. Mus. Vol. 30, p. 722 (1906).
 101. *T. longifasciella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 12 (1863).
 curvistrigella, Chambers, Canad. Ent. Vol. 4, p. 133 (1872).
 obliquifasciella, Chambers, Journ. Cinc. Soc. Nat. Hist. Vol. 2, p. 182 (1880).
 102. *T. latifasciella*, Chambers, Cinc. Quart. Journ. Sc. p. 251 (1875).
 103. *T. limenaea*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 282 (1920).
 104. *T. melanoleuca*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 56, (1911).
 105. *T. alexandriella*, Chambers, Canad. Ent. Vol. 4, p. 149 (1872) (-*driacella*).
 106. *T. bicostimaculella*, Chambers, ibidem, Vol. 4, p. 127 (1872) (-*tomaculella*)
 quercifoliella, Chambers, ibidem, Vol. 4, p. 206 (1872).
 107. *T. probata*, Meyrick, Ann. Transv. Mus. Vol. 2, p. 11, pl. 4, fig. 4 (1909).
 108. *T. inferialis*, Meyrick, Exot. Microlep. Vol. 2, p. 133 (1918).
 109. *T. mariona*, Heinrich, Journ. Agric. Res. Wash. Vol. 20, p. 812 (1921).
 110. *T. ripula*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 57, pl. 2, f. 16 (1911).
- Panama.
 Panama.
 Portuguese E. Africa.
 Brazil.
 Brazil. [naries.
 S. Europe, Algeria, Ca-
 N. & C. Europe, Massa-
 chusetts.
 Asia Minor.
 California.
 Windward Is.
 Pennsylvania, New
 Jersey, Canada.
 Montana.
 Texas, Louisiana.
 E. United States.
- Kentucky, Missouri.
 Cape Colony. [Canada.
 Mexico.
 Kentucky.
 E. United States.
 Tiansvaal.
 Bengal.
 Texas.
 Guatemala.

82. GENUS SYNCOPACMA, NOV. GEN.

Type: *S. acrophylla*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with rather rough scales beneath, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 6 and 7 out of 8, 7 to costa, 9 out of 8 towards base, 11 from middle. Hindwings 1, trapezoidal, termen sinuate; 3-5 approximated towards base, 6 and 7 connate.

Remarks. — Perhaps correlated with the preceding.

Geographical distribution of species. — South African.

Larva unknown.

1. *S. acrophylla*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 65 (1912). Transvaal.

83. GENUS MACRENCHES, MEYRICK

Macrenches, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 306 (1904). — Type: *M. eurybatis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ fasciculate-ciliated, basal joint moderately elongate, without pecten. Labial palpi long, curved, ascending, second joint thickened with dense scales, roughly projecting above towards apex, beneath rough or angularly projecting at apex, terminal joint somewhat shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, termen sinuate, cilia 1 1/2; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated at base.

Remarks. — A development of *Gelechia*.

Geographical distribution of species. — Australian.

Larva unknown.

1. *M. eurybatis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 307 (1904). W. Australia. [nia.
2. *M. clerica*, Rosenstock, Ann. Mag. Nat. Hist. (5), Vol. 16, p. 438 (1885). S. E. Australia, Tasmania.

84. GENUS APOTHETOeca, MEYRICK

Apothetoeeca, Meyrick, Nat. Hist. Juan Fern. Vol. 3, p. 268 (1922). — Type: *A. synaphrista*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrate, ciliated near base, thickened with rough scales beneath from near base to near middle, basal joint elongate, without pecten. (Labial palpi missing in type, presumed as in *Gelechia*.) Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with 1 b furcate, 2 from near angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen faintly sinuate, cilia 4/5; 3 and 4 connate, 5 somewhat approximated, 6 and 7 short-stalked.

Remarks. — Obviously nearly allied to *Gelechia*, from which it differs by the rough scale-thickening of antennae.

Geographical distribution of species. — Juan Fernandez island, associated with the Andean region of South America.

Larva unknown.

1. *A. synaphrista*, Meyrick, Nat. Hist. Juan Fern. Vol. 3, p. 269 (1922). Juan Fernandez.

85. GENUS GELECHIA, HÜBNER

Gelechia, Hübner, Verz. bek. Schmett. p. 415 (1826). — Type: *G. rhombella*, Schiffermüller.

Chionodes, Hübner, ibidem, p. 420 (1826). — Type: *G. lugubrella*, Fabricius.

Lita, Treitschke, Schmett. Eur. Vol. 9 (2), p. 82 (1833). — Type: *G. longicornis*, Curtis.

Ficulea, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 794 (1864). — Type: *G. blandulella*, Walker.

Bryotropha, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 233 (1870). — Type: *G. terrella*, Schiffermüller.

Cirrha, Chambers, Canad. Ent. Vol. 4, p. 146 (1872). — Type: *G. albisparsella*, Chambers.

Pseudochelaria, Dietz, Ent. News, Philad. Vol. 11, p. 252 (1900). — Type: *G. Walsinghami*, Dietz.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 2/3-4/5, in ♂ simple or minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, rough and more or less furrowed beneath, terminal joint as long as second or somewhat shorter or longer, rather slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings rarely with rough scales; 1b furcate, 2 from 3/4 or towards angle, rarely 3 and 4 short-stalked, 7 and 8 stalked, 7 to costa, 6 sometimes out of 7 towards base, 11 from middle. Hindwings 1 or over 1, trapezoidal, apex pointed or obtuse, termen more or less sinuate, cilia 2/3-1 1/2; 3 and 4 connate or seldom short-stalked, 5 somewhat approximated, 6 and 7 closely approximated at base or connate or stalked.

Remarks. — Correlated with *Phthorimaea* as a parallel line of development. Such variation of structure as occurs is gradual and indefinite, and does not allow of further natural subdivision. The greater part of the species are obscurely and similarly coloured, and their large number renders their study difficult; but there is also a considerable proportion with conspicuous black and white colouring. Many are reared in habit, and only obtained readily by rearing the larvae.

Geographical distribution of species. — Summarised approximately (several species occur in both Europe and North America, and it is uncertain with which they should be classed) as follows : American 170, Palaearctic 146, South African 41, India 3, China 1, Australia 11, New Zealand 9. This interesting distribution indicates an origin and extensive development in North America and the Palaearctic region at a time (not very remote) when these two zoological areas had free interconnection by the north, and whilst the Indian peninsula was an island; thence the genus, constitutionally habituated to temperate climates, spread to South America and South Africa mainly by the high lands; it is apparently ill adapted to face the climate of India. The New Zealand species are interesting, since they form a single connected group unquestionably allied to the North American species of the *variabilis* group, and must (like most of the larger genera of New Zealand Lepidoptera) have travelled thither by way of the Andes and the Antarctic lands. The Australian species (which have certainly no connection with the New Zealand group) will probably be found eventually to have come from Eastern Asia by way of the Philippines and New Guinea.

Larva (known for 76 species) feeding usually in spun leaves or shoots, very rarely mining in leaves or fruits.

Foodplants : Leguminosae (14), Salicaceae (10), Polygonaceae (7), Rosaceae (6), Tamaricaceae (5), Ericaceae (4), Cupuliferae (4), Anacardiaceae (3), Compositae (3), Musci (3) and 14 other orders (usually low plants or shrubs).

- | | |
|--|---|
| 1. <i>G. basaltinella</i> , Zeller, Isis, p. 198 (1839). | Europe. |
| 2. <i>G. dryadella</i> , Zeller, Stett. Ent. Zeit. Vol. 11, p. 152 (1850).
<i>phoebeusella</i> , Millière, Cat. Lép. Alp. Mar. p. 328 (1875). | S. Europe, Asia Minor. |
| 3. <i>G. domestica</i> , Haworth, Lep. Brit. p. 551 (1828).
<i>salmonis</i> , Walsingham, Proc. Zool. Soc. Lond. p. 937 (1907). | C. & S. Europe, Madeira,
N. Africa, Palestine. |
| 4. <i>G. umbrosella</i> , Zeller, Isis, p. 201 (1839). | C. Europe. |
| 5. <i>G. affinis</i> , Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 17 (1850).
<i>tegulella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 182 (1855). | C. Europe, Syria. |
| 6. <i>G. thuleella</i> , Staudinger, Stett. Ent. Zeit. Vol. 18, p. 276 (1857). | Iceland. |
| 7. <i>O. nigricella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 316 (1915). | Algeria. |
| 8. <i>G. fuliginosella</i> , Snellen, Vlind. Nederl. Vol. 2, p. 645 (1882). | Holland. |
| 9. <i>G. similis</i> , Stainton, Ins. Brit. Tin. p. 115 (1854).
<i>obscurecinerea</i> , Nolcken, Lep. Faun. Livl. Vol. 2, p. 573 (1870).
<i>confinis</i> , Stainton, Ent. Annual, p. 98 (1871).
<i>stolidella</i> , Gregson, Morris, Brit. Moths, pl. 108, f. 1 (1872). | England, Holland,
Latvia. |

10. *G. tectella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 182 (1855). Switzerland.
 11. *G. mundella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 64 (1850). England, Germany.
 portlandicella, Richardson, Ent. M. Mag. Vol. 26, p. 29 (1890).
 12. *G. Peterseni*, Teich, Korr.-Blatt Naturf. Ver. Riga, Vol. 44, p. 13 (1901). Latvia.
 13. *G. cinerosella*, Tengström, Not. Sällsk. Faun. Fenn. Förh. p. 129 (1847). N. Germany, Finland.
 ? *serratulella*, Tengström, ibidem, p. 128 (1847).
 14. *G. plantariella*, Tengström, ibidem, p. 128 (1847). Finland.
 ? *flavipalpella*, Tengström, ibidem, p. 127 (1847).
 15. *G. glebicolorella*, Erschoff, Lep. Turk. Fedtsch. p. 101, pl. 6, f. 114 (1874). Turkestan.
 16. *G. senectella*, Zeller, Isis, p. 199 (1839). N. & C. Europe.
 obscurella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 239 (1870).
 ? *glabrella*, Heinemann, ibidem, p. 239 (1870).
 ? *minorella*, Heinemann, ibidem, p. 240 (1870).
 17. *G. desertella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 62 (1850). N. C. Europe, Sicily.
 18. *G. brevipalpella*, Rebel, Stett. Ent. Zeit. Vol. 54, p. 47 (1893). Latvia. [Palestine].
 19. *G. plebeiella*, Zeller, Isis, p. 850 (1847). S. Europe, N. Africa,
 20. *G. imperitella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 242 (1859). S. France, Spain.
 21. *G. indignella*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 308 (1880). Asia Minor.
 22. *G. ciliatella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 174, f. 590 (1855). Germany.
 23. *G. politella*, Stainton, Cat. Brit. Tin. Suppl. p. 4 (1851). England.
 24. *G. figulella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 242 (1859). S. Europe, England.
 capnella, Constant, Ann. Soc. Ent. Fr. p. 196, pl. 7, f. 13 (1865).
 25. *G. decrepidella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 177, f. 508, 533 (1855). N. & C. Eur.
 lutescens, Constant, Ann. Soc. Ent. Fr. p. 196, pl. 7, f. 12 (1865).
 26. *G. terrella*, Hübner, Samml. Eur. Schmett. Tin. f. 170 (1796). Europe, Asia Minor.
 inulella, Hübner, ibidem, f. 286 (1805).
 zephyrella, Treitschke, Schmett. Eur. Vol. 9 (1), p. 241 (1832).
 ? *latella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 174, f. 513 (1855).
 ? *alpicolella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 235 (1870).
 27. *G. branella*, Busck, Proc. Ent. Soc. Wash. Vol. 9, p. 87 (1908). Maryland.
 28. *G. neptica*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 65 (1911). Mexico.
 29. *G. bufo*, Walsingham, ibidem, Vol. 4, p. 66 (1911). Mexico.
 30. *G. metallica*, Braun, Proc. Acad. Nat. Sc. Philad. p. 9 (1921). Montana.
 31. *G. clandestina*, Meyrick, Exot. Microlep. Vol. 3, p. 19 (1923). Ontario.
 32. *G. synthetica*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 67 (1911). Mexico.
 33. *G. litigiosa*, Meyrick, Trans. Ent. Soc. Lond. p. 49 (1917). Ecuador.
 34. *G. ceanothiella*, Busck, Proc. U. S. Mus. Vol. 27, p. 760 (1904). Brit. Columbia.
 35. *G. paralogella*, Busck, Proc. Ent. Soc. Wash. Vol. 18, p. 149 (1916). California.
 36. *G. trialbimaculella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 250 (1875) (-bam-). E. United States.
 epigaeella, Chambers, Journ. Cinc. Soc. Nat. Hist. Vol. 3, p. 289 (1881).
 37. *G. minimaculella*, Chambers, Canad. Ent. Vol. 6, p. 235 (1874). Texas.
 38. *G. argentipunctella*, Ely, Proc. Ent. Soc. Wash. Vol. 12, p. 70 (1910). Connecticut.
 39. *G. thoracalbellula*, Chambers, Canad. Ent. Vol. 6, p. 235 (1874) (-cea-). Texas.
 40. *G. argosema*, Meyrick, Trans. Ent. Soc. Lond. p. 50 (1917). Ecuador.
 41. *G. leucaniella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 180 (1909) (-ieella). California.
 42. *G. paraplatulella*, Busck, ibidem, Vol. 11, p. 181 (1909). California.
 43. *G. lutraula*, Meyrick, Exot. Microlep. Vol. 3, p. 20 (1923). Ontario.
 44. *G. consona*, Meyrick, Trans. Ent. Soc. Lond. p. 50 (1917). Peru.
 45. *G. infracta*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 61 (1911). Mexico.
 46. *G. magnetella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 310 (1870). Asia Minor.
 47. *G. platydoxa*, Meyrick, Exot. Microlep. Vol. 3, p. 20 (1923). Fr. Guiana.

48. *G. traducella*, Busck, Proc. U. S. Mus. Vol. 47, p. 12 (1914). Panama, Brazil.
49. *G. compositella*, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1030 (1864). ? C. America.
50. *G. suspensa*, Meyrick, Exot. Microlep. Vol. 3, p. 19 (1923). Brazil.
51. *G. lacticeps*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 61 (1911) (-caput). Mexico.
52. *G. cerussata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 61, pl. 2, f. 18 (1911). Mexico.
53. *G. blandulella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 795 (1864). S. India, Ceylon.
— Pl. 2, Fig. 29.
54. *G. horiaula*, Meyrick, Exot. Microlep. Vol. 2, p. 133 (1918). N. W. India.
55. *G. tetralenca*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 18 (1918). Zululand.
56. *G. psimythota*, Meyrick, ibidem, Vol. 3, p. 293 (1913). Transvaal.
57. *G. triscelis*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 3 (1917). Natal.
58. *G. devia*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 290 (1913). Transvaal.
59. *G. ophitis*, Meyrick, ibidem, Vol. 3, p. 291 (1913). Transvaal.
60. *G. rescissella*, Zeller, Lep. Micr. Caffr. p. 110 (1852). Natal.
61. *G. trisignis*, Meyrick, Proc. Zool. Soc. Lond. p. 725 (1908). — Pl. 2, Fig. 28. Transvaal, Zululand, Cape Colony.
62. *G. isochorda*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 72 (1921). Transvaal.
63. *G. stasiarcha*, Meyrick, ibidem, Vol. 3, p. 290 (1913). Transvaal.
64. *G. prioleuca*, Meyrick, ibidem, Vol. 3, p. 66 (1912). Transvaal.
65. *G. naufragia*, Meyrick, ibidem, Vol. 3, p. 66 (1912). Transvaal.
66. *G. Dolbyi*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 98, pl. 3, f. 22 (1911). Panama.
67. *G. cercerella*, Chambers, Canad. Ent. Vol. 4, p. 108 (1872) (*cercerisella*). — Pl. 2, Fig. 26.
olympiadella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 259, pl. 3, f. 15 (1873).
68. *G. quinella*, Zeller, ibidem, Vol. 23, p. 260, pl. 3, f. 14 (1873). Texas.
69. *G. arizonella*, Busck, Proc. U. S. Mus. Vol. 25, p. 856 (1903). Arizona.
70. *G. lipatiella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 88 (1909). Colorado.
71. *G. bimaculella*, Chambers, Canad. Ent. Vol. 4, p. 108 (1872). Kentucky, Texas.
ternariella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 264, pl. 3, fig. 19 (1873).
- sylvestrella*, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 86 (1878).
72. *G. coloradensis*, Busck, Proc. U. S. Mus. Vol. 25, p. 857 (1903). Colorado, S. Carolina,
73. *G. tessella*, Hübner, Samml. Vög. Schmett. f. 3 (1793). Europe. [Florida.
quadrella, Fabricius, Ent. Syst. Vol. 3 (2), p. 298 (1794).
scopolella, Hübner, Samml. Eur. Schmett. Tin. f. 145 (1796).
funestellia, Hübner, ibidem, f. 466 (1832).
74. *G. luctuella*, Hübner, Samml. Vög. Schmett. f. 5 (1793). N. & C. Europe.
sauteriella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 18, p. 612 (1868).
75. *G. trauniella*, Zeller, Stett. Ent. Zeit. Vol. 29, p. 147 (1868). Carinthia. [dor.
76. *G. viduella*, Fabricius, Ent. Syst. Vol. 3 (2), p. 299 (1794). N. & C. Europe, Labra-
leucomella, Quensel, Acerbi Trav. Swed. Vol. 2, p. 254, pl. 3, f. 2 (1802).
luctuella, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, pl. 298, f. 9 (1838).
luctiferella, Herrich-Schäffer, Neue Schmett. p. 6, f. 42 (1856).
labradoriella, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 12 (1863).
77. *G. prorepta*, Meyrick, Exot. Microlep. Vol. 3, p. 19 (1923). California.
- fulminella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 178 (1909) (praeocc.).
78. *G. albilocella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 261, pl. 3, f. 16 (1873). Colorado, Arizona,
Texas.
- trifasciella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 252 (1875).
79. *G. feralella*, Zeller, Stett. Ent. Zeit. Vol. 33, p. 110 (1872). Switzerland, Austria.
80. *G. lugubrella*, Fabricius, Ent. Syst. Vol. 3 (2), p. 299 (1794). N. & C. Europe, Kam-
luctifcella, Hübner, Samml. Eur. Schmett. Tin. f. 312 (1817). schatka, Maine.
lunatella, Zetterstedt, Ins. Lapp. p. 1005 (1840).

81. *G. dentella*, Busck, Proc. U. S. Mus. Vol. 25, p. 862 (1903). . Arizona.
 82. *G. sistrella*, Busck, ibidem, p. 862 (1903). Arizona, California.
 83. *G. abdominalis*, Busck, ibidem, p. 863 (1903). Arizona.
 84. *G. xanthophilella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 228 (1920). California.
 85. *G. coticola*, Busck, Journ. Ent. Zool. Claremont, Vol. 5, p. 97 (1913). California.
 86. *G. depuratella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 177 (1909). New Mexico.
 87. *G. dromicella*, Busck, ibidem, Vol. 11, p. 177 (1909). Colorado, California.
 88. *G. triangulella*, Busck, Proc. Ent. Soc. Wash. Vol. 8, p. 91 (1907). Arizona.
 89. *G. packardella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 143 (1877). Colorado.
 90. *G. unifasciella*, Busck, Proc. U. S. Mus. Vol. 25, p. 865 (1903). Arizona.
 91. *G. paulella*, Busck, ibidem, Vol. 25, p. 865 (1903). Colorado, Arizona.
 92. *G. kincaidella*, Busck, Proc. Ent. Soc. Wash. Vol. 8, p. 91 (1907). Wyoming.
 93. *G. catalinella*, Busck, Journ. N. York Ent. Soc. Vol. 15, p. 136 (1907). Arizona.
 94. *G. panella*, Busck, Proc. U. S. Mus. Vol. 25, p. 889 (1903). Arizona, California.
 95. *G. exclarella*, Möschler, Abhandl. Senckenb. Naturf. Ges. Vol. 15, p. 343 (1900). Portorico.
96. *G. aristella*, Busck, Proc. U. S. Mus. Vol. 25, p. 866 (1903). Arizona.
 97. *G. morenella*, Busck, Ent. News, Philad. Vol. 19, p. 317 (1908). California.
 98. *G. ribeella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 290 (1875) (*ribesella*). Colorado.
 99. *G. mandella*, Busck, Proc. U. S. Mus. Vol. 27, p. 759 (1904). Brit. Columbia.
 100. *G. catharodes*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 284 (1920). Cape Colony.
 101. *G. leucodoxa*, Meyrick, ibidem, Vol. 17, p. 283 (1920). Cape Colony.
 102. *G. lactiflora*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 71 (1921). Portuguese E. Africa.
 103. *G. albiflora*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 283 (1920). Cape Colony.
 104. *G. triplacopsis*, Neyrick, ibidem, Vol. 10, p. 61 (1912). Cape Colony.
 105. *G. nanodella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 179 (1909). California.
 106. *G. nigrorosea*, Walsingham, Ent. M. Mag. Vol. 40, p. 266 (1904). Algeria.
 107. *G. lunariella*, Walsingham, Proc. Zool. Soc. Lond. p. 939, pl. 51, f. 13 (1907). Canaries.
108. *G. tristis*, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 307 (1880). Asia Minor.
 109. *G. diffinis*, Haworth, Lep. Brit. p. 551 (1828). Europe.
- dissimilella*, Treitschke, Schmett. Eur. Vol. 9 (2), p. 80 (1833).
scabidella, Zeller, Isis, p. 199 (1839)
? bagriotella, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, p. 608, pl. 312, f. 2 (1838).
110. *G. terebinthinella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, f. 597 (1855). S.E. Europe, Asia Minor.
 111. *G. cinctipunctella*, Erschoff, Hor. Soc. Ent. Ross. Vol. 12, p. 344 (1876). E. Siberia.
 112. *G. pyrenaica*, Petry, Iris, Vol. 17, p. 3 (1904). Pyrenees.
 113. *G. Dzieduszycki*, Nowicki, Microlep. Sp. Nov. p. 20, pl. 1, f. 4 (1864). Alps, Hungary, Balkans.
- melaleucella*, Constant, Ann. Soc. Ent. Fr. p. 197, pl. 7, f. 14 (1865)
mariae, Frey, Mitth. Schweiz. Ent. Ges. Vol. 2, p. 302 (1868).
114. *G. perspersella*, Wocke, Stett. Ent. Zeit. Vol. 23, p. 236 (1862). Lapland, Norway,
 115. *G. elatella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 168, f. 499 (1855). Alps. [Latvia.
 116. *G. rosalbella*, Fologne, Ann. Soc. Ent. Belg. p. 167, pl. 2, f. 3 (1862). Belgium, Germany.
 117. *G. albifemorella*, Hofmann, Stett. Ent. Zeit. Vol. 28, p. 204 (1867). Alps.
 118. *G. melanypella*, Mann, Verh. Zool.-bot. Ges. Wien. Vol. 27, p. 498 (1877). Tyrol.
 119. *G. tarandella*, Wocke, Stett. Ent. Zeit. Vol. 25, p. 212 (1864). Norway.
 120. *G. interalbicella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 187 (1855). Alps, S. France, Corsica.
 121. *G. Klosi*, Rebel, Verh. Zool.-bot. Ges. Wien. Vol. 67, p. (30) (1917). Carinthia.
 122. *G. apocleella*, Walsingham, Ent. M. Mag. Vol. 36, p. 216 (1900). Corsica.
 123. *G. perpetuella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 180, f. 511 (1855). Alps, Hungary.
 124. *G. dyariella*, Busck, Proc. U. S. Mus. Vol. 25, p. 877 (1903). Colorado.
 125. *G. albisparsella*, Chambers, Canad. Ent. Vol. 4, p. 92 (1872). Kentucky.
- platanella*, Chambers, ibidem, Vol. 4, p. 146 (1872).
126. *G. periculella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 178 (1909). Oregon, California.
 127. *G. neotrophella*, Heinrich, Journ. Agric. Res. Wash. Vol. 20, p. 811 (1921). Texas.

128. *G. trophella*, Busck, Proc. U. S. Mus. Vol. 25, p. 860 (1903). Colorado.
129. *G. discontinuella*, Rebel, Verh. Zool.-bot. Ges. Wien. Vol. 49, p. 178 (1899). Tyrol.
130. *G. continuaella*, Zeller, Isis, p. 198 (1839).
- trimaculella*, Packard, Proc. Boston Soc. Nat. Hist. p. 61 (1867).
 - nebulosella*, Heinemann, Schmett. Deutsch. (2) Vol. 2, p. 218 (1870).
 - albomaculella*, Chambers, Canad. Ent. Vol. 7, p. 209 (1875).
131. *G. selectella*, Caradja, Iris, Vol. 34, p. 99 (1920). Uralsk.
132. *G. electella*, Zeller, Isis, p. 198 (1839). C. Europe.
133. *G. istrella*, Mann, Verh. Zool.-bot. Ges. Wien. Vol. 16, p. 354, pl. 1, f. 9 (1866). Roumania.
134. *G. impurgata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 67, pl. 2, fig. 23 (1911). Mexico.
135. *G. unistrigella*, Chambers, Canad. Ent. Vol. 5, p. 176 (1873). Kentucky.
136. *G. anthochra*, Lower, Trans. Roy. Soc. S. Australia, Vol. 20, p. 168 (1896). Queensland, New South
137. *G. hyoscyamella*, Rebel, Iris, Vol. 26, p. 89 (1912). Egypt. [Wales.]
138. *G. frankenivorella*, Chrétien, Ann. Soc. Ent. Fr. 1916, p. 474 (1917). Algeria. [Syria, India.]
139. *G. tamariciella*, Zeller, Stett. Ent. Zeit. Vol. 11, p. 153 (1850). S. Europe, N. Africa,
140. *F. heligmatodes*, Walsingham, Ent. M. Mag. Vol. 40, p. 267 (1904) Algeria, Sokotra.
(? = praec.)
141. *G. sonella*, Chrétien, Ann. Soc. Ent. Fr. 1916, p. 474 (1917). Algeria.
142. *G. flammulella*, Walsingham, Proc. Zool. Soc. Lond. p. 72 (1897). Virgin Islands.
143. *G. chloroschema*, Meyrick, Exot. Microlep. Vol. 3, p. 21 (1923). California.
144. *G. discostrigella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 248 (1875). California.
145. *G. elaboratella*, Braun, Proc. Calif. Acad. Sc. (4), Vol. 12, p. 119 (1923). California.
146. *G. lacticoma*, Meyrick, Trans. Ent. Soc. Lond. p. 48 (1917). Peru.
147. *G. trachycosma*, Meyrick, Exot. Microlep. Vol. 3, p. 21 (1923). California.
148. *G. maculatella*, Hübner, Samml. Eur. Schmett. Tin. f. 162 (1796). C. Europe.
149. *G. rhodoptera*, Mann, Verh. Zool.-bot. Ges. Wien. Vol. 16, p. 353, pl. 1, f. 10 (1866). Roumania, Greece.
150. *G. scissella*, Chrétien, Ann. Soc. Ent. Fr. p. 319 (1915). Algeria.
151. *G. cytisella*, Treitschke, Schmett. Eur. Vol. 9 (2), p. 99 (1833). C. & S. Europe.
152. *G. biforella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 89 (1909). Texas.
153. *G. pseudofondella*, Busck, ibidem, Vol. 9, p. 87 (1908). Pennsylvania.
154. *G. fondella*, Busck, Canad. Ent. Vol. 38, p. 122 (1906). Maryland, Pennsylvania,
155. *G. terminimaculella*, Kearfott, Journ. N. York Ent. Soc. Vol. 16, p. 184 Canada. [Ontario.] (1908).
156. *G. anisectis*, Meyrick, Exot. Microlep. Vol. 3, p. 19 (1923). California, Mexico.
- inaequalis*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 66 (1911) (praecocc.)
157. *G. clistrodoma*, Meyrick, Exot. Microlep. Vol. 3, p. 21 (1923). Arizona.
158. *G. inaequalis*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 180 (1909). Mexico, New Mexico.
159. *G. benitella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 229 (1920). Texas.
160. *G. malindella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 179 (1909). New Mexico.
161. *G. lindenella*, Busck, Proc. U. S. Mus. Vol. 25, p. 876 (1903). Texas, Colorado, Ari-
162. *G. crudescens*, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 70 (1920). Kenya Colony. [zona.]
163. *G. creberrima*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 67 (1911). Mexico.
164. *G. concinna*, Walsingham, ibidem, Vol. 4, p. 66, pl. 2, f. 22 (1911). Mexico.
165. *G. cuneifera*, Walsingham, ibidem, Vol. 4, p. 64 (1911). Mexico.
166. *G. mediofuscella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 11 (1863). Canada, E. United States,
ragella, Walker, List Lep. Brit. Mus. Vol. 29, p. 596 (1864). Mexico, Panama.
fuscoochrella, Chambers, Canad. Ent. Vol. 4, p. 106 (1872).
- lituroSELLA*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 265 (1873).
- rhedaria*, Meyrick, Exot. Microlep. Vol. 3, p. 20 (1923).
167. *G. repentina*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 65 Mexico.
(1911).

168. *G. Walsinghami*, Dietz, Ent. News, Philad. Vol. 11, p. 352, pl. 1, f. 3 (1900). Pennsylvania.
169. *G. pennsylvanica*, Dietz, ibidem, Vol. 11, p. 353, pl. 1, f. 4 (1900). Pennsylvania, Arizona.
170. *G. exoenota*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 52 (1918). Transvaal.
171. *G. matutina*, Meyrick, ibidem, Vol. 3, p. 290 (1913). Transvaal.
172. *G. peronectis*, Meyrick, ibidem, Vol. 2, p. 12, pl. 4, f. 9 (1910). Transvaal.
173. *G. sarcographa*, Meyrick, Ann. S Afr. Mus. Vol. 17, p. 3 (1917). Cape Colony.
174. *G. ochnias*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 291 (1913). — Pl. 2, Fig. 27. Transvaal.
175. *G. mesacta*, Meyrick, ibidem, Vol. 2, p. 12, pl. 4, f. 8 (1910). Transvaal.
176. *G. rhodochra*, Meyrick, ibidem, Vol. 3, p. 290 (1913). Transvaal.
177. *G. hutchinsonella*, Walsingham, Trans. Ent. Soc. Lond. p. 93, pl. 4, f. 30 (1891). E. & S. Africa.
178. *G. panula*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 370 (1909). Cape Colony.
179. *G. ferax*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 289 (1913). Transvaal.
180. *G. opaula*, Meyrick, ibidem, Vol. 3, p. 65 (1912). Transvaal.
181. *G. phoenaula*, Meyrick, ibidem, Vol. 3, p. 289 (1913). Transvaal.
182. *G. zetterstedtiella*, Zeller, Lep. Micr. Caffr. p. 111 (1852). Transvaal, Natal.
183. *G. erubescens*, Walsingham, Ent. M. Mag. Vol. 40, p. 265 (1904). Algeria.
184. *G. sinuatella*, Walsingham, ibidem, Vol. 40, p. 223 (1904). (? = seq.). Algeria.
185. *G. plutelliformis*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 239 (1859). S. Europe, N. Africa, *olbiaella*, Millière, Icon. Descr. Lép. Vol. 1, p. 193, pl. 21, f. 1-6 (1864). Canaries, Syria. *siewersiellus*, Christoph, Stett. Ent. Zeit. Vol. 28, p. 239 (1867).
186. *G. Sieversi*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 309 (1870). S. Russia.
187. *G. sesostrella*, Rebel, Iris, Vol. 26, p. 88 (1912). Egypt.
188. *G. caminariella*, Fuchs, Stett. Ent. Zeit. Vol. 63, p. 323 (1902). Germany.
189. *G. flavonigrella*, Chrétien, Ann. Soc. Ent. Fr. p. 318 (1915). Algeria.
190. *G. mulinella*, Zeller, Isis, p. 199 (1839). C. Europe, Spain.
interruptella, Stephens, Ill. Brit. Ent. Haust. Vol. 4, p. 214 (1835).
191. *G. interruptella*, Hübner, Samml. Vög. Schmett. f. 88 (1793). C. Europe, Spain.
192. *G. hetaeria*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 68, pl. 2, f. 24 (1911). Mexico.
193. *G. angulatella*, Chrétien, Ann. Soc. Ent. Fr. p. 318 (1915). Tunis.
194. *G. cerostomella*, Walsingham, Ent. M. Mag. Vol. 40, p. 266 (1904) (- *stomatella*). Algeria.
195. *G. longicornis*, Curtis, Brit. Ent. Vol. 4, p. 189 (1827). N. & C. Europe, Asia Minor, Canada, Colorado.
? virgella, Thunberg, Diss. Ent. Vol. 7, p. 92 (1794).
histriionella, Hübner, Samml. Eur. Schmett. Tin. f. 464 (1832).
zebrella, Treitschke, Schmett. Eur. Vol. 9 (2), p. 82 (1835).
alpicola, Frey, Mitt. Schweiz. Ent. Ges. Vol. 2, p. 299 (1870).
alternatella, Kearnott, Journ. N. York Ent. Soc. Vol. 16, p. 185 (1908).
196. *G. ericotella*, Hübner, Samml. Eur. Schmett. Tin. f. 470 (1832). Europe, Asia Minor, Algeria.
? betulea, Haworth, Lep. Brit. p. 549 (1828).
gallinella, Treitschke, Schmett. Eur. Vol. 9 (2), p. 79 (1835).
? simplicella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 213 (1870).
197. *G. caespitella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 335, pl. 4, f. 107 (1877). Colombia.
198. *G. infernalis*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 177, f. 584 (1855). N. & C. Europe.
? violacea, Tengström, Not. Sällsk. Faun. Fenn. Förh. Vol. 1, p. 125 (1847).
199. *G. tannuolella*, Rebel, Iris, Vol. 30, p. 193 (1917). Colombia.
200. *G. fuscantella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 213 (1870). N. & C. Europe.
201. *G. invenustella*, Berg, Bull. Soc. Nat. Mosc. Vol. 49 (2), p. 240 (1875). Caucasus, Asia Minor.
202. *G. rhypodes*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 69 (1911). Mexico.

203. *G. lapidescens*, Meyrick, Exot. Microlep. Vol. 1, p. 509 (1916). Mexico.
lithodes, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 69 (1911)
(praeocc.).
204. *G. nucifera*, Walsingham, ibidem, Vol. 4, p. 69 (1911). Mexico.
205. *G. rebeliella*, Hauder, Verh. Zool.-bot. Ges. Wien, Vol. 67, p. (29) (1917). Austria.
206. *G. pascuicola*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 301 (1870). Spain.
207. *G. eremella*, Chrétien, Ann. Soc. Ent. Fr. p. 318 (1915). Algeria.
208. *G. pergrandella*, Rebel, Iris, Vol. 30, p. 193 (1917). Mongolia.
209. *G. velocella*, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, pl. 297, f. 3 (1838). Europe, Asia Minor.
subsequilla, Treitschke, Schmett. Eur. Vol. 9 (2), p. 83 (1835).
nebulea, Stephens, Ill. Brit. Ent. Haust. Vol. 4, p. 206 (1835).
brunnea, Schöyen, Troms. Mus. Aars. Vol. 5, p. 53 (1882).
aterrimella, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 39, p. 312, pl. 8, f. 6 (1889).
210. *G. rumicivorella*, Millière, Ann. Soc. Sc. Nat. Cannes, p. 11, pl. 10, f. 13 (1880). S. France.
211. *G. petasitis*, Pfaffenzeller, Stett. Ent. Zeit. Vol. 28, p. 79 (1867). Alps, E. United States.
212. *G. lutilabrella*, Mann, Wien. Ent. Monatschr. Vol. 1, p. 179 (1857). S. France, Hungary, Dalmatia.
- monochromella*, Constant, Bull. Soc. Ent. Fr. p. 52 (1895).
213. *G. bergiella*, Teich, Stett. Ent. Zeit. Vol. 47, p. 170 (1886). Latvia.
214. *G. monophragma*, Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 169 (1886). New Zealand.
215. *G. dividua*, Philpot, ibidem, Vol. 53, p. 340 (1921). New Zealand.
216. *G. aerobatis*, Meyrick, ibidem, Vol. 55, p. 204 (1924). New Zealand.
217. *G. pharetria*, Meyrick, ibidem, Vol. 18, p. 169 (1886). New Zealand.
218. *G. parapleura*, Meyrick, ibidem, Vol. 18, p. 168 (1886). New Zealand.
219. *G. schematica*, Meyrick, ibidem, Vol. 18, p. 168 (1886). New Zealand.
220. *G. neglecta*, Philpot, ibidem, Vol. 55, p. 665 (1924). New Zealand.
221. *G. lithodes*, Meyrick, ibidem, Vol. 18, p. 170 (1886). New Zealand.
222. *G. lapillosa*, Meyrick, ibidem, Vol. 55, p. 203 (1924). New Zealand.
223. *G. sonorensis*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 69, pl. 2, f. 26 (1911). Mexico.
224. *G. abella*, Busck, Proc. U. S. Mus. Vol. 25, p. 889. Colorado.
225. *G. fuscotaeniella*, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 89 (1878) (-iaella). Texas, Colorado.
226. *G. texanella*, Chambers, Journ. Cinc. Soc. Nat. Hist. Vol. 2, p. 179 (1880). Texas.
chambersella, Dyar, List. N. Amer. Lep. p. 524 (1902).
227. *G. petrata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 63, pl. 2, f. 20 (1911). Guatemala.
228. *G. invariabilis*, Kearnott, Journ. N. York Ent. Soc. Vol. 16, p. 184 (1908). Utah.
229. *G. barnesiella*, Busck, Proc. U. S. Mus. Vol. 25, p. 875 (1903). Colorado, Arizona.
230. *G. puertella*, Busck, Proc. Ent. Soc. Wash. Vol. 18, p. 148 (1916). California.
231. *G. variabilis*, Busck, Proc. U. S. Mus. Vol. 25, p. 871 (1903). Colorado, California.
232. *G. diversella*, Busck, Proc. Ent. Soc. Wash. Vol. 18, p. 149 (1916). California.
233. *G. striatella*, Busck, Proc. U. S. Mus. Vol. 25, p. 868 (1903). Arizona.
234. *G. rectistrigella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 229 (1920). California.
235. *G. figurella*, Busck, Rep. Lagun. Mar. Lab. Vol. 1, p. 168 (1913). California.
236. *G. bigella*, Busck, Journ. Ent. Zool. Claremont, Vol. 5, p. 99 (1913). California.
237. *G. bispiculata*, Meyrick, Exot. Microlep. Vol. 3, p. 23 (1923). Arizona.
238. *G. bistrigella*, Chambers, Canad. Ent. Vol. 4, p. 92 (1872). Canada.
239. *G. dryobathra*, Meyrick, Trans. Ent. Soc. Lond. p. 49 (1917). Colombia.
240. *G. xyloglypta*, Meyrick, Exot. Microlep. Vol. 3, p. 22 (1923). California.
241. *G. flavigorporella*, Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 177 (1882). Massachusetts.
242. *G. desiliens*, Meyrick, Exot. Microlep. Vol. 3, p. 23 (1823). California.
243. *G. cacoderma*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 62, (1911). Mexico.

244. *G. ochreosuffusella*, Chambers, Canad. Ent. Vol. 6, p. 236 (1874).
depressostrigella, Chambers, ibidem, Vol. 6, p. 236 (1874). Texas.
245. *G. ochreofuscella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 249 (1875) California.
(ocherf-).
246. *G. ochreostrigella*, Chambers, ibidem, Vol. 2, p. 247 (1875). California.
247. *G. mimella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 163 (1860). Pennsylvania.
248. *G. spilotella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 228 (1929) (-*losella*). California.
249. *G. hibiscella*, Busck, Proc. U. S. Mus. Vol. 25, p. 869 (1903). District of Columbia.
250. *G. wacoella*, Chambers, Canad. Ent. Vol. 6, p. 237 (1874). Texas.
251. *G. Cockerelli*, Busck, Proc. U. S. Mus. Vol. 25, p. 871 (1903). New Mexico, Arizona.
252. *G. discocellella*, Chambers, Canad. Ent. Vol. 4, p. 194 (1872) (-*coocell-*). E. United States.
violaceofusca, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 258 (1873).
253. *G. capitochrella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 252 (1875) Texas.
(-teochr-).
254. *G. pravinomimella*, Chambers, Canad. Ent. Vol. 10, p. 50 (1878). Colorado.
quadrimaculella, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 290 (1875)
(praeocc.).
255. *G. inquilinella*, Busck, Canad. Ent. Vol. 42, p. 168 (1910). New York.
256. *G. pleroma*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 68 (1911). Mexico.
257. *G. abjunctella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 629 (1864). Cape Colony.
258. *G. liberata*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 414 (1910). Cape Colony.
259. *G. xylophaea*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 71 (1921). Rhodesia, Transvaal.
260. *G. arotrias*, Meyrick, Proc. Zool. Soc. Lond. p. 725 (1908). Natal.
261. *G. obruta*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 72 (1921). Natal.
262. *G. anagramma*, Meyrick, ibidem, Vol. 8, p. 72 (1921). Cape Colony.
263. *G. ametris*, Meyrick, ibidem, Vol. 8, p. 72 (1921). Natal.
264. *G. polygramma*, Meyrick, ibidem, Vol. 4, p. 192 (1914). Natal.
265. *G. bletrias*, Meyrick, ibidem, Vol. 3, p. 292 (1913). Transvaal.
266. *G. retiniella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 228 (1920). Nevada.
267. *G. euryanthes*, Meyrick, Ark. f. Zool. Vol. 14, no 15, p. 3 (1922). N. W. Australia.
268. *G. desmanthes*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 23, p. 51 (1898). Queensland, New South Wales.
269. *G. chalcotora*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 123 (1919). Queensland.
270. *G. nephelombra*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 309 (1904). Queensland.
271. *G. dictyomorpha*, Lower, ibidem, Vol. 25, p. 49 (1900). New South Wales.
272. *G. plinthodes*, Lower, Trans. Roy. Soc. S. Australia, Vol. 44, p. 66 (1920). New South Wales.
273. *G. epactaea*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 312 (1904). S. Australia.
274. *G. pycnoda*, Lower, ibidem, Vol. 24, p. 97 (1899). New South Wales.
275. *G. bathropis*, Meyrick, ibidem, Vol. 29, p. 310 (1904). New South Wales, S. Aus-
C. Europe. [tralia.
276. *G. rhombella*, Schiffermüller, Syst. Verz. Schmett. Wien, p. 139 (1776). Colorado.
277. *G. decemmaculella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 290 (1875). Germany, Austria,
278. *G. rhombelliformis*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 303 (1870). China. [S. Russia.
279. *G. anthracopa*, Meyrick, Exot. Microlep. Vol. 2, p. 501 (1922). Algeria.
280. *G. allotria*, n. sp.
anarsiella, Chrétien, Ann. Soc. Ent. Fr. p. 316 (1915) (praeocc.). C. Europe.
281. *G. hippophaella*, Schranck, Faun. Boic. Vol. 2, p. 115 (1802). Switzerland, Tyrol.
- basipunctella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 164, f. 530 (1855). Austria.
282. *G. acupediella*, Frey, Mitth. Schweiz. Ent. Ges. Vol. 3, p. 250 (1870). C. Europe.
283. *G. albicans*; Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 199 (1870). Canada, Kentucky.
284. *G. basiguttella*, Heinemann, ibidem, Vol. 2, p. 199 (1870). Algeria.
285. *G. rileyella*, Chambers, Canad. Ent. Vol. 4, p. 106 (1872). Tunis.
286. *G. aurantiella*, Chrétien, Ann. Soc. Ent. Fr. p. 317 (1915).
287. *G. ocellinella*, Chrétien, ibidem, p. 317 (1915).

288. *G. tragicella*, Heyden, Stett. Ent. Zeit. Vol. 26, p. 380 (1865). Alps, Saxony.
libidinosa, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 229 (1870)
289. *G. nigristrigella*, Wocke, Iris, Vol. 10, p. 374 (1897). Carinthia.
290. *G. incomptella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 178, f. 536 (1855). Germany, Latvia.
turbidella, Nolcken, Lep. Faun. Livi. Vol. 2, p. 561 (1870).
291. *G. tephriditella*, Duponchel, Hist. Nat. Lép. Fr. Suppl. Vol. 4, pl. 84, f. 11 (1842). C. Europe. [Minor.]
292. *G. spurcella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 179, f. 538 (1855). C. & S. W. Europe, Asia
293. *G. reprerella*, Zeller, Stett. Ent. Zeit. Vol. 32, p. 64 (1871). C. Europe, Corsica.
294. *G. superfetella*, Peyerimhoff, Pet. Nouv. Ent. Paris, Vol. 9, p. 102 (1877). Alsace.
295. *G. textorella*, Chrétien, Naturaliste, Vol. 30, p. 59 (1908). France.
296. *G. Hungariae*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 300 (1870). Hungary.
297. *G. suspectella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 202 (1870). Germany.
298. *G. Norvegiae*, Strand, Berl. Ent. Zeitschr. Vol. 46, p. 155 (1902). Norway.
299. *G. holosericella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 176 (1855). Alps.
cognatella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 218 (1870).
300. *G. albifaciella*, Heinemann, ibidem, Vol. 2, p. 205 (1870). Germany.
301. *G. distinctella*, Zeller, Isis, p. 199 (1839). Europe, Asia Minor, Syria, Turkestan, N. Africa.
fumatella, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 67 (1850).
celerella, Stanton, Cat. Brit. Tin. Suppl. p. 5 (1851).
striolatella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 204 (1870).
praeclarrella, Heinemann, ibidem, Vol. 2, p. 204 (1870).
tenebrosella, Teich, Stett. Ent. Zeit. Vol. 47, p. 170 (1886).
? tristella, Teich, Arb. Naturf. Ver. Riga, Vol. 32, p. 108 (1889).
indistinctella, Rebel, Cat. Lep. Palaeart. Faun. Vol. 2, p. 143 (1900).
302. *G. farinosa*, Teich, Arb. Naturf. Ver. Riga Vol. 42, p. 75 (1899). Latvia.
303. *G. oppletella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 180, f. 582 (1855). Germany, Switzerland, Austria.
nigricans, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 205 (1870).
304. *G. pallipalpella*, Snellen, Tijdschr. v. Ent. Vol. 27, p. 167, pl. 9, f. 2 (1884). E. Siberia.
305. *G. praeclarrella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 177 (1855). Alps.
terrestrella, Zeller, Stett. Ent. Zeit. Vol. 33, p. 111 (1872).
306. *G. ochripalpella*, Frey, Lep. Schweiz. p. 358 (1880). Alps.
307. *G. conspurcatella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 206 (1870). Germany.
308. *G. confusella*, Heinemann, ibidem, Vol. 2, p. 207 (1870). Germany.
309. *G. pallidigriseella*, Chambers, Canad. Ent. Vol. 6, p. 237 (1874) (*pallide-*). Texas.
310. *G. glycyrrhizella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 124 (1877) (-saeella). Colorado.
311. *G. fuscoluteella*, Chambers, Canad. Ent. Vol. 4, p. 106 (1872). Kentucky.
312. *G. nundinella*, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 256 (1873). E. United States.
beneficentella, Murtfeldt, Canad. Ent. Vol. 13, p. 245 (1881).
313. *G. monumentella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 125 (1877). Colorado.
314. *G. obscurosuffusella*, Chambers, ibidem, Vol. 4, p. 90 (1878). Texas.
? canofulvella, Chambers, ibidem, Vol. 4, p. 91 (1878).
315. *G. conspersa*, Braun, Proc. Acad. Nat. Sc. Philad. p. 9 (1921). Montana.
316. *G. montivaga*, Walsingham, Ent. M. Mag. Vol. 40, p. 221 (1904). Algeria.
317. *G. limitanella*, Rebel, Ann. Hofmus. Wien. Vol. 19, p. 349, pl. 5, f. 22 (1904). Herzegovina.
318. *G. scotinella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 181, f. 505 (1855). C. Europe.
? Kiesenwetteri, Heuäcker, Stett. Ent. Zeit. Vol. 34, p. 94 (1873).
319. *G. ruptella*, Constant, Ann. Soc. Ent. Fr. p. 192, pl. 7, f. 6 (1865). S. France.
320. *G. psammitella*, Snellen, Tijdschr. v. Ent. Vol. 27, p. 168, pl. 9, f. 3 (1884). E. Siberia.
321. *G. flavigomella*, Zeller, Isis, p. 198 (1839). C. Europe.
322. *G. longipalpella*, Teich, Arb. Naturf. Ver. Riga, Vol. 42, p. 75 (1899). Latvia.
323. *G. Jakovlevi*, Krulikowsky, Rev. Russe Ent. Vol. 5, p. 20 (1905). N. E. Russia

324. *G. sororculella*, Hübner, Samml. Eur. Schmett. Tin. f. 440 (1818). N. & C. Europe.
 325. *G. fulminella*, Millière, Ann. Soc. Linn. Lyon, p. 161, pl. 2, f. 4 (1883). S. France.
 326. *G. pinguinella*, Treitschke, Schmett. Eur. Vol. 9 (1), p. 244 (1832). Europe.
 populella, Hübner, Samml. Eur. Schmett. Tin. f. 21 (1776) (praeocc.).
 turpella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 182 (1855).
 327. *G. cuneatella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 242 (1850). C. Europe.
 328. *G. muscosella*, Zeller, Isis, p. 197 (1839). C. Europe.
 ? *griseella*, Caradja, Iris, Vol. 34, p. 96 (1920).
 329. *G. reuttiella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 196 (1870). Germany.
 syrticola, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 304 (1870).
 330. *G. nigra*, Haworth, Lep. Brit. p. 550 (1828). N. & C. Europe.
 hortuella, Wood, Index Ent. f. 1189 (1837).
 cautella, Zeller, Isis, p. 200 (1839).
 331. *G. intermedia*, Braun, Proc. Calif. Acad. Sc. (4), Vol. 12, p. 120 (1923). California.
 332. *G. paphlactis*, Meyrick, Trans. Ent. Soc. Lond. 1911, p. 693 (1912). Brazil.
 333. *G. versutella*, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 253 (1873). Texas, Wyoming, Colorado.
 334. *G. lynceella*, Zeller, ibidem, Vol. 23, p. 255 (1873). Texas. [rado, Canada.
 335. *G. obscurocellella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 254 (1875) (-rooceella). Texas.
 336. *G. occidentella*, Chambers, ibidem, Vol. 2, p. 246 (1875). California.
 337. *G. negundella*, Heinrich, Proc. U. S. Mus. Vol. 57, p. 63 (1920). N. Dakota.
 338. *G. nigromaculella*, Busck, ibidem, Vol. 25, p. 880 (1903). New York, New Jersey.
 339. *G. flexurella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 163 (1860). Pennsylvania.
 340. *G. maculimarginella*, Chambers, Canad. Ent. Vol. 6, p. 241 (1874). E. United States.
 341. *G. gilvomaculella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 12 (1863). Pennsylvania.
 biminimaculella, Chambers, Cinc. Journ. Nat. Hist. Vol. 2, p. 183 (1880).
 342. *G. veneranda*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 62 (1911). Mexico.
 343. *G. nigripectus*, Walsingham, ibidem, Vol. 4, p. 63 (1911). Mexico.
 344. *G. albipectus*, Walsingham, ibidem, Vol. 4, p. 63 (1911). California, Mexico.
 345. *G. serotinella*, Busck, Proc. U. S. Mus. Vol. 25, p. 882 (1903). Colorado, District of Co.
 346. *G. pseudocacciella*, Chambers, Canad. Ent. Vol. 4, p. 107 (1872) (-seuda-). E. United States. [lumbia.
 caecella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 252 (1873).
 347. *G. thoracocherella*, Chambers, Canad. Ent. Vol. 4, p. 169 (1872). Kentucky.
 obscurella, Chambers, ibidem, Vol. 4, p. 170 (1872).
 perobscurella, Walsingham, Ent. M. Mag. Vol. 39, p. 260 (1903).
 348. *G. thoracistrigella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 245 (1875) (-cest-). California.
 349. *G. fuscopulvella*, Chambers, Canad. Ent. Vol. 4, p. 170 (1872). Kentucky, Canada.
 obscurella, Chambers, ibidem, Vol. 4, p. 106 (1872) (-rusella) (praeocc.).
 350. *G. fuscomaculella*, Chambers, ibidem, Vol. 4, p. 170 (1872). Kentucky.
 351. *G. vernella*, Murtfeldt, ibidem, Vol. 15, p. 139 (1883). E. United States.
 formosella, Murtfeldt, ibidem, Vol. 13, p. 243 (1881) (praeocc.).
 352. *G. brumella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 416 (1864). Labrador.
 353. *G. trilineella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 125 (1877). Colorado, Arizona.
 354. *G. natalis*, Heinrich, Proc. U. S. Mus. Vol. 57, p. 62 (1920). Colorado, Oregon.
 355. *G. trichostola*, Meyrick, Exot. Microlep. Vol. 3, p. 22 (1923). Brit. Columbia.
 356. *G. melanoptila*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 22, p. 272 (1897). New South Wales.
 357. *G. planodes*, Meyrick, Exot. Microlep. Vol. 2, p. 134 (1918). S. India.
 358. *G. boreella*, Douglas, Trans. Ent. Soc. Lond. (2) Vol. 1, p. 105 (1850). N. Europe.
 359. *G. ilmatariella*, Hoffmann, Stett. Ent. Zeit. Vol. 54, p. 138 (1893). Finland.
 360. *G. galbanella*, Zeller, Isis, p. 200 (1839). N. & C. Europe.
 angustella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 217 (1870).
 361. *G. solutella*, Zeller, Isis, p. 199 (1839). C. & S. Europe, Asia
 Pribitzeri, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 49, p. 313, pl. 8, f. 8, 9 (1889). Minor.

LEPIDOPTERA HETEROCEA

362. *G. suppeliella*, Walsingham, Ent. M. Mag. Vol. 32, p. 250 (1896). England, Holland, Germany.
 363. *G. pelliella*, Treitschke, Schmett. Eur. Vol. 10 (3), p. 198 (1835). Europe. [many].
 364. *G. ignorantella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 180, f. 510 (1855). N. Germany, Latvia.
ochrisignella, Nolcken, Lep. Faun. Livl. Vol. 2, p. 548 (1870).
 365. *G. decolorella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 209 (1870). Austria, Tyrol, Yugoslavia.
 366. *G. aristotelis*, Millière, Ann. Soc. Sc. Cannes, pl. 1 f. 1, 2 (1875). S. Europe, Asia Minor, Persia, N. Africa.
astragali, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 304 (1880).
lacertella, Walsingham, Ent. M. Mag. Vol. 40, p. 222 (1904).
 367. *G. ornatissimbiella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 420 (1864). Illinois, Texas, Colorado, Arizona.
unctulella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 257 (1873).
 368. *G. amorphella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 124 (1877) Colorado.
(-phaeella).
 369. *G. abradescens*, Braun, Proc. Acad. Nat. Sc. Philad. p. 9 (1921). Montana.
 370. *G. lentiginosella*, Zeller, Isis, p. 198 (1839). C. & S. E. Europe.
 371. *G. psiloptera*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 230 (1920). Canada.
 372. *G. eburata*, Meyrick, Trans. Ent. Soc. Lond. p. 50 (1917). Colombia.
 373. *G. scotodes*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 68, pl. 2, f. 25 (1911). Mexico.
 374. *G. persicella*, Murtfeldt, Canad. Ent. Vol. 32, p. 164 (1900) (-caeella). Michigan, Canada.
confusella, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 251 (1875) (praeocc.).
 375. *G. fluvialella*, Busck, Canad. Ent. Vol. 40, p. 194 (1908). — Pl. 5, Fig. 117, a, b, c. Pennsylvania, Canada.
 376. *G. cophias*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 292 (1913). Transvaal.
 377. *G. epiphloea*, Meyrick, ibidem, Vol. 3, p. 292 (1913). Transvaal.

UNCLASSIFIED

378. *G. tehuacana*, Busck, Insec. Inscit. Menstr. Vol. 1, p. 140 (1913). Mexico.
 379. *G. scabrella*, Busck, Journ. Ent. Zool. Claremont, Vol. 5, p. 98 (1913). California.
 380. *G. heluanensis*, Debski, Bull. Soc. Ent. Egypte, p. 111 (1913). Egypt.

86. GENUS PLATYEDRA, MEYRICK

Platyedra, Meyrick, Handb. Brit. Lep. p. 605 (1895). — Type : *P. vilella*, Zeller.

Pectinophora, Busck, Journ. Agric. Res. Washington, Vol. 9, p. 347 (1917). — Type : *P. gossypiella*, Saunders.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 2/3, in ♂ serrulate, simple, basal joint elongate, with small distinct basal pecten of long hairscales. Labial palpi very long, recurved, second joint roughened and furrowed beneath, terminal joint as long as second or longer, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Anterior tibiae somewhat thickened with scales; posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from beyond middle. Hindwings somewhat over 1, trapezoidal, termen sinuate, cilia 3:4; 3 and 4 connate, 5 slightly approximated, 6 and 7 closely approximated towards base.

Remarks. — Correlated with *Gelechia*, from which it is distinguished by the antennal pecten. The spontaneous appearance of this positive character in an undoubted member of a group characterised by its absence is a curious feature worthy of study. The well-known cotton-pest, *P. gossypiella*, causes immense damage, and is the subject of much entomological literature; it probably comes from India,

whence it was first described, but has successfully established itself, in spite of the most strenuous resistance of agricultural science, in almost every region where cotton is grown.

Geographical distribution of species. — Naturally centred round the Mediterranean.

Larva (3 known) feeding especially in seed-capsules.

Foodplants always *Malvaceae*.

- | | |
|---|--|
| 1. <i>P. virella</i> , Zeller, Isis, p. 845 (1847). | C. & S. Europe, Canaries, |
| <i>argillosella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, f. 514 (1855). | N. Africa, W.C. & S.W. |
| 2. <i>P. malvella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 281 (1805). | C. & S. Europe. [Asia. |
| 3. <i>P. cruenta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 298 (1920). | Palestine. |
| 4. <i>P. sarcochroma</i> , Walsingham, Bull. Liverpool Mus. Vol. 3, p. 3 (1900). | Sokotra. |
| 5. <i>P. gossypiella</i> , Saunders, Trans. Ent. Soc. Lond. (1), Vol. 3, p. 285 (1843). — | India, Ceylon, Egypt, |
| <i>umbribennis</i> , Walsingham, Proc. Zool. Soc. Lond. p. 884 (1885). | China, Japan, Philip-
pines, New Guinea,
Australia, Hawaii, Fiji,
Texas, Mexico, S. Amer-
ica. |

Pl. 2, Fig. 30.

87. GENUS MOMETA, DURRANT

Mometa, Durrant, Bull. Ent. Research, London, Vol. 5, p. 243 (1914). — Type : *M. semiodes*, Durrant.

Characters. — Head with smoothly appressed scales; ocelli very small, posterior; tongue developed. Antennae 3/4, in ♂ shortly ciliated, basal joint elongate, with slight pecten. Labial palpi long, recurved, second joint thickened with smoothly appressed scales, somewhat furrowed beneath, terminal joint as long as second, thickened with projecting scales posteriorly, apex slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with rough hairscales above. Forewings with 1b furcate, 2 from towards angle, 3-5 closely approximated from angle, 7 and 8 stalked, 7 to costa, 11 from somewhat beyond middle. Hindwings 1, trapezoidal, termen sinuate, cilia 4/5; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated towards base.

Remarks. — Closely allied to *Ptalydrea* in all respects, though of different colouring.

Geographical distribution of species. — Equatorial African.

Larva feeding in seed-capsules.

Foodplant *Gossypium* (*Malvaceae*).

1. *M. semiodes*, Durrant, Bull. Ent. Research, London, Vol. 5, p. 243 (1914). Nigeria, Kenya Colony.
Pl. 2, Fig. 33.

88. GENUS TABERNILLAEA, WALSINGHAM

Tabernillaea, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 53 (-laia). — Type : *T. ephialtes*, Walsingham.

Characters. — Head smooth; tongue developed. Antennae 3/4, in ♂ simple, basal joint without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint nearly as long as second, thickened, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings in ♀ 1, trapezoidal, termen sinuate, cilia nearly 1, in ♂ narrowed, with median area much folded and reduced; 3 and 4 connate, 5 approximated, 6 and 7 almost connate, in ♂ 2-5 concealed in the folded area.

Remarks. — A derivative of *Parastega*.

Geographical distribution of species. — Central American.

Larva unknown.

1. *T. ephialtes*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 54, pl. 2, Panama.
f. 12 (1911).

89. GENUS LOCHARCHA, MEYRICK

Locharcha, Meyrick, Exot. Microlep. Vol. 3, p. 18 (1923). — Type : *L. emicans*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ somewhat stout, simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with very long rough projecting hairscales beneath, terminal joint as long as second, thickened with rough scales projecting posteriorly, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with 2-4 near together, parallel, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen faintly sinuate, cilia 1; 3 and 4 connate, 5 rather approximated, 6 and 7 nearly approximated towards base.

Remarks. — Allied to *Parastega*.

Geographical distribution of species. — South American.

Larva unknown.

1. *L. emicans*, Meyrick, Exot. Microlep. Vol. 3, p. 18 (1923). Peru.

90. GENUS PARASTEGA, MEYRICK

Parastega, Meyrick, Trans. Ent. Soc. Lond. 1911, p. 693 (1912). — Type : *P. niveisignella*, Zeller.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint beneath with brush of rough projecting scales, terminal joint as long as second, expanded posteriorly with loose projecting scales except at apex, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 3 and 4 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 2/3, 3 and 4 nearly approximated at base, 5 approximated, 6 and 7 nearly approximated towards base or stalked.

Remarks. — Apparently related to *Stegasta*.

Geographical distribution of species. — Central and South American.

Larva unknown.

1. *P. chionostigma*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 57, Panama, Ecuador,
pl. 2, f. 17 (1911). — PI. 2, Fig. 32. Guiana, Brazil.
ochropis, Meyrick, Trans. Ent. Soc. Lond. p. 236 (1914).
2. *P. trichella*, Busck, Proc. U. S. Mus. Vol. 47, p. 16 (1914). Panama.
3. *P. niveisignella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 333, pl. 4, f. 106 Panama, Guiana,
(1877). Colombia, Peru.
curvatella, Busck, Proc. U. S. Mus. Vol. 47, p. 16 (1914).

91. GENUS STEGASTA, MEYRICK

Stegasta, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 313 (1904). — Type : *S. variana*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennæ 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, rather rough beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1 $\frac{1}{2}$ furcate, 2 from towards angle, 3 and 4 stalked or connate. 7 and 8 stalked, 7 to costa, 11 from middle; in ♂ beneath with a reflexed fringe of scales from anterior half of costa protecting an expansible pencil of long hairs from base. Hindwings 1, elongate-trapezoidal, termen more or less sinuate, cilia 1-1 $\frac{1}{2}$; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked or approximated.

Remarks. — Allied to *Gelechia*.

Geographical distribution of species. — Tropical American and Australian, but one species has been widely distributed with cultivated plants throughout the Indo-Malayan and African regions. The home of the genus is America, and its connection with Australia may be contemporaneous with that of the Oecophorid genus *Machimia*.

Larva feeding between spun leaves.

Foodplants *Leguminosæ*, especially *Cassia*.

1. <i>S. tenebricosa</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 124 (1919).	Queensland.
2. <i>S. allactis</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 314 (1904).	S. & W. Australia.
3. <i>S. cosmodes</i> , Lower, ibidem, Vol. 24, p. 98 (1899).	New South Wales.
4. <i>S. conjugella</i> , Caradja, Iris, Vol. 34, p. 102 (1920).	W. Turkestan.
5. <i>S. cephalella</i> , Caradja, Iris, Vol. 34, p. 101 (1920).	Uralsk.
6. <i>S. capitella</i> , Fabricius, Ent. Syst. Vol. 3 (2), p. 330 (1794).	Cuba, Jamaica, Portorico, Virgin Islands.
robustella, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 597 (1864).	
• rivulella, Möschler, Abhandl. Senckenb. Naturf. Ges. Vol. 15, p. 344 (1900),	
7. <i>S. zygotoma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 48 (1917).	Colombia, Ecuador,
8. <i>S. comissata</i> , Meyrick, Exot. Microlep. Vol. 3, p. 18 (1923).	Brazil. [Peru.]
9. <i>S. variana</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 314 (1904). — PI. 2, Fig. 31.	Australia, Indo-Malayan region, Africa.
10. <i>S. bosquella</i> , Chambers, Canad. Ent. Vol. 7, p. 92 (1875) (<i>basqueella</i>). costipunctella, Möschler, Abhandl. Senckenb. Naturf. Ges. Vol. 15, p. 344 (1900)	E. United States, Jamaica, Portorico, Antilles.
11. <i>S. biniveipunctata</i> , Walsingham, Proc. Zool. Soc. Lond. p. 75 (1897).	Grenada.
12. <i>S. postpallescens</i> , Walsingham, ibidem, p. 76 (1897)	Grenada.
13. <i>S. phalacra</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 61 (1911).	Mexico.
14. <i>S. donatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 596 (1864).	Cuba, Jamaica.

92. GENUS THYMOSOPHA, MEYRICK

Thymosopha, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 244 (1914). — Type : *T. antileuca*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennæ 3/4, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, slightly roughened anteriorly, terminal joint shorter, thickened with scales projecting posteriorly

above middle, apex slender, acute. Maxillary palpi very short. Posterior tibiae with scanty appressed hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked. 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex tolerably pointed, termen slightly sinuate, cilia 1; 3 and 4 connate, 5 slightly approximated, 6 and 7 parallel.

Remarks. — Seemingly transitional between *Phthorimaea* and the preceding genera.

Geographical distribution of species. — South African.

Larva unknown.

1. *T. antileuca*, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 245 (1914).

Cape Colony.

93. GENUS PHLOEOCECIS, CHRÉTIEN

Phloeocecis, Chrétien, Bull. Soc. Ent. Fr. p. 91 (1908). — Type : *P. fagoniae*, Meyrick.

Characters. — Head smooth. Basal joint of antennae without pecten. Labial palpi long, curved, ascending, second joint with appressed scales, slightly tufted at apex, terminal joint shorter than second, slender, acute. Posterior tibiae clothed with long dense hairs. Forewings with 2 very short, 3 and 4 approximated, 5 somewhat approximated, 7 and 8 stalked, 7 to costa, cell long, narrow. Hindwings under 1, trapezoidal, apex shortly produced, pointed, termen oblique; 2 short, approximated to 3, 3 and 4 connate, 5 slightly curved, 6 and 7 parallel.

Remarks. — Not known to me. I alter the specific name, for which Arabic vernacular is no more permissible than French would be.

Geographical distribution of species. — North African.

Larva feeding in a swelling in stems.

Foodplant *Fagonia* (*Zygophyllaceae*).

1. *P. fagoniae*, nov. sp.

cherregella, Chrétien, Bull. Soc. Ent. Fr. p. 92, (1908) (van.).

Algeria.

94. GENUS THOLEROSTOLA, MEYRICK

Tholerostola, Meyrick, Trans. Ent. Soc. Lond. p. 40 (1917). — Type : *T. omphalopa*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue apparently obsolete. Antennae 3/4, in ♂ serrulate, simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint rather thickened and roughened with scales beneath, terminal joint hardly shorter than second, slightly thickened with scales, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1b long-furcate, 2 from towards angle, 4 and 5 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 4/5, trapezoidal, apex rather produced, acute, termen obtusely emarginate beneath it, cilia 2; 2 from 2/3, 3 approximated to 4, 4 and 5 connate or closely approximated from angle, transverse vein mostly obsolete except near lower extremity, 6 obsolete, represented only by a weak fold parallel to 7.

Remarks. — A peculiar form, of which the affinity may be uncertain.

Geographical distribution of species. — South American.

Larva unknown.

1. *T. omphalopa*, Meyrick, Trans. Ent. Soc. Lond. p. 40 (1917).

Ecuador.

95. GENUS TECIA, STRAND

Tecia, Strand, Berl. Ent. Zeitschr. Vol. 55, p. 165 (1911). — Type : *T. mendozella*, Strand.

Lata, Strand, ibidem, Vol. 55, p. 167 (1911). — Type : *T. Kiefferi*, Strand.

Fapua, Strand, ibidem, Vol. 55, p. 168 (1911). — Type : *T. albinervella*, Strand.

Characters. — Head with appressed scales. Antennae 4/5, in ♂ simple. Labial palpi very long, obliquely ascending, second joint very long, above with long dense rough projecting hairs diminishing to apex, sometimes also tufted beneath, terminal joint less than half second, slender, acute. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from angle, 2-6 parallel, 7 and 8 stalked, 7 to costa, 11 from beyond middle. Hindwings somewhat over 1, rather elongate-ovate, cilia 1; 3 and 4 connate, 5 slightly approximated, 6 and 7 parallel.

Remarks. — Not known to me; probably related to *Gnorimoschema*. The three alleged genera are discriminated only by differences in the scaling of the second joint of palpi, and do not seem distinct.

Geographical distribution of species. — South American.

Larva (all 3 known) feeding in stem-galls.

Foodplants *Compositae*.

1. <i>T. mendozella</i> , Strand, Berl. Ent. Zeitschr. Vol. 55, p. 166 (1911)	Argentina.
2. <i>T. Kiefferi</i> , Strand, ibidem, Vol. 55, p. 167 (1911).	Argentina.
3. <i>T. albinervella</i> , Strand, ibidem, Vol. 55, p. 168 (1911).	Argentina.

96. GENUS GNORIMOSCHEMA, BUSCK

Gnorimoschema, Busck, Proc. U. S. Mus. Vol. 23, p. 227 (1900). — Type : *G. gallaesolidaginis*, Riley.

Tuta, Strand, Berl. Ent. Zeitschr. Vol. 55, p. 169 (1911). — Type : *G. atriplicella*, Strand.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint much thickened with dense scales, rough and furrowed beneath, terminal joint much shorter than second, thickened with scales, compressed, pointed, anterior edge serrate. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long rough hairs above. Forewings sometimes with tufts of scales; 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen more or less sinuate, cilia 1 — 2; 3 and 4 connate, 5 somewhat approximated, 6 and 7 parallel.

Remarks. — A derivative of *Phthorimaea*, from which it is distinguished by the different terminal joint of palpi, and more truly parallel veins 6 and 7 of hindwings.

Geographical distribution of species. — Strictly American.

Larva (21 known) feeding especially in gall-like swellings in stems, but also sometimes mining in leaves or fruits, or within spun shoots or terminal gall.

Foodplants especially *Compositae* (16), but also *Chenopodiaceae*, *Solanaceae*, *Labiatae*.

1. <i>G. serratipalpella</i> , Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 123 (1877).	Colorado, New Mexico.
2. <i>G. dudiella</i> , Busck, Proc. U. S. Mus. Vol. 25, p. 828 (1903).	Arizona, Mexico.

3. *G. septentrionella*, Fyles, Canad. Ent. Vol. 43, p. 422 (1911). Quebec.
 4. *G. gibsoniella*, Busck, Proc. Ent. Soc. Wash. Vol. 17, p. 82 (1915). Manitoba.
 5. *G. gallaesolidaginis*, Riley, Rep. Ins. Missouri, Vol. 1, p. 170 (1869). — E. United States.
Pl. 2, Fig. 36. Canada.
 6. *G. salinaris*, Busck, Canad. Ent. Vol. 43, p. 4 (1911). Massachusetts.
 7. *G. laguna*, Busck, Rep. Laguna Mar. Lab. Vol. 1, p. 167 (1913?). California.
 8. *G. gallaeasteris*, Kellicott, Canad. Ent. Vol. 10, p. 203 (1878) (-*terella*). E. United States, Canada.
 9. *G. gallaeadioplopappi*, Fyles, ibidem, Vol. 22, p. 248 (1890). Quebec.
 10. *G. washingtoniella*, Busck, Proc. U. S. Mus. Vol. 27, p. 757 (1904). Distr. Columbia.
 11. *G. bacchariella*, Busck, ibidem, Vol. 25, p. 825 (1903) (-*isella*). California.
 12. *G. petrella*, Busck, Proc. Ent. Soc. Wash. Vol. 17, p. 83 (1915). New Hampshire.
 13. *G. conifera*, Meyrick, Exot. Microlep. Vol. 1, p. 582 (1916). Ecuador.
 14. *G. aquilina*, Meyrick, Trans. Ent. Soc. Lond. p. 44 (1917). — **Pl. 1,** Peru.
Fig. 36.
 15. *G. subterranea*, Busck, Canad. Ent. Vol. 43, p. 5 (1911). Massachusetts.
 16. *G. coquillettiella*, Busck, Proc. U. S. Mus. Vol. 25, p. 405 (1902). California, Colorado.
 17. *G. buschiella*, Kearfott, Journ. N. York Ent. Soc. Vol. 11, p. 158, pl. 9, f. 7 (1903). New Jersey.
 18. *G. chiquitella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 176 (1909). New Mexico.
 19. *G. tetradymiella*, Busck, Proc. U. S. Mus. Vol. 25, p. 834 (1903). California.
 20. *G. alaricella*, Busck, Can. Ent. Vol. 40, p. 193 (1908). Pennsylvania.
 21. *G. banksiella*, Busck, Proc. U. S. Mus. Vol. 25, p. 832 (1903). N. E. United States,
 22. *G. contraria*, Braun, Proc. Acad. Nat. Sc. Philad. p. 8 (1921). Montana. [Canada].
 23. *G. albimarginella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 291 (1875). Colorado.
 24. *G. semicyclella*, Busck, Proc. U. S. Mus. Vol. 25, p. 828 (1903) (-*clonella*). Colorado, New Mexico.
 25. *G. terracoccella*, Busck, ibidem, Vol. 23, p. 227 (1900) (-*cottella*). Florida.
 26. *G. pedmontella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 123 (1877). Colorado.
 27. *G. faustella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 176 (1909). New Mexico.
 28. *G. collinella*, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 128 (1877) (-*nusella*). Colorado.
 29. *G. atriplex*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 175 (1909). California.
 30. *G. pallidochrella*, Chambers, Canad. Ent. Vol. 4, p. 126 (1872). Kentucky.
 ? *versicolorella*, Chambers, ibidem, Vol. 4, p. 127 (1872).
 31. *G. henshawiella*, Busck, Proc. U. S. Mus. Vol. 25, p. 831 (1903). Montana, Colorado, New
 ochrostrigella, Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 126 (1877) Mexico.
 (praeocc.).
 32. *G. triocellella*, Chambers, ibidem, Vol. 3, p. 127 (1877). Colorado.
 33. *G. octomaculella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 291 (1875). California.
 34. *G. princeps*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 175 (1909). California.
 35. *G. radiatella*, Busck, Proc. U. S. Mus. Vol. 27, p. 758 (1904). Distr. Columbia.
 36. *G. saphirinella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 250 (1875). Texas, Mexico.
 37. *G. splendoriferella*, Busck, Proc. U. S. Mus. Vol. 27, p. 758 (1904). Distr. Columbia.
 38. *G. florella*, Busck, ibidem, Vol. 25, p. 832 (1903). Colorado.
 39. *G. illyella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 337 (1877). Colombia.
 40. *G. batanella*, Busck, Proc. U. S. Mus. Vol. 25, p. 833 (1903). New Jersey.
 41. *G. erigeronella*, Braun, Proc. Acad. Nat. Sc. Philad. p. 7 (1921). Montana.
 42. *G. brackenridgella*, Busck, Dyar, List. N. Amer. Lep. p. 516 (1902). Pennsylvania.
 detersella, Clemens, Proc. Acad. Nat. Sc. Philad. p. 164 (1860) (praeocc.).
 43. *G. scutellariella*, Chambers, Canad. Ent. Vol. 5, p. 175 (1873) (-*iaeella*). Kentucky.
 44. *G. lavernella*, Chambers, ibidem, Vol. 6, p. 242 (1874). E. United States.
 physalivorella, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 238 (1875). Argentina.
 45. *G. atriplicella*, Strand. Berl. Ent. Zeitschr. Vol. 55, p. 169 (1911). Massachusetts.
 46. *G. Chenopodiella*, Busck, Proc. Ent. Soc. Wash. Vol. 18, p. 143 (1916). Kentucky.
 47. *G. ambrosiella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 239 (1875) (-*iaeella*).

48. <i>G. artemisiella</i> , Kearfott, Journ. N. York Ent. Soc. Vol. 11, p. 160, pl. 9, f. 5 (1903).	E. United States, California.
<i>axenopis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 234 (1919).	
49. <i>G. lenta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 41 (1917).	Peru.
50. <i>G. atrifascis</i> , Meyrick, ibidem, p. 45 (1917).	Peru.
51. <i>G. ventralella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 348, pl. 4, f. 116 (1877).	Colombia.
52. <i>G. loquax</i> , Meyrick, Trans. Ent. Soc. Lond. p. 45 (1917).	Peru.
53. <i>G. trichinaspis</i> , Meyrick, ibidem, p. 41 (1917).	Peru.
54. <i>G. crustaria</i> , Meyrick, ibidem, p. 42 (1917).	Peru.
55. <i>G. gregalis</i> , Meyrick, ibidem, p. 43 (1917).	Peru.
56. <i>G. densata</i> , Meyrick, ibidem, p. 42 (1917).	Peru.
57. <i>G. absoluta</i> , Meyrick, ibidem, p. 44 (1917).	Peru.

97. GENUS PHTHORIMAEA, MEYRICK

Phthorimaea, Meyrick, Ent. M. Mag. Vol. 38, p. 103 (1902). — Type : *T. operculella*, Zeller.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple or minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, rough and furrowed beneath, terminal joint as long as second or slightly shorter or longer, with appressed scales, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings seldom with slight scale-tufts; 1 b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, seldom 6 out of 7 near base or 7 and 8 out of 6, 11 from middle. Hindwings 1 or nearly, trapezoidal, apex more or less pointed, termen sinuate, cilia 2/3 to nearly 2; 3 and 4 connate, 5 somewhat approximated, 6 and 7 subparallel, sometimes rather approximated or bent inwards at base.

Remarks. — Probably correlated with *Gelechia*. The species are often obscure, variable, and similar, requiring careful study, with knowledge of the larval habits : *P. operculella* on potato and *P. heliopa* on tobacco are well-known and very injurious pests. The name *Lita*, Treitschke, was erroneously applied to this genus by Heinemann and subsequent European authors.

Geographical distribution of species. — Summarised thus : Palaearctic 133, American 24, South African 20, Indian 9, Australian 13, New Zealand 7. Hence it appears that Europe is the home of this genus, where it may have originated at the same time that *Gelechia* appeared in North America; it is now spreading over the world, but in America, probably the scene of its first invasion, it has been largely supplanted by its own derivative *Gnorimoschema*, specially adapted to the large Composite flora of that region.

Larva (74 known) feeding in spun shoots, or often mining in leaves or stems, sometimes forming galls; almost always on low plants or herbs, rarely shrubs, never trees.

Foodplants : *Caryophyllaceae* 25, *Chenopodiaceae* 14, *Compositae* 13, *Solanaceae* 13, and 8 other Orders with only 2 or 1 species each, showing an extraordinary preference for the four first-named, of which moreover the *Caryophyllaceae* and *Chenopodiaceae* are genetically closely related; connected with this preference is a fondness for dry and sandy localities, often also for sea-coasts. Further, when the species are grouped by their natural affinities, it is found that, in the case of each of these four Orders, all the species which feed on plants of one Order fall naturally together, without the intrusion of a single species known to feed on another of the four; all the Caryophyllaceous feeders fall together, all the Solanaceous (no other genus shows an equal attachment to *Solanaceae*) and so on. I am not acquainted

with any other large genus of *Tineina* in which this holds. Hence the discovery of larval habits in this genus is often easy.

1. *P. subcaerulea*, Meyrick, Exot. Microlep. Vol. 2, p. 136 (1918).
2. *P. atalopis*, Meyrick, ibidem, Vol. 2, p. 135 (1918).
3. *P. infirma*, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 61 (1912).
4. *P. geomicta*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 18 (1918).
5. *P. extorris*, Meyrick, Exot. Microlep. Vol. 3, p. 50 (1923).
6. *P. mixolitha*, Meyrick, ibidem, Vol. 2, p. 135 (1918).
7. *P. suasaria*, Meyrick, ibidem, Vol. 2, p. 135 (1918).
8. *P. ochrodetata*, Meyrick, ibidem, Vol. 3, p. 23 (1923).
9. *P. jamaicensis*, Walsingham, Proc. Zool. Soc. Lond. p. 76 (1897).
10. *P. salva*, n. sp.
leucoccephala, Walsingham, Proc. Zool. Soc. Lond. p. 74 (1897) (praeocc.).
11. *P. sphenophora*, Walsingham, ibidem, p. 73 (1897).
12. *P. textifera*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 288 (1913).
13. *P. fanatica*, Meyrick, ibidem, Vol. 8, p. 73 (1921).
14. *P. dispensata*, Meyrick, ibidem, Vol. 8, p. 73 (1921).
15. *P. synecta*, Meyrick, ibidem, Vol. 2, p. 12, pl. 4, f. 10 (1911). — **Pl. 2,**
Fig. 34.
16. *P. ericinista*, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 245 (1914).
17. *P. cecidiella*, Chrétien, Ann. Soc. Ent. Fr. p. 322 (1915).
18. *P. zygophyllella*, Rebel, Iris, Vol. 26, p. 89 (1912).
19. *P. diminutella*, Zeller, Isis, p. 855 (1847).
subdiminutella, Stainton, Tin. Syr. p. 45 (1867).
20. *P. pusillella*, Rebel, Stett. Ent. Zeit. Vol. 54, p. 47 (1893).
21. *P. promptella*, Staudinger, ibidem, Vol. 20, p. 241 (1859).
22. *P. porcella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 253 (1870).
23. *P. pallidella*, Heinemann, ibidem, Vol. 2, p. 252 (1870).
24. *P. sibila*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 73 (1921).
25. *P. chersophila*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 351 (1909).
26. *P. nomias*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 74 (1921).
27. *P. intestina*, Meyrick, ibidem, Vol. 8, p. 74 (1921).
28. *P. colasta*, Meyrick, ibidem, Vol. 8, p. 74 (1921).
29. *P. pendens*, Meyrick, ibidem, Vol. 6, p. 18 (1918).
30. *P. vicaria*, Meyrick, ibidem, Vol. 8, p. 74 (1921).
31. *P. phelotris*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 371 (1909).
32. *P. eremaula*, Meyrick, Ent. M. Mag. Vol. 27, p. 57 (1891).
33. *P. parvipulex*, Walsingham, ibidem, Vol. 47, p. 13 (1911).
34. *P. gecko*, Walsingham, ibidem, Vol. 47, p. 12 (1911).
35. *P. halymella*, Millière, Icon. Descr. Lép. Vol. 1, p. 352, pl. 42, f. 4-8 (1864).
36. *P. molitor*, Walsingham, Proc. Zool. Soc. Lond. p. 278 (1896).
37. *P. spargulariella*, Chrétien, Naturaliste, Vol. 32, p. 261 (1910).
38. *P. crocoleuca*, Meyrick, Exot. Microlep. Vol. 3, p. 51 (1923).
39. *P. suaedivorella*, Chrétien, Ann. Soc. Ent. Fr. p. 320 (1915).
40. *P. suaedella*, Richardson, Ent. M. Mag. Vol. 29, p. 241 (1893).
41. *P. plantaginella*, Stainton, ibidem, Vol. 19, p. 253 (1883).
42. *P. Brunhildae*, Schawerda, Iris, Vol. 35, p. 134 (1921).
43. *P. submissella*, Stainton, Ann. Mag. Nat. Hist. (3), Vol. 3, p. 212 (1859).
44. *P. saltinella*, Schöyen, Tröms. Mus. Aars. Vol. 5, p. 55 (1882).
Stangei, Hering, Stett. Ent. Zeit. Vol. 50, p. 299 (1889).
45. *P. traganella*, Chrétien, Ann. Soc. Ent. Fr. p. 321 (1915).
46. *P. salicorniae*, Hering, Stett. Ent. Zeit. Vol. 50, p. 302 (1889).
47. *P. omachella*, Oberthür, Etudes Ent. Vol. 12, p. 43, pl. 6, f. 28 (1888).
desertella, Rebel, Iris, Vol. 13, p. 165 (1900).

Kanara.	
Ceylon, India, Burma.	
Rhodesia, Transvaal,	
Natal. [Natal.	
Egypt.	
Bengal.	
India, Palestine.	
Palestine.	
Jamaica.	
Virgin Islands.	
Grenada.	
Transvaal.	
Rhodesia.	
Transvaal.	
Transvaal, Cape Colony.	
Cape Colony.	
Algeria.	
Egypt.	
S. Europe, Palestine.	
Spain.	
Spain, Sicily.	
Bavaria.	
Germany.	
Port. E. Africa.	
Cape Colony.	
Zululand.	
Transvaal.	
Transvaal.	
Zululand.	
Transvaal.	
Cape Colony.	
Algeria.	
Algeria.	
Algeria.	
S. France, Algeria.	
Arabia.	
S. France.	
Egypt.	
Algeria.	
England, France.	
England, Moravia.	
Croatia.	
S. France, Madeira.	
Norway, Germany.	
Algeria.	
England, Germany.	
Algeria.	

48. *P. ocellatella*, Boyd, Ent. Weekl. Intell. Vol. 4, p. 143 (1858).
portosanctana, Stainton, Ann. Mag. Nat. Hist. (3), Vol. 3, p. 212 (1859).
horticolella, Rössler, Verz. Schmett. Nassau, p. 240 (1866).
49. *P. vasconiella*, Rössler, Stett. Ent. Zeit. Vol. 38, p. 377 (1877).
50. *P. lagunella*, Chrétien, Naturaliste, Vol. 32, p. 272 (1910).
51. *P. obsoletella*, Fischer von Röslerstamm, Abbild. Schmett. p. 225, pl. 79 (1840).
52. *P. samadensis*, Pfaffenzeller, Stett. Ent. Zeit. Vol. 31, p. 321 (1870).
instabilella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 249 (1870).
53. *P. instabilella*, Douglas, Zoologist, p. 1270, f. 10 (1846).
54. *P. nitentella*, Fuchs, Stett. Ent. Zeit. Vol. 63, p. 324 (1902).
55. *P. atriplicella*, Fischer van Röslerstamm, Abbild. Schmett. p. 223, pl. 78 (1839).
56. *P. bucolica*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 317 (1904).
57. *P. petrinodes*, Meyrick, ibidem, Vol. 29, p. 318 (1904).
58. *P. nonyma*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 125 (1919).
59. *P. eschatopis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 321 (1904).
60. *P. frequens*, Meyrick, Exot. Microlep. Vol. 2, p. 426 (1921).
61. *P. silignitis*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 125 (1919).
62. *P. pyrrhanthes*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 318 (1904).
63. *P. marina*, Meyrick, ibidem, Vol. 29, p. 319 (1904).
64. *P. chersochlora*, Meyrick, Ark. f. Zool. Vol. 14, no 15, p. 3 (1922).
65. *P. xerophylla*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 320 (1904).
66. *P. thyraula*, Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 167 (1886).
67. *P. brontophora*, Meyrick, ibidem, Vol. 18, p. 168 (1886).
68. *P. plaeisosema*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 126 (1919).
69. *P. leucocephala*, Lower, Trans. Roy. Soc. S. Australia, Vol. 35, p. 169 (1893).
70. *P. perdita*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 24, p. 96 (1899).
71. *P. ferella*, Berg, Bull. Soc. Nat. Mosc. Vol. 49 (2), p. 241 (1875).
72. *P. perfidiosa*, Meyrick, Trans. Ent. Soc. Lond. p. 41 (1917).
73. *P. melanocampta*, Meyrick, ibidem, p. 44 (1917).
74. *P. urosema*, Meyrick, ibidem, p. 43 (1917).
75. *P. milleriella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 253 (1875).
76. *P. subtractella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 592 (1864).
77. *P. impudica*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 51 (1911).
78. *P. epithymella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 242 (1859).
79. *P. inexperta*, n. sp.
simplicella, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 238 (1875) (praeocc.).
80. *P. striatella*, Murtfeldt, Canad. Ent. Vol. 22, p. 163 (1900).
81. *P. minor*, Busck, Proc. U. S. Mus. Vol. 30, p. 726 (1906).
82. *P. glochinella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 263, pl. 3, f. 18 (1873).
solanella, Chambers, Canad. Ent. Vol. 5, p. 176 (1873).
cinerella, Murtfeldt, ibidem, Vol. 13, p. 244 (1881).
inconspicuella, Murtfeldt, ibidem, Vol. 15, p. 139 (1883).
83. *P. gudmannella*, Walsingham, Proc. Zool. Soc. Lond. p. 77 (1897).
84. *P. operculella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 262, pl. 3, f. 17 (1873).
terrella, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1024 (1864) (praeocc.).
solanella, Boisduval, Bull. Soc. Ent. Fr. p. 35 (1875).
tabacella, Ragonot, ibidem, p. 147 (1879).
sedata, Butler, Cistula Ent. Vol. 2, p. 560 (1880).
- W. C. & S. Europe, Asia Minor, Madeira, N. Africa, Ceylon. Spain. S. France. C. Europe. Switzerland. England. Latvia. Germany. C. Europe, Asia Minor, Turkestan. E. & W. Australia. Queensland, New South Victoria. [Wales. West Australia. Queensland. Queensland. West Australia. New South Wales. N. W. Australia. New South Wales, Victoria. New Zealand. New Zealand. New South Wales. New South Wales, S. Australia. New South Wales. Patagonia. Colombia. Peru. Peru. Texas. Nova Scotia. Panama. Spain, S. France. Kentucky. Missouri, California. Texas. Texas, Missouri. San Domingo, Virgin Is. N. America, Portorico, Hawaii, Tahiti, Australia, New Zealand, India, Africa, S. Europe.

85. *P. emancipata*, nov. sp.
marmorella, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 239 (1875) (praeocc.)
86. *P. discomaculella*, Chambers, Canad. Ent. Vol. 4, p. 172 (1872).
aurimaculella, Chambers, ibidem, Vol. 4, p. 172 (1872).
87. *P. lyciella*, Walsingham, Ent. M. Mag. Vol. 36, p. 217 (1900).
88. *P. micradelpha*, Walsingham, ibidem, Vol. 36, p. 217 (1900).
89. *P. heliopa*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 417 (1900).
? *aptatella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 636 (1864).
90. *P. daturae*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 359 (1877).
91. *P. costella*, Westwood, Brit. Moths, Vol. 2, p. 192 (1851).
92. *P. hyoscyamella*, Stainton, Tin. S. Eur. p. 233 (1869).
93. *P. ergasima*, Meyrick, Exot. Microlep. Vol. 1, p. 568 (1916).
94. *P. blapsigona*, Meyrick, ibidem, Vol. 1, p. 569 (1916).
95. *P. cretigena*, Meyrick, Ann. Transv. Mus. Vol. 4, p. 191 (1914).
96. *P. concreta*, Meyrick, ibidem, Vol. 4, p. 191 (1914).
97. *P. epicentra*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 370 (1909).
98. *P. fornacaria*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 289 (1913).
99. *P. phalacrodes*, Meyrick, ibidem, Vol. 3, p. 293 (1913).
100. *P. singula*, Staudinger, Stett. Ent. Zeit. Vol. 37, p. 145 (1876).
101. *P. streliciella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 171, f. 495 (1855).
stralitziella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 245 (1870)
102. *P. Hoefneri*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. (331) (1909).
103. *P. valesiella*, Staudinger, Stett. Ent. Zeit. Vol. 38, p. 205 (1877).
104. *P. steudeliella*, Frey, Lep. Schweiz. p. 363 (1880).
105. *P. Passiezkyi*, Rebel, Rov. Lap. Vol. 20, p. 173 (1913).
106. *P. fiumella*, Krone, Jahrb. Ent. Ver. Wien, Vol. 21, p. 39 (1911).
107. *P. inustella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 171, f. 498 (1855).
108. *P. ustulatella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 307 (1870).
109. *P. psilella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 171, f. 496 (1855).
? *Herbichi*, Nowicki, Microlep. Sp. Nov. p. 17, pl. 1, f. 6 (1864).
110. *P. gallicella*, Constant, Ann. Soc. Ent. Fr. p. 253, pl. 10, f. 14 (1884).
111. *P. tengstroemiella*, Joannis, Bull. Soc. Ent. Fr. p. 296 (1910).
112. *P. thymifoliella*, Constant, Ann. Soc. Ent. Fr. p. 394, pl. 11, f. 5 (1893).
113. *P. cistiflorella*, Constant, Bull. Soc. Ent. Fr. p. 25 (1889).
114. *P. oreocyrniella*, Petry, Stett. Ent. Zeit. Vol. 65, p. 249 (1904).
115. *P. artemisiella*, Treitschke, Schmett. Eur. Vol. 9 (2), p. 97 (1832).
116. *P. proclivella*, Fuchs, Stett. Ent. Zeit. Vol. 47, p. 68 (1886).
117. *P. phagnalella*, Constant, Bull. Soc. Ent. Fr. p. 52 (1895).
118. *P. gallincolella*, Mann, Verh. Zool.-bot. Ges. Wien, Vol. 22, p. 37 (1872).
119. *P. dissectella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 241 (1859).
120. *P. tussilaginella*, Heinemann, Schmett. Deutschl. (2) Vol. 2, p. 251 (1870).
121. *P. furfurella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 308 (1870).
122. *P. ochraceella*, Chrétien, Ann. Soc. Ent. Fr. p. 320 (1915).
123. *P. gypsetta*, Constant, ibidem, p. 396, pl. 11, f. 6 (1893).
124. *P. praticolella*, Christoph, Hor. Soc. Ent. Ross. Vol. 9, p. 25, pl. 2, f. 22 (1872).
125. *P. salinella*, Zeller, Isis, p. 853 (1847).
126. *P. nigromaculella*, Millière, Pet. Nouv. Ent. Paris, Vol. 4, p. 172 (1872).
127. *P. punctata*, Staudinger, Stett. Ent. Zeit. Vol. 37, p. 149 (1876).
128. *P. palermittella*, Laharpe, Contrib. Faun. Sic. p. 8 (1860).
129. *P. insulella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 251 (1870).
insularis, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 304 (1870).

Kentucky.

Kentucky.

S. France, Algeria. [ries.

S. France, Algeria, Cana-

India, Ceylon, Burma,

Java, Australia, S. Afri-

Colombia. [ca.

England, Holland.

S. France.

India, Mesopotamia.

India.

Transvaal.

Transvaal, Natal.

Cape Colony.

Transvaal.

Transvaal.

Sicily.

France, Holland,

Germany.

Carinthia.

Switzerland, Tyrol, Cau-

Switzerland. [casus.

Hungary.

Croatia.

Silesia, Hungary.

S. Russia.

Germany.

S. France.

France.

S. France.

S. France.

Corsica.

C. & S. Europe.

Germany, Austria.

S. France, Corsica, Alge-

Dalmatia, Algeria. [ria.

Spain, Algeria.

Germany, Switzerland,

Hungary.

Austria, S. Russia.

Algeria.

S. France.

S. Russia.

Spain, S. France, Sicily.

S. France.

Sicily.

Sicily.

Germany.

130. *P. deserticolella*, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 305 (1870). S. Russia.
131. *P. trochilella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 253 (1870). Germany.
132. *P. grisella*, Chambers, Canad. Ent. Vol. 4, p. 171 (1872). Kentucky.
133. *P. gregariella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 339, pl. 4, f. 109 (?) Colombia.
(1877).
134. *P. altisona*, Meyrick, Trans. Ent. Soc. Lond. p. 46 (1917). Peru.
135. *P. hippelis*, Meyrick, ibidem, p. 573 (1901). New Zealand.
136. *P. plemochoa*, Meyrick, Trans. N. Zeal. Inst. Vol. 48, p. 415 (1916). New Zealand.
137. *P. glaucoptera*, Meyrick, ibidem, Vol. 43, p. 63 (1911). New Zealand.
138. *P. heterospora*, Meyrick, ibidem, Vol. 55, p. 204 (1924). New Zealand.
139. *P. cheradias*, Meyrick, ibidem, Vol. 41, p. 12 (1909). New Zealand.
140. *P. diffuella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 247 (1870). Alps.
141. *P. cacuminum*, Frey, Mitth. Schweiz. Ent. Ges. Vol. 3, p. 252 (1870). Switzerland.
culminicolella, Staudinger, Berl. Ent. Zeitschr. Vol. 14, p. 306 (1870).
142. *P. excelsa*, Frey, Lep. Schweiz. p. 363 (1880). Switzerland.
143. *P. murinella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 178, f. 535 (1855). C. Europe.
144. *P. nocturnella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 241 (1859). Spain.
145. *P. montanella*, Chrétien, Naturaliste, Vol. 32, p. 272 (1910). S. France.
146. *P. suasella*, Constant, Bull. Soc. Ent. Fr. p. 53 (1895). S. France.
147. *P. rancidella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 176, f. 534 (1855). Germany, Austria.
pauperella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 256 (1870).
? luridella, Teich, Stett. Ent. Zeit. Vol. 47, p. 170 (1886). Galicia.
148. *P. chrysanthemella*, Hofmann, Stett. Ent. Zeit. Vol. 28, p. 202 (1867). Germany.
149. *P. opificella*, Mann, Verh. Zool.-bot. Ges. Wien, Vol. 27, p. 499 (1877). Austria.
150. *P. sabulosella*, Rebel, Ann. Hofmus. Wien, Vol. 20, p. 23 (1905). Asia Minor.
151. *P. halonella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 178 (1855). Germany, Austria.
nigripalpella, Heinemann, ibidem, Vol. 2, p. 258 (1870). Galicia.
152. *P. acuminatella*, Sircom, Zoologist, App. p. 72 (1850). C. Europe, Asia Minor.
? gracilella, Stainton, Ent. Annual, p. 97 (1871).
153. *P. voltinella*, Chrétien, Naturaliste, p. 178 (1898). France.
154. *P. melanella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 258 (1870). Bavaria.
nigripalpella, Heinemann, ibidem, Vol. 2, p. 258 (1870).
155. *P. pygmaeella*, Heinemann, ibidem, Vol. 2, p. 259 (1870). Switzerland, Tyrol.
156. *P. exacta*, Meyrick, Trans. Ent. Soc. Lond. p. 46 (1917). Guiana.
157. *P. epitricha*, Meyrick, ibidem, p. 47 (1917). Guiana.
158. *P. involuta*, Meyrick, ibidem, p. 47 (1917). Guiana.
159. *P. ingloriella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 259 (1870). E. C. Europe, Corsica.
160. *P. brahmiella*, Heyden, Stett. Ent. Zeit. Vol. 23, p. 175 (1862). Germany, S. France.
161. *P. moritzella*, Treitschke, Schmett. Eur. Vol. 10 (3), p. 214 (1835). Germany, Switzerland,
162. *P. roseella*, Zetterstedt, Ins. Lapp. p. 1005 (1840). Norway. [Galicia.]
163. *P. laceratella*, Zeller, Stett. Ent. Zeit. Vol. 29, p. 143 (1868). Carinthia.
164. *P. Huebneri*, Haworth, Lep. Brit. p. 551 (1828). C. Europe.
kroesmanniella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 166, f. 581 (1855).
165. *P. knaggsiella*, Stainton, Ent. Annual, p. 167 (1866). England, Germany.
junctella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 184, f. 594 (1855..
166. *P. maculea*, Haworth, Lep. Brit. p. 552 (1828). C. Europe.
nivella, Wood, Index Ent. f. 1224 (1839).
blandella, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 2, p. 77, pl. 10, f. 2 (1852).
167. *P. blandulella*, Tutt, Ent. M. Mag. Vol. 24, p. 105 (1887). England.
168. *P. fraternella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 101 (1850). England.
169. *P. laudatella*, Walsingham, Proc. U. S. Mus. Vol. 33, p. 197 (1907). California.
170. *P. viscarienda*, Stainton, Ent. Annual, p. 43 (1855). England.

171. *P. tricolorella*, Haworth, Trans. Ent. Soc. Lond. p. 338 (1812). N. & C. Europe.
contigua, Haworth, Lep. Brit. p. 552 (1828).
acerella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 185, f. 580 (1855).
172. *P. jaspidella*, Chrétien, Bull. Soc. Ent. Fr. p. 231 (1908). Algeria. [Greece.]
173. *P. maculifera*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 102 (1850). C. Europe, Sardinia,
proxima, Haworth, Lep. Brit. p. 552 (1828) (non Hübner).
horticola, Peyerimhoff, Mith. Schweiz. Ent. Ges. Vol. 3, p. 411 (1871).
174. *P. semidecandra*, Threlfall, Ent. M. Mag. Vol. 23, p. 233 (1887). England, Holland,
maculifera, Stainton, Nat. Hist. Tin. Vol. 10, p. 154, pl. 12, f. 1 (1867). Germany.
175. *P. Rougemonti*, Rebel, Iris, Vol. 20, p. 236 (1906). Switzerland.
176. *P. pullatella*, Tengström, Not. Sällsk. Faun. Fenn. Förh. p. 126 (1847). Finland.
177. *P. junctella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 103 (1850). N. & C. Europe.
vicinella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 184, f. 471 (1855).
marmorea, Herrich-Schäffer, ibidem, Vol. 5, f. 592 (1855).
178. *P. sciurella*, Walsingham, Proc. Zool. Soc. Lond. p. 941, pl. 51, f. 14 Madeira, Canaries.
(1907).
179. *P. provinciella*, Stainton, Tin. S. Eur. p. 221 (1869). S. France.
180. *P. marmorea*, Haworth, Lep. Brit. p. 553 (1828). C. Europe.
181. *P. rubidella*, Chrétien, Bull. Mus. Hist. Nat. Paris, Vol. 14, p. 361 Canaries.
(1908).
182. *P. pulchra*, Wollaston, Ann. Mag. Nat. Hist. (3), Vol. 1, p. 121 (1858). Madeira.
183. *P. crepusculella*, Teich, Balt. Lep. Faun. p. 111 (1889). Latvia.
184. *P. capsophilella*, Chrétien, Bull. Soc. Ent. Fr. p. 223 (1900). France.
185. *P. behenella*, Constant, ibidem, p. 125 (1889). France. [Austria.]
186. *P. alsinella*, Zeller, Stett. Ent. Zeit. Vol. 29, p. 145 (1868). Latvia, Yugoslavia.
albifrontella, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 266 (1870).
? *tristella*, Heinemann, ibidem, Vol. 2, p. 267 (1870).
? *livoniella*, Teich, Arb. Nat. Ver. Riga, Vol. 41, p. 88 (1888).
187. *P. kiningerella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 186, f. 471 E. Europe.
(1855).
188. *P. fischerella*, Treitschke, Schmett. Eur. Vol. 9 (2), p. 84 (1833). E. Europe.
189. *P. Schleichi*, Christoph, Hor. Soc. Ent. Ross. Vol. 9, p. 22, pl. 1, f. 19 S. Russia.
(1872).
190. *P. tachyptilella*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 66, p. (44) (1916). Bulgaria.
191. *P. melanotephrella*, Erschoff, Hor. Soc. Ent. Ross. Vol. 12, p. 345 (1877). E. Siberia.
192. *P. caulinella*, Schmid, Beil. Ent. Zeitschr. Vol. 7, p. 63 (1863). C. Europe.
193. *P. saginella*, Zeller, Stett. Ent. Zeit. Vol. 29, p. 146 (1868). Carinthia.
194. *P. repentella*, Chrétien, Naturaliste, p. 258 (1908). France, Spain.
195. *P. mucronatella*, Chrétien, Bull. Soc. Ent. Fr. p. 138 (1900). France.
196. *P. gypsophilae*, Stainton, Tin. S. Eur. p. 210 (1869). S. France.
197. *P. trinella*, Fuchs, Stett. Ent. Zeit. Vol. 64, p. 9 (1903). Armenia.
198. *P. Petryi*, Hoffmann, ibidem, Vol. 60, p. 139 (1899). Germany.
199. *P. lakatensis*, Rebel, Ann. Hofmus. Wien, Vol. 19, p. 351, pl. 5, f. 24 Herzegovina.
(1904).
200. *P. coussonella*, Chrétien, Naturaliste, p. 245 (1908). France.
201. *P. Baueri*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 67, p. (33) (1917). Germany.
202. *P. inflatella*, Chrétien, Naturaliste, p. 17 (1901). France.
203. *P. vicinella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 102 (1850). C. Europe.
204. *P. leucomelanella*, Zeller, Isis, p. 198 (1839). N. & C. Europe, Dalmatia.
205. *P. tischeriella*, Zeller, Isis, p. 199 (1839). C. Europe. [Greece.]
206. *P. delphinatella*, Constant, Bull. Soc. Ent. Fr. p. 125 (1889). France.
207. *P. sestertiella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 186, f. 487 Norway, Germany, Austria, Greece.
(1855).
208. *P. pseudolella*, Christoph, Hor. Soc. Ent. Ross. Vol. 22, p. 313 (1888). S. Russia.

98. GENUS POGOCHAETIA, STAUDINGER

Pogochaetia, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 310 (1880). — Type : *P. solitaria*, Staudinger.

Characters. — Head smooth; tongue short. Antennae 4/5, basal joint elongate, without pecten. Labial palpi long, recurved, second joint beneath with rough projecting scales, with 6 or 7 long laterally projecting bristles, terminal joint as long as second, slender, acute. Maxillary palpi very short, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to costa. Hindwings somewhat over 1, trapezoidal, apex pointed, termen sinuate; 3 and 4 connate, 5 somewhat approximated, 6 and 7 subparallel.

Remarks. — A derivative from the Caryophyllaceous-feeding group of *Phthorimaea*, only differing from it by the curious character of the lateral bristles of the labial palpi, which is a feature quite abnormal in this family.

Geographical distribution of species. — Mediterranean.

Larva (both known) feeding in flowers and seeds.

Foodplants *Caryophyllaceae*.

- | | |
|---|-------------|
| 1. <i>P. solitaria</i> , Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 310 (1880). | Asia Minor. |
| 2. <i>P. ocyroidella</i> , Walsingham, Ent. M. Mag. Vol. 36, p. 218 (1900). | S. France. |

99. GENUS SAROTORNA, MEYRICK

Sarotorna, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 322 (1904). — Type : *S. eridora*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, basal joint moderately elongate, without pecten. Labial palpi rather long, recurved, second joint considerably dilated with scales roughly projecting anteriorly beneath, almost tufted, laterally compressed, terminal joint as long as second or much shorter, thickened with scales rather rough anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 from near angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex produced, round-pointed, termen sinuate, cilia 1; 3 and 4 connate, 5 parallel, 6 and 7 remote, nearly parallel.

Remarks. — Apparently related to *Phthorimaea* and *Gnorimoschema*, yet with palpi quite different from both, and also peculiar in superficial appearance.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|---|------------------|
| 1. <i>S. eridora</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 323 (1904). — | New South Wales. |
|---|------------------|

Pl. 2, Fig. 37.

- | | |
|--|-------------|
| 2. <i>S. myrrhina</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 126 (1919). — | Queensland. |
|--|-------------|

100. GENUS NOTHRIS, HÜBNER

Nothris, Hübner, Verz. bek. Schmett. p. 411 (1826). — Type : *N. verbascella*, Schiffermüller.

Oeselis, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 255 (1875). — Type : *N. sabinella*, Zeller.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, forming a compact rough tuft at apex beneath, terminal joint longer than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 3 and 4 approximated or connate, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 2/3; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 stalked.

Remarks. — Correlated with early forms of *Gelechia*, which is its nearest ally. This genus has been much misapprehended, partly through the inclusion of heterogeneous material, partly through the erroneous assumption that it was close to *Dichomeris*, with which there is in fact no near relationship.

Geographical distribution of species. — Essentially European, but extending into Northwest America, Siberia, and India.

Larva (6 known) feeding in spun shoots.

Foodplants : *Scrophulariaceae* (2), *Coniferae* (2), *Salicaceae*, *Rhamnaceae*.

1. <i>N. bilbainella</i> , Rössler, Stett. Ent. Zeit. Vol. 38, p. 378 (1877).	Spain.
2. <i>N. lemniscella</i> , Zeller, Isis, p. 190 (1839).	Germany, Austria, Hun-
3. <i>N. hastata</i> , Meyrick, Exot. Micolep. Vol. 2, p. 152 (1918).	Bengal. [gary.
4. <i>N. monella</i> , Busck, Proc. U. S. Mus. Vol. 27, p. 759 (1904).	Brit. Columbia.
5. <i>N. anarsiella</i> , Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 126 (1877).	Colorado.
6. <i>N. notandella</i> , Busck, Proc. Ent. Soc. Wash. Vol. 18, p. 149 (1916).	California.
7. <i>N. sabinella</i> , Zeller, Isis, p. 190 (1839).	C. Europe, Asia Minor,
<i>biannullella</i> , Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 225 (1875).	Colorado.
<i>ocellella</i> , Chambers, Bull. U. S. Geol. Surv. Vol. 3, p. 126 (1877).	
8. <i>N. senticetella</i> , Staudinger, Stett. Ent. Zeit. Vol. 20, p. 238 (1859).	S. Europe, Asia Minor.
9. <i>N. chinganella</i> , Christoph, Bull. Soc. Nat. Mosc. p. 32 (1882).	E. Siberia, Punjab, Asia
10. <i>N. hoffmanniella</i> , Strand, Norsk. Selsk. Skr. no 8, p. 4 (1901).	Norway. [Minor.
11. <i>N. obscuripennis</i> , Frey, Lep. Schweiz. p. 372 (1880).	Switzerland.
12. <i>N. asinella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 166 (1796).	C. Europe, Corsica.
<i>aurorella</i> , Frey, Mitt. Schweiz. Ent. Ges. Vol. 6, p. 365 (1882).	
13. <i>N. discretella</i> , Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 39, p. 318, pl. 8, f. 14 (1889).	Styria.
14. <i>N. congressariella</i> , Bruand, Ann. Soc. Ent. Fr. p. 171, pl. 11, f. 7 (1858). <i>declaratella</i> , Staudinger, Stett. Ent. Zeit. Vol. 20, p. 238 (1859).	S. France, Spain.
15. <i>N. sulcella</i> , Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 328 (1880).	Asia Minor.
16. <i>N. verbascella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 98 (1796).	Europe, Asia Minor,
	Algeria, Turkestan.

101. GENUS ANISOPLACA, MEYRICK

Anisoplaca, Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 171 (1886). — Type : *A. phyoptera*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ moderately ciliated, basal joint elongate, rather stout, without pecten. Labial palpi long, recurved, second joint with rough projecting scales beneath, sometimes prominent below apex, terminal joint nearly as long as second or longer, stout, sometimes roughened anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae roughly haired above. Forewings with 1b furcate, 2 from about 3/4, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen rounded, cilia 2/5; 3 and 4 connate or short-stalked, 5 nearly parallel to 4, 6 and 7 approximated and parallel on basal half, then diverging.

Remarks. — Though varying considerably in the development of the rough scaling of palpi, these species are naturally associated together. The genus represents an archaic form of this group, probably indicating the origin of *Nothris*, *Gelechia* and *Platyedra*.

Geographical distribution of species. — Represented by scattered species through the Southern hemisphere, presumably remnants of a larger number.

Larva unknown.

1. <i>A. achyrota</i> , Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 170 (1886).	New Zealand.
2. <i>A. acrodactyla</i> , Meyrick, ibidem, Vol. 39, p. 118 (1907).	New Zealand.
3. <i>A. pycnoides</i> , Meyrick, Ann. Transv. Mus. Vol. 2, p. 17, pl. 5, f. 9 (1911).	Transvaal.
4. <i>A. viatrix</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 165 (1921).	Java.
5. <i>A. ptyoptera</i> , Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 171 (1886).	New Zealand.
6. <i>A. praesignis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 175 (1913).	Peru.

Group 4 (Anacampsis type)

In this group the labial palpi have the second joint characteristically smooth beneath, and those forms which in other respects are most like *Gelechia* and its allies can always be separated from them by this character; but there are also exceptional genera (as *Sennostoma* and *Ethirostoma*) in which the palpi have the second joint tufted or roughly haired beneath. The singular fine black and white transverse ribbing of this joint displayed in many species of *Compsolechia* and certain other genera is confined to this group. Instead of the dark rings of the previous group, the terminal joint of palpi tends to be marked with longitudinal lines. In the forewings the apex shows a general tendency to be hooked or falcate, sometimes very pronounced; vein 2 is usually separate and remote, but in the earlier forms sometimes stalked with 3; and vein 7 usually runs to costa when present, but in the earlier forms sometimes to apex, and is rather frequently absent (coincident with 8). In the hindwings veins 3 and 4 are normally connate, 6 and 7 connate or stalked; and rather frequently there is a cubital pecten (fringe of hairs on lower margin of cell towards base) which is not found elsewhere in the family except in the *Dichomeris* group, and is undoubtedly evidence of phylogenetic relationship with that group, supported by the other structures noted above as primitive in the group. This group contains the most ornamental and highly decorated members of the family, many of the tropical species being adorned with bright-coloured or metallic markings; these probably fly in sunshine. The greatest development of the group is in South America, which abounds in genera and species; there is also a marked but smaller development in the Indian region; elsewhere the group is not prominent.

102. GENUS THYRSOSTOMA, MEYRICK

Thrysostoma, Meyrick, Journ. Bomb. Nat. Hist. Soc. Vol. 17, p. 736 (1907). — Type: *T. glaucitis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae $\frac{3}{4}$, in ♂ moderately or strongly ciliated, basal joint very long, without pecten, second joint also often enlarged and elongate. Labial palpi very long, recurved, smooth, second joint in ♂ with expanded fringe or long expandible tuft of hairs enclosed in a furrow beneath, terminal joint as long as second or longer, sometimes in ♂ thickened, more or less pointed, in ♀ slender, acute. Maxillary palpi rudimentary. Posterior tibiae with three expanded whorls of stiff scales. Forewings with 1b furcate, 2 from towards

angle, 4 absent, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings 1/2, very narrowly elongate-trapezoidal, acutely pointed, cilia 4; 2-4 parallel, 5 approximated, 6 and 7 long-stalked.

Remarks. — A development of *Thiotricha*. The species are very similar, but good distinguishing characters are found in the secondary sexual structures of the male palpi and antennae. In repose the posterior legs are thrust forwards beneath the wings and erected on each side of the shoulders. This singular habit at once suggests *Stathmopoda* (*Heliodinidae*), to which there is also an extraordinary resemblance, not only in superficial appearance but in many details of structure, and hence Stainton was led to place the European *Guerini* actually in *Stathmopoda* (where it still remains in Staudinger's Catalogue), though he correctly noted and emphasized the different form of hindwings, but without recognising their Gelechiad type. No adequate explanation of these curious analogies in abnormal characters can be given, but it is possible that local research in India would show that there is a third (non-Lepidopterous) group of insects, of which both these genera are mimics.

Geographical distribution of species. — Indo-Malayan, with one species on the Mediterranean.

Larva (2 known) feeding in Aphid-galls on leaves.

Foodplants *Anacardiaceae*.

- | | |
|---|----------------|
| 1. <i>T. chelophora</i> , Meyrick, Exot. Microlep. Vol. 2, p. 121 (1918). | Assam. |
| 2. <i>T. tanyrrhina</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 162 (1921). | Java. |
| 3. <i>T. glaucitis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 736 (1907). | India, Ceylon. |

— Pl. 2, Fig. 38.

- | | |
|--|------------|
| 4. <i>T. Guerini</i> , Stainton, Ent. Annual, p. 152, f. 5 (1858). | S. Europe. |
| 5. <i>T. oxyprora</i> , Meyrick, Exot. Microlep. Vol. 2, p. 501 (1922). | China. |
| 6. <i>T. diplobathra</i> , Meyrick, ibidem, Vol. 2, p. 120 (1918). | Assam. |
| 7. <i>T. fissilis</i> , Meyrick, ibidem, Vol. 2, p. 121 (1918). | Assam. |
| 8. <i>T. macrodelta</i> , Meyrick, ibidem, Vol. 2, p. 121 (1918). | Assam. |
| 9. <i>T. pylartis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 441 (1908). | Assam. |

103. GENUS HIERANGELA, MEYRICK

Hierangela, Meyrick, Trans. Ent. Soc. Lond. p. 14 (1894). — Type : *H. erythrogramma*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ serrulate, simple, basal joint slender, without pecten. Labial palpi very long, recurved, smooth, slender, second joint somewhat thickened, terminal joint longer than second, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs. Forewings with 1b fuscate, 2 from towards angle, 7 absent, 8 and 9 stalked, 10 approximated, 11 from beyond middle. Hindwings under 1, narrow-trapezoidal, apex strongly produced, acute, termen emarginate, cilia 3; 3 and 4 connate, 5 rather approximated, 6 and 7 connate.

Remarks. — Correlated with *Thiotricha*. An elegant insect, with crimson and yellow colouring.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

1. *H. erythrogramma*, Meyrick, Trans. Ent. Soc. Lond. p. 15 (1894). — Pl. 2, Burma.
Fig. 42.

104. GENUS SYMPHANACTIS, NOV. GEN.

Type : *S. hetaera*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ rather thick, simple, basal joint elongate, rather swollen towards apex, without pecten. Labial palpi long, curved, ascending, second joint somewhat thickened towards apex, terminal joint longer than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1^b furcate, 2 from towards angle, 3 and 5 connate from angle, 4 absent, 6 and 7 out of 8, 7 to costa, 11 from middle. Hindwings 2/3, narrow-trapezoidal, apex long-produced, acute, termen sinuate-emarginate, cilia 4; 3 and 4 connate, 5 curved, rather approximated to 4, 6 and 7 stalked.

Remarks. — Correlated with *Thiotricha*.

Geographical distribution of species. — South American.

Larva unknown.

1. *S. hetaera*, Meyrick, Trans. Ent. Soc. Lond. p. 231 (1914). — **Pl. I, Fig. 17.** Guiana.

105. GENUS THIOTRICA, MEYRICK

Thiotricha, Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 164 (1886). — Type : *T. thorybodes*, Meyrick.

Reutlla, Hofmann, Iris, Vol. 10, p. 228 (1897). — Type : *T. subocellea*, Stephens.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrate, with long fine ciliations (3-6), basal joint elongate, without pecten. Labial palpi long, recurved, smooth-scaled, second joint hardly thickened, terminal joint as long as second or longer, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1^b furcate, 2 from towards angle, 4 absent, 6 out of 8 or separate, 7 absent, 9 approximated or out of 8, 11 from beyond middle. Hindwings 1 or under 1, elongate-trapezoidal, apex more or less produced, pointed, termen emarginate, cilia 1 1/2-6; 3 and 4 connate, 5 bent, approximated, 6 and 7 stalked.

Remarks. — The long antennal ciliations of ♂ are a constant feature, very unusual in this family. The species are often elegantly marked, sometimes with orange or rosy colouring. The genus is correlated with *Polyhymno*.

Geographical distribution of species. — Characteristically Indo-Malayan, with a considerable Australian section (mainly Queensland), and scattered elements in Europe, Africa, South America and New Zealand.

Larva (3 known) always feeding in a portable case (very exceptional in this family) on flowers, seeds, or leaves.

Foodplants : *Labiatae*, *Proteaceae*, *Oleaceae*.

- | | |
|---|-------------------------------|
| 1. <i>T. animosella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1022 (1864).
<i>acrocelea</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 116 (1919). | India, Ceylon.
Queensland. |
| 2. <i>T. xanthaspis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 122 (1918). | |
| Assam. | |
| 3. <i>T. glenias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 439 (1908). — Pl. 2, Fig. 40. | |
| Ceylon. | |

4. *T. prososteia*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 116 (1919).
 5. *T. pyrphora*, Meyrick, Exot. Microlep. Vol. 2, p. 122 (1918).
 6. *T. Godmani*, Walsingham, Proc. Zool. Soc. Lond. p. 525 (1891).
 7. *T. sciurella*, Walsingham, ibidem, p. 78 (1897).
 argoxantha, Meyrick, Trans. Ent. Soc. Lond. p. 237 (1914).
 8. *T. laterestriata*, Walsingham, Proc. Zool. Soc. Lond. p. 78 (1897).
 9. *T. cleodrella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 355, pl. 5, f. 120
 (1877).
 10. *T. centritis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 439 (1908).
 11. *T. janitrix*, Meyrick, Exot. Microlep. Vol. 1, p. 64 (1912).
 12. *T. atractodes*, Meyrick, ibidem, Vol. 2, p. 502 (1922).
 13. *T. oxytheces*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 293
 (1904).
 14. *T. panglycera*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 117 (1919).
 15. *T. epiclista*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 440 (1908).
 16. *T. complicata*, Meyrick, Exot. Microlep. Vol. 2, p. 122 (1918).
 17. *T. clidias*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 439 (1908).
 18. *T. chrysopa*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 293
 (1904).
 19. *T. saulotis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 138 (1906).
 20. *T. clinopeda*, Meyrick, Exot. Microlep. Vol. 2, p. 124 (1918).
 21. *T. tenuis*, Walsingham, Trans. Ent. Soc. Lond. p. 96, pl. 4, f. 33 (1891).
 22. *T. paltobola*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 75 (1921).
 23. *T. cuneiformis*, Meyrick, Exot. Microlep. Vol. 2, p. 124 (1918).
 24. *T. acrantha*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 440
 (1908).
 25. *T. leucothona*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 294
 (1904).
 26. *T. margarodes*, Meyrick, ibidem, Vol. 29, p. 294 (1904).
 27. *T. pteropis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 443 (1908).
 28. *T. paraconta*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 295
 (1904).
 29. *T. hamulata*, Meyrick, Zool. Med. Leid. Vol. 6, p. 162 (1921).
 30. *T. balanopa*, Meyrick, Exot. Microlep. Vol. 2, p. 123 (1918).
 31. *T. coleella*, Constant, Ann. Soc. Ent. Fr. p. 255, pl. 10, f. 16 (1884).
 32. *T. subocellea*, Stephens, Ill. Brit. Ent. Haust. Vol. 4, p. 214 (1835).
 internella, Zeller, Isis, p. 291 (1846).
 dissimella, Herrich Schäffer, Schmett. Eur. Vol. 5, p. 200, f. 553 (1855).
 33. *T. majorella*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 60, p. (28) (1910).
 34. *T. delacma*, Meyrick, Exot. Microlep. Vol. 3, p. 24 (1923).
 35. *T. galactaea*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 441
 (1908).
 36. *T. hoplomacha*, Meyrick, ibidem, Vol. 18, p. 441 (1908).
 37. *T. acronipha*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 118 (1919).
 38. *T. argyrea*, Turner, ibidem, Vol. 31, p. 118 (1919).
 39. *T. hemiphaea*, Turner, ibidem, Vol. 31, p. 118 (1919).
 40. *T. rhodopa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 442
 (1908).
 41. *T. chrysantha*, Meyrick, ibidem, Vol. 18, p. 443 (1908).
 42. *T. oxygramma*, Meyrick, Exot. Microlep. Vol. 2, p. 123 (1918).
 43. *T. characias*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 440
 (1908).
 44. *T. galenaea*, Meyrick, ibidem, Vol. 18, p. 443 (1908).
 45. *T. grammatis*, Meyrick, ibidem, Vol. 18, p. 442 (1908).
 Queensland.
 Coorg. [ti.]
 Windward Islands, Hay-
 Mexico, Virgin Islands,
 Guiana, Brazil, Peru.
 Virgin Islands.
 Colombia.
 S. India.
 Coorg, Bengal.
 Queensland.
 New South Wales,
 Queensland.
 Queensland.
 Assam.
 Coorg, Queensland.
 Assam, Ceylon.
 Queensland.
 Ceylon.
 Coorg, Ceylon.
 Gambia, Seychelles.
 Transvaal.
 Coorg.
 Assam.
 New South Wales.
 Queensland.
 Assam, Ceylon.
 New South Wales.
 Java.
 Assam, Borneo.
 S. France.
 C. & S. Europe.
 Herzegovina.
 S. India.
 S. India.
 Assam.
 Queensland.
 Queensland.
 Queensland.
 Assam.
 Assam.
 Assam.
 S. India.
 Ceylon.
 Assam.

46. <i>T. synacma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 124 (1918).	Assam.
47. <i>T. polyaula</i> , Meyrick, ibidem, Vol. 2, p. 125 (1918). — PI. 2, Fig. 41.	Assam.
48. <i>T. rhodomicta</i> , Meyrick, ibidem, Vol. 2, p. 126 (1918).	Assam.
49. <i>T. obvoluta</i> , Meyrick, ibidem, Vol. 2, p. 126 (1918).	Assam.
50. <i>T. scioplecta</i> , Meyrick, ibidem, Vol. 2, p. 123 (1918).	Assam.
51. <i>T. pancratiasitis</i> , Meyrick, ibidem, Vol. 2, p. 426 (1921).	Assam.
52. <i>T. nephodesma</i> , Meyrick, ibidem, Vol. 2, p. 125 (1918).	Assam.
53. <i>T. operaria</i> , Meyrick, ibidem, Vol. 2, p. 125 (1918).	Assam.
54. <i>T. xanthodora</i> , Meyrick, ibidem, Vol. 3, p. 24 (1921).	Burma.
55. <i>T. scotaea</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 442 (1908).	Ceylon.
56. <i>T. orthiastis</i> , Meyrick, ibidem, Vol. 16, p. 591 (1905).	Punjab.
57. <i>T. parthenica</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 297 (1904).	S.E. Australia, Tasmania.
58. <i>T. arthrodies</i> , Meyrick, ibidem, Vol. 29, p. 295 (1904).	New South Wales.
59. <i>T. bullata</i> , Meyrick, ibidem, Vol. 29, p. 296 (1904).	New South Wales.
60. <i>T. niphastis</i> , Meyrick, ibidem, Vol. 29, p. 296 (1904).	West Australia.
61. <i>T. tetraphala</i> , Meyrick, Trans. N. Zeal. Inst. Vol. 18, p. 164 (1886).	New Zealand.
62. <i>T. thorybodes</i> , Meyrick, ibidem, Vol. 18, p. 164 (1886).	New Zealand.
63. <i>T. anticentra</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 296 (1904).	Queensland.

106. GENUS SEMNOSTOMA, MEYRICK

Semnostama, Meyrick, Exot. Microlep. Vol. 2, p. 127 (1918). — Type : *S. leucochalca*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 5/6, in ♂ simple or moderately ciliated, basal joint very elongate, slender, without pecten. Labial palpi very long, recurved, second joint thickened with scales roughly expanded at apex above and forming a rough projecting apical tuft beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with 2 from towards angle, 7 absent, 8 and 9 short-stalked, 11 from middle. Hindwings 1, narrow-trapezoidal, apex acutely produced, termen sinuate, cilia 2/3; 2 remote, 3 and 4 nearly approximated from angle, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Correlated with *Thiotricha*, which it resembles in general appearance; the development of a tuft on palpi is exceptional in this group.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|---|--------|
| 1. <i>S. leucochalca</i> , Meyrick, Exot. Microlep. Vol. 2, p. 127 (1918). | Assam. |
| 2. <i>S. poecilopa</i> , Meyrick, ibidem, Vol. 2, p. 128 (1918). | Assam. |
| 3. <i>S. barathrota</i> , Meyrick, ibidem, Vol. 2, p. 128 (1918). — PI. 2, Fig. 39. | Assam. |
| 4. <i>S. scabrosa</i> , Meyrick, ibidem, Vol. 2, p. 128 (1918). | Assam. |

107. GENUS PLECTROCOSMA, MEYRICK

Plectrocosma, Meyrick, Ann. Transv. Mus. Vol. 8, p. 75 (1921). — Type : *P. centrophora*, Meyrick.

Characters. — Head smooth-scaled; ocelli posterior; tongue developed. Antennae 3/4, in ♂ moderately ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second

joint with appressed scales, terminal joint longer than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 from towards angle, 7 absent, 8-10 approximated, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex acute, produced, termen sinuate, cilia 2 1/2; 3 and 4 connate, 5 parallel, 6 and 7 stalked.

Remarks. — Allied to the preceding.

Geographical distribution of species. — South African.

Larva unknown.

1. *P. centrophora*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 75 (1921).

Transvaal.

108. GENUS CNAPHOSTOLA, MEYRICK

Cnaphostola, Meyrick, Exot. Microlep. Vol. 2, p. 131 (1918). — Type : *C. adamantina*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 6 and 8 stalked, 7 absent, 11 from beyond middle. Hindwings nearly 1, narrow-trapezoidal, apex produced, acute, termen sinuate, cilia 2; 2 remote, 3 from before angle, 4 from angle, 5 approximated, 6 and 7 stalked.

Remarks. — Also correlated with the two preceding.

Geographical distribution of species. — Indian.

Larva unknown.

1. *C. adamantina*, Meyrick, Exot. Microlep. Vol. 2, p. 132 (1918).

Assam.

109. GENUS CRAMBODOXA, MEYRICK

Crambodoxa, Meyrick, Trans. Ent. Soc. Lond. p. 174 (1913). — Type : *C. platyaula*, Meyrick.

Characters. — Head loosely rough-haired on crown; ocelli posterior; tongue short. Antennae 3/4, in ♂ moderately ciliated, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint rather short, thickened with scales, in ♂ with very long expandible projecting apical pencil of fine hairs above, terminal joint in ♂ much longer than second, thickened with appressed scales, hardly pointed. Maxillary palpi imperceptible. Posterior tibiae clothed with long fine hairs above. Forewings with 2 from 4/5, 3 from angle, 7 absent, 8 and 9 stalked, 11 from 3/4. Hindwings 1, elongate-trapezoidal, termen faintly sinuate beneath apex, cilia 1; 2 widely remote, 3-5 slightly approximated towards base, 6 and 7 parallel.

Remarks. — A genus of quite uncertain position, but possibly an aberrant member of this group, and at least equally abnormal elsewhere.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. platyaula*, Meyrick, Trans. Ent. Soc. Lond. p. 174 (1913). — PI. 2. Colombia.

Fig. 43.

110. GENUS POLYHYMNO, CHAMBERS

Polyhymno, Chambers, Canad. Ent. Vol. 6, p. 246 (1874). — Type : *P. luteostrigella*, Chambers.

Copocercia, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 374 (1877). — Type : *P. crambinella*, Zeller.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ shortly ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint smooth-scaled or seldom rough at apex beneath, terminal joint as long as second or longer, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with apex more or less produced and subfalcate; 1b furcate, 2 from angle, 7 and 8 very long-stalked or coincident, 7 (if present) to costa, 11 from beyond middle. Hindwings under 1, elongate-trapezoidal, apex produced, acute, termen sinuate-emarginate, cilia 2-3; 3 and 4 connate or short-stalked, 5 approximated, 6 and 7 long-stalked.

Remarks. — Structurally and superficially it would seem natural that this characteristic genus should be the ancestor of *Thiotricha* and its allies, yet the geographical distribution makes it highly improbable. It is likely therefore that all are alike descended from a more ancient extinct form, of which *Polyhymno* preserves the main features.

Geographical distribution of species. — Summarised thus : African 16, American 10, Indian 1. This is therefore one of the comparatively few genera (as *Tiquadra* and *Ceromitia*) exhibiting direct faunal connection between Africa and America.

Larva (2 known) feeding in spun leaves (?).

Foodplants *Leguminosae*.

1. <i>P. conficta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 51 (1917).	Peru.
2. <i>P. colleta</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 71 (1911).	Mexico.
3. <i>P. convergens</i> , Walsingham, ibidem, Vol. 4, p. 71, pl. 2, f. 27 (1911).	Mexico.
4. <i>P. subaequalis</i> , Walsingham, ibidem, Vol. 4, p. 70 (1911).	Mexico.
5. <i>P. acaciella</i> , Busck, Journ. N. York Ent. Soc. Vol. 8, p. 235, pl. 9, f. 1 (1900).	Texas.
6. <i>P. luteostrigella</i> , Chambers, Canad. Ent. Vol. 6, p. 247 (1874). <i>fuscostrigella</i> , Chambers, ibidem, Vol. 8, p. 30 (1876).	S. E. United States, Jamaica, Virgin Islands.
7. <i>P. crambinella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 375, pl. 5, f. 129 (1877).	Colombia.
8. <i>P. sexstrigella</i> , Chambers, Canad. Ent. Vol. 6, p. 248 (1874).	Texas.
9. <i>P. leucocras</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 71 (1911).	Mexico.
10. <i>P. gladiata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 51 (1917).	Colombia.
11. <i>P. cemistostomella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 357 (1877).	Zanzibar.
12. <i>P. tropaea</i> , Meyrick, Proc. Zool. Soc. Lond. p. 728 (1908).	Transvaal.
13. <i>P. hieracitis</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 283 (1913).	Transvaal.
14. <i>P. alcimacha</i> , Meyrick, Exot. Microlep. Vol. 2, p. 129 (1918).	S. India, Assam.
15. <i>P. palinorsa</i> , Meyrick, Ann. Transv. Mus. Vol. 2, p. 15, pl. 5, f. 4 (1911) — PI. 2, Fig. 44.	Transvaal.
16. <i>P. deuteraula</i> , Meyrick, ibidem, Vol. 4, p. 193	Transvaal.
17. <i>P. tetraphrapha</i> , Meyrick, ibidem, Vol. 3, p. 283 (1913).	Transvaal.
18. <i>P. oxystola</i> , Meyrick, ibidem, Vol. 3, p. 284 (1913).	Transvaal.
19. <i>P. cleodorella</i> , Walsingham, Trans. Ent. Soc. Lond. p. 95, pl. 4, f. 32 (1891).	Gambia.
20. <i>P. intorta</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 19 (1918).	Natal.

21. *P. multifida*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 4 (1917). Zululand.
 22. *P. pausimacha*, Meyrick, Ann. Transv. Mus. Vol. 2, p. 14, pl. 5, f. 3 Transvaal.
 (1911).
 23. *P. hostilis*, Meyrick, ibidem, Vol. 6, p. 19 (1918). Transvaal.
 24. *P. paracma*, Meyrick, ibidem, Vol. 2, p. 15, pl. 5, f. 5 (1911). Transvaal.
 25. *P. eurydoxa*, Meyrick, ibidem, Vol. 2, p. 15, pl. 5, f. 6 (1911). Transvaal.
 26. *P. chionarcha*, Meyrick, ibidem, Vol. 3, p. 282 (1913). Transvaal, Port. E. Afri-
 27. *P. inermis*, Meyrick, ibidem, Vol. 3, p. 284 (1913). Transvaal, Natal. [ca.

III. GENUS CALLIPRORA, MEYRICK

Calliprora, Meyrick, Trans. Ent. Soc. Lond. p. 242 (1914). — Type: *C. pentagramma*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint somewhat thickened with appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with appressed scales, with whorls of projecting scales on origin of spurs. Forewings with apex falcate; 1b furcate, 2 from 3/4, 7 absent, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, produced, termen concave beneath apex, cilia 1; 3 and 4 connate, 5 rather approximated, 6 and 7 connate.

Remarks. — Probably correlated with the preceding, from which it is distinguished by the falcate apex of forewings, and veins 6 and 7 of hindwings not long-stalked; it is always easily recognisable by the characteristic and elegant superficial appearance.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|--|-----------------------|
| 1. <i>C. erethistis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 70 (1922). | Peru. |
| 2. <i>C. peritura</i> , Meyrick, ibidem, p. 70 (1922). | Brazil. |
| 3. <i>C. rhodogramma</i> , Meyrick, ibidem, p. 67 (1922). | Brazil. |
| 4. <i>C. centrocrossa</i> , Meyrick, ibidem, p. 67 (1922). | Brazil. |
| 5. <i>C. tetraplecta</i> , Meyrick, ibidem, p. 68 (1922). | Peru. |
| 5. <i>C. pentagramma</i> , Meyrick, ibidem, p. 243 (1914). — Pl. 2, Fig. 48. | Guiana, Brazil, Peru. |
| 7. <i>C. trigramma</i> , Meyrick, ibidem, p. 243 (1914). | Guiana, Brazil, Peru. |
| 8. <i>C. eurydelta</i> , Meyrick, ibidem, p. 69 (1922). | Peru. |
| 9. <i>C. platyxipha</i> , Meyrick, ibidem, p. 69 (1922). | Brazil, Peru. |

112. GENUS SOPHRONIA, HÜBNER

Sophronia, Hübner, Verz. bek. Schmett. p. 407 (1826). — Type: *S. illustrata*, Hübner.

Characters. — Head with dense appressed scales; ocelli conspicuous, posterior; tongue developed. Antennae 2/3, in ♂ shortly ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with very long dense rough projecting tuft beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long fine hairs above. Forewings with apex prominent or subfalcate; 1b furcate, 2 from 5/6, 3 from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen sinuate beneath apex, cilia 4/5-1; without cubital pecten; 3 and 4 connate or stalked, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — This somewhat puzzling but natural genus is probably referable here, notwithstanding the strong tuft of palpi; certainly it has no near relation to the *Dichomeris* group, with which it has been usually associated.

Geographical distribution of species. — Especially characteristic of the European region, but extending into America. I have also an unpublished species from South Africa.

Larva (4 known) feeding between spun leaves.

Foodplants *Compositae*.

- | | |
|---|---------------------------|
| 1. <i>S. mediatrix</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 377, pl. 5, f. 130 (1877). | Colombia. |
| 2. <i>S. roseicrinella</i> , Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 87 (1909). | Texas. |
| 3. <i>S. primella</i> , Busck, ibidem, Vol. 8, p. 89 (1907). | Arizona, Colorado. |
| 4. <i>S. semicostella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 396 (1817).
<i>parenthesella</i> , Haworth, Lep. Brit. p. 540 (1828). | C. Europe, Spain. |
| 5. <i>S. alaicella</i> , Caradja, Iris, Vol. 34, p. 116 (1920). | W. Turkestan. |
| 6. <i>S. catharurga</i> , Meyrick, Exot. Microlep. Vol. 3, p. 34 (1923). | Palestine. |
| 7. <i>S. chilonella</i> , Treitschke, Schmett. Eur. Vol. 9 (2), p. 36 (1833). — Pl. 3, | S. C. & E. Europe. |
| Fig. 70. | |
| 8. <i>S. consanguinella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 156, f. 369 (1855). | E. C. Europe, Asia Minor. |
| 9. <i>S. acaudella</i> , Rebel, Ann. Hofmus. Wien, Vol. 18, p. 333, pl. 3, f. 16 (1903). | Bulgaria. |
| 10. <i>S. finitimella</i> , Rebel, ibidem, Vol. 20, p. 25 (1905). | Asia Minor. |
| 11. <i>S. sagittans</i> , Meyrick, Exot. Microlep. Vol. 3, p. 33 (1923). | Palestine. |
| 12. <i>S. cosmella</i> , Constant, Ann. Soc. Ent. Fr. p. 258, pl. 10, f. 19 (1884). | Corsica. |
| 13. <i>S. exustella</i> , Zeller, Isis, p. 820 (1847). | Spain, Italy. |
| 14. <i>S. curonella</i> , Standfuss, Stett. Ent. Zeit. Vol. 45, p. 193 (1884). | Italy. |
| 15. <i>S. humerella</i> , Schiffermüller, Syst. Verz. Schmett. Wien. p. 137 (1776). | Europe, Asia Minor. |
| 16. <i>S. sicariella</i> , Zeller, Isis, p. 189 (1839). | C. & E. Europe. |
| 17. <i>S. santolinae</i> , Staudinger, Stett. Ent. Zeit. Vol. 24, p. 270 (1863). | Spain. [Asia Minor. |
| 18. <i>S. illustrella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 158 (1796). | E. C. & S. E. Europe, |

113. GENUS ANASPHALTIS, NOV. GEN.

Type : *A. renigerella*, Zeller.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with dense appressed scales, forming a projecting apical tuft beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired beneath. Forewings with apex prominent, subfalcate; 1b furcate, 2 and 3 separate, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings somewhat over 1, trapezoidal, termen slightly sinuate, cilia 2/3; 3 and 4 connate, 5 somewhat approximated, 6 and 7 approximated towards base.

Remarks. — Doubtfully related to the preceding.

Geographical distribution of species. — European.

Larva feeding in spun leaves.

Foodplant *Melissa* (*Labiatae*).

- | | |
|---|--------------------|
| 1. <i>A. renigerella</i> , Zeller, Isis, p. 189 (1839). | C. & S. E. Europe. |
|---|--------------------|

114. GENUS IDIOPHANTIS, MEYRICK

Idiophantis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 298 (1904). — Type : *I. habrias*, Meyrick.

Colobodes, Meyrick, ibidem, Vol. 29, p. 297 (1904). — Type : *I. insomnis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple or shortly ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, sometimes expanded towards apex above, terminal joint as long as second or longer, slender or thickened with sometimes loose scales anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with termen deeply excavated beneath slender produced falcate apex, tornus often appearing as a strong prominent lobe; 1b furcate, 2 and 3 separate or stalked or coincident, 6 sometimes absent, 7 absent, 9 absent, 11 from middle. Hindwings 1 or over 1, elongate-trapezoidal, termen more or less sinuate, cilia 2/3-1 2/3; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 somewhat approximated.

Remarks. — A singular and interesting form of this group, with exaggerated wing-characters.

Geographical distribution of species. — Especially Indo-Malayan, but extending into the Australasian and African regions.

Larva (*chiridota*) feeding in galls or seed-capsules.

Foodplant *Eugenia* (Myrtaceae).

1. *I. habrias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 298 (1904). Queensland.
2. *I. chiridota*, Meyrick, Exot. Microlep. Vol. 1, p. 201 (1914). — Pl. 2, Ceylon, Java, Malay States, Fiji.
- Fig. 45.**
3. *I. soreuta*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 139 (1906). Ceylon.
4. *I. hemiphaea*, Meyrick, ibidem, Vol. 18, p. 149 (1907). Assam.
5. *I. butyraula*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 285 (1913). Transvaal.
6. *I. spectrata*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 272 (1911). Seychelles.
7. *I. chalcura*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 148 (1907). Assam.
8. *I. anisosticta*, Meyrick, Exot. Microlep. Vol. 1, p. 566 (1916). Ceylon, Burma.
9. *I. disparata*, Meyrick, ibidem, Vol. 3, p. 24 (1923). Fiji.
10. *I. insomnis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 297 (1904). New South Wales.
11. *I. paraptila*, Meyrick, Exot. Microlep. Vol. 1, p. 566 (1916). Ceylon.
12. *I. melanosacta*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 148 (1907). Coorg.
13. *I. carpotoma*, Meyrick, Exot. Microlep. Vol. 1, p. 567 (1916). S. India.
14. *I. croconota*, Meyrick, ibidem, Vol. 2, p. 129 (1918). Madagascar.
15. *I. discura*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 148 (1907). Ceylon.
16. *I. stoica*, Meyrick, ibidem, Vol. 18, p. 149 (1907). S. India.

115. GENUS PAURONEURA, TURNER

Pauroneura, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 120 (1919). — Type : *P. brachysticha*, Turner.

Characters. — Head smooth; ocelli minute, posterior; tongue developed. Antennae 5/6, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint smooth, terminal joint longer than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with rough

scales above. Forewings with 2 from 5/6, 4 absent, 3 and 5 approximated at base, 7 absent, 11 from middle. Hindwings over 1, subovate, apex obtuse, termen not sinuate, cilia 1/4; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 approximated at base.

Remarks. — Doubtless a derivative of *Chaliniastis*.

Geographical distribution of species. — Australian.

Larva unknown.

1. *P. brachysticha*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 121 (1919). Queensland.

116. GENUS CHALINIASTIS, MEYRICK

Chaliniastis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 301 (1904). — Type : *C. astrapaea*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint smooth, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough scales above. Forewings with 2 from near angle, 7 absent, 11 from middle. Hindwings considerably over 1, trapezoidal-ovate, apex obtuse, termen not sinuate, cilia 2/3; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 rather approximated.

Remarks. — An early form of this sub-group.

Geographical distribution of species. — Australian.

Larva unknown.

1. *C. astrapaea*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 302 (1904). Queensland.

117. GENUS HYPELICTIS, MEYRICK

Hypelictis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 600 (1905). — Type : *H. acrochlora*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with dense scales, somewhat rough beneath and with rough projecting scales above, terminal joint as long as second, variably thickened with scales more or less roughly projecting posteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 2 and 3 stalked, 6 sometimes to apex, 7 to costa, 8 out of 7 or absent, 11 from middle. Hindwings over 1, trapezoidal, apex obtuse, termen hardly sinuate, cilia 1/2; without cubital pecten; 3 and 4 connate, 5 parallel, 6 and 7 stalked.

Remarks. — Notwithstanding the peculiar but variable palpi, this genus is correctly referable here, and appears to be a collateral development from the presumed ancestral form of all the preceding genera of the group. The forewings have a tendency to be curiously crumpled at the apex, hinting at the potential origin of the falcate apex and excised termen of some of these.

Geographical distribution of species. — Indian.

Larva (1 known) feeding between spun leaves.

Foodplant *Salix*.

- | | |
|---|---------|
| 1. <i>H. frenigera</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 171 (1913). | Assam. |
| 2. <i>H. acrochlora</i> , Meyrick, ibidem, Vol. 16, p. 600 (1905). | Ceylon. |
| 3. <i>H. lupata</i> , Meyrick, ibidem, Vol. 22, p. 171 (1913). | Assam. |
| 4. <i>H. albiscricta</i> , Meyrick, ibidem, Vol. 22, p. 773 (1914). | Kanara. |
| 5. <i>H. charonaea</i> , Meyrick, ibidem, Vol. 22, p. 172 (1913). | Ceylon. |

118. GENUS SCINDALMOTA, TURNER

Scindalmota, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 121 (1919). — Type : *S. limata*, Turner.

Characters. — Head smooth; tongue developed. Antennae 3/4, in ♂ minutely ciliated, basal joint without pecten. Labial palpi long, recurved, second joint smooth-scaled, terminal joint nearly as long as second, slender, acute. Maxillary palpi very short. Forewings with 2 and 3 parallel, 6 and 7 out of 8, 6 to costa, Hindwings nearly 1 1/2, trapezoidal, apex acute, produced, termen strongly sinuate, 3 and 4 long-stalked, 5 parallel, 6 and 7 long-stalked.

Remarks. — I have not seen this genus, which is considered by its author to be allied to *Stomopteryx*; its true affinity seems very uncertain.

Geographical distribution of species. — Australian.

Larva unknown.

1. *S. limata*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 122 (1919). Queensland.

119. GENUS STIPHROSTOLA, MEYRICK

Stiphrostola, Meyrick, Exot. Microlep. Vol. 3, p. 25 (1923). — Type : *S. longinqua*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint shorter than second, moderate, pointed. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from angle, 2-5 approximated, 7 and 8 stalked, 7 to costa, 11 from beyond middle. Hindwings hardly 1, narrow-trapezoidal, apex produced, termen emarginate, cilia 2; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — A modification of *Stomopteryx*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *S. longinqua*, Meyrick, Exot. Microlep. Vol. 3, p. 25 (1923). Assam.

120. GENUS INOTICA, MEYRICK

Inotica, Meyrick, Exot. Microlep. Vol. 1, p. 65 (1913). — Type : *I. gaesata*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, slender, without pecten. Labial palpi very long,

recurved, second joint thickened with dense appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with dense long hairs. Forewings with 2 from $3/4$, 3 from before angle, 4 from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex long-produced, acute, termen emarginate, cilia 1; without cubital pecten; 3 and 4 approximated towards base, 5 rather approximated, 6 and 7 stalked.

Remarks. — Derived from an early form of *Stomopteryx*.

Geographical distribution of species. — Asia Minor.

Larva unknown.

1. *I. gaeata*, Meyrick, Exot. Microlep. Vol. 1, p. 66 (1913). — Pl. 2, Fig. 47. Asia Minor.

121. GENUS STOMOPTERYX, HEINEMANN

Stomopteryx, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 324 (1870). — Type : *S. detersella*, Zeller.

Aproaerema, Durrant, Ent. M. Mag. Vol. 33, p. 221 (1897). — Type : *S. anthyllidella*, Hübner.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint smooth-scaled, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 6 sometimes out of 7 near base, 11 from middle. Hindwings 1 or under 1, elongate-trapezoidal, apex pointed, produced, termen sinuate-emarginate, cilia 1 1/2-2; without cubital pecten; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — Derived from a form approaching *Melitoxestis*. The genus, as here limited, is quite natural, but has been subject to much misapprehension, and also to much confusion of nomenclature, the name *Anacampsis* being often wrongly applied to it.

Geographical distribution of species. — Perhaps Mediterranean in origin; about half the species are Palaearctic, the rest occurring in diminishing numbers in all the other regions, but barely represented in Australia and in New Zealand only by one introduced species. Certain species have been widely introduced with cultivated foodplants.

Larva (16 species known) feeding usually between spun leaves, seldom in stems or mining in leaves or fruits.

Foodplants usually *Leguminosae* (13), but also *Linaceae*, *Rubiaceae*, *Umbelliferae*.

- | | |
|--|---|
| 1. <i>S. oxyspila</i> , Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 351 (1909). | Transvaal, Cape Colony. |
| 2. <i>S. coracina</i> , Meyrick, Exot. Microlep. Vol. 2, p. 427 (1921). | Queensland. |
| 3. <i>S. simplicella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1024 (1864) (<i>simplexella</i>).
<i>isoscelixantha</i> , Lower, Proc. Linn. Soc. N. S. Wales, Vol. 22, p. 272 (1897). | India, China, E. Australia, New Zealand. |
| 4. <i>S. elachistella</i> , Stainton, Ann. Mag. Nat. Hist. (3) Vol. 3, p. 213 (1859). | Madeira, Canaries. |
| 5. <i>S. subsecivella</i> , Zeller, Micr. Caffr. p. 113 (1852).
<i>modicella</i> , Deventer, Tijdschr. v. Ent. Vol. 47, p. 4, pl. 1, f. 2 (1904).
<i>nerteria</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 139 (1906). | S. Africa, India, Ceylon, Malay States, Java. |
| 6. <i>S. anthyllidella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 330 (1817).
<i>nigritella</i> , Stainton, Ins. Brit. Tin. p. 133 (1854).
<i>sparsiciliella</i> , Barrett, Ent. M. Mag. Vol. 27, p. 7 (1891). | C. & S. Europe, Asia Minor, N. & S. Africa. |
| 7. <i>S. phaeopa</i> , Meyrick, Exot. Microlep. Vol. 2, p. 136 (1918). | Peru. |

LEPIDOPTERA HETEROCEA

8. *S. nigrella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 250 (1875).
 9. *S. palpilineella*, Chambers, ibidem, Vol. 2, p. 252 (1875).
 10. *S. crotalariella*, Busck, Proc. U. S. Mus. Vol. 23, p. 226 (1900).
 11. *S. vinella*, Banks, Ent. M. Mag. Vol. 34, p. 242 (1898).
 12. *S. albipalpella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 195 (1855).
leucopalpella, Herrich-Schäffer, ibidem, Vol. 5, f. 523 (1855).
 13. *S. submissella*, Frey, Lep. Schweiz. p. 367 (1880).
 14. *S. balcanica*, Rebel, Ann. Hofmus. Wien, Vol. 18, p. 330 (1903).
 15. *S. sanguella*, Stanton, Ent. Annual, p. 149 (1863).
 16. *S. splendens*, Staudinger, Hor. Soc. Ent. Ross. Vol. 16, p. 90 (1881).
 17. *S. patruella*, Mann, Wien. Ent. Monatsschr. Vol. 1, p. 180 (1857).
 18. *S. ignobiliella*, Heinemann, Schmett. Deutschl. (2) Vol. 2, p. 313 (1870).
 19. *S. fulvistillella*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 41, p. 632 (1891).
 20. *S. nigrinella*, Zeller, Isis, p. 857 (1847).
 21. *S. coronillella*, Treitschke, Schmett. Eur. Vol. 9 (2), p. 87 (1833).
 22. *S. biguttella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 192, f. 521 (1855).
 23. *S. psoralella*, Millière, Icon. Descr. Lép. Vol. 2, p. 83, pl. 61, f. 1-6 (1865).
infestella, Rebel, Ann. Hofmus. Wien, Vol. 11, p. 128 (1896).
linella, Chrétien, Naturaliste (2), Vol. 26, p. 151 (1904).
 24. *S. prolapsa*, Meyrick, Exot. Microlep. Vol. 2, p. 137 (1918).
 25. *S. rastrifera*, Meyrick, ibidem, Vol. 2, p. 137 (1918).
 26. *S. praecipitata*, Meyrick, ibidem, Vol. 2, p. 137 (1918).
 27. *S. azosterella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 194 (1855).
 28. *S. melagonella*, Constant, Bull. Soc. Ent. Fr. p. 53 (1895).
 29. *S. lachtenensis*, Erschoff, Hor. Soc. Ent. Ross. Vol. 12, p. 345 (1876).
 30. *S. embrocha*, Meyrick, Ann. Transv. Mus. Vol. 4, p. 192 (1914).
 31. *S. sarothamnella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 18, p. 615
 (1868).
? albifrontella, Heinemann, Schmett. Deutschl. (2) Vol. 2, p. 319 (1870).
 32. *S. ussuricella*, Caradja, Iris, Vol. 34, p. 108 (1920).
 33. *S. vorticella*, Scopoli, Ent. Carn. no 651 (1763).
? cinctella, Hübner, Samml. Vög. Schmett. f. 11 (1793).
ligulella, Zeller, Isis, p. 201 (1839).
? obliquella, Ragonot, Ann. Soc. Ent. Fr. p. 586 (1874).
 34. *S. cincticulella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 194, f. 527
 (1855).
 35. *S. taeniola*, Zeller, Isis, p. 201 (1839).
 36. *S. captivella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 194, f. 579 (1855).
 37. *S. genistae*, Walsingham, Proc. Zool. Soc. Lond. p. 933, pl. 51, f. 8
 (1907).
 38. *S. zonariella*, Walsingham, Ent. M. Mag. Vol. 41, p. 39 (1905).
 39. *S. mitrella*, Walsingham, ibidem, Vol. 41, p. 39 (1905).
 40. *S. acanthyllidis*, Walsingham, ibidem, Vol. 41, p. 40 (1905) (? — seq.).
 41. *S. polychromella*, Rebel, Iris, Vol. 15, p. 109 (1902). — **Pl. 2, Fig. 46.**
faceta, Meyrick, Ann. Transv. Mus. Vol. 4, p. 192 (1914).
 42. *S. biangulata*, Meyrick, ibidem, Vol. 8, p. 77 (1921).
 43. *S. exsulata*, Meyrick, ibidem, Vol. 6, p. 20 (1918).
 44. *S. elaeocoma*, Meyrick, ibidem, Vol. 6, p. 19 (1918).
 45. *S. maledicta*, Meyrick, Zool. Med. Leid. Vol. 6, p. 162 (1921).
 46. *S. inumbrata*, Meyrick, Ann. Transv. Mus. Vol. 4, p. 193 (1914).
 47. *S. oncodes*, Meyrick, ibidem, Vol. 3, p. 285 (1913).
 48. *S. circea*, Meyrick, ibidem, Vol. 3, p. 67 (1912).
 49. *S. bathrarcha*, Meyrick, ibidem, Vol. 8, p. 76 (1921).
- E. United States.
 E. United States.
 Florida.
 England.
 England, Belgium,
 Holland, Germany.
 Switzerland.
 Bulgaria.
 C. Europe, Asia Minor.
 Macedonia, Asia Minor.
 S. C. & S. E. Europe, Asia
 Minor.
 Germany, Austria.
 Dalmatia.
 Spain, Sicily, Dalmatia.
 Europe, Asia Minor.
 C. & S. Europe, Asia Mi-
 France, Spain, [nor.
 Canaries, Madeira.
 Ceylon.
 Ceylon.
 India.
 Silesia, Austria.
 S. France.
 N. Russia.
 Natal.
 Germany.
 E. Siberia.
 Europe, Asia Minor.
 E. C. & S. Europe.
 C. & S. E. Europe, Asia
 Minor.
 Hungary, Dalmatia.
 Canaries.
 Algeria.
 Algeria.
 Algeria.
 India, S. W. Asia,
 Egypt, S. Africa.
 Tanganyika Protect.
 Zululand.
 Transvaal.
 Java.
 Transvaal.
 Transvaal.
 Transvaal.
 Rhodesia.

50. <i>S. cirrhocoma</i> , Meyrick, Ann. Transv. Mus. Vol. 4, p. 193 (1914).	Natal.
51. <i>S. thoracica</i> , Meyrick, ibidem, Vol. 3, p. 67 (1912).	Transvaal.
52. <i>S. hyperythra</i> , Meyrick, ibidem, Vol. 8, p. 76 (1921).	Transvaal.
53. <i>S. alaopis</i> , Meyrick, ibidem, Vol. 8, p. 76 (1921).	Natal.
54. <i>S. basalis</i> , Staudinger, Stett. Ent. Zeit. Vol. 37, p. 147 (1876).	Sicily.
55. ? <i>S. Wollastoni</i> , Walsingham, Trans. Ent. Soc. Lond. p. 545 (1894).	Madeira.
56. <i>S. nugatricella</i> , Rebel, Stett. Ent. Zeit. Vol. 54, p. 50 (1893).	Spain.
57. <i>S. deverrae</i> , Walsingham, Ent. M. Mag. Vol. 41, p. 124 (1905).	Algeria.
58. <i>S. maraschella</i> , Caradja, Iris, Vol. 34, p. 107 (1920).	Asia Minor.
59. <i>S. geryella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 323 (1915).	Algeria.
60. <i>S. bivittella</i> , Chrétien, ibidem, p. 324 (1915).	Tunis.
61. <i>S. detersella</i> , Zeller, Isis, p. 846 (1847). <i>egenella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, f. 340 (1855).	S. Europe, Asia Minor, Algeria.
62. <i>S. kermella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 328 (1915).	Algeria
63. <i>S. quadripunctella</i> , Chrétien, ibidem, p. 327 (1915).	Algeria.

122. GENUS GLAPHYRERGA, NOV. GEN.

Type : *G. mauricaudella*, Oberthür.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 2/3, in ♂ shortly ciliated, basal joint elongate, without pecten. Labial palpi long, recurved, second joint smooth-scaled, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 from towards angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen hardly sinuate, cilia 1/2; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 rather approximated towards base.

Remarks. — A development of *Melitoxestis*.

Geographical distribution of species. — North African.

Larva unknown.

1. *G. mauricaudella*, Oberthür, Et. Ent. Vol. 12, p. 43, pl. 6, f. 34 (1888). Algeria.

123. GENUS MELITOXESTIS, MEYRICK

Melitoxestis, Meyrick, Ann. Transv. Mus. Vol. 8, p. 75 (1921). — Type : *M. centrotypa*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ moderately ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint rather thickened with appressed scales, terminal joint longer than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, apex obtuse, termen hardly sinuate, cilia 2/3; without cubital pecten; 3 and 4 connate, 5 rather approximated, 6 and 7 closely approximated towards base.

Remarks. — Derived from a form approaching *Compsolechia*.

Geographical distribution of species. — South African.

Larva unknown.

1. *M. centrotypa*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 76 (1921). Rhodesia.

124. GENUS ANTERETHISTA, MEYRICK

Anterethista, Meyrick, Trans. Ent. Soc. Lond. p. 237 (1914). — Type : *A. heteractis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, slender, second joint with scales somewhat expanded towards apex above, terminal joint longer than second, acute. Maxillary palpi minute, filiform, appressed. Posterior tibiae with appressed hairs above. Forewings with 1b furcate, 2 from towards angle, 7 absent, 11 from 2/3. Hindwings 1, elongate-trapezoidal, apex tolerably pointed, termen somewhat sinuate, cilia 3; with slight cubital pecten; 3 and 4 rather approximated towards base, 5 nearly parallel, 6 absent, 7 to apex.

Remarks. — Perhaps a specialised derivative of *Commatica*.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|--|-----------------------|
| 1. <i>A. phosphoropa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 66 (1922). | Brazil, Peru. |
| 2. <i>A. heteractis</i> , Meyrick, ibidem, p. 237 (1914). | Guiana, Brazil, Peru. |

125. GENUS STAGMATURGIS, MEYRICK

Stagmaturgis, Meyrick, Exot. Microlep. Vol. 3, p. 25 (1923). — Type : *S. catharosema*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, slender, smooth, terminal joint longer than second, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 2-6 parallel, 7 absent, 11 from middle. Hindwings 2/3, narrow-trapezoidal, apex pointed, termen sinuate, cilia 1 1/4; with slight cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 approximated towards base.

Remarks. — Probably derived from *Commatica*.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|---|---------|
| 1. <i>S. catharosema</i> , Meyrick, Exot. Microlep. Vol. 3, p. 25 (1923). | Brazil. |
|---|---------|

126. GENUS SIMONEURA, WALSINGHAM

Simoneura, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 72 (1911). — Type : *S. ophitis*, Walsingham.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, somewhat expanded above towards apex, terminal joint nearly as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from 3/4, 3 and 4 closely approximated, 5 rather approximated, 6 remote, to costa, 7 absent, 11 from

middle. Hindwings 1, trapezoidal, termen slightly sinuate; with cubital pecten (?); 3 and 4 connate, 5 approximated, 6 and 7 connate.

Remarks. — A derivative of *Commatica*.

Geographical distribution of species. — North American.

Larva unknown.

1. *S. ophitis*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 73, pl. 2, Mexico.
f. 29 (1911).

127. GENUS COMMATICA, MEYRICK

Commatica, Meyrick, Trans. Ent. Soc. Lond. p. 18 (1909). — Type : *C. eremna*, Meyrick.

Apopira, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 73 (1911). — Type : *C. falcatella*, Walker.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, slender, without pecten. Labial palpi long, recurved, second joint thickened with scales, somewhat roughly expanded towards apex above, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, apex more or less pointed, termen from emarginate to hardly sinuate, cilia 1; with cubital pecten; 3 and 4 connate or almost, 5 nearly parallel, 6 and 7 separate at base, diverging.

Remarks. — This and the four following genera form an associated group derived from *Compsolechia*.

Geographical distribution of species. — South American, extending up to Mexico, with a straggler in South Africa.

Larva unknown.

- | | |
|---|---|
| 1. <i>C. parvula</i> , Meyrick, Trans. Ent. Soc. Lond. p. 241 (1914). | Guiana, Brazil. |
| 2. <i>C. falcatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 625 (1864).
<i>rostella</i> , Felder, Reis. Novar. Lep. pl. 140, f. 12 (1875). | Mexico, Guatemala,
Guiana, Braz., Colom-
bia. |
| 3. <i>C. cyanorrhoea</i> , Meyrick, Trans. Ent. Soc. Lond. p. 241 (1914). — PI. 2,
Fig. 49. | Guiana, Brazil. [bia.] |
| 4. <i>C. chionura</i> , Meyrick, ibidem, p. 240 (1914). | Guiana, Brazil, Peru. |
| 5. <i>C. extremella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 625 (1864). | Brazil. |
| 6. <i>C. xanthocarpa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 73 (1922). | Peru. |
| 7. <i>C. acropelta</i> , Meyrick, ibidem, p. 238 (1914). | Guiana, Brazil, Peru. |
| 8. <i>C. stygia</i> , Meyrick, ibidem, p. 71 (1922). | Brazil. |
| 9. <i>C. eremna</i> , Meyrick, ibidem, p. 19 (1909). | Bolivia. |
| 10. <i>C. cryptina</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 76 (1911). | Mexico. |
| 11. <i>C. nerterodes</i> , Meyrick, Trans. Ent. Soc. Lond. p. 239 (1914). | Guiana. |
| 12. <i>C. hexacentra</i> , Meyrick, ibidem; p. 73 (1922). | Brazil. |
| 13. <i>C. servula</i> , Meyrick, ibidem, p. 72 (1922). | Peru. |
| 14. <i>C. placoterra</i> , Meyrick, Exot. Microlep. Vol. 2, p. 146 (1918). | Colombia. |
| 15. <i>C. compsotoma</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 77 (1921). | Rhodesia, Port. E. Africa. |
| 16. <i>C. palirrhoa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 72 (1922). | Brazil, Peru. |
| 17. <i>C. lupata</i> , Meyrick, ibidem, p. 239 (1914). | Guiana, Peru. |
| 18. <i>C. phanocrossa</i> , Meyrick, ibidem, p. 72 (1922). | Brazil. |
| 19. <i>C. emplasta</i> , Meyrick, ibidem, p. 240 (1914). | Guiana, Brazil, Peru. |
| 20. <i>C. metochra</i> , Meyrick, ibidem, p. 238 (1914). | Guiana, Brazil, Peru. |

128. GENUS UNTOMIA, BUSCK

Untomia, Busck, Proc. U. S. Mus. Vol. 30, p. 727 (1906). — Type : *U. untomiella*, Busck.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with smooth scales, somewhat expanded at apex above, apex truncate, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 3 and 4 stalked, 7 absent, 11 from middle. Hindwings nearly 1, elongate-trapezoidal, apex produced, termen sinuate, cilia nearly 1; with cubital pecten; 3 and 4 connate, 5 curved, transverse vein partially obsolete, 6 and 7 stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — North and Central America to Ecuador.

Larva unknown.

- | | |
|---|------------------|
| 1. <i>U. alticolans</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 74 (1911) (-lens). | Mexico. |
| 2. <i>U. rotundata</i> , Walsingham, ibidem, Vol. 4, p. 75 (1911). | Mexico. |
| 3. <i>U. latistriga</i> , Walsingham, ibidem, Vol. 4, p. 75, pl. 2, f. 30 (1911). | Mexico. |
| 4. <i>U. juvenella</i> , Walsingham, Proc Zool. Soc. Lond. p. 86 (1897).
<i>horista</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 76 (1911). | Mexico, Jamaica. |
| 5. <i>U. albistrigella</i> , Chambers, Canad. Ent. Vol. 4, p. 171 (1872). | Kentucky. |
| 6. <i>U. untomiella</i> , Busck, Proc. U. S. Mus. Vol. 30, p. 727 (1906). | Texas. |
| 7. <i>U. acicularis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 146 (1918). | Ecuador. |
| 8. <i>U. melanobathra</i> , Meyrick, ibidem, Vol. 2, p. 146 (1918). | Ecuador. |
| 9. <i>U. acuminata</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 75, pl. 2, f. 31 (1911). | Mexico. |

129. GENUS HAPALONOMA, MEYRICK

Hapalonoma, Meyrick, Trans. Ent. Soc. Lond. p. 244 (1914). — Type : *H. sublustricella*, Walker.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae almost 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint expanded with rough hairs above towards apex, beneath with fringe of long rough projecting hairs, terminal joint longer than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 6 to apex, 7 absent, 11 from beyond middle. Hindwings 1, elongate trapezoidal, apex rounded, termen hardly sinuate, cilia over 1; with cubital pecten; 3 and 4 connate, 5 rather approximated, 6 and 7 closely approximated toward base.

Remarks. — This and the next genus differ curiously from their allies by the long fringe-tuft of palpi, but are certainly referable here, but perhaps separately evolved from *Compsolechia*.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|---|-----------------------|
| 1. <i>H. sublustricella</i> , Walker, List Lep. Brit. Mus. Vol. 29, p. 623 (1864).
<i>argyracta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 244 (1914). | Guiana, Brazil, Peru. |
|---|-----------------------|

130. GENUS ETHIROSTOMA, MEYRICK

Ethirostoma, Meyrick Trans. Ent. Soc. Lond. p. 244 (1914). — Type : *E. semiacma*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint clothed above with dense scales expanded towards apex, beneath tufted with long rough projecting hairs, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse-pointed, termen hardly sinuate, cilia over 1; with cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated towards base.

Remarks. — Except in the palpi, closely allied to *Battaristis*.

Geographical distribution of species. — South American.

Larva unknown.

1. *E. semiacma*, Meyrick, Trans. Ent. Soc. Lond. p. 245 (1914).

Guiana, Brazil.

2. *E. interpolata*, Meyrick, ibidem, p. 71 (1922).

Brazil, Peru.

131. GENUS BATTARISTIS, MEYRICK

Battaristis, Meyrick, Trans. Ent. Soc. Lond. p. 245 (1914). — Type : *B. ichnota*, Meyrick.

Duvita, Busck, Proc. Ent. Soc. Wash. Vol. 18, p. 147 (1916). — Type : *B. vittella*, Busck.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, thickened with appressed scales, second joint with scales somewhat expanded at apex above, truncate, terminal joint as long as second, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from 4/5-5/6, 6 sometimes to apex, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse, termen faintly sinuate, cilia 1; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated towards base or stalked.

Larva (*vittella*) feeding in seed-cones and galls.

Foodplants *Coniferae*.

- | | |
|--|--|
| 1. <i>B. emissurella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 620 (1864). | Panama, Guiana, Brazil,
Peru. |
| <i>severella</i> , Walker, ibidem, Vol. 30, p. 1035 (1864). | |
| <i>fuliginosa</i> , Felder, Reis. Novar, Lep. pl. 138, f. 43 (1875). | |
| <i>brunniceps</i> , Felder, ibidem, pl. 140, f. 25 (1875). | |
| <i>dorsalis</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 8 (1914). | |
| <i>astroconis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 139 (1918). | |
| 2. <i>B. cycella</i> , Busck, Proc. U. S. Mus. Vol. 25, p. 848 (1903). | Arizona. |
| 3. <i>B. vittella</i> , Busck, Proc. Ent. Soc. Wash. Vol. 18, p. 147 (1916). | New York, Maryland. |
| 4. <i>B. nigratomella</i> , Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 11 (1863). | Canada, E. United
States, Colorado. |
| <i>apicilinella</i> , Clemens, ibidem, Vol. 2, p. 120 (1863). | |
| <i>apicistrigella</i> , Chambers, Canad. Ent. Vol. 4, p. 66 (1872). | |
| <i>concinnusella</i> , Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 253 (1875). | |
| 5. <i>B. bistrigella</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 9 (1914). | Panama. |
| 6. <i>B. unistrigella</i> , Busck, ibidem, Vol. 47, p. 9 (1914). | Panama. |
| 7. <i>B. curtella</i> , Busck, ibidem, Vol. 47, p. 10 (1914). | Panama. |

8. *B. perinaeta*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 40 Mexico.
(1910).
9. *B. syngraphopa*, Meyrick, Trans. Ent. Soc. Lond. p. 73 (1922). Brazil, Peru.
10. *B. amphiscolia*, Meyrick, ibidem, p. 248 (1914). Guiana, Brazil, Peru.
11. *B. prismatopa*, Meyrick, ibidem, p. 246 (1914). Guiana.
12. *B. ardiophora*, Meyrick, ibidem, p. 247 (1914). — PI. 2, Fig. 50. Guiana, Brazil, Peru.
13. *B. orthocampta*, Meyrick, ibidem, p. 246 (1914). Guiana, Brazil.
14. *B. coniosema*, Meyrick, ibidem, p. 74 (1922). Brazil, Peru.
15. *B. symphora*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 76, pl. 2, f. 32 (1911). Mexico.
16. *B. ichnota*, Meyrick, Trans. Ent. Soc. Lond. p. 247 (1914). Guiana, Brazil.
17. *B. synocha*, Meyrick, ibidem, p. 74 (1922). Peru.
18. *B. sphenodelta*, Meyrick, ibidem, p. 75 (1922). Brazil.
19. *B. atelesta*, Meyrick, ibidem, p. 248 (1914). Guiana.
20. *B. melanamba*, Meyrick, ibidem, p. 249 (1914). Guiana, Brazil, Peru.
21. *B. stereogramma*, Meyrick, ibidem, p. 249 (1914). Guiana, Brazil, Peru.

132. GENUS MOLOPOSTOLA, MEYRICK

Molopostola, Meyrick, Exot. Microlep. Vol. 2, p. 298 (1920). — Type : *M. rufitincta*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, curved, ascending, or in ♀ nearly straight, porrected, second joint much thickened with dense scales roughly projecting above, terminal joint shorter than second, in ♀ sometimes much shorter, thickened with appressed scales, slightly rough anteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from 3/4, 3 from before angle, 4 and 5 stalked or connate from angle, 7 absent, 11 from middle. Hindwings 1, rounded-trapezoidal, termen hardly sinuate, cilia 3/5; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate.

Remarks. — A development from *Promolopica*, differing from it by the loss of vein 7 of forewings, and of the cubital pecten of hindwings.

Geographical distribution of species. — South American.

Larva unknown.

1. *M. rufitincta*, Meyrick, Exot. Microlep. Vol. 2, p. 299 (1920). Guiana, Brazil.
2. *M. calumnians*, Meyrick, ibidem, Vol. 3, p. (ined.). Brazil.

133. GENUS PROMOLOPICA, NOV. GEN.

Type : *P. epiphanta*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, basal joint elongate, without pecten. Labial palpi (♀) long, porrected, second joint long, thickened with dense scales, above with rough projecting hairs diminishing to apex, terminal joint hardly 1/3 of second, thickened with scales rather rough beneath, pointed (in ♂ perhaps more as *Molopostola*). Maxillary palpi very short, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 from 3/4, 3 from before angle, 4 and 5 connate from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1,

trapezoidal, termen slightly sinuate, cilia 3/5; with strong cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate.

Remarks. — Derived from *Compsolechia*, with curious modification of palpi.

Geographical distribution of species. — South American.

Larva unknown.

1. *P. epiphanta*, Meyrick, Exot. Microlep. Vol 3, p. (ined.). Brazil.

134. GENUS COLEOSTOMA, MEYRICK

Coleostoma, Meyrick, Trans. Ent. Soc. Lond. p. 99 (1922). — Type : *C. entryphopa*, Meyrick.

Characters. — Head with appressed scales, sidetufts somewhat raised; ocelli small, posterior; tongue developed. Antennae nearly 1, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint broadly thickened with dense scales, slightly expanded at apex above, terminal joint as long as second, thickened with dense scales roughly projecting posteriorly except at apex, pointed. Maxillary palpi short, loosely scaled, appressed to tongue. Posterior tibiae shortly rough-scaled above. Forewings with 1b furcate, 2 from towards angle, 3 from before angle, 4 and 5 approximated, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings somewhat over 1, trapezoidal ovate, apex obtuse, termen faintly bisinuate, cilia 1/2; with cubital pecten; 3 and 4 connate, 5 approximated, 6 and 7 closely approximated towards base.

Remarks. — Also a development from *Compsolechia* with specialised palpi.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. entryphopa*, Meyrick, Trans. Ent. Soc. Lond. p. 99 (1922). Brazil.

135. GENUS COMPSOLECHIA, MEYRICK

Compsolechia, Meyrick, Exot. Microlep. Vol. 2, p. 137 (1918). — Type : *C. repandella*, Walker.

Characters. — Head smooth; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ simple or shortly ciliated, basal joint elongate, slender, without pecten. Labial palpi very long, recurved, second joint smooth-scaled, compressed, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae more or less shortly rough-scaled above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa (in *eurygypsa* 7 absent), 11 from middle. Hindwings 1 or over 1, trapezoidal, apex obtuse, termen not or slightly sinuate, cilia 1/3-1; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 nearly approximated at base.

Remarks. — Correlated with *Anacampsis*. The species are elegantly marked but sometimes very similar, and require careful attention to detail.

Geographical distribution of species. — Characteristic of South America, where the genus is numerous developed, apparently spreading thence to North America and Europe, with stragglers in Japan and South Africa.

Larva (15 known) feeding between spun leaves.

Foodplants : *Leguminosae* (4), *Rosaceae* (3), *Salicaceae* (2), and six other Orders.

1. *C. suffectella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 610 (1864). Mexico to Brazil & Peru.
 2. *C. stelliferella*, Walker, ibidem, Vol. 29, p. 613 (1864). Guatemala to Brazil & Peru.
 3. *C. pentastra*, Meyrick, Trans. Ent. Soc. Lond. p. 81 (1922). Brazil, Peru.
 4. *C. trochilea*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 41 (1910). Mexico.
 5. *C. perlatella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 612 (1864). — Guiana, Brazil, Peru.
- Pl. 3, Fig. 52.**
- | | |
|---|-----------------------|
| smaragdulella, Walker, ibidem, Vol. 29, p. 614 (1864). | Brazil. |
| secundella, Walker, ibidem, Vol. 29, p. 615 (1864). | Brazil. |
| 6. <i>C. chelidonia</i> , Meyrick, Trans. Ent. Soc. Lond. p. 82 (1922). | Guiana, Brazil. |
| 7. <i>C. hemileucas</i> , Meyrick, ibidem, p. 82 (1922). | Guiana, Brazil, Peru. |
| 8. <i>C. cassidata</i> , Meyrick, ibidem, p. 257 (1914). | Guiana, Brazil, Peru. |
| 9. <i>C. secretella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 616 (1864). | Guiana, Brazil, Peru. |
| <i>cistulata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 258 (1914). | |
| <i>trimolybda</i> , Meyrick, ibidem, p. 258 (1914). | |
| 10. <i>C. mesodelta</i> , Meyrick, ibidem, p. 83 (1922). | Brazil. |
| 11. <i>C. peculella</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 7 (1914). | Panama. |
| 12. <i>C. siderophaea</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 39 (1910). | Mexico. |
| 13. <i>C. metadupa</i> , Walsingham, ibidem, Vol. 4, p. 40 (1910). | Mexico. |
| 14. <i>C. ferreata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 257 (1914). | Guiana. |
| 15. <i>C. quadrifascia</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 617 (1864). | Brazil. |
| <i>superella</i> , Walker, ibidem, Vol. 29, p. 617 (1864). | |
| 16. <i>C. orthophracta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 257 (1914). | Guiana. |
| 17. <i>C. refracta</i> , Meyrick, ibidem, p. 258 (1914). | Guiana, Brazil, Peru. |
| 18. <i>C. platiastis</i> , Meyrick, ibidem, p. 84 (1922). | Brazil. |
| 19. <i>C. praenirea</i> , Meyrick, ibidem, p. 260 (1914). | Guiana. |
| 20. <i>C. sporozona</i> , Meyrick, ibidem, p. 259 (1914). | Guiana, Brazil, Peru. |
| 21. <i>C. blepharopa</i> , Meyrick, ibidem, p. 259 (1914). | Guiana. |
| 22. <i>C. petromorpha</i> , Meyrick, ibidem, p. 84 (1922). | Brazil, Peru. |
| 23. <i>C. lithomorpha</i> , Meyrick, ibidem, p. 256 (1914). | Guiana, Brazil, Peru. |
| 24. <i>C. sciomima</i> , Meyrick, ibidem, p. 84 (1922). | Brazil, Peru. |
| 25. <i>C. phaeotoxa</i> , Meyrick, ibidem, p. 85 (1922). | Brazil. |
| 26. <i>C. religata</i> , Meyrick, ibidem, p. 85 (1922). | Peru. |
| 27. <i>C. cognatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 606 (1864). | Brazil. |
| 28. <i>C. ischnoptera</i> , Meyrick, Trans. Ent. Soc. Lond. p. 85 (1922). | Brazil. |
| 29. <i>C. eupecta</i> , Meyrick, ibidem, p. 265 (1914). | Guiana, Brazil. |
| 30. <i>C. canofusella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 617 (1864). | Brazil. |
| 31. <i>C. suspectella</i> , Walker, ibidem, Vol. 29, p. 615 (1864). | Brazil. |
| 32. <i>C. diplodichna</i> , Meyrick, Trans. Ent. Soc. Lond. p. 86 (1922). | Brazil. |
| 33. <i>C. crocodilopa</i> , Meyrick, ibidem, p. 86 (1922). | Brazil, Peru. |
| 34. <i>C. coverdalella</i> , Kearfott, Journ. N. York Ent. Soc. Vol. 11, p. 162, pl. 9, f. 13 (1903). | Louisiana. |
| 35. <i>C. fasciella</i> , Felder, Reis. Novara Lep. pl. 140, f. 1 (1875). | Brazil. |
| 36. <i>C. scitella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 605 (1864). | Guiana, Brazil, Peru. |
| 37. <i>C. pallodoriella</i> , Busck, Proc. U. S. Mus. Vol. 25, p. 848 (1903). | Montana, New Mexico. |
| 38. <i>C. lingulata</i> , Meyrick, Exot. Microlep. Vol. 2, p. 140 (1918). | Colombia. |
| 39. <i>C. salebrosa</i> , Meyrick, ibidem, p. 140 (1918). | Colombia, Guiana. |
| 40. <i>C. levipedella</i> , Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 4 (1863). | E. United States. |
| 41. <i>C. recta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 86 (1922). | Brazil. |
| 42. <i>C. argyrahma</i> , Meyrick, ibidem, p. 83 (1922). | Brazil. |

43. *C. amazonica*, Meyrick, Exot. Microlep. Vol. 2, p. 139 (1918).
suffusella, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 623 (1864) (praeocc.).
 44. *C. scopulata*, Meyrick, Trans. Ent. Soc. Lond. p. 260 (1914).
 45. *C. zebrina*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 37 (1910).
 46. *C. versatella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 604 (1864).
 47. *C. seductella*, Walker, ibidem, Vol. 29, p. 602 (1864).
 48. *C. ambusta*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 40,
 pl. 1, f. 33 (1910).
 brochospila, Meyrick, Trans. Ent. Soc. Lond. p. 265 (1914).
 49. *C. campalea*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 36
 (1910).
 50. *C. succincta*, Walsingham, ibidem, Vol. 4, p. 38, pl. 2, f. 1 (1910).
 51. *C. titanota*, Walsingham, ibidem, Vol. 4, p. 38, pl. 1, f. 32 (1910).
 52. *C. caryoterma*, Meyrick, Trans. Ent. Soc. Lond. p. 88 (1922).
 53. *C. scholias*, Meyrick, ibidem, p. 88 (1922).
 54. *C. minicosma*, Meyrick, ibidem, p. 89 (1922). — **Pl. 3, Fig. 54.**
 55. *C. trachycnemis*, Meyrick, ibidem, p. 89 (1922).
 56. *C. sesamodes*, Meyrick, ibidem, p. 90 (1922).
 57. *C. subapicalis*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 599 (1864).
 58. *C. leucorrhapha*, Meyrick, Trans. Ent. Soc. Lond. p. 261 (1914).
 59. *C. niphocentra*, Meyrick, ibidem, p. 90 (1922).
 60. *C. amaurola*, Meyrick, ibidem, p. 262 (1914).
 61. *C. tardella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 606 (1864).
 sublatella, Walker, ibidem, Vol. 29, p. 607 (1864).
 collocatella, Walker, ibidem, Vol. 29, p. 616 (1864).
 62. *C. volubilis*, Meyrick, Trans. Ent. Soc. Lond. p. 91 (1922).
 63. *C. halmyra*, Meyrick, ibidem, p. 262 (1914).
 64. *C. eurygypsa*, Meyrick, ibidem, p. 91 (1922).
 65. *C. antiplaca*, Meyrick, ibidem, p. 92 (1922).
 66. *C. ptochogramma*, Meyrick, ibidem, p. 93 (1922).
 67. *C. inusta*, Meyrick, ibidem, p. 264 (1914).
 68. *C. scutella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 363, pl. 5, f. 123 (1877).
 69. *C. repandella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 602 (1864).
 subscriptella, Walker, ibidem, Vol. 29, p. 603 (1864).
 episema, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 37, pl. 1, f. 34
 (1910).
 diortha, Meyrick, Trans. Ent. Soc. Lond. p. 263 (1914).
 70. *C. tetrortha*, Meyrick, ibidem, p. 92 (1922).
 71. *C. picticornis*, Walsingham, Proc. Zool. Soc. Lond. p. 68 (1897).
 72. *C. stillata*, Meyrick, Trans. Ent. Soc. Lond. p. 93 (1922).
 73. *C. anthracura*, Meyrick, ibidem, p. 263 (1914).
 74. *C. superfusella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 600 (1864).
 75. *C. diazeucta*, Meyrick, Exot. Microlep. Vol. 2, p. 138 (1918).
 trajectella, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 599 (1864) (praeocc.).
 76. *C. phepsalitis*, Meyrick, Trans. Ent. Soc. Lond. p. 94 (1922).
 77. *C. solidella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 597 (1864). —
 Pl. 3, Fig. 53.
 78. *C. drachmaea*, Meyrick, Trans. Ent. Soc. Lond. p. 94 (1922).
 79. *C. thysanora*, Meyrick, ibidem, p. 261 (1914).
 80. *C. binotatella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 601 (1864).
 81. *C. abruptella*, Walker, ibidem, Vol. 29, p. 601 (1864).
 sectella, Walker, ibidem, Vol. 29, p. 605 (1864).
 82. *C. suspectella*, Walker, ibidem, Vol. 29, p. 604 (1864).
 83. *C. trapezias*, Meyrick, Trans. Ent. Soc. Lond. p. 87 (1922).
 84. *C. rhombica*, Meyrick, ibidem, p. 95 (1922).

85. *C. pungens*, Meyrick, Trans. Ent. Soc. Lond. p. 95 (1922). Peru.
 86. *C. erebodelta*, Meyrick, ibidem, p. 96 (1922). Peru.
 87. *C. incurva*, Meyrick, ibidem, p. 264 (1914). Guiana, Brazil, Peru.
 88. *C. accinctella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 600 (1864). Brazil, Peru.
 89. *C. transjectella*, Walker, ibidem, Vol. 29, p. 598 (1864). Brazil, Peru.
 90. *C. corymbas*, Meyrick, Exot. Microlep. Vol. 2, p. 139 (1918). Guiana.
 91. *C. par mata*, Meyrick, ibidem, Vol. 2, p. 139 (1918). Colombia.
 92. *C. tornoptila*, Meyrick, Trans. Ent. Soc. Lond. p. 96 (1922). Brazil.
 93. *C. stasiogastra*, Meyrick, ibidem, p. 97 (1922). Brazil.
 94. *C. loxogramma*, Meyrick, ibidem, p. 97 (1922). Brazil.
 95. *C. chrysoplaca*, Meyrick, ibidem, 1911, p. 694 (1912). Venezuela.
 96. *C. glaphyra*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 39, Mexico.
 pl. 1, f. 31 (1910).
 97. *C. neurophora*, Meyrick, Trans. Ent. Soc. Lond. p. 98 (1922). Brazil.
 98. *C. percnospila*, Meyrick, ibidem, p. 266 (1914). Guiana.
 99. *C. abolitella*, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1025 (1864). ? S. America.
 100. *C. terrenella*, Busck, Proc. U. S. Mus. Vol. 47, p. 10 (1914). Panama.
 101. *C. ocelligera*, Butler, Trans. Ent. Soc. Lond. p. 77 (1883). Chile.
 102. *C. monochromella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 607 Brazil, Peru.
 (1864).
 displacitella, Walker, ibidem, Vol. 29, p. 609 (1864).
 103. *C. permagna*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 284 (1920). Transvaal.
 104. *C. fullonella*, Zeller, Verh. Zool.-Bot. Ges. Wien, Vol. 23, p. 276 (1873). Texas.
 rufusella, Chambers, Canad. Ent. Vol. 6, p. 240 (1874).
 subrubrella, Chambers, ibidem, p. 240 (1874).
 rubescens, Walsingham, Proc. Zool. Soc. Lond. p. 319, pl. 36, f. 9 (1881).
 105. *C. dicax*, Meyrick, Trans. Ent. Soc. Lond. p. 266 (1914). Guiana, Peru.
 106. *C. lagunculariella*, Busck, Proc. U. S. Mus. Vol. 23, p. 230 (1900). Florida, Panama.
 107. *C. lupinella*, Busck, Canad. Ent. Vol. 33, p. 14 (1901). Canada.
 108. *C. scintillella*, Fischer von Röslerstamm, Abbild. Schmett. p. 221, pl. 77, S. C. & S. Europe, Asia
 f. 3 (1839). Minor.
 brunneella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 193, f. 578 (1855).
 contubernella, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 240 (1859).
 109. *C. acosmeta*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 34 Mexico.
 (1910).
 110. *C. mangelivora*, Walsingham, Proc. Zool. Soc. Lond. p. 80 (1897). Virgin Islands.
 111. *C. balia*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 34 Mexico.
 (1910).
 112. *C. plumbeolata*, Walsingham, Proc. Zool. Soc. Lond. p. 79 (1897). Leeward Islands.
 113. *C. dyocrossa*, Meyrick, Trans. Ent. Soc. Lond. p. 98 (1922). Brazil.
 114. *C. desectella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 362, pl. 5, f. 122 Cuba.
 (1877).
 115. *C. temerella*, Zeller, Isis, p. 284 (1846). C. Europe.
 fernigrella, Douglas, Trans. Ent. Soc. Lond. (2) Vol. 1, p. 64 (1850).
 116. *C. molybdina*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 36 Mexico.
 (1910).
 117. *C. fragariella*, Busck, Proc. U. S. Mus. Vol. 27, p. 760 (1904). Distr. Columbia.
 118. *C. subsequella*, Hübner, Samml. Eur. Schmett. Tin. f. 161 (1796). E. Europe, Asia Minor.
 obscurella, Fischer von Röslerstamm, Abbild. Schmett. p. 220 (1839).
 119. *C. psoraliella*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 226, pl. 28, Iowa.
 f. 10 (1920).
 120. *C. hirsutella*, Constant, Ann. Soc. Ent. Fr. p. 256, pl. 10, f. 17 (1884). S. France.
 121. *C. timidella*, Wocke, Bresl. Ent. Zeit. p. 63 (1887). Silesia.
 122. *C. panormitella*, Caradja, Iris, Vol. 34, p. 106 (1920). Asia Minor.
 123. *C. trifoliella*, Constant, Bull. Soc. Ent. Fr. p. 125 (1889). France.

124. *C. crescentifasciella*, Chambers, Canad. Ent. Vol. 6, p. 237 (1874). Texas.
 125. *C. epibola*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 34 (1910). Mexico, Panama.
 126. *C. rhoifruetella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 163 (1860). E. United States.
consoneella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 251 (1873).
quadrimaculella, Chambers, Canad. Ent. Vol. 6, p. 237 (1874).
ochreocostella, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 91 (1878).
 127. *C. lacteochrella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 244 (1875) (-teusochrella). California.
 128. *C. argyrothamniella*, Busck, Proc. U. S. Mus. Vol. 23, p. 231 (1900). Florida.
 129. *C. niveopulvella*, Chambers, Canad. Ent. Vol. 7, p. 210 (1875). Canada, Arizona.
 130. *C. elephas*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 39 (1910). Mexico, Guatemala.
 131. *C. nonstrigella*, Busck, Canad. Ent. Vol. 38, p. 121 (1906). Pennsylvania.
 132. *C. tristrigella*, Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 181 (1882). Illinois, Kansas.
 133. *C. metagramma*, Meyrick, Exot. Microlep. Vol. 2, p. 138 (1918). Japan.
 134. *C. agrimonella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 162 (1860). E. United States, Canada,
aduncella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 18, p. 614 (1868). Siberia, Asia Minor.
solemnella, Christoph, Bull. Soc. Nat. Mosc. p. 27 (1882).
 135. *C. kearfottella*, Busck, Proc. U. S. Mus. Vol. 25, p. 842 (1903). New Jersey.

136. GENUS CATALEXIS, WALSHAM

Catalexis, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 19 (1909). — Type : *C. tapinota*, Walsingham.

Characters. — Head smooth; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, rather rough towards apex beneath, terminal joint somewhat shorter than second, moderate, acute. Maxillary palpi rudimentary. Posterior tibiae with appressed scales. Forewings with 1b furcate, 2 from 2/3, 3 from before angle, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings hardly 1, trapezoidal, termen sinuate, cilia 3/5; 3 and 4 separate, 4 from angle, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — I have not seen this genus, of which the affinity is perhaps uncertain.

Geographical distribution of species. — Central American.

Larva unknown.

1. *C. tapinota*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 20, pl. 1, f. 18 (1909). Guatemala.

137. GENUS ACANTHOPHILA, HEINEMANN

Acanthophila, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 320 (1870). — Type : *A. alacella*, Duponchel.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, slightly projecting above towards apex, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 from towards angle, 3 and 4 closely approximated from angle, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, termen sinuate, cilia 1; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Certainly a derivative of *Anacampsis*.

Geographical distribution of species. — European.

Larva feeding in silken galleries on tree-trunks.

Foodplants *Lichenes*.

1. *A. alacella*, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, pl. 297, f. 12 (1838). Europe.

138. GENUS SOROTACTA, MEYRICK

Sorotacta, Meyrick, Trans. Ent. Soc. Lond. p. 253 (1914). — Type : *S. viridans*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, considerably expanded towards apex, somewhat projecting angularly above and beneath, terminal joint as long as second, with rough projecting scales posteriorly except towards apex, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with tufts of scales; 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse-pointed, termen slightly bisinuate, cilia 1 1/2; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated towards base.

Remarks. — Probably derived from *Alsodryas*.

Geographical distribution of species. — South American.

Larva unknown; but considering the strongly marked adaptation of the imago in this and the next genus to lichen-covered tree-trunks, it is likely that the larval habit may be similar to that of the preceding genus.

- | | |
|--|-----------------|
| 1. <i>S. viridans</i> , Meyrick, Trans. Ent. Soc. Lond. p. 254 (1914). | Guiana, Brazil. |
| 2. <i>S. bryochlora</i> , Meyrick, ibidem, p. 76 (1922). | Brazil. |

139. GENUS ALSODRYAS, MEYRICK

Alsodryas, Meyrick, Trans. Ent. Soc. Lond. p. 250 (1914). — Type : *A. lactaria*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint densely scaled, with rough projecting scales or strong triangular tuft at apex beneath, terminal joint as long as second or longer, thickened with scales and slightly roughened anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with tufts of scales; 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse, termen hardly sinuate, cilia 2/3-1 1/2; with cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated towards base.

Remarks. — A derivative of *Anacampsis*.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|--|---------|
| 1. <i>A. lactaria</i> , Meyrick, Trans. Ent. Soc. Lond. p. 250 (1914). | Guiana. |
| 2. <i>A. prasinoptila</i> , Meyrick, ibidem, p. 75 (1922). | Brazil. |
| 3. <i>A. deltochlora</i> , Meyrick, ibidem, p. 76 (1922). | Brazil. |

140. GENUS ANACAMPSIS, CURTIS

Anacampsis, Curtis, Brit. Ent. Vol. 4, pl. 189 (1827). — Type : *A. populella*, Clerck.

Tachyptilia, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 321 (1870). — Type : *A. populella*, Clerck.

Agriastis, Meyrick, Trans. Ent. Soc. Lond. p. 251 (1914). — Type : *A. peloptila*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, more or less expanded towards apex above, terminal joint as long as second or longer, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with slight tufts of scales; 1b furcate. 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse, termen hardly sinuate, cilia over 1; with cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated towards base.

Remarks. — The species are generally obscurely mottled insects of similar appearance, adapted for concealment on tree-trunks, where they habitually rest.

Geographical distribution of species. — Characteristically American, but with stragglers in Europe, India and Africa.

Larva (4 known) feeding between spun leaves.

Foodplants : *Salicaceae*, *Cupuliferae*, *Euphorbiaceae*.

1. <i>A. lithodelta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 77 (1922).	Peru.
2. <i>A. diplodelta</i> , Meyrick, ibidem, p. 76 (1922).	Brazil.
3. <i>A. viretella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 340, pl. 4, f. 110 (1877).	Guiana, Brazil.
4. <i>A. petrographa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 79 (1922).	Brazil.
5. <i>A. pomaceella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 620 (1864).	Brazil.
6. <i>A. prasina</i> , Meyrick, Trans. Ent. Soc. Lond. p. 251 (1914). — Pl. 3,	Guiana.
Fig. 51.	
7. <i>A. phytomiella</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 8 (1914).	Panama.
8. <i>A. considerata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 78 (1922).	Brazil, Peru.
9. <i>A. cenelpis</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 77, pl. 2, f. 34 (1910).	Mexico.
10. <i>A. scalata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 252 (1914).	Guiana.
11. <i>A. multinotata</i> , Meyrick, Exot. Microlep. Vol. 2, p. 134 (1918).	Guiana.
12. <i>A. perquisita</i> , Meyrick, Trans. Ent. Soc. Lond. p. 78 (1922).	Brazil.
13. <i>A. nocturna</i> , Meyrick, ibidem, p. 252 (1914).	Guiana.
14. <i>A. subactella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1026 (1864).	? S. America.
15. <i>A. conclusella</i> , Walker, ibidem, Vol. 29, p. 593 (1864). <i>tephriasiella</i> , Chambers, Canad. Ent. Vol. 4, p. 68 (1872). <i>grisefasciella</i> , Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 253 (1875).	E. United States. Canada.
16. <i>A. insularis</i> , Walsingham, Proc. Zool. Soc. Lond. p. 81 (1897).	Virgin Islands.
17. <i>A. poliombra</i> , Meyrick, Trans. Ent. Soc. Lond. p. 77 (1922).	Brazil.
18. <i>A. canevodes</i> , Meyrick, ibidem, p. 79 (1922).	Brazil.
19. <i>A. peloptila</i> , Meyrick, ibidem, p. 251 (1914).	Guiana.
20. <i>A. triangularis</i> , Braun, Proc. Calif. Acad. Sc. (4) Vol. 12, p. 118 (1923).	California.
21. <i>A. inquieta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 253 (1914).	Guiana.
22. <i>A. languens</i> , Meyrick, Exot. Microlep. Vol. 2, p. 142 (1918).	Ecuador.
23. <i>A. lapidella</i> , Walsingham, Proc. Zool. Soc. Lond. p. 81 (1897).	Windward Islands.
24. <i>A. cornifera</i> , Walsingham, ibidem, p. 79 (1897).	Virgin Islands.
25. <i>A. specularis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 142 (1918).	S. India, Ceylon.

26. *A. rivalis*, Meyrick, Exot. Microlep. Vol. 2, p. 141 (1918). S. India, Ceylon.
 27. *A. rhabdodes*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 36, Mexico.
 pl. 1, f. 30 (1910).
 28. *A. cosmia*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 77 (1921). Natal.
 29. *A. primigenia*, Meyrick, Exot. Microlep. Vol. 2, p. 141 (1918). Colombia, Ecuador.
 30. *A. ursula*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 35 (1910). Mexico.
 31. *A. conistica*, Walsingham, ibidem, Vol. 4, p. 35 (1910). Mexico.
 32. *A. quinquepunctella*, Walsingham, Proc. Zool. Soc. Lond. p. 80 (1897). Windward Islands.
 33. *A. tridentella*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 28, Mexico.
 pl. 1, f. 24 (1909).
 34. *A. innocuella*, Zeller, Verh. Zool.-Bot. Ges. Wien, Vol. 23, p. 249 (1873). Colorado, Texas.
 35. *A. populella*, Clerck, Icon. Ins. pl. 11, f. 5 (1760). Europe, Mongolia,
blattariella, Hübner, Samml. Eur. Schmett. Tin. f. 148 (1796). E. Siberia.
tremulella, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, pl. 296, f. 5 (1838).
laticinctella, Wood, Ind. Ent. f. 1188 (1838).
 36. *A. lugens*, Caradja, Iris, Vol. 34, p. 105 (1920). E. Siberia.
 37. *A. querella*, Lafaury, Naturaliste, Vol. 29, p. 250 (1907). France, Germany.
Disquei, Meess, Mitt. Karlsr. Zool. Ver. Vol. 18, p. 125 (1910).
suberiella, Caradja Iris, Vol. 34, p. 105 (1920).
 38. *A. capyrodes*, Meyrick, Trans. Ent. Soc. Lond. p. 80 (1922). Brazil.
 39. *A. flexiloqua*, Meyrick, ibidem, p. 80 (1922). Peru.
 40. *A. idiocentra*, Meyrick, ibidem, p. 80 (1922). Brazil.

141. GENUS ANTHINORA, MEYRICK

Anthinora, Meyrick, Trans. Ent. Soc. Lond. p. 255 (1914). — Type : *A. xanthophanes*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, curved, ascending, in ♂ second joint very long, basal half slender, apical half thickened and roughened anteriorly, terminal joint very short, pointed, in ♀ second joint slightly thickened, hardly roughened, terminal joint 2/3 of second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 3/4, elongate-trapezoidal, apex produced, teumen obliquely sinuate, cilia 2 1/2; without cubital pecten; lower margin of cell very near dorsum, 3 and 4 connate, 5 absent, transverse vein absent, 6 and 7 stalked.

Remarks. — This peculiar insect is of quite exceptional appearance, and its immediate affinity seems entirely problematical.

Geographical distribution of species. — South American.

Larva unknown.

1. *A. xanthophanes*, Meyrick, Trans. Ent. Soc. Lond. p. 256 (1914). Guiana.

142. GENUS HARPOGRAPTIS, NOV. GEN.

Type : *H. eucharacta*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, smooth, slender, terminal joint longer than second, acute. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above.

Forewings with 1^b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex pointed, termen hardly sinuate, cilia 3/4; without cubital pecten; 3 and 4 connate, 5 closely approximated at base, 6 and 7 rather approximated towards base.

Remarks. — This again is a striking but very puzzling insect, conjecturally referred here; doubtless many diverse forms of this group remain to be discovered.

Geographical distribution of species. — South American.

Larva unknown.

1. *H. eucharacta*, Meyrick, Trans. Ent. Soc. Lond. p. 66 (1922). Brazil.

143. GENUS DESMAUCHA, MEYRICK

Desmaucha, Meyrick, Exot. Microlep. Vol. 2, p. 146 (1918). — Type : *D. chrysostoma*, Meyrick.

Characters. — Head smooth; ocelli small, posterior; tongue developed. Antennae 4/5, serrulate, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint smooth, terminal joint shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae shortly haired above. Forewings with 2-5 approximated, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings 1, trapezoidal, rather expanded posteriorly, apex rounded-obtuse, termen not sinuate, cilia 3/4; without cubital pecten; 3 and 4 connate, 5 parallel, 6 and 7 separate, diverging.

Remarks. — Probably derived from a form approaching *Strobisia*.

Geographical distribution of species. — South American.

Larva unknown.

1. *D. chrysostoma*, Meyrick, Exot. Microlep. Vol. 2, p. 147 (1918). — PI. 3, Guiana, Brazil.

Fig. 55.

144. GENUS MERIMNETRIA, WALSINGHAM

Merimnetria, Walsingham, Faun. Hawaiensis, Vol. 1, p. 482 (1907). — Type : *M. flaviterminella*, Walsingham.

Characters. — Head smooth; tongue developed. Antennae 1, in ♂ stout, slightly compressed, serrulate, simple, basal joint without pecten. Labial palpi long, recurved, smooth, terminal joint as long as second, acute. Maxillary palpi obsolete. Posterior tibiae clothed with hairs above. Forewings with 2 from near angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia over 1; 3 and 4 connate, 6 and 7 rather approximated towards base.

Remarks. — Probably representing an immigrant of the *Strobisia* sub-group from America.

Geographical distribution of species. — North Pacific.

Larva unknown.

1. *M. flaviterminella*, Walsingham, Faun. Hawaiensis, Vol. 1, p. 482, pl. 13, Hawaiian Islands. f. 26 (1907).

145. GENUS DIASTALTICA, WALSINGHAM

Diastattica, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 32 (1910). — Type : *D. separabilis*, Walsingham.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 2/3, basal joint without pecten. Labial palpi long, recurved, second joint thickened with scales, compressed, smooth, terminal joint as long as second, slender, acute. Maxillary palpi very short. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen slightly sinuate; 3 and 4 connate, 5 slightly approximated, 6 and 7 almost connate.

Remarks. — A modified form of the *Strobisia* sub-group.

Geographical distribution of species. — Central American.

Larva unknown.

1. *D. separabilis*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 33, Guatemala.
pl. 1, f. 29 (1910).

146. GENUS EUNOMARCHA, MEYRICK

Eunomarcha, Meyrick, Exot. Microlep. Vol. 3, p. 26 (1923). — Type : *E. glycinopis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with dense whorled scales, somewhat roughened anteriorly, terminal joint shorter than second, stout, somewhat roughened anteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked, 6 to costa, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 4/5; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 almost connate.

Remarks. — An interesting form allied to *Strobisia*, with specialised palpi and neuration.

Geographical distribution of species. — South American.

Larva unknown.

1. *E. glycinopis*, Meyrick, Exot. Microlep. Vol. 3, p. 26 (1923). Brazil.

147. GENUS SPHENOGRYPA, MEYRICK

Sphenogrypa, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 71 (1920). — Type : *S. syncosma*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5 (?), basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint thickened with dense scales, rather rough anteriorly, terminal joint shorter than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely haired above. Forewings with 2 and 3 very short-stalked from angle, 4 and 5 approximated, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 4/5; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — Apparently allied to *Strobisia*.

Geographical distribution of species. — African.

Larva unknown.

1. *S. syncosma*, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 71 (1920). Kenya Colony.

148. GENUS DREPANOTERMA, WALSINGHAM

Drepanotermes, Walsingham, Proc. Zool. Soc. Lond. p. 84 (1897). — Type : *D. lacticaudella*, Walsingham.

Characters. — Head with appressed scales; tongue developed. Antennae 1, in ♂ biciliated (1). Labial palpi long, recurved, smooth, second joint strongly compressed, terminal joint scarcely half second, acute. Maxillary palpi very short. Posterior tibiae rather rough-scaled. Forewings with 2 and 3 stalked, 7 absent. Hindwings slightly over 1, trapezoidal; 3 and 4 short-stalked, 5 approximated, 6 and 7 connate.

Remarks. — Probably a development of *Onebala*.

Geographical distribution of species. — Caribbean.

1. *D. lacticaudella*, Walsingham, Proc. Zool. Soc. Lond. p. 85 (1897). Windward Islands.

149. GENUS ERIPNURA, MEYRICK

Eripnura, Meyrick, Trans. Ent. Soc. Lond. p. 242 (1914). — Type : *E. criodes*, Meyrick.

Characters. — Head smooth; ocelli very small, posterior; tongue absent. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, rather thickened with appressed scales throughout, terminal joint shorter than second, pointed. Maxillary palpi rudimentary. Posterior tibiae with long hairs above. Forewings with 1b furcate, 2 from towards angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen hardly sinuate, cilia 3/4; without cubital pecten; 3 and 4 connate, 5 remote, parallel, 6 and 7 rather approximated towards base.

Remarks. — Probably related to *Strobisia*.

Geographical distribution of species. — South American.

Larva unknown.

1. *E. criodes*, Meyrick, Trans. Ent. Soc. Lond. p. 242 (1914). Guiana.

150. GENUS PARELECTRA, NOV. GEN.

Type : *P. helicopis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long; recurved, second joint thickened with scales, strongly compressed, with hairs loosely expanded above posteriorly, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely

haired above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 11 from beyond middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 3/5; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate or short-stalked.

Remarks. — Nearly related to *Strobisia*.

Geographical distribution of species. — American.

Larva unknown.

- | | |
|--|---------------|
| 1. <i>P. selectella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 614 (1864). | Brazil. |
| 2. <i>P. ithycosma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 267 (1914). | Guiana. |
| 3. <i>P. subvectella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 610 (1864). | Brazil. |
| 4. <i>P. scintillula</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 81, pl. 3, f. 1 (1911). | Mexico. |
| 5. <i>P. helicopis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 101 (1922). | Brazil, Peru. |

151. GENUS STROBISIA, CLEMENS

Strobisia, Clemens, Proc. Acad. Nat. Sc. Philad. p. 164 (1860). — Type : *S. iridipennella*, Clemens.

Systasiota, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 28 (1910). — Type : *S. leucura*, Walsingham.

Characters. — Head smooth; ocelli rather large, posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint smooth-scaled, compressed, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae more or less rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 11 from beyond middle. Hindwings nearly 1, trapezoidal, termen slightly sinuate, cilia 2/3-1; with cubital pecten; 3 and 4 connate or short-stalked, 5 somewhat approximated, 6 and 7 closely approximated at base.

Remarks. — The imagoes are described as restless, turning in circles on leaves in shaded places, a habit probably shared by the nearly allied genera, which are characterised by bright blue-metallic or silvery markings, apparently adapted for display.

Geographical distribution of species. — American.

Larva unknown.

- | | |
|--|-------------------|
| 1. <i>S. argentifrons</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 81, pl. 3, f. 2 (1911). | Mexico. |
| 2. <i>S. iridipennella</i> , Clemens, Proc. Acad. Nat. Sc. Philad. p. 165 (1860).
<i>aphroditella</i> , Chambers, Canad. Ent. Vol. 4, p. 88 (1872). | E. United States. |
| 3. <i>S. proserpinella</i> , Frey, Stett. Ent. Zeit. Vol. 39, p. 251 (1878). | Missouri, Texas. |
| 4. <i>S. sapphiritis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 267 (1914). — PI. 5, Fig. 121a, b, c. | Guiana, Brazil. |
| 5. <i>S. spintheropsis</i> , Meyrick, ibidem, p. 101 (1922) — PI. 3, Fig. 57. | Brazil. |
| 6. <i>S. leucura</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 29, pl. 1, f. 25 (1910). | Mexico. |

152. GENUS HOLOPHYSIS, WALSHAM

Holophysis, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 29 (1910). — Type : *H. emblemella*, Clemens.

Characters. — Head smooth; ocelli moderate, posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint smooth-scaled, compressed, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae loosely haired above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia nearly 1; without cubital pecten; 3 and 4 connate, 5 rather approximated or nearly parallel, 6 and 7 connate or nearly approximated at base.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — American.

Larva unknown.

- | | |
|---|-------------------|
| 1. <i>H. tentatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 611 (1864). | Brazil. |
| 2. <i>H. barydesma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 142 (1918). | Ecuador. |
| 3. <i>H. autodesma</i> , Meyrick, ibidem, Vol. 2, p. 143 (1918). | Colombia. |
| 4. <i>H. auxiliaris</i> , Meyrick, ibidem, Vol. 2, p. 143 (1918). | Colombia. |
| 5. <i>H. anomia</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 30 (1910). | Mexico. |
| 6. <i>H. emblemella</i> , Clemens, Proc. Acad. Nat. Sc. Philad. p. 164 (1860).
<i>venustella</i> , Chambers, Canad. Ent. Vol. 4, p. 90 (1872). | E. United States. |
| 7. <i>H. stigmatophoria</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 30 (1910). | Mexico. |
| 8. <i>H. quadrimaculata</i> , Walsingham, ibidem, Vol. 4, p. 30, pl. 1, f. 26 (1910). | Mexico. |
| 9. <i>H. xanthostoma</i> , Walsingham, ibidem, Vol. 4, p. 31, pl. 1, f. 27 (1910). | Mexico. |

153. GENUS TRICYANAULA, NOV. GEN.

Type : *T. aurantiaca*, Walsingham.

Characters. — Head smooth; ocelli moderate, posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint strongly compressed, smooth-scaled, above with scales more or less roughly expanded posteriorly, terminal joint longer than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 11 from beyond middle. Hindwings slightly over 1, trapezoidal, termen slightly sinuate, cilia 1/2; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate.

Remarks. — Probably a derivative from *Zalithia*.

Geographical distribution of species. — Indian and African.

Larva (*amethystias*) feeding in the fungus-bed formed within the nest of *Termites*.

- | | |
|--|--------------------|
| 1. <i>T. metallica</i> , Walsingham, Trans. Ent. Soc. Lond. p. 97, pl. 4, f. 34 (1891). | Gambia, Transvaal. |
| 2. <i>T. aurantiaca</i> , Walsingham, Moore's Lep. Ceyl. Vol. 3, p. 518, pl. 209, f. 6 (1886). | Ceylon, S. India. |
| 3. <i>T. amethystias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 140 (1906). | Ceylon, S. India. |
| 4. <i>T. cyanozona</i> , Meyrick, Exot. Microlep. Vol. 3, p. 26 (1923). | Coorg. |
| 5. <i>T. augusta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 727 (1911). | Assam. |

154. GENUS ZALITHIA, MEYRICK

Zalithia, Meyrick, Trans. Ent. Soc. Lond. p. 18 (1894). — Type : *Z. uranopis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ moderately ciliated, basal joint elongate, without pecten. Labial palpi long, recurved, smooth, second joint rather thickened, terminal joint as long as second or shorter, slender or moderate, acute. Maxillary palpi very short, slender. Posterior tibiae shortly rough-scaled. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, oblong-ovate, cilia 2/5-2/3; with cubital pecten; 3 and 4 connate, 5 parallel, 6 and 7 closely approximated at base.

Remarks. — Probably a rather early form of this group.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

- | | |
|---|--------|
| 1. <i>Z. uranopis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 18 (1894). | Burma. |
| 2. <i>Z. doxarcha</i> , Meyrick, Exot. Microlep. Vol. 1, p. 578 (1916). | Burma. |

155. GENUS HYPERECTA, NOV. GEN.

Type : *H. enoptrias*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, strongly compressed, terminal joint as long as second, moderate or slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae shortly rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 6 to costa, 7 and 8 stalked, 11 from middle. Hindwings considerably over 1, trapezoidal, termen hardly sinuate, cilia 1/2; without cubital pecten; 3 and 4 connate, 5 parallel, 6 and 7 stalked.

Remarks. — Correlated with early forms of *Onebala*.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|---|--------|
| 1. <i>H. viridescens</i> , Meyrick, Exot. Microlep. Vol. 2, p. 143 (1918). | Assam. |
| 2. <i>H. enoptrias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 728 (1911). | Assam. |

156. GENUS SATRAPODOXA, NOV. GEN.

Type : *S. regia*, Meyrick.

Characters. — Head smooth, shining; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderate, without pecten. Labial palpi very long, recurved, smooth, slender, terminal joint nearly twice as long as second, acute. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above. Forewings with 2 and 3 stalked, 7 absent, 11 from middle. Hindwings somewhat over 1, trapezoidal,

termen slightly sinuate, cilia 3/5; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 connate.

Remarks. — A splendidly decorated insect, probably related to the two following genera.

Geographical distribution of species. — South American.

Larva unknown.

1. *S. regia*, Meyrick, Trans. Ent. Soc. Lond. p. 267 (1914).

Guiana, Peru.

157. GENUS EUZONOMACHA, NOV. GEN.

Type : *E. subjectella*, Walker.

Characters. — Head smooth, shining; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderate, without pecten. Labial palpi very long, recurved, smooth, slender, terminal joint much longer than second, acute. Maxillary palpi rudimentary. Posterior tibiae with appressed scales. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 2/3; without cubital pecten; 3 and 4 connate, 5 approximated, 6 and 7 connate.

Remarks. — Correlated with the following.

Geographical distribution of species. — South American.

Larva unknown.

1. *E. subjectella*, Walker, List. Lep. Het. Brit. Mus. Vol. 29, p. 611 (1864). Brazil, Peru.

158. GENUS CHARISTICA, NOV. GEN.

Type : *C. rhodopetala*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6 to almost 1, in ♂ sometimes rather thick, simple, basal joint elongate, without pecten. Labial palpi very long, recurved, slender, smooth, terminal joint longer than second, acute. Maxillary palpi rudimentary. Posterior tibiae shortly rough-scaled or almost smooth above. Forewings with 1 b furcate, 2 and 3 stalked, 6 to apex, 7 and 8 stalked, 7 to costa, or 7 absent, 11 from beyond middle. Hindwings considerably over 1, trapezoidal, termen hardly sinuate, cilia 1/2-3/4; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate.

Remarks. — These elegant and graceful insects are conspicuous from their strikingly polychromatic adornment, sometimes displaying all the colours of the rainbow.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. caeligena*, Meyrick, Trans. Ent. Soc. Lond. p. 101 (1922).

Brazil.

2. *C. sandaracota*, Meyrick, ibidem, p. 269 (1914).

Guiana.

3. *C. callichroma*, Meyrick, ibidem, p. 269 (1914). — Pl. 3, Fig. 58.

Guiana, Brazil, Peru.

4. *C. exteriorella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 610 (1864).

Brazil.

5. *C. iriantha*, Meyrick, Trans. Ent. Soc. Lond. p. 270 (1914).

Guiana, Peru.

6. *C. rhodopetala*, Meyrick, ibidem, p. 102 (1922).

Brazil.

7. *C. Walkeri*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 80 Panama, Colombia,
(1911).
Guiana, Brazil, Peru.
cuphracta, Meyrick, Trans. Ent. Soc. Lond. p. 268 (1914).
8. *C. porphyraspis*, Meyrick, ibidem, p. 23 (1909). Bolivia.
9. *C. ioploca*, Meyrick, ibidem, p. 103 (1922). Brazil.

159. GENUS CATOPTRISTIS, NOV. GEN.

Type *C. trissoxantha*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint somewhat thickened, smooth-scaled, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 3/5; with cubital pecten; 3 and 4 connate, 5 rather approximated, 6 and 7 connate.

Remarks. — Probably derived from the following.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. trissoxantha*, Meyrick, Trans. Ent. Soc. Lond. p. 100 (1922). Brazil, Peru.

160. GENUS CERYCANGELA, NOV. GEN.

Type : *C. sacricola*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint rather thickened, smooth-scaled, compressed, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae somewhat rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings rather over 1, trapezoidal, termen hardly sinuate, cilia 1/2; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate.

Remarks. — Perhaps correlated with *Onebala*.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. sacricola*, Meyrick, Trans. Ent. Soc. Lond. p. 102 (1922). Brazil, Peru.

161. GENUS HARPAGIDIA, RAGONOT

Harpagidia, Ragonot, Bull. Soc. Ent. Fr. p. 107 (1895). — Type : *H. pallidibasella*, Ragonot.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ stout, very shortly ciliated. Labial palpi very long, recurved, smooth-scaled, second joint somewhat thickened,

terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with appressed scales. Forewings with 2-4 parallel, curved, 5 and 6 straight, 7 absent. Hindwings over 1, trapezoidal, apex produced, termen emarginate; 3 and 4 connate, 5 remote, 6 from near angle, 7 from angle, to apex.

Remarks. — Of dubious affinity, but on general characters would be referable here.

Geographical distribution of species. — Asia Minor.

Larva unknown.

1. *H. pallidibasella*, Ragonot, Bull. Soc. Ent. Fr. p. 107 (1895). Asia Minor.

162. GENUS ZELOSYNE, WALSINGHAM

Zelosyne, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 50 (1911). — Type : *Z. poecilosoma*, Walsingham.

Characters. — Head smooth; ocelli very small, posterior; tongue developed. Antennae 3/4, in ♂ shortly ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint beneath with rough projecting scales whorled so as to form a series of acute teeth, terminal joint as long as second or somewhat longer, thickened with scales, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1 b furcate, 2 from 4/5, 3 from near angle, 6 to apex, 7 and 8 stalked or coincident, 11 from before middle. Hindwings under 1, trapezoidal, termen not sinuate, cilia over 1; without cubital pecten; 3 and 4 connate or closely approximated, 5 somewhat approximated, 6 and 7 separate, nearly parallel.

Remarks. — A remarkable genus of abnormal structure and facies; it may not belong here, but goes no better anywhere else. The two species differ in neuration, though extremely similar in superficial appearance; but Walsingham's description and diagram are incorrect in several particulars. The typical species possesses a beautiful coloured hyaline eye-like spot in the hindwings.

Geographical distribution of species. — Central and South America.

Larva unknown.

1. *Z. poecilosoma*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 51, Panama.
pl. 2, f. 11 (1911).
2. *Z. olga*, Meyrick, Trans. Ent. Soc. Lond. p. 201 (1915). — Pl. I, Fig. 22. Guiana.

163. GENUS TEUCHOPHANES, MEYRICK

Teuchophanes, Meyrick, Trans. Ent. Soc. Lond. p. 274 (1914). — Type : *T. leucopleura*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, shortly or strongly ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, compressed, terminal joint as long as second, with projecting scales posteriorly except near apex, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1 b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, apex obtuse, termen hardly

sinuate, cilia $1/2$; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated towards base.

Remarks. — Probably related to *Onebala*.

Geographical distribution of species. — Central and South America.

Larva unknown.

- | | |
|---|---------|
| 1. <i>T. cornuta</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 20 (1914). | Panama. |
| 2. <i>T. luminosa</i> , Busck, ibidem, Vol. 47, p. 19 (1914). | Panama. |
| 3. <i>T. perceptella</i> , Busck, ibidem, Vol. 47, p. 19 (1914). | Panama. |
| 4. <i>T. leucopleura</i> , Meyrick, Trans. Ent. Soc. Lond. p. 274 (1914). — Pl. 3, Guiana. | |

Fig. 59.

164. GENUS ATASTHALISTIS, MEYRICK

Atasthalistis, Meyrick, Trans. Ent. Soc. Lond. p. 279 (1886). — Type : *A. pyrocosma*, Meyrick.

Croesopola, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 410 (1904). — Type : *A. euchroa*, Lower.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae $3/4$, in ♂ serrulate, moderately ciliated, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint much thickened with dense scales, rather rough beneath and more or less roughly expanded above towards apex, terminal joint as long as second or shorter, thickened with dense slightly rough scales at least at base, apex slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 and 3 stalked or coincident, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia $1/6-2/5$; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated or stalked.

Remarks. — Correctly referred here, and probably correlated with *Onebala*. All the species are brightly coloured, usually with orange hindwings, the larger ones being amongst the most conspicuous of the *Tineina*.

Geographical distribution of species. — Polynesian and Malayan.

Larva unknown.

- | | |
|---|-------------------------------|
| 1. <i>A. gorgopa</i> , Meyrick, Exot. Microlep. Vol. 2, p. 115 (1918). | New Guinea. |
| 2. <i>A. concinalis</i> , Feisthamel, Mag. Zool. pl. 26, f. 2 (1839). | Moluccas. |
| 3. <i>A. gnophrina</i> , Felder, Reise Novara, Lep. pl. 139, f. 38 (1875). — Pl. 3, Fig. 74. | Ternate, Moluccas. |
| 4. <i>A. pyrocosma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 280 (1886). | New Guinea. |
| 5. <i>A. tricolor</i> , Felder, Reis. Novara Lep. pl. 139, f. 18 (1875). | Celebes, Solomon Islands. |
| 6. <i>A. hieropla</i> , Meyrick, Exot. Microlep. Vol. 2, p. 235 (1919). | Fiji. [Islands.] |
| 7. <i>A. euchroa</i> , Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 47 (1900). — Pl. 3, Fig. 75. | Queensland, Bismarck Islands. |
| <i>ochreoviridella</i> , Pagenstecher, Zoologica, Vol. 29, p. 236 (1900). | |
| 8. <i>A. viridella</i> , Snellen, Tijdschr. v. Ent. Vol. 44, p. 86, pl. 5, f. 10 (1901). | Java, Philippines. |

165. GENUS PERIORISTICA, WALSINGHAM

Perioristica, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 31 (1910). — Type : *P. chalcoptera*, Walsingham.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened, smooth-scaled, terminal joint about half second, slender, acute. Maxillary palpi very short, drooping. Posterior tibiae somewhat rough-haired above. Forewings with 1b furcate, 2 and 3 approximated from about angle, 7 and 8 stalked, 7 to costa, 11 from beyond middle. Hindwings slightly over 1, trapezoidal, termen hardly sinuate; 3 and 4 connate, 5 parallel, 6 and 7 approximated towards base.

Remarks. — Apparently allied to *Onebala*.

Geographical distribution of species. — North American.

Larva unknown.

1. *P. chalcoptera*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 32, Mexico.
pl. 1, f. 28 (1910).

166. GENUS SCHEMATASPIS, MEYRICK

Schemataspis, Meyrick, Exot. Microlep. Vol. 2, p. 144 (1918). — Type : *S. gradata*, Meyrick.

Characters. — Head smooth; ocelli moderate, posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint smooth-scaled, strongly compressed laterally, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 8 and 9 out of 7, 7 to apex, or seldom 8 and 9 stalked, 7 absent, 11 from middle. Hindwings 1 or somewhat over 1, trapezoidal, termen sinuate, cilia 1; without cubital pecten; 3 and 4 connate, 5 nearly approximated, 6 and 7 stalked.

Remarks. — A small group of very similar species, derived from *Onebala*.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

1. *S. epicentra*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 730 (1911). Ceylon.
2. *S. immeritella*, Walker, List. Lep. Brit. Mus. Vol. 29, p. 634 (1864). Ceylon, Java.
3. *S. gradata*, Meyrick, Rec. Ind. Mus. Vol. 5, p. 221 (1910). Assam.
4. *S. bicunea*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 731 (1911). Assam.
5. *S. rhabducha*, Meyrick, ibidem, Vol. 20, p. 730 (1911). S. India, Ceylon.

167. GENUS ONEBALA, WALKER

Onebala, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 792 (1864). — Type : *O. blandiella*, Walker.

Helcystogramma, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 371 (1877). — Type : *O. hibisci*, Stainton.

Dectobathra, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 299 (1904). — Type : *O. choristis*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, or seldom nearly 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, smooth, second joint thickened, laterally compressed, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings over 1,

trapezoidal, termen hardly sinuate, cilia 1/2-1; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate or closely approximated at base.

Remarks. — An early form of this group, indicating the connection between various more specialised genera; certain species have ribbed palpi such as are found in *Compsolechia*. The name *Onebala*, through wrong identification, has been misapplied by me to another genus.

Geographical distribution of species. — American, Indo-Malayan, Australian, and African, in warm regions only, but nowhere very numerous; one widely spread species reaches Europe.

Larva (4 known) feeding in rolled or spun leaves.

Foodplants : *Convolvulaceae* (2), *Malvaceae*, *Leguminosae*.

- | | |
|---|--|
| 1. <i>O. carycastis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 104 (1922). | Guiana, Brazil. |
| 2. <i>O. euargyra</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 120 (1919). | Queensland. |
| 3. <i>O. amethystina</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 300 (1904). | Queensland, New South Wales. |
| 4. <i>O. iridosoma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 144 (1918). | N. Queensland. |
| 5. <i>O. choristis</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 300 (1904). | E. & W. Australia. |
| 6. <i>O. zapyrodes</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 119 (1919). | Queensland. |
| 7. <i>O. gypsaspis</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 163 (1921). | Java. |
| 8. <i>O. infibulata</i> , Meyrick, Exot. Microlep. Vol. 1, p. 577 (1916). | India, Ceylon. |
| 9. <i>O. mercedella</i> , Walsingham, Proc. Zool. Soc. Lond. p. 934, pl. 11, f. 11 (1907). | Canaries. |
| 10. <i>O. thaumalea</i> , Walsingham, Ent. M. Mag. Vol. 41, p. 41 (1905). | N. Africa, Canaries. |
| 11. <i>O. lamprostoma</i> , Zeller, Isis, p. 851 (1847).
<i>sulu</i> , Walsingham, Trans. Ent. Soc. Lond. p. 261, pl. 12, f. 30 (1881).
<i>sentata</i> , Meyrick, ibidem, p. 14 (1894). | S. Europe, Canaries, N. & S. Africa, Asia Minor, India, Burma, Ceylon, Brazil. [Java.] Mexico. |
| 12. <i>O. chalyburga</i> , Meyrick, ibidem, p. 103 (1922). | |
| 13. <i>O. daedalea</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 100, pl. 3, f. 17 (1911). | |
| 14. <i>O. leucoplecta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 729 (1911). | S. India, Ceylon, Java. |
| 15. <i>O. brabylitis</i> , Meyrick, ibidem, Vol. 20, p. 729 (1911). | S. India, Java. |
| 16. <i>O. armata</i> , Meyrick, ibidem, Vol. 20, p. 728 (1911). | Assam. |
| 17. <i>O. hoplophora</i> , Meyrick, Exot. Microlep. Vol. 1, p. 577 (1911). | India, Burma, Ceylon. |
| 18. <i>O. victrix</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 727 (1911). | Coorg. |
| 19. <i>O. lithostrota</i> , Meyrick, Exot. Microlep. Vol. 1, p. 578 (1916). | Malay States. |
| 20. <i>O. balteata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 732 (1911). | Assam. |
| 21. <i>O. septella</i> , Zeller, Micr. Caffr. p. 108 (1852).
<i>digitata</i> , Meyrick, Exot. Microlep. Vol. 1, p. 200 (1914). | Nyassaland, Natal. |
| 22. <i>O. hibisci</i> , Stainton, Trans. Ent. Soc. Lond. (2) Vol. 5, p. 117 (1859).
<i>obseratella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 371, pl. 5, f. 127 (1877).
<i>eudela</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 160 (1919). | India, Ceylon, Java, N. Queensland. |
| 23. <i>O. blandiella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 792 (1864). | S. India, Ceylon. |
| 24. <i>O. ribbeella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 369, pl. 5, f. 126 (1877). | Panama. |
| 25. <i>O. symbolica</i> , Meyrick, Trans. Ent. Soc. Lond. p. 270 (1914). — PI. 3, Fig. 58. | Guiana. |
| 26. <i>O. adaequata</i> , Meyrick, ibidem, p. 271 (1914). | Guiana. |
| 27. <i>O. rusticella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 621 (1864). | Brazil. |
| 28. <i>O. sertigera</i> , Meyrick, Exot. Microlep. Vol. 3, p. 27 (1923). | Peru. |
| 29. <i>O. stellatella</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 20 (1914). | Panama. |

30. *O. alacella*, Clemens, Proc. Ent. Soc. Philad. Vol. 1, p. 132 (1862). E. United States, Canada.
ochripalpella, Zeller, Verh. Zool.-bot. Ges. Wien. Vol. 23, p. 279 (1873).
goodellella, Chambers, Journ. Cinc. Soc. Nat. Hist. Vol. 3, p. 289 (1881).
31. *O. tegulella*, Walsingham, Proc. Zool. Soc. Lond. p. 83 (1897). Windward Islands,
servilis, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 103, pl. 3, Panama, Guiana.
f. 25 (1911).
32. *O. meconitis*, Meyrick, Trans. Ent. Soc. Lond. p. 176 (1913). Argentina.

168. GENUS MYCONITA, MEYRICK

Myconita, Meyrick, Exot. Microlep. Vol. 3, p. 27 (1923). — Type : *M. plutelliformis*, Snellen.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint much thickened with dense appressed scales somewhat expanded at apex above, laterally compressed, terminal joint shorter than second, moderate, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia 2/3; without cubital pecten; 3 and 4 connate, 5 approximated at base, 6 and 7 stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Indo-Malayan and Australian.

Larva feeding in rolled leaves.

Foodplant *Ipomoea* (*Convolvulaceae*).

1. *M. plutelliformis*, Snellen, Tijdschr. v. Ent. Vol. 44, p. 84, pl. 6, f. 4 (1901). India, Ceylon, Java, Queensland.

169. GENUS THELYASCETA, MEYRICK

Thelyasceta, Meyrick, Exot. Microlep. Vol. 3, p. 27 (1923). — Type : *T. nonstrigella*, Chambers.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ moderately ciliated, in ♀ thickened with dense rough projecting scales above on basal 2/3, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, laterally compressed, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae in ♂ smooth, in ♀ hairy above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings slightly over 1, trapezoidal, termen slightly sinuate, cilia 3/4; without cubital pecten; 3 and 4 connate, 5 slightly approximated, 6 and 7 connate.

Remarks. — A distinct early form of this group.

Geographical distribution of species. — North American.

Larva feeding in spun leaves.

Foodplant *Aster* (*Compositae*).

1. *T. nonstrigella*, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 92 (1878). E. United States, Canada.
purpureofusca, Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 184 (1882).

170. GENUS PROSTOMEUS, BUSCK

Prostomeus, Busck, Proc. U. S. Mus. Vol. 25, p. 837 (1903). — Type : *P. brunneus*, Busck.

Characters. — Head with appressed scales; tongue developed. Antennae 3/4, in ♂ simple. Labial palpi long, recurved, strongly laterally compressed, sharp-edged anteriorly, second joint thickened with scales, truncate, terminal joint as long as second, thickened with smooth appressed scales terminating abruptly just below acute apex, sharp-edged posteriorly. Maxillary palpi obsolete. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — I am not acquainted with this genus, supposed to be characterised by the peculiar palpi; it is possibly related to *Teuchophanes*.

Geographical distribution of species. — North American.

Larva unknown.

1. *P. brunneus*, Busck, Proc. U. S. Mus. Vol. 25, p. 838 (1903).

Florida.

171. GENUS AEOLOTROCHA, MEYRICK

Aeolotrocha, Meyrick, Ann. Transv. Mus. Vol. 8, p. 78 (1921). — Type : *A. generosa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ serrulate, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint considerably thickened with appressed scales, strongly compressed, terminal joint rather shorter than second, moderate, acute. Maxillary palpi obsolete. Posterior tibiae clothed with rough hairs above. Forewings with 1b furcate, 2 from near angle, 4 and 5 approximated at base, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen faintly sinuate, cilia 4/5; 3 and 4 connate, 5 nearly parallel, 6 and 7 stalked.

Remarks. — Apparently an early form of this group, but differing in the terminal ending of vein 7 of forewings.

Geographical distribution of species. — South African.

Larva unknown.

1. *A. generosa*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 78 (1921).

Natal.

172. GENUS DECATOPSEUSTIS, NOV. GEN.

Type : *D. xanthastis*, Lower.

Characters. — Head smooth, sidetuffs slightly raised; ocelli posterior; tongue developed. Antennae 3/4, in ♂ minutely ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint somewhat thickened with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, in ♂ 10 absent, 11 from towards base, with a subhyaline streak beneath it, in ♀ 10 present, 11 from middle. Hindwings 1, trapezoidal,

apex produced, termen emarginate, cilia 1; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 connate.

Remarks. — Quite a peculiar genus, whose relationships must be considered very doubtful.

Geographical distribution of species. — Australian.

Larva unknown.

1. *D. xanthastis*, Lower, Trans. Roy. Soc. S. Australia, Vol. 20, p. 168 (1896). Queensland.

173. GENUS AREGHA, CHRÉTIEN

Aregha, Chrétien, Ann. Soc. Ent. Fr. Vol. 84, p. 333 (1915). — Type : *A. abhaustella*, Chrétien.

Characters. — Head smooth; tongue obsolete. Antennae 4/5, in ♂ stout, simple. Labial palpi long, recurved, slender, second joint with appressed scales, terminal joint as long as second, acute. Maxillary palpi obsolete. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1, elongate-trapezoidal, termen slightly sinuate; 3 and 4 connate, 5 approximated, 6 and 7 approximated towards base.

Remarks. — Unknown to me; considered by its author to be near *Epidola*, but on the hindwings this is unlikely.

Geographical distribution of species. — North African.

Larva unknown.

1. *A. abhaustella*, Chrétien, Ann. Soc. Ent. Fr. p. 334 (1915). Algeria.

174. GENUS LEOBATUS, WALSINGHAM

Leobatus, Walsingham, Ent. M. Mag. Vol. 40, p. 220 (1904). — Type : *L. fagoniae*, Walsingham.

Characters. — Head smooth; tongue developed. Antennae 2/3, in ♂ shortly ciliated, basal joint without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint as long as second, pointed. Maxillary palpi very short. Posterior tibiae clothed with rough hairs. Forewings with 2 separate, 6 and 7 out of 8, 7 to costa. Hindwings slightly over 1, ovate-trapezoidal, termen slightly sinuate, cilia 1; 3 and 4 connate, 5 hardly approximated, 6 and 7 stalked.

Remarks. — Also unknown to me.

Geographical distribution of species. — North African.

Larva feeding in loose web amongst trailing stems.

Foodplant *Fagonia* (*Zygophyllaceae*).

1. *L. fagoniae*, Walsingham, Ent. M. Mag. Vol. 40, p. 221 (1904). Algeria.

175. GENUS ACOMPSSIA, HÜBNER

Acompssia, Hübner, Verz. bek. Schmett. p. 409 (1826) — Type : *A. cinerella*, Clerck.

Brachycrossata, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 323 (1870). — Type : *A. cinerella*, Clerck.

Cathegesis, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 27 (1910). — Type : *A. vinitincta*, Walsingham.

Oxyptoryx, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 61, p. (151) (1911). — Type : *A. jordanella*, Rebel.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 3/4. in ♂ very shortly ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint rather thickened with appressed scales, terminal joint as long as second or somewhat shorter, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, separate, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen faintly sinuate, cilia 1/2-4/5; without cubital pecten; 3 and 4 connate or short-stalked, 5 rather approximated, 6 and 7 connate or approximated towards base. Wings in ♀ sometimes abbreviated.*

Remarks. — The simple and unspecialised characters of this genus cause its precise affinity to be rather indefinite.

Geographical distribution of species. — European, extending into America and Africa.

Larva (2 known) feeding in spun leaves.

Foodplants *Plantaginaceae*, *Scrophulariaceae*.

- | | |
|---|---|
| 1. <i>A. jordanella</i> , Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 61, p. (151) (1911). | Palestine. |
| 2. <i>A. vinitincta</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 27 (1910). | Mexico. |
| 3. <i>A. oenochyta</i> , Meyrick, Ann. Trans. Mus. Vol. 8, p. 78 (1921). | Rhodesia. |
| 4. <i>A. formosella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 160 (1796). — | C. & S. Europe, Asia Minor, Syria, N. Africa. |
| Pl. 3, Fig. 60. | |
| <i>flammella</i> , Treitschke, Schmett. Eur. Vol. 9 (2), p. 20 (1833). | |
| 5. <i>A. flavella</i> , Duponchel, Hist. Nat. Lép. Fr. Suppl. Vol. 4, pl. 89, f. 7 (1842). | S. Europe, Asia Minor, Syria, N. Africa. |
| <i>segetella</i> , Zeller, Isis, p. 847 (1847). | |
| 6. <i>A. pallidipulchra</i> , Walsingham, Ent. M. Mag. Vol. 40, p. 269 (1904). | Algeria, Tunis, Egypt. |
| 7. <i>A. psoricopterella</i> , Walsingham, Proc. Zool. Soc. Lond. p. 523 (1891). | Windward Islands. |
| 8. <i>A. angulifera</i> , Walsingham, ibidem, p. 82 (1897). | Virgin & Windward Rhodesia. [Islands. |
| 9. <i>A. sphenopis</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 79 (1921). | Spain. |
| 10. <i>A. dimorpha</i> , Petry, Iris, Vol. 17, p. 4 (1904). | S. France. |
| 11. <i>A. antirrhinella</i> , Millière, Icon. Descr. Lép. Vol. 2, p. 274, pl. 80, f. 6-8 (1868). | Alps, France, Galicia. |
| 12. <i>A. tripunctella</i> , Schiffermüller, Syst. Verz. Schmett. Wien, p. 319 (1776). | Bohemia, Austria, Tyrol. |
| <i>maculosella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 202, f. 463 (1855). | Europe, Asia Minor. |
| 13. <i>A. minorella</i> , Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 49, p. 180 (1899). | |
| 14. <i>A. cinerella</i> , Clerck, Icon. Ins. pl. 11, fig. 6 (1760). | |
| <i>ardelliella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 437 (1818). | |
| <i>spodiella</i> , Treitschke, Schmett. Eur. Vol. 9 (2), p. 78 (1833). | |
| 15. <i>A. labradorica</i> , Möschler, Wien. Ent. Monatsschr. Vol. 8, p. 200, pl. 5, f. 17 (1864). | Labrador. |

Group 5 (Protolechia type)

The species of this group have the general aspect of *Gelechia*, though tending to be broader-winged, especially in the earlier forms, but are usually characterised by having veins 2 and 3 of the forewings stalked, and 6 and 7 of hindwings parallel. Vein 7 of forewings when present always runs to costa (except in *Phloeographis* to apex); 3 and 4 of hindwings are always connate, and the cubital pecten is absent. The group is almost entirely confined to Australia, where it must have been developed in seclusion.

176. GENUS METEORISTIS, MEYRICK

Meteoristis, Meyrick, Exot. Microlep. Vol. 3, p. 27 (1923). — Type : *M. religiosa*, Meyrick.

Characters. — Head with appressed scales, sidetufts raised; ocelli posterior; tongue developed. Antennae 2/3, in ♂ minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint much thickened with appressed scales, somewhat roughened beneath towards apex, terminal joint much shorter than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1^b furcate, 2 and 3 out of 4, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, narrowed posteriorly, termen slightly sinuate, cilia 3/4; 3 and 4 stalked, 5 nearly approximated at base, 6 and 7 connate.

Remarks. — Seems best regarded as a rather aberrant straggler of this group.

Geographical distribution of species. — Indian.

Larva boring in aerial roots.

Foodplant *Ficus religiosa*.

1. *M. religiosa*, Meyrick, Exot. Microlep. Vol. 3, p. 28 (1923). Bengal.

177. GENUS TRITADELPHA, MEYRICK

Tritadelpha, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 323 (1904). — Type : *T. microptila*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint with scales somewhat rough beneath, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1^b furcate, 2 and 3 coincident, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex pointed, termen sinuate, cilia 2; 3 and 4 connate, 5 somewhat approximated to 4, 6 and 7 approximated.

Remarks. — This and the next genus are superficially extremely similar; they are correlated derivatives of *Protolechia*. I have amended my original interpretation of the neuration.

Geographical distribution of species. — Australian.

Larva unknown.

1. *T. microptila*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 323 (1904). Queensland.

178. GENUS EPIBRONTIS, MEYRICK

Epibrontis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 324 (1904). — Type : *E. hemichlaena*, Lower.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint with scales somewhat roughened beneath, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1^b furcate,

2 and 3 connate, 6 to apex, 7 absent, 11 from middle. Hindwings 1, trapezoidal, apex pointed, termen slightly sinuate, cilia 1 1/4; 3 and 4 connate, 5 somewhat approximated to 4, 6 and 7 tolerably parallel.

Remarks. — These small white and black insects possibly mimic the excrement of birds.

Geographical distribution of species. — Australian.

Larva unknown.

1. *E. hemichlaena*, Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 55 (1897). E. Australia, Tasmania.
2. *E. pallacopa*, Meyrick, Exot. Microlep. Vol. 2, p. 502 (1922). Victoria.

179. GENUS EPIMIMASTIS, MEYRICK

Epimimastis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 325 (1904). — Type : *E. porphyroloma*, Lower.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with scales, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked or connate, 7 absent, 11 from middle. Hindwings 1, or over 1, rounded-trapezoidal, apex somewhat pointed, termen sinuate, cilia 2/5-1; 3 and 4 connate, 5 approximated, 6 and 7 tolerably parallel.

Remarks. — A simple derivative of *Protolechia*.

Geographical distribution of species. — Australian and Indo-Malayan. This genus, an undoubted member of the group, is the only one that appears to be well established outside Australia. Note that the natural order of the species agrees with their geographical order.

Larva unknown.

1. *E. escharitis*, Meyrick, Exot. Microlep. Vol. 1, p. 589 (1916). Ceylon.
2. *E. glaucodes*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 461 (1910). Ceylon.
3. *E. emblematica*, Meyrick, Exot. Microlep. Vol. 1, p. 589 (1916). Assam, Borneo.
4. *E. calopta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 127 (1919). Queensland.
5. *E. tegminata*, Meyrick, Exot. Microlep. Vol. 1, p. 590 (1916). Queensland.
6. *E. porphyroloma*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 22, p. 22 (1897). E. Australia, Tasmania.

180. GENUS CRASPEDOTIS, MEYRICK

Craspedotis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 326 (1904). — Type : *C. pragmatica*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with scales somewhat roughened beneath, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 connate or stalked, 6 and 8 stalked, 7 absent, 11 from middle. Hindwings 1, rounded-trapezoidal, apex pointed, termen slightly sinuate, cilia 1; 3 and 4 connate, 5 approximated to 4, 6 and 7 approximated or stalked.

Remarks. — A development of *Sphaleractis*, to which it is very similar.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|---|------------------|
| 1. <i>C. soloeca</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 326 (1904). | New South Wales. |
| 2. <i>C. pragmatica</i> , Meyrick, ibidem, Vol. 29, p. 327 (1904). | New South Wales. |
| 3. <i>C. thinodes</i> , Meyrick, ibidem, Vol. 29, p. 327 (1904). | E. Australia. |
| 4. <i>C. diasticha</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 129 (1919). | New South Wales. |

181. GENUS SPHALERACTIS, MEYRICK

Sphaleractis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 328 (1904). — Type: *S. platyleuca*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, rough beneath, terminal joint shorter than second, thickened with scales and somewhat roughened anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 11 from middle. Hindwings over 1, rounded-trapezoidal, apex round-pointed, termen sinuate, cilia 4/5; 3 and 4 connate, 5 somewhat approximated to 4, 6 and 7 tolerably parallel.

Remarks. — A development from *Protolechia*.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|--|-------------------------------|
| 1. <i>S. parasticta</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 328 (1904). — PI. 3, Fig. 61. | E. & W. Australia. |
| 2. <i>S. platyleuca</i> , Lower, ibidem, Vol. 22, p. 22 (1897). | S. E. Australia, Tasmania. |
| 3. <i>S. eurysema</i> , Meyrick, ibidem, Vol. 29, p. 329 (1904). | New South Wales. |
| 4. <i>S. epiclysta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 299 (1920). | New South Wales,
Tasmania. |

182. GENUS LEXIARCHA, MEYRICK

Lexiarcha, Meyrick, Exot. Microlep. Vol. 1, p. 590 (1916). — Type: *L. galactopa*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough scales above. Forewings with 1b furcate, 2 and 3 connate, 7 and 8 out of 6, 7 to costa. 11 from middle. Hindwings somewhat over 1, trapezoidal, apex pointed, termen somewhat sinuate, cilia 1; 3 and 4 connate, 5 curved, approximated, 6 and 7 remote, parallel, transverse vein very oblique from 6 to 7.

Remarks. — A development of *Protolechia*.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|--|---------------|
| 1. <i>L. galactopa</i> , Meyrick, Exot. Microlep. Vol. 1, p. 590 (1916). | N. Australia. |
|--|---------------|

183. GENUS PRODOSIARCHA, MEYRICK

Prodosiarcha, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 330 (1904). — Type : *P. loxodesma*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with scales, somewhat rough beneath, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 coincident, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, apex rounded, termen not sinuate, cilia 3/4; 3 and 4 connate, 5 parallel, 6 and 7 nearly parallel.

Remarks. — A derivative of *Protolechia*.

Geographical distribution of species. — Australian.

Larva unknown.

1. *P. glagera*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 129 (1919). N. Queensland.
2. *P. loxodesma*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 330 (1904). S. Australia.
3. *P. thanatodes*, Lower, Trans. Roy. Soc. S. Australia, Vol. 17, p. 170 (1893). S. Australia.

184. GENUS CORYNAEA, TURNER

Corynaea, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 129 (1919). — Type : *C. dilechia*, Turner.

Characters. — Head smooth; tongue developed. Antennae 3/4, in ♂ simple, basal joint without pecten. Labial palpi long, recurved, second joint greatly thickened towards apex with long appressed hairs, terminal joint half second, slender, acute. Maxillary palpi very short. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to costa. Hindwings over 1, trapezoidal, apex pointed, termen scarcely sinuate; 3 and 4 stalked, 5 approximated, 6 and 7 stalked.

Remarks. — Not known to me.

Geographical distribution of species. — Australian.

Larva unknown.

1. *C. dilechia*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 130 (1919). N. Queensland.

185. GENUS HEMIARCHA, MEYRICK

Hemiarcha, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 331 (1904). — Type : *H. thermochroa*, Lower.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/5, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with scales, somewhat roughened beneath, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings 1b furcate, 2 and 3 stalked, 7 and 8 stalked, or sometimes out of 6, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, apex rounded, termen hardly sinuate, cilia 2/3; 3 and 4 connate, 5 somewhat approximated to 4, 6 and 7 stalked.

Remarks. — A derivative of *Protolechia*.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|--|-------------------------|
| 1. <i>H. macroplaca</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 17, p. 170 (1893). | Victoria. |
| 2. <i>H. bleptodes</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 132 (1919). | Queensland, New South |
| 3. <i>H. tetrasticta</i> , Turner, ibidem, Vol. 31, p. 133 (1919). | Queensland. [Wales. |
| 4. <i>H. thermochroa</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 17, p. 169 (1893). | Victoria, S. Australia. |
| 5. <i>H. polioleuca</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 133 (1919). | Queensland. |
| 6. <i>H. caliginosa</i> , Turner, ibidem, Vol. 31, p. 133 (1919). | Queensland. |

186. GENUS SEMOCHARISTA, MEYRICK

Semocharista, Meyrick, Ark. f. Zool. Vol. 14, no. 15, p. 4 (1922). — Type : *S. idiospila*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint nearly as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal-ovate, termen slightly sinuate, cilia 3/5; 3 and 4 connate, 5 parallel, 6 absent, 7 to apex, 8 anastomosing with central point of upper margin of cell.

Remarks. — A modification of *Protolechia*.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|---|------------------|
| 1. <i>S. idiospila</i> , Meyrick, Ark. f. Zool. Vol. 14, no. 15, p. 4 (1922). | N. W. Australia. |
|---|------------------|

187. GENUS PROTOLECHIA, MEYRICK

Protolechia, Meyrick, Ent. M. Mag. Vol. 39, p. 291 (1903). — Type : *P. mesochra*, Lower.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint more or less thickened with appressed scales, loose or somewhat rough beneath, terminal joint as long as second or shorter, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 connate or stalked, or seldom nearly approximated and then 2 curved from angle, 7 and 8 stalked, seldom 6 out of 7 near base, 7 to costa, 11 from middle. Hindwings 1 or over 1, trapezoidal-ovate, apex obtuse or pointed, termen sometimes sinuate, cilia 3/5-4/5; 3 and 4 connate, 5 rather approximated to 4, 6 and 7 nearly parallel, rarely 8 anastomosing shortly with upper margin of cell.

Remarks. — This characteristic genus may be regarded as originating from a form of the *Brachmia* group. The numerous species are in general obscurely coloured and adapted to concealment, especially perhaps on tree-trunks.

Geographical distribution of species. — Exclusively Australian.

Larva (7 known) usually feeding in spun leaves, but one mining blotches in leaves.

Foodplants : *Myrtaceae* (5), *Sapindaceae*, *Rutaceae*; judging from habits, a large number of the species are attached to particular species of *Eucalyptus*.

1. *P. tetraploa*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 338 (1904). Victoria.
2. *P. telopis*, Meyrick, ibidem, Vol. 29, p. 338 (1904). New South Wales.
3. *P. desmata*, Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 56 (1897). S. E. Australia.
4. *P. hilara*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 135 (1919). Victoria.
5. *P. exarista*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 339 (1904). W. Australia. [toria.]
6. *P. acroleuca*, Meyrick, ibidem, Vol. 29, p. 340 (1904). New South Wales, Vic-
7. *P. caminopsis*, Meyrick, ibidem, Vol. 29, p. 340 (1904). New South Wales.
8. *P. temenitis*, Meyrick, ibidem, Vol. 29, p. 341 (1904). Queensland.
9. *P. voluta*, Meyrick, ibidem, Vol. 29, p. 341 (1904). New South Wales.
10. *P. pacifica*, Meyrick, ibidem, Vol. 29, p. 342 (1904). S. E. & W. Australia.
- leptoictica*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 136 (1919).
11. *P. trachyphanes*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 343 (1904). W. Australia.
12. *P. prisca*, Meyrick, ibidem, Vol. 29, p. 343 (1904). New South Wales.
13. *P. diplonesa*, Meyrick, ibidem, Vol. 29, p. 344 (1904). W. Australia.
14. *P. classopis*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 136 (1919). Queensland.
15. *P. crypsibatis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 344 (1904). S. E. Australia, Tasmania.
16. *P. flexilis*, Meyrick, ibidem, Vol. 29, p. 345 (1904). New South Wales.
17. *P. frugalis*, Meyrick, ibidem, Vol. 29, p. 345 (1904). New South Wales, W. Australia.
18. *P. microdora*, Meyrick, ibidem, Vol. 29, p. 346 (1904). W. Australia.
19. *P. invalida*, Meyrick, ibidem, Vol. 29, p. 346 (1904). Queensland, New South Tasmania. [Wales.]
20. *P. cladara*, Meyrick, ibidem, Vol. 29, p. 346 (1904). Tasmania.
21. *P. sciodes*, Meyrick, ibidem, Vol. 29, p. 347 (1904). W. Australia.
22. *P. euryanya*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 135 (1919). Queensland.
23. *P. trichosema*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 347 (1904). W. Australia.
24. *P. aclera*, Meyrick, ibidem, Vol. 29, p. 348 (1904). South Australia.
25. *P. autopis*, Meyrick, ibidem, Vol. 29, p. 348 (1904). W. Australia.
26. *P. elpistis*, Meyrick, ibidem, Vol. 29, p. 349 (1904). W. Australia.
27. *P. lithina*, Lower, ibidem, Vol. 24, p. 98 (1899). New South Wales.
28. *P. argocentra*, Meyrick, ibidem, Vol. 29, p. 350 (1904). W. Australia.
29. *P. liota*, Meyrick, ibidem, Vol. 29, p. 350 (1904). W. Australia.
30. *P. hypoleuca*, Meyrick, ibidem, Vol. 29, p. 351 (1904). New South Wales.
31. *P. cosmotis*, Meyrick, ibidem, Vol. 29, p. 351 (1904). W. Australia.
32. *P. psephias*, Meyrick, ibidem, Vol. 29, p. 352 (1904). W. Australia.
33. *P. actinota*, Meyrick, ibidem, Vol. 29, p. 352 (1904). W. Australia.
34. *P. mitophora*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 138 (1919). Queensland.
35. *P. creperrima*, Turner, ibidem, Vol. 31, p. 137 (1919). Queensland.
36. *P. nyctias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 353 (1904). Queensland.
37. *P. hypocneca*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 137 (1919). Queensland.
38. *P. melicrata*, Turner, ibidem, Vol. 31, p. 137 (1919). N. Queensland.
39. *P. iochlaena*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 353 (1904). W. Australia.
40. *P. xestolitha*, Meyrick, ibidem, Vol. 29, p. 354 (1904). New South Wales.
41. *P. nephelota*, Meyrick, ibidem, Vol. 29, p. 354 (1904). E. Australia.
42. *P. anthracina*, Meyrick, ibidem, Vol. 29, p. 355 (1904). New South Wales.
43. *P. phloeodes*, Meyrick, ibidem, Vol. 29, p. 355 (1904). Queensland.
44. *P. thyridota*, Meyrick, ibidem, Vol. 29, p. 355 (1904). New South Wales, W. Australia.
45. *P. sarisias*, Meyrick, ibidem, Vol. 29, p. 356 (1904). Queensland.

46. *P. enchotypha*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 138 (1919).
 47. *P. aversella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 649 (1864).
 48. *P. penthicodes*, Meyrick, Exot. Microlep. Vol. 2, p. 427 (1921).
 49. *P. englypta*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 357 (1904).
 50. *P. bistrigata*, Meyrick, Exot. Microlep. Vol. 2, p. 427 (1921).
 51. *P. obeliscota*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 358 (1904).
 52. *P. sisyraea*, Meyrick, ibidem, Vol. 29, p. 359 (1904).
 53. *P. phloeopola*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 140 (1919).
 54. *P. chenias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 359 (1904).
 55. *P. mechanitis*, Meyrick, ibidem, Vol. 29, p. 360 (1904).
 56. *P. furcifera*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 139 (1919).
 57. *P. xuthias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 361 (1904).
 58. *P. chalazodes*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 140 (1919).
 59. *P. vacatella*, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1026 (1864).
 60. *P. orthanotis*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 50 (1900).
 61. *P. pyrrhica*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 141 (1919).
 62. *P. trimetropis*, Meyrick, Ark. f. Zool. Vol. 14, no 15, p. 5 (1922).
 63. *P. ceramica*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 362 (1904). — **Pl. 3, Fig. 62.**
64. *P. hormodes*, Meyrick, ibidem, Vol. 29, p. 363 (1904).
 65. *P. plinthactis*, Meyrick, ibidem, Vol. 29, p. 363 (1904).
 66. *P. decaspila*, Lower, ibidem, Vol. 24, p. 99 (1899).
 67. *P. mesochra*, Lower, Trans. Roy. Soc. S. Australia, Vol. 18, p. 107 (1894).
 68. *P. stratifera*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 366 (1904).
 69. *P. catarrhacta*, Meyrick, ibidem, Vol. 29, p. 366 (1904).
 70. *P. tabulata*, Meyrick, ibidem, Vol. 29, p. 367 (1904).
 71. *P. pelogramma*, Meyrick, ibidem, Vol. 29, p. 367 (1904).
 72. *P. amblopis*, Meyrick, ibidem, Vol. 29, p. 368 (1904).
 73. *P. aeolopis*, Meyrick, ibidem, Vol. 29, p. 369 (1904).
 74. *P. hylas*, Meyrick, ibidem, Vol. 29, p. 369 (1904).
 75. *P. compsochroa*, Meyrick, ibidem, Vol. 29, p. 370 (1904).
 76. *P. odorifera*, Meyrick, ibidem, Vol. 29, p. 371 (1904).
 77. *P. secta*, Meyrick, Exot. Microlep. Vol. 2, p. 428 (1921).
 78. *P. acricula*, Meyrick, ibidem, Vol. 1, p. 198 (1914).
 79. *P. cryptosperma*, Meyrick, ibidem, Vol. 2, p. 428 (1921).
 80. *P. micropa*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 371 (1904).
 81. *P. deltodes*, Lower, Trans. Roy. Soc. S. Australia, Vol. 20, p. 169 (1896).
 82. *P. diplanetis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 373 (1904).
 83. *P. involuta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 142 (1919).
 84. *P. cephalota*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 373 (1904).
 85. *P. scytina*, Meyrick, ibidem, Vol. 29, p. 374 (1904).
 86. *P. trochias*, Meyrick, Exot. Microlep. Vol. 2, p. 428 (1921).
 87. *P. megalosticta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 141 (1919).
 88. *P. ananeura*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 374 (1904).
 89. *P. gypsocrana*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 143 (1919).
 90. *P. blacica*, Turner, ibidem, Vol. 31, p. 141 (1919).
 91. *P. albifrons*, Turner, ibidem, Vol. 31, p. 142 (1919).
 92. *P. haemaspila*, Lower, Trans. Roy. Soc. S. Australia, Vol. 18, p. 107 (1894).
 nana, Lower, ibidem, Vol. 18, p. 107 (1894).
 93. *P. tridecta*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 48 (1900).
 94. *P. xanthocephala*, Meyrick, ibidem, Vol. 29, p. 376 (1904). — **Pl. 3, Fig. 63; Pl. 5, Fig. 120.**
 95. *P. trichalina*, Meyrick, ibidem, Vol. 29, p. 377 (1914).
 96. *P. erudita*, Meyrick, Exot. Microlep. Vol. 1, p. 591 (1916).
- Victoria.
 E. & W. Australia.
 Queensland.
 Victoria.
 Queensland.
 Victoria.
 Victoria.
 Victoria.
 Victoria.
 New South Wales,
 Tasmania. [Victoria.
 Victoria.
 South Australia.
 Queensland.
 Queensland.
 Victoria.
 Queensland.
 Victoria.
 N. W. Australia.
 New South Wales.
 New South Wales.
 New South Wales.
 Victoria.
 E. Australia.
 E. Australia.
 New South Wales.
 S. E. Australia.
 Victoria.
 W. Australia.
 New South Wales.
 Queensland, New South
 S. E. Australia. [Wales.
 Queensland, New South
 Queensland. [Wales.
 New South Wales.
 Victoria. [lia.
 Victoria, South Austra-
 N. South Wales, Victoria.
 N. South Wales, Victoria.
 N. Queensland.
 W. Australia.
 S. E. Australia.
 Queensland.
 New South Wales.
 New South Wales.
 Queensland.
 Queensland.
 N. Australia.
 S. E. Australia.
 South Australia.
 Queensland, New South
 Wales.
 W. Australia.
 Victoria.

97. *P. arganthes*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 377 (1904). Queensland.
 98. *P. selenia*, Meyrick, ibidem, Vol. 29, p. 378 (1904). New South Wales.
 99. *P. sodalella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 646 (1864). Queensland, New South
 100. *P. gorgonias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 380 (1904). Queensland. [Wales.
 101. *P. celidophora*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 145 (1919). Queensland.
 102. *P. chiradia*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 380 (1904). Queensland.
 103. *P. hedana*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 145 (1919). Queensland.
 104. *P. thyrsopelta*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 381 (1904). Queensland.
 105. *P. annularia*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 143 (1919). Queensland.
 106. *P. phasianis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 382 (1904). S.E. Australia,
 Tasmania.
 107. *P. taracta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 143 (1919). Queensland.
 108. *P. eustephana*, Turner, ibidem, Vol. 31, p. 146 (1919). Queensland.
 109. *P. nothrodes*, Meyrick, Exot. Microlep. Vol. 2, p. 429 (1921). Queensland.
 110. *P. eumela*, Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 59 (1897). S.E. Australia, Tasmania.
 111. *P. semigrapha*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 146 (1919). Queensland.
 112. *P. loemias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 383 (1904). New South Wales,
 Victoria.
 113. *P. proscripta*, Meyrick, Exot. Microlep. Vol. 2, p. 429 (1921). Queensland.
 114. *P. megalommata*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 384 (1904). Queensland.
 115. *P. banausodes*, Meyrick, ibidem, Vol. 29, p. 384 (1904). Queensland.
 116. *P. mesopsamma*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 144 (1919). Queensland.
 117. *P. crotalodes*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 385 (1904). Queensland.
 118. *P. sporodeta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 144 (1919). Queensland.
 119. *P. molynnis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 385 (1904). Victoria, South Australia.
 120. *P. lechriosema*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 147 (1919). Queensland.
 121. *P. aspetodes*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 386 (1904). Victoria.
 122. *P. subnexella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 645 (1864). Queensland, New South
 bistellella, Walker, ibidem, Vol. 30, p. 1025 (1864). Wales.
 monoleuca, Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 55 (1897).

188. GENUS HETEROZANCLA, TURNER

Heterozancla, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 134 (1919). — Type : *H. rubida*, Turner.

Characters. — Head smooth; tongue developed. Antennae 3/4, in ♂ subserrate, basal joint without pecten. Labial palpi long, recurved, second joint very long, much thickened with loosely appressed scales throughout, scales loosely expanded towards apex above, terminal joint about 1/4 of second, slender, acute. Maxillary palpi very short. Forewings with 2 and 3 connate, 7 and 8 stalked, 7 to costa. Hindwings 1, trapezoidal-ovate, apex obtuse, termen not sinuate; 3 and 4 connate, 5 approximated, 6 and 7 parallel.

Remarks. — Derived from an early form of the preceding.

Geographical distribution of species. — Australian.

Larva unknown.

1. *H. rubida*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 134 (1919). Victoria.

189. GENUS EPHELICTIS, MEYRICK

Ephelictis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 387 (1904). — Type : *E. neochalca*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi very long, curved, ascending, second joint very long, thickened with rough scales above and beneath, terminal joint much shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 approximated or connate or stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal-ovate, apex obtuse, termen hardly sinuate, cilia 1/2-3/4; 3 and 4 connate, 5 somewhat approximated to 4, 6 and 7 parallel.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|---|---------------|
| 1. <i>E. megalarthra</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 388 (1904). | W. Australia. |
| 2. <i>E. neochalca</i> , Meyrick, ibidem, Vol. 29, p. 388 (1904). | W. Australia. |

190. GENUS PANCOENIA, MEYRICK

Pancoenia, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 389 (1904). — Type : *P. periphora*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ rather stout, serrulate, simple, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint with appressed scales, somewhat rough beneath, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 11 from middle. Hindwings 1, rounded-trapezoidal, termen sometimes sinuate, cilia 1/2-1; 3 and 4 connate, 5 somewhat approximated to 4, 6 and 7 parallel.

Remarks. — Modified from an early form of *Protolechia*; inconspicuous insects.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|---|------------------|
| 1. <i>P. periphora</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 389 (1904). | New South Wales. |
| 2. <i>P. pelota</i> , Meyrick, ibidem, Vol. 29, p. 390 (1904). | New South Wales. |
| 3. <i>P. pygmaea</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 148 (1919). | N. Queensland. |

191. GENUS ORTHOPTILA, MEYRICK

Orthoptila, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 392 (1904). — Type : *O. abruptella*, Walker.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ shortly ciliated, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint thickened with appressed scales, slightly rough beneath, terminal joint shorter than second, acute.

Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with subdorsal tuft of scales before middle; 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal-ovate, apex obtuse, termen not sinuate, cilia 2/3; 3 and 4 connate, 5 somewhat approximated to 4, 6 and 7 stalked.

Remarks. — Allied to early forms of *Protolechia*, some of which show raised scales corresponding to the subdorsal tuft of this genus, but much less developed; but distinguished also by the stalking of veins 6 and 7 of hindwings.

Geographical distribution of species. — Australian.

Larva unknown.

1. *O. abruptella*, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1032 (1864). — S. E. Australia,
Pl. 3, Fig. 64.
triforella, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 336, pl. 4, f. 108 (1877).

192. GENUS PHLOEOGRAPTIS, MEYRICK

Phloeoegraptis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 393 (1904). — Type: *P. macrynta*,
Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ moderately ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, somewhat roughened towards apex beneath, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked, 4 and 5 nearly connate, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings 1 or somewhat over 1, trapezoidal, termen not sinuate, cilia 2/5; 3 and 4 connate, 5 slightly approximated to 4 at base, 6 and 7 tolerably parallel.

Remarks. — A peculiar type, but allied to early forms of *Protolechia*. The species are adapted to concealment on *Eucalyptus* trunks.

Geographical distribution of species. — Australian.

Larva unknown.

1. *P. macrynta*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 394 (1904). Victoria.
2. *P. brachynta*, Meyrick, ibidem, Vol. 29, p. 394 (1904). Queensland.
3. *P. zopherota*, Meyrick, ibidem, Vol. 29, p. 394 (1904). Victoria.

Group 6 (Chelaria type)

Specialised forms constituting a rather recent line of development from a common origin with the *Compsolechia* group. They are commonly marked by more or less tufting of the palpi, especially by posterior scale-projections and several dark rings on a long terminal joint, but in *Anarsia* the terminal joint is usually aborted in the ♂. The forewings often have tufts of scales; vein 2 is separate, 3 tending rather to unite with 4, 7 to costa; in the hindwings veins 3 and 4 are connate or nearly approximated towards base, 6 and 7 nearly approximated or stalked (except *Thriophora*), the cubital pecten never present. The species are generally narrow-winged, and a small dark median costal spot is a very usual and characteristic feature.

193. GENUS ANARSIA, ZELLER

Anarsia, Zeller, Isis, p. 190 (1839). — Type : *A. spartiella*, Schrank.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi with scales of second joint forming a dense projecting tuft beneath, terminal joint in ♂ very short, usually concealed, in ♀ longer than second, ascending, acute, or rarely as in ♂. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 separate, parallel, 7 and 8 stalked, 7 to costa, 6 sometimes out of 8 near base, 11 from middle. Hindwings 1, trapezoidal, termen hardly sinuate, cilia 3/4-1 1/4; 3 and 4 connate or approximated, 5 rather approximated to 4, 6 and 7 stalked or seldom approximated.

Remarks. — A derivative of *Chelaria*. The species are often similar in appearance, but then generally possess easy distinctions in the secondary sexual hair-pencils and scale-patches of the wings in the ♂.

Geographical distribution of species. — Essentially Indo-Malayan and African, but with a few representatives in Australia and the Palaearctic region, one of which has been introduced into America.

Larva (18 known) feeding in spun leaves, or sometimes in fruits or shoots.

Foodplants by strong preference *Leguminosae* (11), but also *Rosaceae*, *Anacardiaceae*, *Rhamnaceae*, *Capparidaceae*, *Tamaricaceae*, *Loranthaceae*, *Salicaceae*.

1. <i>A. gravata</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 69 (1912).	Transvaal.
2. <i>A. ambitiosa</i> , Meyrick, ibidem, Vol. 3, p. 299 (1913).	Transvaal.
3. <i>A. carbonaria</i> , Meyrick, ibidem, Vol. 3, p. 299 (1913).	Transvaal.
4. <i>A. inculta</i> , Walsingham, Trans. Ent. Soc. Lond. p. 112, pl. 5, f. 49 (1891).	Gambia.
5. <i>A. leucophora</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 417 (1904).	New South Wales.
6. <i>A. acrotoma</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 169 (1913).	Coorg.
7. <i>A. phortica</i> , Meyrick, ibidem, Vol. 22, p. 167 (1913).	Ceylon, S. India, Borneo.
8. <i>A. tricornis</i> , Meyrick, ibidem, Vol. 22, p. 168 (1913).	Ceylon.
9. <i>A. vectaria</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 21 (1918).	Natal.
10. <i>A. euphorodes</i> , Meyrick, Exot. Microlep. Vol. 2, p. 503 (1922).	China.
11. <i>A. molybdota</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 417 (1904). — Pl. 3, Fig. 66.	S. & W. Australia.
12. <i>A. isogona</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 169 (1913).	S. India.
13. <i>A. sagmatica</i> , Meyrick, Exot. Microlep. Vol. 1, p. 582 (1916).	India.
14. <i>A. pensilis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 168 (1913).	Ceylon. [Wales.
15. <i>A. epiula</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 418 (1904).	Queensland, New South Rhodesia, Port.E.Africa.
16. <i>A. semnopha</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 79 (1921).	Ceylon, S. India,
17. <i>A. patulella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 635 (1864).	Java. [Queensland.
18. <i>A. eutacta</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 163 (1921).	Rhodesia.
19. <i>A. balionearia</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 79 (1921).	Port. E. Africa.
20. <i>A. citromitra</i> , Meyrick, ibidem, Vol. 8, p. 80 (1921).	Bengal.
21. <i>A. altercata</i> , Meyrick, Exot. Microlep. Vol. 2, p. 148 (1918).	S. Africa, Kenya Colony.
22. <i>A. agricola</i> , Walsingham, Trans. Ent. Soc. Lond. p. 111, pl. 5, f. 48 (1891).	Bengal.
23. <i>A. ephippias</i> , Meyrick, Ent. M. Mag. Vol. 44, p. 197 (1908).	Bengal.
24. <i>A. idioptila</i> , Meyrick, Exot. Microlep. Vol. 1, p. 582 (1916).	Queensland.
25. <i>A. hippocoma</i> , Meyrick, ibidem, Vol. 2, p. 429 (1921).	India, Tonkin.
26. <i>A. acerata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 169 (1913).	

27. <i>A. subfulvescens</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 21 (1918).	Natal.
28. <i>A. didymopa</i> , Meyrick, Exot. Microlep. Vol. 1, p. 583 (1916).	Bengal.
29. <i>A. veruta</i> , Meyrick, ibidem, Vol. 2, p. 148 (1918).	Bengal.
30. <i>A. mitescens</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 299 (1913).	Transvaal.
31. <i>A. omoptila</i> , Meyrick, Exot. Microlep. Vol. 2, p. 147 (1918).	India.
32. <i>A. lineatella</i> , Zeller, Isis, p. 190 (1839). ? <i>pallatella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 118 (1796). <i>pruniella</i> , Clemens, Proc. Acad. Nat. Sc. Philad. p. 169 (1860).	C. & S. Europe, W. C. Asia, Mesopotamia, N. Africa, N. America. Kenya Colony. Transvaal. Algeria. Turkestan. N. Africa, Syria, Arabia. S. India, Burma. S. India. E. Australia. N. Persia.
33. <i>A. arsenopa</i> , Meyrick, Voyage All. Jean. Lép. Vol. 2, p. 72 (1920).	C. & S. Europe, Asia Minor.
34. <i>A. amalleuta</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 298 (1913).	
35. <i>A. retamella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 331 (1915).	
36. <i>A. eburnella</i> , Christoph, Mém. Lép. Roman. Vol. 3, p. 122, pl. 5, f. 14 (1887).	
37. <i>A. acaciae</i> , Walsingham, Proc. Zool. Soc. Lond. p. 278 (1896).	
38. <i>A. triaenota</i> , Meyrick, Journ. Bomb. Nat. Hist. Soc. Vol. 22, p. 169 (1913).	
39. <i>A. reciproca</i> , Meyrick, Exot. Microlep. Vol. 2, p. 300 (1918).	
40. <i>A. trichodeta</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 415 (1904).	
41. <i>A. halimodendri</i> , Christoph, Hor. Soc. Ent. Ross. Vol. 12, p. 297, pl. 8, f. 69 (1877).	
42. <i>A. spartiella</i> , Schranck, Faun. Boic. Vol. 2, p. 104 (1802). <i>robertsonella</i> , Curtis, Brit. Ent. Vol. 14, p. 655 (1838). <i>genistae</i> , Stainton, Ins. Brit. Tin. p. 144 (1854).	S. E. Australia, Tasmania. Transvaal. [nia. Mesopotamia.
43. <i>A. dryinopa</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 57 (1897).	Bengal.
44. <i>A. nimboza</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 300 (1913).	Transvaal.
45. <i>A. aleurodes</i> , Meyrick, Exot. Microlep. Vol. 2, p. 502 (1922).	Algeria.
46. <i>A. sagittaria</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 774 (1914).	Bengal.
47. <i>A. spicata</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 21 (1918).	
48. <i>A. luteostella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 332 (1915).	
49. <i>A. melanoplecta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 774 (1914).	
50. <i>A. epotias</i> , Meyrick, Exot. Microlep. Vol. 1, p. 583 (1916).	Bengal.
51. <i>A. stylota</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 168 (1913).	Ceylon.

194. GENUS DOLEROTRICHA, NOV. GEN.

Type : *D. flabellifera*, Rebel.

Characters. — Head with appressed scales; tongue developed. Antennae $\frac{3}{4}$, in ♂ simple, basal joint without pecten. Labial palpi with second joint densely scaled, with long dense projecting tuft beneath, terminal joint moderate, pointed, strongly reflexed, wholly concealed in scales of second. Maxillary palpi very short. Forewings with 2 separate, 7 and 8 stalked, 7 to costa. Hindwings over 1, trapezoidal, apex pointed, termen somewhat sinuate; 3 and 4 connate, 6 and 7 connate (?).

Remarks. — The typical species (described from a single specimen), which I have not seen, is placed by its author in *Nothris*, but with the admission that the peculiar form of the palpi should justify a new genus. This is now called for; I give the characters as they seem to have been conceived by Rebel, but the neuration was apparently imperfectly ascertained.

Geographical distribution of species. — North African.

Larva unknown.

1. *C. flabellifera*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 46, p. 175 (1896). Morocco.

195. GENUS PALINTROPA, MEYRICK

Palintropa, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 160 (1913). — Type : *P. hippica*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales above, beneath with broad rough tuft of projecting scales, terminal joint longer than second, thickened with scales, posteriorly with dense rough projecting scales except at apex, acute. Maxillary palpi short, filiform, appressed to tongue. Posterior tibiae with rough scales above. Forewings with 2 from angle, 3 absent, 6 to apex, 7 and 8 stalked, 11 from middle. Hindwings 2/3, narrowly elongate-trapezoidal, apex pointed, produced, termen beneath apex bisinuate, very oblique, cilia 3; 3 absent, 4 and 5 connate, 6 and 7 stalked.

Remarks. — A development of *Chelaria*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *P. hippica*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 160 (1913). Ceylon.

196. GENUS CHELARIA, HAWORTH

Chelaria, Haworth, Lep. Brit. p. 526 (1828). — Type : *C. conscriptella*, Hübner.

Psoricoptera, Stainton, Inst. Brit. Tin. p. 100 (1854). — Type : *C. gibbosella*, Zeller.

Cymatomorpha, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 411 (1904). — Type : *C. euplecta*, Meyrick.

Deuteroptila, Meyrick, ibidem, Vol. 29, p. 418 (1904). — Type : *C. sphenophora*, Meyrick.

Allocota, Meyrick, ibidem, Vol. 29, p. 419 (1904). — Type : *C. simulacrella*, Meyrick.

Semodictis, Meyrick, Ann. Transv. Mus. Vol. 2, p. 16 (1909). — Type : *C. tetraptera*, Meyrick.

Episacta, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 161 (1919). — Type : *C. discissa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with double or single tuft beneath or rough-scaled, terminal joint longer than second, thickened with scales often forming one or two projections posteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings usually with more or less developed tufts of scales; 1 b furcate, 2 from towards angle, 6 to apex or termen, 7 and 8 stalked, 7 to costa, or 7 and 8 out of 6, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse, termen more or less bisinuate beneath apex, oblique, cilia about 1; 3-5 more or less approximated at base, 6 and 7 stalked or closely approximated at base.

Remarks. — I am now satisfied that the variations in scaling of palpi and in neuration included within this genus are only specific (they are reliable for discriminating species); so treated the genus forms a natural and recognisable whole, of no excessive extent. Some of the Australian species are very similar to one another and require observation of the genitalia, which happen to be remarkably diverse and striking in character.

Geographical distribution of species. — Indo-Malayan, Australasian, and African, with stragglers in Europe and America.

- Larva (6 known) feeding in spun leaves or inflorescence, or sometimes in shoots or Psyllid galls.
- Foodplants: *Anacardiaceae* (2), *Betulaceae*, *Cupuliferae*, *Euphorbiaceae*, *Lecythidaceae* (all trees or shrubs).
1. *C. scopulosa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 165 Kanara.
(1913).
 2. *C. iophana*, Meyrick, ibidem, Vol. 22, p. 162 (1913). Ceylon, Java.
 3. *C. isoptila*, Meyrick, ibidem, Vol. 22, p. 163 (1913). Assam, Ceylon.
 4. *C. corynetis*, Meyrick, ibidem, Vol. 22, p. 162 (1913). — Pl. 3, Fig. 67. Ceylon.
 5. *C. isotricha*, Meyrick, Zool. Med. Leid. Vol. 6, p. 164 (1921). Java.
 6. *C. eriozona*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 80 (1921). Port. E. Africa.
 7. *C. syncrypta*, Meyrick, Exot. Microlep. Vol. 1, p. 580 (1916). Ceylon.
 8. *C. scriniata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 163 Ceylon.
(1913).
 9. *C. simulacrella*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 420 New South Wales.
(1904).
 10. *C. tortuosa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 165 (1913). Ceylon.
 11. *C. caryodora*, Meyrick, ibidem, Vol. 22, p. 164 (1913). — Pl. 6, Fig. 126. Assam.
 12. *C. tonsa*, Meyrick, ibidem, Vol. 22, p. 164 (1913). Assam, Coorg.
 13. *C. lactifera*, Meyrick, ibidem, Vol. 22, p. 161 (1913). Assam.
 14. *C. spathota*, Meyrick, ibidem, Vol. 22, p. 165 (1913). India, Queensland.
 15. *C. indica*, Walsingham, Proc. Zool. Soc. Lond. p. 884 (1885). Bombay.
 16. *C. pilosella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 640 (1864). Borneo.
 17. *C. dissidens*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 301 (1913). Transvaal.
 18. *C. albogrisea*, Walsingham, Trans. Ent. Soc. Lond. p. 264, pl. 12, f. 34 Natal, Zululand.
(1881).
 19. *C. probolaea*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 298 (1913). Transvaal.
 20. *C. formidolosa*, Meyrick, Exot. Microlep. Vol. 1, p. 581 (1916). Natal.
 21. *C. antaris*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 297 (1913). Transvaal.
 22. *C. improba*, Meyrick, ibidem, Vol. 3, p. 297 (1913). Transvaal.
 23. *C. mancipata*, Meyrick, ibidem, Vol. 3, p. 297 (1913). Transvaal.
 24. *C. procax*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 274 (1911). Seychelles.
 25. *C. tetraptera*, Meyrick, Ann. Transv. Mus. Vol. 2, p. 16, pl. 5, f. 7 (1910). Transvaal.
 26. *C. silvestris*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 164 (1913). Assam.
 27. *C. discissa*, Meyrick, Exot. Microlep. Vol. 1, p. 581 (1916). N. Queensland.
 28. *C. toreuta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 162 (1919). N. Queensland.
 29. *C. dermatica*, Meyrick, Exot. Microlep. Vol. 2, p. 432 (1921). Queensland.
 30. *C. sphenophora*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 419 (1904). Queensland.
 31. *C. conscriptella*, Hübner, Samml. Eur. Schmett. Tin. f. 283 (1805). C. Europe.
 - huebnerella*, Donovan, Nat. Hist. Brit. Ins. Vol. 11, pl. 382, f. 2 (1806).
 - rhomboidea*, Curtis, Brit. Ent. Vol. 8, p. 368 (1832).
 32. *C. arignota*, Meyrick, Exot. Microlep. Vol. 1, p. 579 (1916). Burma.
 33. *C. instaurata*, Meyrick, Zool. Med. Leid. Vol. 6, p. 165 (1921). Java.
 34. *C. obtruncata*, Meyrick, Exot. Microlep. Vol. 3, p. 30 (1923). Assam.
 35. *C. verticosa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 166 (1913). Coorg.
 36. *C. demonstrata*, Meyrick, Exot. Microlep. Vol. 2, p. 303 (1920). New Guinea.
 37. *C. ericta*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 162 (1913). Ceylon.
 38. *C. trachydyta*, Meyrick, Exot. Microlep. Vol. 2, p. 304 (1920). Kanara.
 39. *C. rhinota*, Meyrick, ibidem, Vol. 1, p. 580 (1916). S. India.
 40. *C. brachyrrhiza*, Meyrick, ibidem, Vol. 2, p. 431 (1921). Fiji.
 41. *C. apicipuncta*, Busck, Proc. U. S. Mus. Vol. 40, p. 206, pl. 9, f. 35 Panama, Costa Rica,
(1911) (apice-). Guiana.

42. <i>C. hora</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 14 (1914).	Panama.
43. <i>C. euchorda</i> , Meyrick, Exot. Microlep. Vol. 3, p. 31 (1923).	Brazil.
44. <i>C. speciosella</i> , Teich, Stett. Ent. Zeit. Vol. 53, p. 358 (1892).	Latvia.
45. <i>C. gibbosella</i> , Zeller, Isis, p. 202 (1839).	C. Europe.
46. <i>C. cyrtopleura</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 165 (1919).	N. Australia, Queensland.
47. <i>C. anguinea</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 161 (1913).	Assam.
48. <i>C. apparitrix</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 164 (1921).	Java.
49. <i>C. melanecta</i> , Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 246 (1914).	Transvaal.
50. <i>C. austerodes</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 22 (1918).	Transvaal.
51. <i>C. loxosaris</i> , Meyrick, ibidem, Vol. 6, p. 21 (1918).	Natal.
52. <i>C. cirrhospila</i> , Meyrick, Exot. Microlep. Vol. 2, p. 302 (1920).	Assam.
53. <i>C. isosema</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 81 (1921).	Rhodesia.
54. <i>C. phacelota</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 166 (1913).	Ceylon.
55. <i>C. tessulata</i> , Meyrick, Exot. Microlep. Vol. 2, p. 431 (1921).	N. Queensland.
56. <i>C. baliodes</i> , Lower, Trans. Roy. Soc. S. Australia Vol. 44, p. 66 (1920).	Queensland.
57. <i>C. paroctas</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 166 (1913).	Ceylon.
58. <i>C. tephroplintha</i> , Meyrick, Exot. Microlep. Vol. 3, p. 30 (1923).	Fiji.
59. <i>C. triannulata</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 69 (1912).	S. Africa.
60. <i>C. solutrix</i> , Meyrick, ibidem, Vol. 3, p. 69 (1912).	Transvaal.
61. <i>C. sciograpta</i> , Meyrick, ibidem, Vol. 8, p. 80 (1921).	Transvaal.
62. <i>C. scotia</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 160 (1919).	N. Queensland.
63. <i>C. tenebrosa</i> , Meyrick, Exot. Microlep. Vol. 2, p. 301 (1920).	South Australia.
64. <i>C. euplecta</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 412 (1904).	New South Wales, Queensland.
65. <i>C. microgramma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 301 (1920).	New South Wales.
66. <i>C. attenuata</i> , Meyrick, ibidem, Vol. 2, p. 300 (1920).	New South Wales.
67. <i>C. ammonura</i> , Meyrick, ibidem, Vol. 2, p. 430 (1921).	Queensland.
68. <i>C. metaphorica</i> , Meyrick, ibidem, Vol. 2, p. 430 (1921).	Queensland.
69. <i>C. orthostathma</i> , Meyrick, ibidem, Vol. 2, p. 429 (1921).	Queensland.
70. <i>C. harpophora</i> , Meyrick, ibidem, Vol. 2, p. 431 (1921).	Queensland.
71. <i>C. particulata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 167 (1913).	Ceylon, Java.
72. <i>C. aridella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 639 (1864).	Borneo.
73. <i>C. levata</i> , Meyrick, Exot. Microlep. Vol. 2, p. 304 (1920).	Bengal.
74. <i>C. stictocosma</i> , Meyrick, ibidem, Vol. 2, p. 303 (1920).	Coorg, Kanara.

197. GENUS HAPLOCHELA, MEYRICK

Haplochela, Meyrick, Exot. Microlep. Vol. 3, p. 32 (1923). — Type : *H. mundana*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint much thickened with dense appressed scales, terminal joint longer than second, thickened with scales projecting posteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with tufts of scales; 2 from towards angle, 3-5 approximated, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen rounded, cilia 2/3; 3 and 4 connate, 5 rather approximated, 6 and 7 closely approximated towards base.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — South American.

Larva unknown.

1. *H. mundana*, Meyrick, Trans. Ent. Soc. Lond. p. 254 (1914).

Guiana, Brazil, Peru.

198. GENUS SCLEROGRAPTIS, MEYRICK

Sclerogaptis, Meyrick, Exot. Microlep. Vol. 3, p. 31 (1923). — Type : *S. oxytypa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint much thickened with dense appressed scales, somewhat expanded at apex above, terminal joint longer than second, thickened with scales roughly projecting posteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae somewhat rough-scaled above. Forewings with 1b furcate, 2 from towards angle, 3 and 4 approximated, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings slightly under 1, trapezoidal, termen slightly bisinuate, cilia 1; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — Also correlated with *Chelaria*.

Geographical distribution of species. — South American.

Larva unknown.

1. *S. oxytypa*, Meyrick, Exot. Microlep. Vol. 3, p. 31 (1923). Guiana.

199. GENUS ENCOLAPTA, MEYRICK

Encolapta, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 167 (1913). — Type : *E. metorcha*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint with short apical tuft beneath, terminal joint longer than second, somewhat thickened with scales except at apex, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 from towards angle, 3 absent, 4 from angle, 6 and 8 stalked, 6 to apex, 7 absent, 11 from middle. Hindwings 4/5, elongate-trapezoidal, apex round-pointed, termen somewhat bisinuate beneath apex, cilia 1 1/2; 3-5 closely approximated at base, 6 and 7 long-stalked.

Remarks. — Correlated with early forms of *Chelaria*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *E. metorcha*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 167 (1913). Ceylon.

200. GENUS PESSOGRAPTIS, MEYRICK

Pessoigraptis, Meyrick, Exot. Microlep. Vol. 3, p. 29 (1923). — Type : *P. thalamias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint with rather short dense projecting apical tuft of scales beneath, terminal joint much longer than second, stout, slightly roughened anteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with tufts of scales; 1b furcate, 2 from near angle, 3-5

approximated, 7 and 8 stalked or coincident, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen slightly bisinuate, cilia 1; 3 and 4 connate, 5 approximated at base, 6 and 7 long-stalked.

Remarks. — Allied to *Chelaria*. The two species, though differing in neuration, are otherwise hardly distinguishable.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|--|-----------------|
| 1. <i>P. cancellata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 255 (1914). | Guiana, Brazil. |
| 2. <i>P. thalamias</i> , Meyrick, Exot. Microlep. Vol. 3, p. 30 (1923). | Brazil. |

201. GENUS APOTACTIS, MEYRICK

Apotactis, Meyrick, Ann. Transv. Mus. Vol. 6, p. 52 (1918). — Type : *A. drimylota*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint with strong dense projecting apical tuft beneath, terminal joint longer than second, somewhat thickened and roughened anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with very long fine hairs above. Forewings with 1b furcate, 2-5 rather approximated, 6 absent, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse, termen scarcely sinuate, cilia 1; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Apparently an early form of this group.

Geographical distribution of species. — African.

Larva unknown.

- | | |
|---|--------------------|
| 1. <i>A. drimylota</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 52 (1918). | Tanganyika Territ. |
|---|--------------------|

202. GENUS PILOCRATES, MEYRICK

Pilocrates, Meyrick, Exot. Microlep. Vol. 2, p. 299 (1920). — Type : *P. prograpta*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4 (?), basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with scales, with dense compact projecting apical tuft beneath, terminal joint longer than second, moderate, acute. Maxillary palpi obsolete. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 3 and 4 connate from angle, 5 approximated, 6 and 8 stalked, 7 absent, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen somewhat sinuate, cilia 1; 3 and 5 closely approximated at base, 4 absent, 6 and 7 stalked.

Remarks. — Somewhat allied to the preceding.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|--|--------|
| 1. <i>P. prograpta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 299 (1920). | Coorg. |
|--|--------|

203. GENUS METABOLAEA, MEYRICK

Metabolaea, Meyrick, Exot. Microlep. Vol. 3, p. 32 (1923). — Type : *M. chlorophthalma*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderate, without pecten. Labial palpi long, recurved, second joint rather rough above and with long rough projecting tuft beneath, terminal joint somewhat shorter than second, rather slender, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with rough scales above. Forewings with tufts of scales; 1b furcate, 2 from towards angle, 3 and 4 approximated from angle, 6 and 7 stalked, 6 to apex, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 1; 3 and 4 connate, 5 rather approximated, 6 and 7 stalked.

Remarks. — Related to the two following.

Geographical distribution of species. — South American.

Larva unknown.

1. *M. chlorophthalma*, Meyrick, Exot. Microlep. Vol. 3, p. 32 (1923). Brazil.

204. GENUS OESTOMORPHA, WALSINGHAM

Oestomorpha, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 107 (1911). — Type : *O. alloea*, Walsingham.

Characters. — Head smooth; ocelli very small, posterior; tongue developed. Antennae 3/4, basal joint moderate, without pecten. Labial palpi moderately long, recurved, second joint much thickened with dense rough scales beneath forming a short tuft, terminal joint shorter than second, moderate, pointed. Maxillary palpi short, filiform appressed to tongue. Posterior tibiae rough-haired above. Forewings with some raised scales; 2 from towards angle, 3 and 4 short-stalked from angle, 5 connate, 6 and 8 long-stalked, 6 to apex, 7 absent, 9 closely approximated, 11 from middle. Hindwings over 1, ovate-trapezoidal, termen not sinuate, cilia 2/5; 3 and 4 connate or short-stalked, 5 rather approximated, 6 and 7 stalked.

Remarks. — A derivative of *Crasimorpha*.

Geographical distribution of species. — Tropical American.

Larva unknown.

1. *O. alloea*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 108, pl. 3, Mexico, Panama, Guiana, f. 29 (1911).

205. GENUS CRASIMORPHA, MEYRICK

Crasimorpha, Meyrick, Exot. Microlep. Vol. 3, p. 33 (1923). — Type : *C. peragrata*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderate, without pecten. Labial palpi rather long, curved, ascending, second joint thickened with dense scales, with rough projecting scales towards apex beneath, terminal joint shorter than second, rather stout, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with some raised scales; 1b furcate, 2 from towards angle, 3 and 4 stalked, 7 and 8 stalked, 7 to costa, 6 and 9 approximated to 8 at base, 11 from middle. Hindwings over 1,

trapezoidal-ovate, termen not sinuate, cilia 1/3; 3 and 4 short-stalked, 5 rather approximated, 6 and 7 stalked.

Remarks. — Nearly related to the following.

Geographical distribution of species. — South American.

Larva unknown.

1. *C. peragrata*, Meyrick, Exot. Microlep. Vol. 3, p. 33 (1923). Guiana.

206. GENUS PORPODRYAS, MEYRICK

Porpodryas, Meyrick, Exot. Microlep. Vol. 2, p. 304 (1920). — Type : *P. prasinantha*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue strong. Antennae 1/3, in ♂ moderately ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint much thickened with dense appressed scales, somewhat expanded towards apex above, terminal joint as long as second, thickened with dense scales somewhat projecting towards apex posteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with dense rough scales above and beneath. Forewings with tufts of scales; 1b furcate, 2 from 5/6, 3-5 approximated, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal-ovate, apex obtuse, termen hardly sinuate, cilia 2/5; 3 and 4 connate, 5 rather approximated, 6 and 7 closely approximated towards base.

Remarks. — This must nearly approach the primitive form of all the preceding genera of the group.

Geographical distribution of species. — South American.

Larva unknown.

1. *P. prasinantha*, Meyrick, Exot. Microlep. Vol. 2, p. 305 (1920). Guiana.

207. GENUS STACHYOSTOMA, MEYRICK

Stachyostoma, Meyrick, Exot. Microlep. Vol. 3, p. 28 (1923). — Type : *S. psilodoxa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, minutely ciliated, basal joint elongate, without pecten. Labial palpi very long, second joint straight, densely clothed with scales, roughly expanded towards apex above and with loose rough spreading apical tuft beneath, terminal joint erect, somewhat longer than second, slender, pointed. Maxillary palpi rudimentary. Posterior tibiae clothed with rough hairs above. Forewings with 1b furcate, 2 from 3/4, 3 and 4 closely approximated from angle, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, termen hardly sinuate, cilia 3/4; 2-4 remote, slightly approximated towards base, 5 parallel, nearly equidistant from 4 and 6, 6 and 7 nearly approximated at base.

Remarks. — A specialised accessory of the group.

Geographical distribution of species. — South American.

Larva unknown.

1. *S. psilodoxa*, Meyrick, Exot. Microlep. Vol. 3, p. 28 (1923). Ecuador.

208. GENUS METOPLEURA, BUSCK

Metopleura, Busck, Proc. Ent. Soc. Wash. Vol. 14, p. 83 (1912). — Type : *M. potosi*, Busck.

Characters. — Head with appressed scales; ocelli very small, inferior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi very long, slightly curved, ascending, second joint very long, much thickened with dense appressed scales roughly expanded towards apex above, terminal joint half second, thickened with dense slightly roughened scales, acute. Maxillary palpi very short, loosely scaled, appressed to tongue. Posterior tibiae loosely scaled above. Forewings with slight tufts of scales; 1b furcate, 2 from 3/4, 3 from angle, 7 absent, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 1/2; 3 and 4 connate or nearly so, 5 parallel, 6 and 7 connate or closely approximated.

Remarks. — A specialised form related to *Crasimorpha*.

Geographical distribution of species. — North American.

Larva unknown.

1. *M. potosi*, Busck, Proc. Ent. Soc. Wash. Vol. 14, p. 83 (1912). Mexico.

209. GENUS TORNODOXA, MEYRICK

Tornodoxa, Meyrick, Exot. Microlep. Vol. 2, p. 432 (1921). — Type : *T. tholochorda*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint densely scaled, with long rough projecting hairscales beneath throughout, terminal joint longer than second, thickened with dense appressed scales, laterally compressed, acute. Maxillary palpi short, loosely scaled, appressed to tongue. Posterior tibiae clothed with rough hair-scales above. Forewings with 2 from towards angle, 3 and 4 short-stalked from angle, 5 closely approximated, 6 to apex, 7 absent, 11 from middle. Hindwings 1, elongate-ovate, costa somewhat sinuate, apex rounded-obtuse, termen rounded, cilia 1/2; 3 and 4 short-stalked, 5 rather approximated, 6 and 7 stalked.

Remarks. — A peculiar form, perhaps allied to *Tituacia*.

Geographical distribution of species. — Japan.

Larva unknown.

1. *T. tholochorda*, Meyrick, Exot. Microlep. Vol. 2, p. 432 (1921). Japan.

210. GENUS TITUACIA, WALKER

Tituacia, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 812 (1864). — Type : *T. devilla*, Walker.

Stomylla, Snellen, Tijdschr. v. Ent. Vol. 22, p. 14 (1878). — Type : *T. devilla*, Walker.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales forming a short obtuse tuft at apex beneath, terminal joint longer than

second, with two projections of rough scales posteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with tufts of scales; 1b furcate, 2 from towards angle, 3 and 4 stalked, 7 and 8 stalked or seldom coincident, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen hardly sinuate, cilia $\frac{3}{4}$; 3 and 4 connate, 5 nearly approximated at base, 6 and 7 stalked.

Remarks. — Probably derived from a form approaching *Dactylethra*.

Geographical distribution of species. — Indo-Malayan and (perhaps by introduction) Australian.

Larva unknown (probably on a cultivated tree or shrub).

- | | |
|--|--------------------------------------|
| 1. <i>T. deviella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 812 (1864).
erossella, Snellen, Tijdschr. v. Ent. Vol. 22, p. 14, pl. 8, f. 1-6 (1878). | Ceylon, Java, Borneo,
Queensland. |
|--|--------------------------------------|

211. GENUS AXYROSTOLA, MEYRICK

Axyrostola, Meyrick, Exot. Microlep. Vol. 3, p. 29 (1923). — Type: *A. acherusia*, Meyrick.

Characters. — Head with loosely appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint beneath with tuft of very long rough projecting scales, whorled towards apex, terminal joint longer than second, slender, acute. Maxillary palpi very short, drooping. Posterior tibiae with rough projecting hairs above. Forewings with 2 from near angle, 2-5 approximated, 6 and 8 stalked, 7 absent, 11 from middle. Hindwings 1, elongate-trapezoidal, termen hardly sinuate, cilia 1; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — Probably derived from *Dactylethra*.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|---|--------|
| 1. <i>A. acherusia</i> , Meyrick, Exot. Microlep. Vol. 3, p. 29 (1923). | Burma, |
|---|--------|

212. GENUS THRIOPHORA, MEYRICK

Thriophora, Meyrick, Ann. Transv. Mus. Vol. 2, p. 231 (1911). — Type: *T. ovulata*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely spreading; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint with very long loosely spreading tuft of scales beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from angle, 3 absent, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings almost 1, trapezoidal, termen somewhat sinuate beneath apex, cilia $\frac{1}{4}$; 3 and 4 connate, 5 somewhat approximated, 6 and 7 parallel.

Remarks. — A curious form, but probably derived from *Dactylethra*.

Geographical distribution of species. — South African.

Larva unknown.

- | | |
|--|------------|
| 1. <i>T. ovulata</i> , Meyrick, Ann. Transv. Mus. Vol. 2, p. 231 (1911). | Transvaal. |
|--|------------|

213. GENUS DACTYLETHRA, MEYRICK

Dactylethra, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 153 (1906). — Type : *D. candida*, Stainton.

Characters. — Head with appressed scales, sidetufts loosely spreading; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, simple, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint with long tuft of loose spreading scales beneath, terminal joint as long as second, loosely scaled, acute, with several black rings. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 1b furcate, 2 from near angle, 2-4 parallel, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen faintly sinuate, cilia 4/5; 2 remote, 3 and 4 connate or nearly approximated, 5 slightly approximated, 6 and 7 stalked.

Remarks. — This genus, which must closely approach the primitive form of the whole group, may be derived from a form of the *Dichomeris* group.

Geographical distribution of species. — Indian and African, with a species in Siberia
Larva (*candida*) feeding in galls formed on seed-pods.
Foodplants Leguminosae.

- | | |
|---|---|
| 1. <i>D. tetrametra</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 300 (1913).
2. <i>D. albidella</i> , Snellen, Tijdschr. v. Ent. Vol. 27, p. 171, pl. 9, f. 6 (1884).
3. <i>D. globulata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 461 (1910).
4. <i>D. candida</i> , Stainton, Trans. Ent. Soc. Lond. (2), Vol. 5, p. 114 (1859). — | Transvaal.
E. Siberia.
Ceylon.
India, Ceylon.
Pl. 3, Fig. 68.
<small><i>plagiferella</i>, Walker, List. Lep. Brit. Mus. Vol. 28, p. 540 (1863).
 <i>tetroctas</i>, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 153 (1906).</small> |
| 5. <i>D. incondita</i> , Meyrick, ibidem, Vol. 22, p. 170 (1913). — Pl. 3, Fig. 69.
6. <i>D. bryophilella</i> , Walsingham, Trans. Ent. Soc. Lond. p. 108, pl. 5, f. 46 (1891).
7. <i>D. siccifolii</i> , Walsingham, ibidem, p. 267, pl. 13, f. 37 (1881).
8. <i>D. chionitis</i> , Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 414 (1910). | Ceylon.
Gambia.
Natal:
Transv., Port. E. Africa. |

Group 7 (Dichomeris type)

A considerable group in general recognisable by the characteristic palpi, which have the second joint usually variously tufted or expanded beneath or above or both, the terminal long and acute; the forewings have veins 2 and 3 as a rule stalked (yet exceptions occur freely in the more advanced forms), 7 running to costa or apex or rarely (probably by reversion) to termen; in the hindwings veins 3 and 4 are usually connate, 6 and 7 generally closely approximated or stalked, and the cubital pecten is more often present. The group is well developed in all main regions, but absent from New Zealand. The larvae are normally leaf-feeders.

214. GENUS HYODECTIS, MEYRICK

Hyodectis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 411 (1904). — Type : *H. crenoides* Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ shortly ciliated, basal joint moderate, without pecten. Labial palpi very long, recurved, second

joint rather short, with very long projecting apical scaletuft beneath, terminal joint twice as long as second, smooth-scaled, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 straight, parallel, 7 and 8 out of 6, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal, apex much produced, pointed, termen sinuate-emarginate, cilia 1 2/3; without cubital pecten; 3 and 4 remote, 5 nearly parallel to 4, 6 and 7 long-stalked.

Remarks. — This curious insect is in some respects so discordant that its true position seems very uncertain, but on the whole I regard it as an aberrant member of this group related to *Symbolistis*; the palpi are characteristic.

Geographical distribution of species. — Australian.

Larva unknown.

1. *H. crenoides*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 411 (1904). — S. E. Australia.
Pl. 3, Fig. 65.

215. GENUS APONOEA, WALSINGHAM

Aponeoa, Walsingham, Ent. M. Mag. Vol. 41, p. 125 (1905). — Type : *A. obtusipalpis*, Walsingham.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ shortly biciliated, basal joint without pecten. Labial palpi very long, porrected, second joint very long, densely scaled, somewhat roughened above and beneath, terminal joint very short, erect, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 almost connate, 7 and 8 stalked, 7 to costa, 9 connate. Hindwings over 1, trapezoidal-ovate, apex obtuse, termen not sinuate, cilia 1/2; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — Regarded by Walsingham as allied to *Holcophora*.

Geographical distribution of species. — North African.

Larva (both known) feeding in slight web on leaves and bark of stems.

Foodplants *Plumbaginaceae*.

1. *A. obtusipalpis*, Walsingham, Ent. M. Mag. Vol. 41, p. 125 (1905). Algeria.
2. *A. pruinosa*, Chrétien, Ann. Soc. Ent. Fr. p. 330 (1915). Algeria.

216. GENUS RHYNCHOPACHA, STAUDINGER

Rynchopacha, Staudinger, Berl. Ent. Zeit. Vol. 14, p. 303 (1870). — Type : *R. spiraeæ*, Staudinger.

Characters. — Head with appressed scales; tongue developed. Antennae 4/5. Labial palpi long, recurved, second joint much thickened with dense scales rough above, terminal joint about 1/3 of second, somewhat thickened with scales towards middle, pointed. Forewings (presumed) with 2 separate, 7 and 8 stalked, 7 to costa. Hindwings trapezoidal, apex somewhat produced, termen sinuate; (presumed) 3 and 4 connate, 6 and 7 connate.

Remarks. — I have not seen this species, and Staudinger, whilst suggesting a new generic name, omitted to describe fully any details of structure except the palpi; these however appear to justify the genus, which I only place here conjecturally.

Geographical distribution of species. — South-East Europe.

Larva observed but unrecorded.

Foodplant *Spiraea* (*Rosaceae*).

1. *R. spiraea*, Staudinger, Berl. Ent. Zeit. Vol. 14, p. 302 (1870).

S. E. Russia.

217. GENUS HOLCOPHORA, STAUDINGER

Holcophora, Staudinger, Berl. Ent. Zeit. Vol. 14, p. 313 (1870). — Type : *H. statice*, Staudinger.

Characters. — Head with appressed scales; tongue developed. Antennae 4/5, in ♂ very shortly ciliated. Labial palpi long, recurved, second joint thickened with appressed scales, somewhat expanded towards apex above, terminal joint 1/5 of second, stout, pointed. Maxillary palpi rudimentary. Forewings with 2 remote from 3, strongly curved, 7 and 8 stalked, 7 to costa. Hindwings trapezoidal, termen slightly sinuate; 3 and 4 nearly approximated, 6 and 7 long-stalked.

Remarks. — Probably related to *Mesophleps*.

Geographical distribution of species. — South-East Europe.

Larva unrecorded.

Foodplant (apparently) *Statice* (*Plumbaginaceae*).

1. *H. statice*, Staudinger, Berl. Ent. Zeit. Vol. 14, p. 313 (1870).

S. E. Russia.

218. GENUS SYMBOLISTIS, MEYRICK

Symbolistis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 413 (1904). — Type : *S. orophota*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ thick, serrate, simple, somewhat roughened with scales above, basal joint moderately elongate, without pecten. Labial palpi long, curved, ascending, second joint with dense appressed scales, forming a slight apical projection beneath, and projecting roughly towards apex above, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 widely remote, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex acutely produced, termen emarginate, cilia 2 1/2; without cubital pecten; 3 and 4 remote, 5 somewhat approximated to 4, 6 and 7 somewhat approximated towards base.

Remarks. — This genus is also discordant structurally, yet seems to be an advanced form of the group.

Geographical distribution of species. — Australian.

Larva unknown.

1. *S. argyromitra*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 414 (1904). N. South Wales, Victoria.

2. *S. orophota*, Meyrick, ibidem, Vol. 29, p. 414 (1904).

Queensland, New South Wales.

219. GENUS STRENIASTIS, MEYRICK

Streniastis, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 428 (1904). — Type : *S. thermaea*, Lower.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ serrulate, shortly ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint with projecting tuft of scales beneath, terminal joint as long as second or shorter, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from about 4/5, 3 from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex round-pointed, termen sinuate, oblique, cilia 1 1/2; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 parallel.

Remarks. — Another curious and abnormal form, probably allied to the preceding.

Geographical distribution of species. — Australian.

Larva unknown.

- | | |
|--|------------------|
| 1. <i>S. thermaea</i> , Lower, Proc. Linn. Soc. N. S. Wales, Vol. 22, p. 271 (1897). — | New South Wales, |
| Pl. 5, Fig. 113. | Tasmania. |
| 2. <i>S. composita</i> , Meyrick, Exot. Microlep. Vol. 2, p. 503 (1922). | Queensland. |

220. GENUS THIOGNATHA, MEYRICK

Thiognatha, Meyrick, Voyage All. Jean. Lép. Vol. 2, p. 74 (1920). — Type : *T. metachalca*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, basal joint elongate, without pecten. Labial palpi with second joint extremely long, porrected, thickened with dense scales, at apex forming a very short tuft beneath, above with rough projecting hairs throughout, terminal joint 1/3 of second, erect, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 2 from near angle, 2-4 rather approximated, 7 absent, 11 from middle. Hindwings 1, elongate-trapezoidal, apex obtuse, termen faintly sinuate, cilia 4/5; without cubital pecten; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — A derivative of *Mesophleps*.

Geographical distribution of species. — African.

Larva unknown.

- | | |
|---|---------------|
| 1. <i>T. metachalca</i> , Meyrick, Voyage All. Jean. Lép. Vol. 2, p. 74 (1920). | Kenya Colony. |
|---|---------------|

221. GENUS CROSSEBELA, MEYRICK

Crossobela, Meyrick, Exot. Microlep. Vol. 3, p. 34 (1923). — Type : *C. barysphena*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint thickened with dense rough projecting scales beneath throughout, above with scales rather expanded towards apex, terminal joint half second, slender, pointed. Maxillary palpi rudimentary. Forewings with

2-6 separate, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, termen emarginate, cilia 2; 3 and 4 rather approximated, 5 remote, 6 and 7 stalked.

Remarks. — Correlated with *Mesophleps*.

Geographical distribution of species. — Mediterranean.

Larva unknown.

1. *C. barysphena*, Meyrick, Exot. Microlep. Vol. 3, p. 34 (1923).

Cyprus.

222. GENUS MESOPHLEPS, HÜBNER

Mesophleps, Hübner, Verz. bek. Schmett. p. 405 (1826). — Type : *M. silacella*, Hübner.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, subascending, second joint long, dilated beneath with dense appressed scales, above with loose rough hairs longest towards base, terminal joint 1/4 of second, obliquely ascending, slender, pointed. Maxillary palpi rudimentary. Posterior tibiae clothed with long fine hairs above. Forewings with 1b furcate, 2 remote, 3 from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings slightly over 1, trapezoidal, termen slightly sinuate, cilia 1; without cubital pecten; 3 and 4 connate or closely approximated, 5 somewhat approximated towards base, 6 and 7 short-stalked.

Remarks. — Probably derived from *Brachyacma*.

Geographical distribution of species. — European.

Larva (both known) feeding in seed-capsules.

Foodplants *Cistaceae*.

1. *M. corsicella*, Herrich-Schäffer, Neue Schmett. p. 7, f. 47 (1856).

S. Fr., Sardinia, Corsica.

2. *M. silacella*, Hübner, Samml. Eur. Schmett. Tin. f. 117 (1796).

C. & S. Europe, Asia

pyropella, Hübner, Samml. Vog. Schmett. f. 95 (1793) (praeocc.).

Minor.

223. GENUS BRACHYACMA, MEYRICK

Brachyacma, Meyrick, Trans. Ent. Soc. Lond. p. 278 (1880). — Type : *B. epiochra*, Meyrick.

Lethontogenus, Walsingham, Proc. Zool. Soc. Lond. p. 87 (1897). — Type : *B. palpigera*, Walsingham.

Paraspistes, Meyrick, Journ. Bomb. Nat. Hist. Soc. Vol. 16, p. 600 (1905). — Type : *B. palpigera*, Walsingham.

Lipatia, Busck, Bull. Trinidad Dep. Agric. Vol. 9, p. 243 (1910). — Type : *B. palpigera*, Walsingham.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5; in ♂ simple, basal joint elongate, without pecten. Labial palpi long, curved, ascending, second joint thickened with dense scales roughly expanded above towards apex, terminal joint shorter than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 remote, 3 and 4 connate or stalked from angle, 6 sometimes out of 7 near base, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings nearly 1, elongate-trapezoidal, apex produced, termen bisinuate or emarginate, cilia 2-2; without cubital pecten; 3 and 4 connate or stalked, 5 nearly parallel, 6 and 7 stalked.

Remarks. — An interesting genus, probably related to *Xerometra*.

Geographical distribution of species. — Chiefly Indian, but with single species in Europe and Fiji, and one species has been artificially widely spread with its foodplant.

Larva (3 known) feeding in seed-pods or seed-cones.

Foodplants : *Leguminosae* (2), *Coniferae*.

1. *B. oxycedrella*, Millière, Icon. Descr. Lép. Vol. 3, p. 177, pl. 118, f. 1-6 (1874). S. France.
2. *B. aprica*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 170 (1913). Coorg.
3. *B. palpigera*, Walsingham, Trans. Ent. Soc. Lond. p. 94, pl. 4, f. 31 (1891). India, Ceylon, Seychelles, *adustipennis*, Walsingham, Proc. Zool. Soc. Lond. p. 88 (1897); *ioloncha*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 600 (1905); *crotulariella*, Busck, Bull. Trinidad Dep. Agric. Vol. 9, p. 244 (1910); *epichorda*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 163 (1919).
4. *B. tabellata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 170 (1913). Coorg.
5. *B. epiochra*, Meyrick, Trans. Ent. Soc. Lond. p. 279 (1886). — **Pl. 3, Fig. 71.** Fiji.

224. GENUS EPIMESOPHLEPS, REBEL

Epimesophleps, Rebel, Denkschr. Kais. Akad. Wiss. Wien, Vol. 71, p. 125 (1907). — Type : *E. symmocella*, Rebel.

Characters. — Head with appressed scales; tongue developed. Antennae 4/5. Labial palpi long, ascending, second joint thickened with dense appressed scales, roughly expanded towards apex above, terminal joint 1/3 of second, slender, somewhat roughened, pointed. Forewings with 1b furcate, 2 and 3 approximated from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings somewhat over 1, trapezoidal, termen slightly sinuate, cilia 3/5; 3 and 4 connate, 5 somewhat approximated, 6 and 7 nearly approximated towards base.

Remarks. — Approaching *Mesophleps* in structure, but of doubtful immediate affinity.

Geographical distribution of species. — Arabian.

Larva unknown.

1. *E. symmocella*, Rebel, Denkschr. Kais. Akad. Wiss. Wien, Vol. 71, p. 125 Sokotra. (1907).

225. GENUS AROTRIA, MEYRICK

Arotria, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 387 (1904). — Type . *A. iophaea*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ strongly fasciculate-ciliated, basal joint moderate, with slight pecten. Labial palpi very long, curved, ascending, second joint extremely long, beneath with appressed scales, above with long projecting scales diminishing to apex, terminal joint less than half second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 1; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Seems to be really allied to *Mesophleps*, yet differing in several important particulars; the basal pecten of antennae is unique in this group.

Geographical distribution of species. — Australian.

Larva unknown.

1. *A. iophaea*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 387 (1904). Queensland.

226. GENUS EMPALACTIS, NOV. GEN.

Type : *E. sporogramma*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue short. Antennae 3 4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint with dense rough projecting apical tuft beneath, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1 b furcate, 2 from towards angle, 3-5 separate, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, narrow-trapezoidal, apex rather produced, termen somewhat sinuate, cilia 2; without cubital pecten; 3-5 closely approximated at base, 6 and 7 closely approximated at base, 6 and 7 closely approximated towards base.

Remarks. — Probably a derivative of *Xerometra*.

Geographical distribution of species. — Australian.

Larva unknown.

1. *E. sporogramma*, Meyrick, Exot. Microlep. Vol. 2, p. 433 (1921).

N. Australia.

227. GENUS XEROMETRA, NOV. GEN.

Type : *X. crocina*, Meyrick.

Characters. — Head with appressed scales, sidetufts somewhat raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with dense projecting tuft beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 1 b furcate, 2 remote, parallel, 6 sometimes out of 7 near base, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, apex often produced, termen more or less sinuate, cilia 4/5-1 1/2; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated or stalked.

Remarks. — This genus, formerly confused by me with *Nothris*, appears to represent a line of development correlated with *Dichomeris*, and replacing it in the true indigenous fauna of Australia.

Geographical distribution of species. — Exclusively Australian.

Larva unknown.

- | | |
|--|------------------|
| 1. <i>X. chloristis</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 421 (1904). | W. Australia. |
| 2. <i>X. centrothetis</i> , Meyrick, ibidem, Vol. 29, p. 422 (1904). | W. Australia. |
| 3. <i>X. chloranthes</i> , Lower, ibidem, Vol. 25, p. 418 (1900). | New South Wales. |
| 4. <i>X. meliphanes</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 18, p. 107 (1894). | South Australia. |
| 5. <i>X. apentheta</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 164 (1919). | New South Wales. |
| 6. <i>X. crocina</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 423 (1904). | S. E. Australia. |
| 7. <i>X. tetrachroa</i> , Lower, ibidem, Vol. 23, p. 49 (1898). | New South Wales. |
| 8. <i>X. ochroloma</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 80 (1901). | New South Wales. |
| 9. <i>X. macrosema</i> , Lower, Proc. Linn. Soc. N. S. Wales, Vol. 25, p. 418 (1900). | New South Wales. |
| 10. <i>X. centrosipa</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 165 (1919). | Queensland. |

11. *X. trichombra*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 23, p. 50 (1898). New South Wales.
 12. *X. mylicotis*, Meyrick, ibidem, Vol. 29, p. 426 (1904). South Australia.
 13. *X. acromelas*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 164 (1919). New South Wales.
 14. *X. cycnobathra*, Lower, Proc. Linn. Soc. N. S. Wales, Vol. 23, p. 50 (1898). New South Wales.
 15. *X. tephrastris*, Meyrick, ibidem, Vol. 29, p. 427 (1904). W. Australia.
 16. *X. mesophracta*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 166 (1919). Victoria. [toria.
 17. *X. dentata*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 427 (1904). New South Wales, Vic-

228. GENUS ACRIBOLOGA, MEYRICK

Acribologa, Meyrick, Exot. Microlep. Vol. 2, p. 622 (1923). — Type : *A. malacodes*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ moderately or very strongly ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint expanded towards apex with dense scales, forming a moderate or long projecting tuft beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 separate, from before angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, apex rather produced, cilia 1 1/2-2; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate or stalked.

Remarks. — A simple derivative of *Dichomeris*.

Geographical distribution of species. — Indo-Malayan and African.

Larva unknown.

1. *A. citharista*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 170 Coorg. (1913). [Borneo.
 2. *A. malacodes*, Meyrick, Trans. Ent. Soc. Lond. p. 451 (1910). India, Ceylon, Java,
 3. *A. cymotrocha*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 300 (1913). Transvaal.

229. GENUS LACHNOSTOLA, MEYRICK

Lachnosta, Meyrick, Ann. Transv. Mus. Vol. 6, p. 22 (1918). — Type : *L. amphizeucta*, Meyrick.

Characters. — Head smooth, sidetuffs somewhat raised; ocelli small, far posterior; tongue developed. Antennae 3/4, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, forming a long projecting triangular apical tuft beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from 4/5, 3 and 4 closely approximated from angle, 6 and 8 stalked, 7 absent, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex produced, pointed, termen sinuate, cilia 1 2/3; with cubital pecten; 3 and 4 connate, 5 nearly approximated, 6 and 7 stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — African.

Larva unknown.

1. *L. amphizeucta*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 22 (1918). Natal.

230. GENUS PAPPOPHORUS, WALSINGHAM

Pappophorus, Walsingham, Trans. Ent. Soc. Lond. p. 39 (1897). — Type : *P. eurynota*, Walsingham.

Characters. — Head smooth; tongue developed. Antennae 3/4, in ♂ dentate, basal joint without pecten. Labial palpi very long, recurved, second joint triangularly expanded with dense scales projecting at apex beneath, terminal joint twice as long as second, slender, acute. Maxillary palpi very short, filiform. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 (probably) to costa. Hindwings over 1, trapezoidal, termen slightly sinuate; in ♂ 2 and 3 absent, 6 and 7 approximated towards base, a long hairpencil from base in submedian groove; probably with cubital pecten.

Remarks. — A derivative of *Dichomeris*.

Geographical distribution of species. — African.

Larva unknown.

1. *P. eurynota*, Walsingham, Trans. Ent. Soc. Lond. p. 40, pl. 2, f. 4 (1897). Sierra Leone, Fr. Congo.

231. GENUS RHYNCHOTONA, MEYRICK

Rhynchotona, Meyrick, Exot. Microlep. Vol. 3, p. 35 (1923). — Type : *R. phaeostrota*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue absent. Antennae 3/4, in ♂ serrulate, shortly ciliated, basal joint elongate, without pecten. Labial palpi extremely long, porrected, second joint thickened with dense scales loosely appressed and projecting at apex, terminal joint short, slender, acute, projecting little from apical scales of second. Maxillary palpi obsolete. Posterior tibiae shortly rough-scaled above. Forewings with 2 and 3 stalked from angle, 8 and 9 out of 7, 7 to costa just above apex, 11 from middle. Hindwings under 1, elongate-trapezoidal, apex rather produced, termen sinuate, cilia 1; with cubital pecten; 3-5 nearly approximated at base, 6 and 7 connate.

Remarks. — Also derived from *Dichomeris*.

Geographical distribution of species. — South American.

Larva unknown.

1. *R. phaeostrota*, Meyrick, Exot. Microlep. Vol. 3, p. 35 (1923). Peru.

232. GENUS AGELIARCHIS, MEYRICK

Ageliarchis, Meyrick, Exot. Microlep. Vol. 2, p. 622 (1923). — Type : *A. rhizogramma*, Meyrick.

Characters. — Head with appressed scales; ocelli far posterior; tongue developed. Antennae 4/5, in ♂ serrulate, ciliated, basal joint elongate, in ♂ stalk thickened near base, with subbasal notch and large scaletuft above it. Labial palpi with second joint extremely long, straight, porrected, with rough projecting hairscales above throughout and at apex, terminal joint less than half second, ascending, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae shortly rough-scaled above. Forewings with 2 and 3 stalked from angle, 6 closely approximated to 7 towards base, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, elongate-trapezoidal, termen faintly sinuate, cilia 3/4; with cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — South American.

Larva unknown.

1. *A. rhizogramma*, Meyrick, Exot. Microlep. Vol. 2, p. 623 (1923). Brazil.

233. GENUS RHADINOPHYLLA, TURNER

Rhadinophylla, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 166 (1919). — Type : *R. siderosema*, Turner.

Characters. — Head with appressed scales, sidetuffs somewhat raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint rough-haired above at apex and with long rough apical projecting tuft beneath, terminal joint longer than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 coincident, 7 and 8 stalked, 7 to costa. Hindwings under 1, narrow-trapezoidal, apex acute, strongly produced, termen obtusely emarginate, cilia 2 1/2; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — A derivative of *Dichomeris*.

Geographical distribution of species. — Australasian.

Larva feeding in spun leaves.

Foodplant *Pongamia* (*Leguminosae*).

1. *R. siderosema*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 166 (1919). Queensland, Fiji.

234. GENUS TELEPHILA, MEYRICK

Telephila, Meyrick, Exot. Microlep. Vol. 2, p. 626 (1923). — Type : *T. schmidiella*, Heyden.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint expanded with dense scales, forming a long projecting apical tuft beneath, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1 or somewhat over 1, trapezoidal, termen slightly sinuate, cilia 1/2-3/4; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated towards base.

Remarks. — Intermediate in characters between *Dichomeris* and *Gaesia*, and probably a derivative of the former.

Geographical distribution of species. — European, North American, and Australian.

Larva (2 known) feeding in spun leaves.

Foodplants *Labiate*, *Ericaceae*.

1. *T. schmidiella*, Heyden, Isis, p. 954 (1848). C. & S. W. Europe.
durhamella, Stainton, Cat. Brit. Tin. p. 12 (1849).
quadrinella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 154, f. 616 (1855).

2. *T. plastica*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 433 (1904). New South Wales.
 3. *T. vacciniella*, Busck, Proc. Ent. Soc. Wash. Vol. 17, p. 83 (1915). New Jersey, Canada.
 4. *T. delotella*, Busck, ibidem. Vol. 11, p. 90 (1909). Arizona.

235. GENUS DICHOMERIS, HÜBNER

- Dichomeris**, Hübner, Samml. Exot. Schmett. Vol. 1, p. 25 (1818). — Type : *D. ligulella*, Hübner.
Oxybella, Hübner, Verz. bek. Schmett. p. 407 (1826). — Type : *D. ustulella*, Fabricius.
Rhinosia, Treitschke, Schmett. Eur. Vol. 9 (2), p. 9 (1833). — Type : *D. ustulella*, Fabricius.
Anorthosia, Clemens, Proc. Acad. Nat. Sc. Philad. p. 161 (1860). — Type : *D. punctipennella*, Clemens.
Rhobonda, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 802 (1864) (praeocc.). — Type : *D. punctatella*, Walker.
Carna, Walker, ibidem, Vol. 30, p. 1038 (1864). — Type : *D. punctatella*, Walker.
Sagaritis, Chambers, Canad. Ent. Vol. 4, p. 225 (1872) (praeocc.). — Type : *D. punctipennella*, Clemens.
Macrozancla, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 130 (1919). — Type : *D. mendica*, Turner.
Euryzanca, Turner, ibidem, Vol. 31, p. 131 (1919). — Type : *D. melanophylla*, Turner.
Eurysara, Turner, ibidem, Vol. 31, p. 167 (1919). — Type : *D. pleurophaea*, Turner.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ shortly ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with scales more or less expanded above towards apex and with projecting tuft beneath, sometimes in ♂ with dense long fine expansible hairs above, terminal joint as long as second or longer (only in *heteracma* ♂ short), slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked from angle (rarely closely approximated), 7 and 8 stalked (or exceptionally coincident), 7 to costa, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen more or less sinuate, cilia 1/2-1; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate or stalked.

Remarks. — Probably derivable from *Cymotricha*. The type of *Eurysara*, Turner, is said to have vein 9 of forewings rising out of 7, but this is erroneous, the veins being closely approximated on lower half but not stalked.

Geographical distribution of species. — Nearly cosmopolitan, but absent from New Zealand, and more numerous in warmer regions.

Larva (20 known) feeding in spun or rolled leaves.

Foodplants : *Leguminosae* (4), *Cistaceae* (3), *Cupuliferae* (3), *Betulaceae* (2), *Rosaceae* (2), *Coniferae* (2), *Euphorbiaceae* (2), and other Orders, showing adaptability, but a general preference for trees and shrubs.

- | | |
|--|--|
| 1. <i>D. zygophora</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 430 (1904). | Queensland. |
| 2. <i>D. adelocentra</i> , Meyrick, Exot. Microlep. Vol. 2, p. 305 (1920). | Java. |
| 3. <i>D. brachygrapha</i> , Meyrick, ibidem, Vol. 2, p. 305 (1920). | Assam. |
| 4. <i>D. attenta</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 84 (1921). | Rhodesia. |
| 5. <i>D. brachyptila</i> , Meyrick, Exot. Microlep. Vol. 1, p. 584 (1916). | Burma, Java. |
| 6. <i>D. lotella</i> , Constant, Ann. Soc. Ent. Fr. p. 398, pl. 11, f. 7 (1893). | S. France. |
| 7. <i>D. ianthes</i> , Meyrick, Trans. Ent. Soc. Lond. p. 273 (1887).
<i>ochrophanes</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 981 (1907). | India, Ceylon, Celebes, Formosa, Egypt, Seychelles, Reunion. |

8. *D. ammoxantha*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 430 (1904). Queensland, Kei Islands.
9. *D. sciastes*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 90, pl. 3, f. 10 (1911). Mexico.
10. *D. trimaculella*, Chambers, Canad. Ent. Vol. 6, p. 243 (1874). Kentucky, Texas.
toucayellus, Busck, Dyar List N. Amer. Lep. p. 508 (1902). [Islands.]
11. *D. rustica*, Walsingham, Proc. Zool. Soc. Lond. p. 525 (1891). Texas, Windward
12. *D. cachrydias*, Meyrick, Trans. Ent. Soc. Lond. p. 283 (1914). Guiana, Brazil.
13. *D. acuminata*, Staudinger, Stett. Ent. Zeit. Vol. 37, p. 148 (1876). Sicily.
14. *D. consertella*, Christoph, Bull. Soc. Nat. Mosc. p. 31 (1882). E. Siberia.
15. *D. dignella*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 91 (1911). Mexico.
16. *D. piperata*, Walsingham, Proc. Zool. Soc. Lond. p. 526 (1891). Windward Islands.
17. *D. lucrifuga*, Meyrick, Exot. Microlep. Vol. 2, p. 620 (1923). Brazil.
18. *D. saturata*, Meyrick, ibidem, Vol. 2, p. 621 (1923). Brazil.
19. *D. lypetica*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 91 (1911). Mexico.
20. *D. horiodes*, Meyrick, Exot. Microlep. Vol. 2, p. 620 (1923). Brazil.
21. *D. stratigera*, Meyrick, Trans. Ent. Soc. Lond. p. 111 (1922). Brazil.
22. *D. percnoptolis*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 93, pl. 3, f. 11 (1911). Guatemala.
23. *D. citrifoliella*, Chambers, Journ. Cinc. Soc. Nat. Hist. Vol. 2, p. 184 (1880). Florida.
24. *D. famulata*, Meyrick, Trans. Ent. Soc. Lond. p. 284 (1914). Colombia, Guiana, Bra-
25. *D. instans*, Meyrick, Exot. Microlep. Vol. 2, p. 619 (1923). Brazil, Peru. [zil, Peru.]
26. *D. thermodynias*, Meyrick, ibidem, Vol. 2, p. 621 (1923). Peru.
27. *D. argentinella*, Berg, An. Soc. Arg. Vol. 19, p. 282 (1885). Argentina.
28. *D. fida*, Meyrick, Exot. Microlep. Vol. 2, p. 620 (1923). Brazil.
29. *D. hirculella*, Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 89 (1909). Connecticut.
30. *D. ligulella*, Hübner, Zutr. Exot. Schmett. f. 143, 144 (1818). — **Pl. 5,**
Fig. 125.
- pometella* (Harris), Fitch, Journ. N. York Agric. Soc. Vol. 4, p. 36 (1853). .
contubernatellus, Fitch, ibidem, Vol. 4, p. 36 (1853).
- malifoliellus*, Fitch, Trans. N. York Agric. Soc. Vol. 13, p. 231 (1854).
- pauciguttellus*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 123 (1863).
- flavivittellus*, Clemens, ibidem, Vol. 2, p. 429 (1863).
- reedella*, Chambers, Canad. Ent. Vol. 4, p. 222 (1872).
- quercipomonella*, Chambers, ibidem, Vol. 4, p. 222 (1872)
- ruderella*, Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 167 (1878).
31. *D. zomias*, Meyrick, Trans. Ent. Soc. Lond. p. 283 (1914). Guiana, Brazil.
32. *D. stratella*, Walsingham, Proc. Zool. Soc. Lond. p. 87 (1897). Trinidad.
33. *D. substratella*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 93 (1911). Mexico, Guiana.
34. *D. hemichrysella*, Walker, List. Lep. Brit. Mus. Vol. 28, p. 536 (1863). Guiana, Brazil, Peru.
excisarella, Walker, ibidem, Vol. 29, p. 665 (1864).
macroptera, Meyrick, Trans. Ent. Soc. Lond. p. 283 (1914).
35. *D. capillata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 87, pl. 3, f. 9 (1911). Guatemala.
36. *C. prensans*, Meyrick, Trans. Ent. Soc. Lond. p. 111 (1922). Guiana, Brazil, Peru.
37. *D. acrolychna*, Meyrick, ibidem, p. 112 (1922). Brazil.
38. *D. punctatella*, Walker, List. Lep. Brit. Mus. Vol. 29, p. 802 (1864). Guiana, Brazil.
39. *D. mexicana*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 96 (1911). California, Mexico.
40. *D. pleurophaea*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 167 (1919). Queensland.
hyalombra, Meyrick, Exot. Microlep. Vol. 2, p. 503 (1922).
41. *D. summata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 172 (1913). Assam.
42. *D. ferruginosa*, Meyrick, ibidem, Vol. 22, p. 173 (1913). Assam.

43. *D. ostracodes*, Meyrick, Exot. Microlep. Vol. 1, p. 583 (1916).
 44. *D. intensa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 173 (1913).
 45. *D. ignorata*, Meyrick, Zool. Med. Leid. Vol. 6, p. 165 (1921).
 46. *D. ceponoma*, Meyrick, Exot. Microlep. Vol. 2, p. 151 (1918).
 47. *D. crepitatrix*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 173 (1913).
 48. *D. harmonias*, Meyrick, Exot. Microlep. Vol. 2, p. 504 (1922).
 49. *D. praevacua*, Meyrick, ibidem, Vol. 2, p. 504 (1922).
 50. *D. querxicola*, Meyrick, ibidem, Vol. 2, p. 433 (1921).
 51. *D. sciodora*, Meyrick, ibidem, Vol. 2, p. 504 (1922).
 52. *D. trinotella*, Coquillet, Papilio Vol. 3, p. 81 (1883).
 53. *D. ventrella*, Fitch, Trans. N. York Agric. Soc. Vol. 13, p. 234 (1853).
 uncipunctellus, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 123 (1863).
 querceella, Chambers, Canad. Ent. Vol. 4, p. 223 (1872).
 54. *D. caryifoliella*, Chambers, ibidem, Vol. 4, p. 224 (1872). (*caryaef...*)
 55. *D. hexasticta*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 94 (1911).
 56. *D. argentaria*, Meyrick, Ann. Transv. Mus. Vol. 3 p. 304 (1913).
 57. *D. eustacta*, Meyrick, ibidem, Vol. 8, p. 84 (1921).
 58. *D. metrodes*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 172 (1913).
 59. *D. chlorophracta*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 82 (1921).
 60. *D. cotifera*, Meyrick, ibidem, Vol. 3, p. 303 (1913).
 61. *D. fluitans*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 284 (1920).
 62. *D. antizyga*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 303 (1913).
 63. *D. marginella*, Fabricius, Spec. Ins. Vol. 2, p. 307 (1781).
 striatella, Hübner, Samml. Eur. Schmett. Tin. f. 154 (1796).
 clarella, Treitschke, Schmett. Eur. Vol. 9 (2), p. 54 (1833).
 64. *D. aulotoma*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 5 (1917).
 65. *D. pleuroleuca*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 169 (1919).
 66. *D. ventosa*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 304 (1913).
 67. *D. chalcophaea*, Meyrick, Exot. Microlep. Vol. 2, p. 434 (1921).
 68. *D. melanophylla*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 131 (1919).
 69. *D. peristylis*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 431 (1904).
 perleva, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 168 (1919).
 70. *D. lativittata*, Meyrick, Exot. Microlep. Vol. 2, p. 434 (1921).
 71. *D. polyommata*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 131 (1919).
 72. *D. melichroa*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 431 (1904).
 73. *D. lygropa*, Lower, Trans. Roy. Soc. S. Australia, Vol. 27, p. 70 (1903).
 74. *D. acrogypsa*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 168 (1919).
 75. *D. xuthochyta*, Turner, ibidem, Vol. 31, p. 168 (1919).
 76. *D. mesoetenis*, Meyrick, Exot. Microlep. Vol. 2, p. 434 (1921).
 77. *D. mendica*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 130 (1919).
 78. *D. achlyodes*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 432 (1904).
 79. *D. dysorata*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 170 (1919).
 80. *D. iodora*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 432 (1904).
 81. *D. holomelas*, Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 57 (1897).
 82. *D. cirrhostola*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 169 (1919).
 83. *D. apludella*, Lederer, Hor. Soc. Ent. Ross. Vol. 6, p. 92, pl. 5, f. 14 (1869).
 84. *D. cisti*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 244 (1859).
 85. *D. limbipunctella*, Staudinger, ibidem, Vol. 20, p. 245 (1859).
 millierellus, Stainton, Nat. Hist. Tin. Vol. 13, p. 336 (1873).
- Burma, Java.
 S. India, Ceylon.
 Java.
 Coorg, Java.
 Coorg, Kanara.
 China.
 China.
 Punjab.
 Assam.
 Illinois.
 E. United States.
- Kentucky, Texas, Mis-
 Mexico. [souri.]
- Transvaal.
 Rhodesia.
 India, Ceylon, S. Africa.
 Rhodesia.
 Transvaal.
 Natal.
 Transvaal.
 Europe, Siberia.
- Cape Colony.
 Queensland.
 Transvaal.
 Queensland.
 Queensland.
 Queensland, W. Aus-
 tralia.
- Queensland.
 New South Wales.
 New South Wales (?).
- South Australia.
 Queensland.
 Queensland.
 Queensland.
 Queensland.
 Queensland.
 New South Wales.
- New South Wales.
 Queensland.
 New South Wales.
 Queensland.
 N. Persia.
- Spain.
 S. France, Spain, Algeria.

86. *D. helianthemi*, Walsingham, Ent. M. Mag. Vol. 39, p. 265 (1903).
limbipunctella, Millière, Ann. Soc. Ent. Fr. p. 118, pl. 2, f. 9 (1885).
87. *D. neatodes*, Meyrick, Exot. Microlep. Vol. 3, p. 35 (1923).
88. *D. juniperella*, Linnaeus, Faun. Suec. no. 1449 (1761).
89. *D. ustulella*, Fabricius, Ent. Syst. Vol. 3 (2), p. 307 (1794).
capucinella, Hübner, Samml. Eur. Schmett. Tin. f. 159 (1796).
90. *D. fasciella*, Hübner, ibidem, f. 111 (1796).
91. *D. limosella*, Schläger, Ber. Lep. Tausch. p. 43 (1849).
deflectivella, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 154, f. 383 (1855).
92. *D. suffusella*, Chambers, Canad. Ent. Vol. 6, p. 243 (1874).
93. *D. ferrata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 174 (1913).
94. *D. clarescens*, Meyrick, ibidem, Vol. 22, p. 174 (1913).
95. *D. excoriata*, Meyrick, ibidem, Vol. 22, p. 174 (1913).
96. *D. oceanis*, Meyrick, Exot. Microlep. Vol. 2, p. 306 (1920). — PI. 3,
- Fig. 72.**
97. *D. mesoglena*, Meyrick, ibidem, Vol. 2, p. 619 (1923).
98. *D. imbricata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 175 (1913).
99. *D. eridantis*, Meyrick, ibidem, Vol. 17, p. 981 (1907).
100. *D. aphanopa*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 83 (1921).
101. *D. impigra*, Meyrick, ibidem, Vol. 3, p. 305 (1913).
102. *D. pladarota*, Meyrick, ibidem, Vol. 8, p. 84 (1921).
103. *D. meridionella*, Walsingham, Trans. Ent. Soc. Lond. p. 268, pl. 13, f. 38 (1881).
104. *D. hylurga*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 83 (1921).
105. *D. erixantha*, Meyrick, Exot. Microlep. Vol. 1, p. 279 (1914).
106. *D. scenites*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 371 (1909).
107. *D. oleata*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 305 (1913).
108. *D. aequata*, Meyrick, Trans. Ent. Soc. Lond. p. 282 (1914).
109. *D. baccata*, Meyrick, Exot. Microlep. Vol. 2, p. 621 (1923).
110. *D. brachymetra*, Meyrick, ibidem, Vol. 2, p. 620 (1923).
111. *D. squalens*, Meyrick, Trans. Ent. Soc. Lond. p. 282 (1914).
112. *D. mollis*, Barnes, Contrib. N. Amer. Lep. Vol. 4, p. 230 (1920).
113. *D. georgiella*, Walker, List Lep. Het. Brit. Mus. Vol. 35, p. 1827 (1866).
roseocostellus, Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 185 (1882).
114. *D. punctidiscella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 123 (1863).
stramineellus, Chambers, Canad. Ent. Vol. 4, p. 224 (1872).
115. *D. punctipennella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 156 (1860).
gracilella, Chambers, Canad. Ent. Vol. 4, p. 226 (1872).
116. *D. bipunctella*, Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 186 (1882).
117. *D. ingloria*, Meyrick, Exot. Microlep. Vol. 2, p. 621 (1923).
118. *D. heteracma*, Meyrick, ibidem, Vol. 2, p. 622 (1923).
119. *D. indigna*, Walsingham, Proc. Zool. Soc. Lond. p. 526 (1891).
120. *D. tactica*, Meyrick, Exot. Microlep. Vol. 2, p. 152 (1918).
121. ? *D. adactella*, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1027 (1864).
- Spain, France, Italy.
- Cyprus.
- Europe, Asia Minor.
- Europe.
- [Minor.]
- C. & S. Europe, Asia
- Germany, Switzerland,
Asia Minor, S.E. Siberia.
- Texas.
- Assam.
- Ceylon.
- Assam.
- China, Japan.
- Coorg.
- Coorg.
- Bengal.
- Rhodesia.
- Transvaal.
- Rhodesia.
- Natal.
- Rhodesia.
- Nyassaland, Rhodesia.
- Cape Colony.
- Transvaal.
- Guiana, Brazil.
- Brazil.
- Peru.
- Guiana, Brazil.
- Arizona.
- Georgia, Missouri.
- E. United States, Canada.
- E. United States, Canada.
- Massachusetts.
- Peru.
- Brazil, Peru.
- Windward Islands.
- Ecuador.
- ? Australia.

236. GENUS HYLOGRAPTIS, MEYRICK

Hylograptis, Meyrick, Trans. Ent. Soc. Lond. p. 450 (1910). — Type : *H. thryptica*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with

dense scales, forming a long expansible tuft towards apex above, terminal joint much shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough scales above. Forewings with 2 and 3 stalked, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, termen sinuate, cilia 4/5; with cubital pecten; 3 and 4 stalked, 5 parallel, 6 and 7 stalked.

Remarks. — Correlated with *Dichomeris*.

Geographical distribution of species. — Papuan.

Larva unknown.

1. *H. thryptica*, Meyrick, Trans. Ent. Soc. Lond. p. 451 (1910). — Pl. 4, New Guinea.
Fig. 96.

237. GENUS VAZUGADA, WALKER

Vazugada, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 803 (1864). — Type : *V. abscessella*, Walker.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, minutely ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with scales, roughly projecting towards apex above and at apex, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 2/3; with cubital pecten; 3 and 4 connate, 5 rather approximated, 6 and 7 closely approximated towards base.

Remarks. — Nearly related to *Dichomeris* and *Cymotricha*.

Geographical distribution of species. — Tropical American.

Larva unknown.

- | | |
|---|-----------------|
| 1. <i>V. costalis</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 18 (1914). | Panama. |
| 2. <i>V. amphicoma</i> , Meyrick, Trans. Ent. Soc. Lond. 1911, p. 695 (1912). | Brazil. |
| 3. <i>V. leucostena</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 94, pl. 3, f. 12 (1911). | Mexico. |
| 4. <i>V. macrosphena</i> , Meyrick, Trans. Ent. Soc. Lond. p. 175 (1913). | Brazil. |
| 5. <i>V. perenacma</i> , Meyrick, Exot. Microlep. Vol. 2, p. 623 (1923). | Brazil, Peru. |
| 6. <i>V. abscessella</i> , Walker, List. Lep. Het. Brit. Mus. Vol. 28, p. 536 (1863).
<i>strigiplenella</i> , Walker, ibidem, Vol. 29, p. 803 (1864);
<i>zonostoma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 281 (1914). | Guiana, Brazil. |

238. GENUS GAESA, WALKER

Gaesia, Walker, List. Lep. Het. Brit. Mus. Vol. 29, p. 803 (1864). — Type : *G. decusella*, Walker.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi long, recurved, second joint triangularly expanded with dense scales, forming a compact projecting apical tuft beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle (rarely approximated), 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen sinuate, cilia 2/5-3/5; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated at base or stalked.

Remarks. — Probably correlated with *Dichomeris*, but indicating a quite distinct branch from a common origin.

Geographical distribution of species. — Indo-Malayan, African, and Australian, with stragglers in Europe and South America.

Larva unknown.

- | | | |
|--|---|---|
| 1. <i>G. marmorata</i> , Walsingham, Trans. Ent. Soc. Lond. p. 108, pl. 5 f. 44 (1891). | <i>basistriata</i> , Walsingham, ibidem, p. 41, pl. 3, f. 13 (1897). | Gambia, French Congo,
Rhodesia, Transvaal. |
| 2. <i>G. olivescens</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 175 (1913). | | Ceylon. |
| 3. <i>G. decusella</i> , Walker, List. Lep. Het. Brit. Mus. Vol. 29, p. 804 (1864). | <i>alternella</i> , Walker, ibidem, Vol. 30, p. 1023 (1864).
<i>Granti</i> , Walsingham, Bull. Liverp. Mus. Vol. 3, p. 2 (1900).
<i>thoracella</i> , Walsingham, ibidem, Vol. 3, p. 3 (1900). | India, Ceylon, Sokotra. |
| 4. <i>G. barbella</i> , Hübner, Samml. Eur. Schmett. Tin. f. 291 (1805). | | Bavaria, Bohemia, Asia |
| 5. <i>G. sparsella</i> , Christoph, Bull. Soc. Nat. Mosc. p. 29 (1882). | | E. Siberia. [Minor.] |
| 6. <i>G. thanatopsis</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 25, p. 82 (1901). | | Queensland. |
| 7. <i>G. capnites</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 435 (1904). | | Queensland, South Aus- |
| 8. <i>G. dryinodes</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 58 (1897). | | Queensland. [tralia.] |
| 9. <i>G. ptychosema</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 175 (1913). | | Assam. |
| 10. <i>G. microsphena</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 166 (1921). | | Assam, Java. |
| 11. <i>G. bisignella</i> , Snellen, Tijdschr. v. Ent. Vol. 28, p. 30, pl. 3, f. 12 (1885). | <i>deltaspis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 601 (1905). | India, Ceylon, Celebes. |
| 12. <i>G. agathopa</i> , Meyrick, Ann Transv. Mus. Vol. 8, p. 85 (1921). | | Rhodesia. |
| 13. <i>G. xanthodeta</i> , Meyrick, ibidem, Vol. 3, p. 305 (1913). | | Transvaal. |
| 14. <i>G. melitura</i> , Meyrick, Exot. Microlep. Vol. 1, p. 585 (1916). | | Kanara. |
| 15. <i>G. rhodophaea</i> , Meyrick, Voyage All. Jean. Lép. Vol. 2, p. 73 (1920). | | Tanganyika Territ. |
| 16. <i>G. leucothicta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 235 (1919). | | Bombay. |
| 17. <i>G. formulata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 110 (1922). | | Brazil. |
| 18. <i>G. ampliata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 175 (1913). | | Assam, Ceylon. |

239. GENUS OXYCRYPTIS, MEYRICK

Oxycryptis, Meyrick, Trans. Ent. Soc. Lond. 1911, p. 692 (1912). — Type : *O. attonita*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint relatively short, clothed with dense scales, rough beneath, terminal joint much longer than second, thickened with dense scales, somewhat roughly expanded posteriorly throughout and concealing apex. Maxillary palpi very short, filiform, appressed to tongue. Forewings 2 from near angle, 3 and 4 closely approximated at base, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia 4/5; with cubital pecten; 3 and 4 closely approximated at base, 5 approximated, 6 and 7 connate or closely approximated.

Remarks. — A peculiarly modified form of uncertain affinity.

Geographical distribution of species. — Tropical American.

Larva unknown.

- | | |
|---|-------------------------|
| 1. <i>O. trigonota</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 58 (1911). | Mexico, Panama, Guiana. |
| 2. <i>O. attonita</i> , Meyrick, Trans. Ent. Soc. Lond. 1911, p. 692 (1912). | Colombia. |

240. GENUS MYROPHILA, MEYRICK

Myrophila, Meyrick, Exot. Microlep. Vol. 2, p. 624 (1923). — Type : *M. carycina*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, shortly ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with scales roughly expanded before apex above, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 1/3-2/5; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate or closely approximated.

Remarks. — A simple derivative of *Cymotricha*.

Geographical distribution of species. — South American and Malayan.

Larva unknown.

- | | |
|--|-----------------|
| 1. <i>M. traumatis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 625 (1923). | Borneo. |
| 2. <i>M. caryophragma</i> , Meyrick, ibidem, Vol. 2, p. 624 (1923). | Guiana, Brazil. |
| 3. <i>M. carycina</i> , Meyrick, Trans. Ent. Soc. Lond. p. 280 (1914). | Guiana, Brazil. |
| 4. <i>M. diacnista</i> , Meyrick, Exot. Microlep. Vol. 2, p. 624 (1923). | Guiana. |

241. GENUS BROCHOMETIS, MEYRICK

Brochometis, Meyrick, Exot. Microlep. Vol. 2, p. 625 (1923). — Type : *B. plexigramma*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli posterior; tongue developed. Antennae 5/6, in ♂ serrate, moderately ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales forming a rough tuft at apex beneath, terminal joint shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia 1/3; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated at base.

Remarks. — A derivative from *Cymotricha*.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|---|-----------------------|
| 1. <i>B. plexigramma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 110 (1922). | Guiana, Brazil, Peru. |
|---|-----------------------|

242. GENUS EUPOLIS, MEYRICK

Eupolis, Meyrick, Exot. Microlep. Vol. 2, p. 625 (1923). — Type : *E. stygnota*, Walsingham.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ moderately ciliate, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense appressed scales, slightly rough beneath, terminal joint as long as second, rather stout, slightly roughened anteriorly, pointed. Maxillary palpi very short,

filiform, appressed to tongue. Posterior tibiae with appressed scales. Forewings with 2 and 3 stalked from near angle, 4 from angle, 7 absent, 11 from middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia 1/4; with cubital pecten; 3 and 4 connate, 5 parallel, 6 and 7 connate.

Remarks. — Related to the preceding.

Geographical distribution of species. — Central and South American.

Larva unknown.

1. *E. stygnotis*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 111, Panama, Brazil, Peru. pl. 3, f. 32 (1911).

243. GENUS PROPHORULA, MEYRICK

Prophorula, Meyrick, Trans. Ent. Soc. Lond. p. 105 (1922). — Type: *P. pyrrhopis*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ moderately ciliated, basal joint elongate, without pecten. Labial palpi extremely long, straight, porrected, second joint extremely long, rough-scaled above throughout with scales longer towards base, and beneath on apical half with rough projecting scales becoming longer towards apex, terminal joint very short, filiform, pointed, almost concealed in scales of second. Maxillary palpi obsolete. Posterior tibiae rough-haired above and beneath. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 2/3; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 short-stalked.

Remarks. — Apparently a development of *Neochrista*.

Geographical distribution of species. — South American.

Larva unknown.

1. *P. pyrrhopis*, Meyrick, Trans. Ent. Soc. Lond. p. 106 (1922). Brazil.

244. GENUS NEOCHRISTA, MEYRICK

Neochrista, Meyrick, Exot. Microlep. Vol. 2, p. 625 (1923). — Type: *N. auritogata*, Walsingham.

Characters. — Head with appressed scales, sidetufts slightly raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ fasciculate-ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint triangularly expanded with dense scales, roughly projecting along terminal edge, terminal joint much shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with appressed scales. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 1/3; with cubital pecten; 3 and 4 connate or short-stalked, 5 parallel, 6 and 7 closely approximated at base.

Remarks. — Derived from *Cymotricha*.

Geographical distribution of species. — Central and South American.

Larva unknown.

1. *N. auritogata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 85, Panama, Guiana, Brazil. pl. 3, f. 6 (1911).

pyropis, Meyrick, Exot. Microlep. Vol. 2, p. 149 (1918).

245. GENUS SEMIOMERIS, MEYRICK

Semiomeris, Meyrick, Exot. Microlep. Vol. 2, p. 626 (1923). — Type : *S. pyretodes*, Meyrick.

Characters. — Head with appressed scales, sidetufts slightly raised; ocelli posterior; tongue developed. Antennae 5/6, in ♂ serrulate, minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with scales roughly expanded above towards apex, and forming a tufted apical projection beneath, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 2/5; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated at base.

Remarks. — Probably correlated with the preceding.

Geographical distribution of species. — South American.

Larva unknown.

1. *S. pyretodes*, Meyrick, Trans. Ent. Soc. Lond. p. 277 (1914).

Guiana, Brazil, Peru.

246. GENUS CATELAPHRIS, NOV. GEN.

Type : *C. torrefacta*, Meyrick.

Characters. — Head smooth; sidetufts slightly raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ rather strongly and evenly ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, somewhat expanded loosely towards apex above, terminal joint somewhat shorter than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely haired above. Forewings with 1b furcate, 2 and 3 stalked, 8 and 9 out of 7, 7 to costa just above apex, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 1; with cubital pecten; 3 and 4 connate or stalked, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — A development of *Cymotricha*.

Geographical distribution of species. — South African.

Larva unknown.

1. *C. torrefacta*, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 245 (1914).

Transvaal.

247. GENUS AULIDIOTIS, NOV. GEN.

Type : *A. phoxopterella*, Snellen.

Characters. — Head with appressed scales, sidetufts raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint in ♂ much longer than second, stout, compressed, pointed, with slight groove on inner side, in ♀ rather longer than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with

1 b furcate, 2 from angle, 3 absent, 4 connate, 5 nearly approximated, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings over 1, trapezoidal, termen sinuate, cilia 2/5; without cubital pecten; 3 and 4 connate, 5 rather approximated, 6 and 7 stalked.

Remarks. — Probably a derivative of *Sphagiocrates*.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

1. *A. phoxopterella*, Snellen, Tijdschr. v. Ent. Vol. 46, p. 41, pl. 4, f. 11 (1903). Assam, Java.

248. GENUS SPHAGIOCRATES, NOV. GEN.

Type : *S. lusoria*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, stout, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint longer than second, stout, compressed, pointed. Maxillary palpi very short, filiform, appressed to tongue. Forewings with 1 b furcate, 2 and 3 long-stalked from angle, 4 connate from angle, 5 closely approximated, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings over 1, trapezoidal, apex pointed, termen sinuate, cilia 1/3; with cubital pecten; cell short, 3 and 4 short-stalked, 5 nearly approximated at base, straight, 6 and 7 short-stalked.

Remarks. — Apparently allied to *Eporastis*.

Geographical distribution of species. — Malayan.

Larva unknown.

1. *S. lusoria*, Meyrick, Zool. Med. Leid. Vol. 7, p. 87 (1922). Java, Sumatra.

249. GENUS DEOCLONA, BUSCK

Deoclona, Busck, Proc. U. S. Mus. Vol. 25, p. 837 (1903). — Type: *D. yuccacella*, Busck.

Proclesis, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 83 (1911). — Type: *D. xanthoselena*, Walsingham.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint with appressed scales, terminal joint much shorter, 1/4-2/3 of second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with dense hairs above and beneath. Forewings with 1 b furcate, 2 from near angle, 7 and 8 stalked, 7 to costa, 9 almost connate or out of 7, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 1; without cubital pecten; 3 and 4 stalked, 5 absent, 6 and 7 stalked.

Remarks. — Of quite uncertain affinity, but possibly referable here.

Geographical distribution of species. — American.

Larva (*yuccacella*) feeding in seed-capsules.

Foodplant *Yucca* (*Liliaceae*).

1. *D. xanthoselena*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 83, Panama, Guiana, Brazil.
pl. 3, f. 4 (1911) (-ne). — **Pl. 4, Fig. 95.**
2. *D. yuccacella*, Busck, Proc. U. S. Mus. Vol. 25, p. 837 (1903) (-asella). California.

250. GENUS IOCHARES, MEYRICK

Iochares, Meyrick, Ann. Transv. Mus. Vol. 8, p. 81 (1921). — Type : *I. festa*, Meyrick.

Characters. — Head with appressed hair-scales; ocelli posterior; tongue developed. Antennae 2/3, basal joint elongate, without pecten. Labial palpi very long, second joint long, subascending, densely clothed with rough projecting scales above and slightly rough scales beneath, terminal joint erect, half second, scaled, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long rough hairs above. Forewings with 2 and 3 connate or approximated from angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen faintly sinuate, cilia 4/5; with cubital pecten; 3 and 4 short-stalked, 5 somewhat approximated to 4, 6 and 7 nearly approximated towards base.

Remarks. — A development of *Eporgastis*.

Geographical distribution of species. — African.

Larva unknown.

- | | |
|---|-------------------------|
| 1. <i>I. straminis</i> , Walsingham, Trans. Ent. Soc. Lond. p. 266, pl. 12, f. 36 (1881). | Transvaal, Natal, Port. |
| 2. <i>I. festa</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 81 (1921). | Transvaal. [E. Africa.] |

251. GENUS EPORGASTIS, MEYRICK

Eporgastis, Meyrick, Ann. Transv. Mus. Vol. 8, p. 81 (1921). — Type : *E. naturata*, Meyrick.

Characters. — Head with appressed scales, sidetuffs projecting over forehead; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ serrulate, rather strongly ciliated, basal joint elongate, without pecten. Labial palpi very long, curved, obliquely ascending, second joint very long, much thickened with dense rather rough scales beneath, and more or less expanded with scales sometimes rough above, terminal joint half second, thickened with scales, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 2 and 3 stalked from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen somewhat sinuate, cilia 3/4; with cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate.

Remarks. — Derived from *Cymotricha*.

Geographical distribution of species. — African.

Larva unknown.

- | | |
|--|-----------|
| 1. <i>E. torrescens</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 82 (1921). | Rhodesia. |
| 2. <i>E. naturata</i> , Meyrick, ibidem, Vol. 8, p. 82 (1921). | Rhodesia. |
| 3. <i>E. conclusa</i> , Meyrick, ibidem, Vol. 6, p. 23 (1918). | Natal. |

252. GENUS CARBATINA, MEYRICK

Carbatina, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 181 (1913). — Type : *C. picrocarpa*, Meyrick.

Characters. — Head with appressed scales, sidetuffs somewhat raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ ciliated, basal joint moderate, without pecten. Labial palpi long, recurved,

second joint clothed with dense appressed scales rather roughly expanded above towards apex, terminal joint rather shorter than second, thickened with loose scales, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen somewhat sinuate, cilia 2/3; with cubital pecten; 3 and 4 connate, 5 approximated, 6 and 7 stalked.

Remarks. — Also a derivative of *Cymotricha*.

Geographical distribution of species. — Indian, extending to Japan.

Larva unknown.

1. *C. levigata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 182 (1913). Ceylon.
 2. *C. picrocarpa*, Meyrick, ibidem, Vol. 22, p. 182 (1913). Assam, Japan.

253. GENUS STRYPHNOCOPA, MEYRICK

Stryphnocopa, Meyrick, Exot. Microlep. Vol. 2, p. 306 (1920). — Type : *S. trinotata*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with scales rather expanded towards apex above and somewhat projecting angularly at apex beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to termen, 9 almost connate with 7, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen faintly sinuate, cilia 3/5; without cubital pecten; 3 and 4 connate, 5 absent, 6 and 7 stalked.

Remarks. — A modified form, of which the immediate relationship is doubtful.

Geographical distribution of species. — Indian.

Larva unknown.

1. *S. trinotata*, Meyrick, Exot. Microlep. Vol. 2, p. 307 (1920). Assam.

254. GENUS MYTHOGRAPHA, MEYRICK

Mythographa, Meyrick, Exot. Microlep. Vol. 2, p. 626 (1923). — Type : *M. chartaria*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, roughly expanded towards apex above, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 2 and 3 stalked, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia 3/5; with cubital pecten; 3 and 4 short-stalked, 5 somewhat approximated, 6 and 7 closely approximated at base.

Remarks. — A development of *Cymotricha*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *M. chartaria*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 178 (1913). Ceylon.

255. GENUS MACHLOTRICA, MEYRICK

Machlotricha, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 61 (1912). — Type : *M. caeca*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi very long, porrected, second joint very long, straight, densely rough-scaled above and with very long rough projecting scales beneath, terminal joint directed obliquely sideways, shorter than second, slender, acute. Maxillary palpi minute, filiform, appressed to tongue. Posterior tibiae smooth-scaled. Forewings with 2 and 3 stalked from angle, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 1; 4 absent, 5 somewhat approximated to 3, 6 and 7 stalked.

Remarks. — Probably referable in this neighbourhood.

Geographical distribution of species. — African.

Larva unknown.

1. *M. caeca*, Meyrick, Ann. S. Afr. Mus. Vol. 10, p. 62 (1912).

Zululand.

256. GENUS PACHYSARIS, MEYRICK

Pachysaris, Meyrick, Trans. Ent. Soc. Lond. p. 276 (1914). — Type : *P. rurigena*, Meyrick.

Characters. — Head with appressed scales, sidetufts roughly spreading; ocelli posterior; tongue developed. Antennae 5/6, in ♂ moderately ciliated, basal joint moderate, without pecten. Labial palpi very long, curved, ascending, second joint much thickened with dense scales, above with long dense projecting hairs on apical half, in ♀ sometimes expandible interiorly, terminal joint much shorter than second, more or less roughened posteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with rough projecting hairs above and beneath. Forewings with 1 b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, apex obtuse, termen faintly sinuate, cilia 1/2; with cubital pecten; 3 and 4 connate, 5 approximated at base, 6 and 7 connate or closely approximated.

Remarks. — A specialised modification of *Cymotricha*.

Geographical distribution of species. — South American.

Larva unknown.

1. *P. contrita*, Meyrick, Trans. Ent. Soc. Lond. p. 105 (1922).

Guiana, Brazil.

2. *P. collina*, Meyrick, ibidem, p. 277 (1914).

Peru.

3. *P. pacifensis*, Meyrick, Exot. Microlep. Vol. 2, p. 626 (1923).

Brazil.

4. *P. rurigena*, Meyrick, Trans. Ent. Soc. Lond. p. 277 (1914).

Guiana.

257. GENUS TOCMIA, WALKER

Tocmia, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 805 (1864). — Type : *T. versicolorella*, Walker.

Characters. — Head smooth; tongue developed. Antennae nearly 1, in ♂ simple. Labial palpi very long, second joint porrected, with rough projecting scales above and beneath, terminal joint less

than half second, erect, slender, acute. Forewings with 2 and 3 stalked (?), 7 and 8 stalked, 7 to apex. Hindwings trapezoidal; 3 and 4 connate, 6 and 7 approximated at base.

Remarks. — This is probably a good genus, but the structure has not been adequately ascertained owing to the deteriorated condition of the typical example, which is still unique.

Geographical distribution of species. — South American.

Larva unknown.

1. *T. versicolorella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 806 (1864). Brazil.

258. GENUS SCHEMATISTIS, MEYRICK

Schematistis, Meyrick, Ann. Transv. Mus. Vol. 3, p. 67 (1912). — Type: *S. analoxa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrate, minutely ciliated, basal joint elongate, without pecten. Labial palpi very long, slightly curved, subascending, second joint with rough projecting scales above and long dense projecting tuft beneath, terminal joint shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1, trapezoidal, termen hardly sinuate, cilia 3/4; 3 from angle, 4-7 tolerably parallel.

Remarks. — A dubious insect, but perhaps a form of this group with exceptionally modified neuration; the specimen is no longer before me.

Geographical distribution of species. — African.

Larva unknown.

1. *S. analoxa*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 68 (1912).

259. GENUS PARISTHmia, MEYRICK

Paristhmia Meyrick, Ann. Transv. Mus. Vol. 2, p. 13 (1909). — Type: *P. barathrodes*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ shortly ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with dense appressed scales, slightly expanded towards apex above, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 1; in ♂ with some fine long expandible hairs at base of lower margin of cell, representing cubital pecten; 2 and 3 stalked from much before angle, 4 from angle, 5 somewhat approximated at base, 6 and 7 stalked.

Remarks. — A modification of *Cymotricha*, of which the peculiar neuration is perhaps sexual only.

Geographical distribution of species. — African.

Larva unknown.

1. *P. barathrodes*, Meyrick, Ann. Transv. Mus. Vol. 2, p. 13, pl. 5, f. 1 (1909). Transvaal.

260. GENUS CYMOTRICA, MEYRICK

Cymotricha, Meyrick, Exot. Microlep. Vol. 2, p. 626 (1923). — Type : *C. miltohrauma*, Meyrick.

Oxysactis, Meyrick, ibidem, Vol. 3, p. 35 (1923). — Type : *C. sciritis*, Meyrick.

Characters. — Head with appressed scales: ocelli posterior; tongue developed. Antennae 4/5, in ♂ shortly or moderately ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, more or less roughly expanded towards apex above and sometimes on terminal edge, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with appressed or rather rough scales above. Forewings with 1 b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to costa, or seldom 7 absent, 11 from middle. Hindwings 1 or over 1, trapezoidal, apex obtuse, termen slightly sinuate, cilia 1/3-2/3; with cubital pecten; 3 and 4 connate or short-stalked, 5 somewhat approximated, 6 and 7 connate or closely approximated at base.

Remarks. — Correlated with *Trichotaphe* as derivatives of a form approaching *Brachmia*.

Geographical distribution of species. — South American, African, and Indian, with stragglers in North America, Europe, and Madagascar; some South American and African species show near affinity. Hence it seems likely that the genus originated in South America, which possesses the largest number of species, and was transmitted thence to South Africa, whence the comparatively limited Indian contingent was derived; compare the case of *Polyhymno*.

Larva (2 known) feeding in spun leaves.

Foodplants *Combretaceae*, *Euphorbiaceae*.

- | | |
|---|------------------|
| 1. <i>C. macroxyla</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 180 (1913). | Assam. |
| 2. <i>C. cinnamicostella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 332, pl. 4, f. 105 (1877). | Panama. |
| 3. <i>C. thalpodes</i> , Meyrick, Trans. Ent. Soc. Lond. p. 111 (1922). | Brazil, Peru. |
| 4. <i>C. procyphodes</i> , Meyrick, ibidem, p. 115 (1922). | Brazil. |
| 5. <i>C. fluctuans</i> , Meyrick, Exot. Microlep. Vol. 3, p. 2 (1923). | Peru. |
| 6. <i>C. miltohrauma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 115 (1922). — PI. 4. | Brazil, Peru. |
| Fig. 77. | |
| 7. <i>C. fracticostella</i> , Walsingham, ibidem, p. 110, pl. 5, f. 45 (1891). | Gambia. |
| 8. <i>C. cymatodes</i> , Meyrick, Exot. Microlep. Vol. 1, p. 584 (1916). | Assam, Tonkin. |
| 9. <i>C. cyclospila</i> , Meyrick, ibidem, Vol. 2, p. 151 (1918). | Guiana. |
| 10. <i>C. cinctella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 619 (1864).
<i>subrutila</i> , Meyrick, Exot. Microlep. Vol. 3, p. 1 (1923). | Brazil, Peru. |
| 11. <i>C. oenombra</i> , Meyrick, Vol. 1, p. 201 (1914). | Nyassaland. |
| 12. <i>C. furvella</i> , Zeller, Micr. Caffr. p. 115 (1852). | Natal, Zululand. |
| 13. <i>C. monococca</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 85 (1921). | Port. E. Africa. |
| 14. <i>C. physeta</i> , Meyrick, ibidem, Vol. 3, p. 302 (1913). | Transvaal. |
| 15. <i>C. oxygrapha</i> , Meyrick, ibidem, Vol. 3, p. 302 (1913). | Transvaal. |
| 16. <i>C. melanosoma</i> , Meyrick, Voyage All. Jean. Lép. Vol. 2, p. 75 (1920). | Kenya Colony. |
| 17. <i>C. coarctata</i> , Walsingham, Trans. Ent. Soc. Lond. p. 252, pl. 11, f. 20 (1881). | Natal. |
| 18. <i>C. famosa</i> , Meyrick, Exot. Microlep. Vol. 1, p. 202 (1914). | Nyassaland. |
| 19. <i>C. pyrrhitis</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 68 (1912). | Transvaal. |
| 20. <i>C. ironica</i> , Meyrick, ibidem, Vol. 2, p. 17, pl. 6, f. 1 (1909). | Transvaal. |
| 21. <i>C. hercogramma</i> , Meyrick, ibidem, Vol. 8, p. 86 (1921). | Rhodesia. |
| 22. <i>C. geochrota</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 775 (1914). | Bombay. |
| 23. <i>C. mochlopis</i> , Meyrick, Exot. Microlep. Vol. 3, p. 1 (1923). | Brazil. |

24. <i>C. pleuropa</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 86 (1921).	Natal.
25. <i>C. pseudometra</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 178 (1913).	Coorg, Kanara.
26. <i>C. sciritis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 149 (1918).	Madras.
27. <i>C. centracma</i> , Meyrick, ibidem, Vol. 3, p. 4 (1923).	Bombay.
28. <i>C. pelitis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 179 (1913).	Assam.
29. <i>C. pudicella</i> , Mann, Wien. Ent. Monatsschr. Vol. 5, p. 190, pl. 3, f. 10 (1861).	Hungary, Dalmatia, Asia
30. <i>C. syngrapta</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 85 (1921).	Rhodesia. [Minor.
31. <i>C. ochroxesta</i> , Meyrick, ibidem, Vol. 8, p. 86 (1921).	Transvaal.
32. <i>C. homaloxesta</i> , Meyrick, ibidem, Vol. 8, p. 86 (1921).	Rhodesia.
33. <i>C. tepeps</i> , Meyrick, Exot. Microlep. Vol. 3, p. 2 (1923).	Madagascar.
34. <i>C. condylodes</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 85 (1921).	Rhodesia.
35. <i>C. claviculata</i> , Meyrick, ibidem, Vol. 2, p. 17, pl. 5, f. 10 (1909).	Transvaal.
36. <i>C. byrsoxantha</i> , Meyrick, ibidem, Vol. 6, p. 23 (1918).	Natal.
37. <i>C. intentella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 627 (1864).	Brazil, Peru.
38. <i>C. directa</i> , Meyrick, Trans. Ent. Soc. Lond. 1911, p. 694 (1912).	Venezuela.
39. <i>C. abortiva</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 98, pl. 3, f. 21 (1911).	Guatemala.
40. <i>C. amauropis</i> , Meyrick, Exot. Microlep. Vol. 3, p. 1 (1923).	Peru.
41. <i>C. tristicta</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 17 (1914).	Panama.
42. <i>C. rubiginosella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 745 (1864).	Guiana, Brazil, Peru.
43. <i>C. ochropyga</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 98 (1911).	Mexico.
44. <i>C. serena</i> , Meyrick, Trans. Ent. Soc. Lond. p. 23 (1909).	Bolivia.
45. <i>C. excepta</i> , Meyrick, Exot. Microlep. Vol. 1, p. 279 (1914).	Nyassaland.
46. <i>C. externella</i> , Zeller, Micr. Caffr. p. 109 (1852).	Transvaal.
47. <i>C. turrita</i> , Meyrick, Trans. Ent. Soc. Lond. p. 279 (1914).	Guiana, Brazil.
48. <i>C. designatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 619 (1864).	Guiana, Brazil, Peru.
49. <i>C. sphycroopa</i> , Meyrick, Exot. Microlep. Vol. 2, p. 150 (1918).	Guiana.
50. <i>C. themelia</i> , Meyrick, Trans. Ent. Soc. Lond. p. 175 (1913).	Brazil.
51. <i>C. subdentata</i> , Meyrick, ibidem, p. 113 (1922).	Brazil.
52. <i>C. ptilocompa</i> , Meyrick, ibidem, p. 113 (1922).	Brazil, Peru.
53. <i>C. sumptella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 609 (1864).	Brazil, Peru.
54. <i>C. permundella</i> , Walker, ibidem, Vol. 29, p. 608 (1864). tactella, Walker, ibidem, Vol. 29, p. 608 (1864).	Guiana, Brazil, Peru.
55. <i>C. thalamopa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 112 (1922).	Brazil.
56. <i>C. ligyra</i> , Meyrick, Ann. Trans. Mus. Vol. 3, p. 301 (1913).	Transvaal.
57. <i>C. ellipsias</i> , Meyrick, Trans. Ent. Soc. Lond. p. 114 (1922).	Peru.
58. <i>C. trigonella</i> , Walsingham, Proc. Zool. Soc. Lond. p. 523 (1891).	Windward Islands.
59. <i>C. jugata</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 97 (1911).	Mexico, Guatemala,
60. <i>C. melissa</i> , Walsingham, ibidem, Vol. 4, p. 97 (1911).	Panama. [Panama.
61. <i>C. hortulana</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 23 (1918).	Transvaal.
62. <i>C. thesmiopa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 114 (1922).	Brazil.
63. <i>C. melanota</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 94, pl. 3, f. 13 (1911).	Mexico.
64. <i>C. memnonia</i> , Meyrick, Trans. Ent. Soc. Lond. p. 176 (1913).	Brazil.
65. <i>C. ostensella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 618 (1864).	Guiana, Brazil.

261. GENUS DEIMNESTRA, MEYRICK

Deimnestra, Meyrick, Exot. Microlep. Vol. 2, p. 150 (1918). — Type : *D. thrysicola*, Meyrick.

Characters. — Head with appressed scales; ocelli moderate, posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi extremely long, second joint very long

straight, porrected, densely scaled, above with rough projecting hair-scales diminishing towards apex, terminal joint much shorter than second, obliquely ascending, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 1^b furcate, 2 and 3 stalked, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 3/5; without cubital pecten; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated at base.

Remarks. — Probably a development of *Trichotaphe*.

Geographical distribution of species. — Indian and African.

Larva unknown.

1. *D. chalybitis*, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 75 (1920). Tanganyika Territ.
2. *D. thysicola*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 171 (1913). Assam.

262. GENUS NOEZA, WALKER

Noeza, Walker, List Lep. Het. Brit. Mus. Vol. 35, p. 1839 (1866). — Type : *N. telegraphella*, Walker.

Characters. — Head with appressed scales, sidetufts somewhat raised; ocelli very small, posterior; tongue developed. Antennae 1, in ♂ rather strongly ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint broadly thickened with dense scales, triangularly expanded above and with projecting apical tuft, terminal joint longer than second, very slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with tolerably appressed hair-scales above. Forewings with 2 and 3 stalked, 6 to apex, 7 absent, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 1/3; without cubital pecten; 3 and 4 connate or stalked, 5 nearly parallel, 6 and 7 closely approximated at base.

Remarks. — A peculiar form of which the immediate affinity is uncertain.

Geographical distribution of species. — South American.

Larva unknown.

1. *N. telegraphella*, Walker, List Lep. Het. Brit. Mus. Vol. 35, p. 1839 (1866). Brazil.

263. GENUS ILINGIOTIS, MEYRICK

Ilingiotis, Meyrick, Trans. Ent. Soc. Lond. p. 275 (1914). — Type : *I. sevestella*, Walker.

Sirogenes, Meyrick, Exot. Microlep. Vol. 3, p. 3 (1923). — Type : *I. thermophaea*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae almost 1, in ♂ serrulate, minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with scales triangularly expanded and rough on terminal edge, sometimes with projecting apical tuft beneath, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1^b furcate, 2 and 3 stalked, 7 absent, 8 and 9 sometimes stalked, 11 from middle. Hindwings 1, trapezoidal, apex obtuse, termen slightly sinuate, cilia 1; without cubital pecten; 3 and 4 connate, 5 approximated towards base, 6 and 7 connate.

Remarks. — Perhaps really allied to *Cymotricha*, but without cubital pecten.

Geographical distribution of species. — South American, with one South African species in all respects very similar.

Larva unknown.

- | | |
|--|-----------------|
| 1. <i>I. tephrodes</i> , Meyrick, Ann. Transv. Mus. Vol. 2, p. 18, pl. 6, f. 2 (1909). | Transvaal. |
| 2. <i>I. thrasynta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 275 (1914). | Guiana. |
| 3. <i>I. vigilans</i> , Meyrick, ibidem, p. 275 (1914). | Guiana. |
| 4. <i>I. thermophaea</i> , Meyrick, Exot. Microlep. Vol. 3, p. 3 (1923). | Brazil, Peru. |
| 5. <i>I. hemeropa</i> , Meyrick, ibidem, Vol. 3, p. 2 (1923). | Brazil. |
| 6. <i>I. sevectella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1020 (1864). | Guiana, Brazil. |

264. GENUS SATHROGENES, MEYRICK

Sathrogenes, Meyrick, Exot. Microlep. Vol. 3, p. 2 (1923). — Type : *S. malachias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, triangularly expanded with dense scales, with rough projecting hairscales along terminal edge, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 3/5; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 connate or closely approximated.

Remarks. — Presumably derived from *Trichotaphe*.

Geographical distribution of species. — Indian and African.

Larva unknown.

- | | |
|---|-----------|
| 1. <i>S. xestobyrsa</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 82 (1921). | Rhodesia. |
| 2. <i>S. planata</i> , Meyrick, Rec. Ind. Mus. Vol. 5, p. 222 (1910). | Punjab. |
| 3. <i>S. malachias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 179 (1913). | Assam. |

265. GENUS HOLAXYRA, MEYRICK

Holaxyra, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 176 (1913). — Type : *H. ampycota*, Meyrick.

Characters. — Head with appressed scales, sidetuffs loose; ocelli posterior; tongue developed. Antennae 3/4, in ♂ ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, porrected, second joint expanded with dense rough projecting scales above and beneath, longest towards base above and towards apex beneath, terminal joint much shorter than second, obliquely ascending, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal-ovate, termen rather sinuate, cilia 2/5; without cubital pecten; 3 and 4 connate, 5 rather approximated, 6 and 7 approximated at base or short-stalked.

Remarks. — A development of *Trichotaphe*.

Geographical distribution of species. — Indian and African; the Australian and North American species, which I have not seen, are conjectural.

Larva (*deflecta*) feeding in spun leaves.

Foodplant *Thurberia* (*Malvaceae*).

- | | |
|--|-------------|
| 1. <i>H. picrophanes</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 304 (1913). | Transvaal. |
| 2. <i>H. latipalpis</i> , Walsingham, Trans. Ent. Soc. Lond. p. 265, pl. 12, f. 35 (1881). | Natal. |
| 3. <i>H. isoclera</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 176 (1913). | Ceylon. |
| 4. <i>H. ampycota</i> , Meyrick, ibidem, Vol. 22, p. 176 (1913). | Ceylon. |
| 5. <i>H. ancylosticha</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 169 (1919). | Queensland. |
| 6. ? <i>H. deflecta</i> , Busck, Proc. Ent. Soc. Wash. Vol. 11, p. 91 (1909). | Arizona. |

266. GENUS EPICORTHYLIS, ZELLER

Epicorthylis, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 248 (1873). — Type : *E. inversella*, Zeller.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ minutely ciliated, basal joint moderate, without pecten. Labial palpi very long, curved, ascending, second joint long, thickened with dense appressed scales, above with long rough projecting hairscales diminishing to apex, terminal joint half second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 long-stalked, 7 to apex, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen faintly sinuate, cilia 2/3-3 4; with cubital pecten; 3 and 4 connate, 5 rather approximated, 6 and 7 closely approximated towards base.

Remarks. — Correlated with *Cymotricha* and *Trichotaphe*.

Geographical distribution of species. — North American.

Larva unknown.

1. *E. inversella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 248, pl. 3, Texas.
f. 13 (1873).

267. GENUS PARANOEA, WALSHAM

Paranoea, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 78 (1911). — Type : *P. latescens*, Walsingham.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ shortly ciliated, basal joint elongate, without pecten. Labial palpi long, recurved, second joint with appressed scales, somewhat expanded at apex above, apex truncate, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 connate or stalked, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia over 1; 3 and 4 connate or short-stalked, 5 rather approximated, 6 and 7 closely approximated at base.

Remarks. — Not known to me.

Geographical distribution of species. — Tropical American.

Larva (*fulvidella*) in a web on leaves.

Foodplant *Bromelia*.

1. *P. fulvidella*, Walsingham, Proc. Zool. Soc. Lond. p. 62 (1897). Virgin Islands.
2. *P. latescens*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 79, pl. 2, Mexico.
f. 28 (1911).

268. GENUS COTYLOSCIA, MEYRICK

Cotyloscia, Meyrick, Exot. Microlep. Vol. 3, p. 3 (1923). — Type : *C. caustonota*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, fasciculate-ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense appressed scales, above with rough projecting hairs except at base, terminal joint shorter than second, rather thickened with scales more or less roughly projecting posteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from beyond middle. Hindwings 1 or over 1, trapezoidal, termen slightly sinuate, cilia 2/3-3/4; without cubital pecten; 3 and 4 connate or short-stalked, 5 nearly parallel, 6 and 7 closely approximated towards base.

Remarks. — A development of *Tricholaphe*.

Geographical distribution of species. — Central and South American.

Larva unknown.

- | | |
|--|-----------------|
| 1. <i>C. terracotta</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 87, Panama.
pl. 3, f. 8 (1911). | |
| 2. <i>C. caustonota</i> , Meyrick, Trans. Ent. Soc. Lond. p. 280 (1914). | Guiana, Brazil. |
| 3. <i>C. triplagella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 618 (1864). | Brazil. |

269. GENUS EUNEBRISTIS, MEYRICK

Eunebristis, Meyrick, Exot. Microlep. Vol. 3, p. 3 (1923). — Type : *E. zachroa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue deve'oped. Antennae 5/6, in ♂ serrulate, simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, above with long erect hairs, terminal joint shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 11 from middle. Hindwings 1 or over 1, rounded-trapezoidal, cilia 2/5-3/5; without cubital pecten; 3 and 4 connate, 5 approximated, 6 and 7 closely approximated towards base.

Remarks. — A derivative of *Tricholaphe*.

Geographical distribution of species. — Tropical American.

Larva (*zingarella*) mining in fleshy leaves.

Foodplant *Coccoloba* (*Polygonaceae*).

- | | |
|--|-----------------|
| 1. <i>E. oncotera</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 85, Mexico.
pl. 3, f. 7 (1911). | |
| 2. <i>E. cinctilias</i> , Meyrick, Exot. Microlep. Vol. 2, p. 150 (1918). | Guiana. |
| 3. <i>E. zingarella</i> , Walsingham, Proc. Zool. Soc. Lond. p. 84 (1897). | Virgin Islands. |
| 4. <i>E. zachroa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 278 (1914). — Pl. 3, | Guiana. |
| Fig. 73. | |
| 5. <i>E. gyralea</i> , Meyrick, ibidem, p. 106 (1922). | Brazil. |

270. GENUS ZOMEUTIS, MEYRICK

Zomeutis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 182 (1913). — Type : *Z. dicausta*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint clothed with dense appressed scales expanded above towards apex, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 2 and 3 stalked, 7 to apex, 8 absent, 11 from middle. Hindwings over 1, oblong-ovate, cilia 2/5; without cubital pecten; 3 and 4 connate, 5 parallel, 6 and 7 approximated towards base.

Remarks. — Also derived from *Trichotaphe*.

Geographical distribution of species. — Indian and Chinese.

Larva unknown.

- | | |
|--|--------|
| 1. <i>Z. praealbescens</i> , Meyrick, Exot. Microlep. Vol. 2, p. 505 (1922). | China. |
| 2. <i>Z. dicausta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 182 (1913). | Assam. |

271. GENUS MUSURGA, MEYRICK

Musurga, Meyrick, Exot. Microlep. Vol. 3, p. 3 (1923). — Type : *M. sandycitis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ evenly ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense appressed scales, triangularly roughly expanded above towards apex, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae somewhat rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen slightly sinuate, cilia 1/2-3 5; without cubital pecten; 3 and 4 connate, 5 parallel or slightly approximated, 6 and 7 closely approximated at base.

Remarks. — A modification of *Trichotaphe*.

Geographical distribution of species. — Indian and African.

Larva unknown.

- | | |
|--|-----------|
| 1. <i>M. turgida</i> , Meyrick, Ann. Transv. Mus. Vol. 6, p. 24 (1918). | Zululand. |
| 2. <i>M. cinnabarina</i> , Meyrick, Exot. Microlep. Vol. 3, p. 3 (1923) | Ceylon. |
| 3. <i>M. polyaema</i> , Meyrick, ibidem, Vol. 3, p. 4 (1923). | Ceylon. |
| 4. <i>M. sandycitis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 150 (1907). — Pl. 4, Fig. 76. | Assam. |

272. GENUS TAPHROSARIS, MEYRICK

Taphrosaris, Meyrick, Trans. Ent. Soc. Lond. p. 104 (1922). — Type : *T. malthacopa*, Meyrick.

Characters. — Head with broad appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ strongly ciliated, basal joint elongate, subclavate, without pecten. Labial palpi

long, recurved, basal joint enlarged and much thickened with dense scales, second joint very long, broad, smooth, on inner side hollowed throughout (σ) into a deep trough filled with a long dense expansible hair-pencil, terminal joint extremely short, filiform. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal-ovate, termen hardly sinuate, cilia 1/5; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — A specialised derivative of *Trichotaphe*.

Geographical distribution of species. — South American.

Larva unknown.

1. *T. malthacopa*, Meyrick, Trans. Ent. Soc. Lond. p. 104 (1922). — Pl. 4, Guiana, Brazil.
Fig. 79.

273. GENUS TRICHOTAPHE, CLEMENS

Trichotaphe, Clemens, Proc. Acad. Nat. Sc. Philad. p. 166 (1860). — Type : *T. setosella*, Clemens.

Begoe, Chambers, Canad. Ent. Vol. 4, p. 209 (1872). — Type : *T. setosella*, Clemens.

Malacotricha, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 282 (1873). — Type : *T. setosella*, Clemens.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5-5/6, in σ ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales beneath, above with scales more or less roughly expanded towards apex, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen slightly sinuate, cilia 2/5-3/4; without cubital pecten; 3 and 4 connate, 5 somewhat approximated, 6 and 7 closely approximated or stalked.

Remarks. — Derived from a form approaching *Brachmia*.

Geographical distribution of species. — Chiefly American, but also fairly represented in the Indo-Malayan region, and more scantily in Africa; probably however of Indian origin.

Larva (10 known) feeding in spun leaves.

Foodplants especially *Compositae* (8), but also *Boraginaceae*, *Rhamnaceae*.

- | | |
|--|-------------------------|
| 1. <i>T. semicuprata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 107 (1922). | Peru. |
| 2. <i>T. euparypha</i> , Meyrick, ibidem, p. 108 (1922). | Peru. |
| 3. <i>T. aurisulcata</i> , Meyrick, ibidem, p. 108 (1922). | Brazil. |
| 4. <i>T. argogastra</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 99, pl. 3, f. 16 (1911) (argi-). | Mexico. |
| 5. <i>T. porphyrogramma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 278 (1914). | Guiana, Brazil, Peru. |
| 6. <i>T. excavata</i> , Busck, Proc. U. S. Mus. Vol. 47, p. 18 (1914). | Panama. |
| 7. <i>T. habrochitona</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 102, pl. 3, f. 26 (1911). | Panama, Guiana, Brazil. |
| 8. <i>T. cyanoneura</i> , Meyrick, Trans. Ent. Soc. Lond. p. 109 (1922). | Guiana, Brazil. |
| 9. <i>T. evitata</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 99, pl. 3, f. 15 (1911). | Panama. |
| 10. <i>T. vetustella</i> , Walker, List Lep. Brit. Mus. Vol. 29, p. 830 (1864). | Brazil. |
| 11. <i>T. mistipalpis</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 102, pl. 3, f. 24 (1911). | Panama, Guiana, Brazil. |
| <i>violaria</i> , Meyrick, Trans. Ent. Soc. Lond. p. 279 (1914). | |

12. *T. xerodes*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 100 (1911). Mexico.
 13. *T. fulvicilia*, Meyrick, Trans. Ent. Soc. Lond. p. 109 (1922). Brazil.
 14. *T. xuthostola*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 101, pl. 3, f. 18 (1911). Mexico.
 15. *T. griseella* Chambers, Canad. Ent. Vol. 6, p. 245 (1874). Texas.
 16. *T. opsonoma*, Meyrick, Trans. Ent. Soc. Lond. p. 281 (1914). Guiana.
 17. *T. bidiscomaculella*, Chambers, Canad. Ent. Vol. 6, p. 241 (1874). Texas.
 18. *T. trimaculella*, Chambers, ibidem, Vol. 6, p. 238 (1874). Texas.
 19. *T. fernaldella*, Busck, Proc. U. S. Mus. Vol. 25, p. 915 (1903). Maine.
 20. *T. serrativittella*, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 280, pl. 4, f. 27 (1873). E. United States.
 platella, Chambers, Canad. Ent. Vol. 6, p. 238 (1874).
 21. *T. barnesiella*, Busck, Proc. Ent. Soc. Wash. Vol. 8, p. 92 (1907). Arizona.
 22. *T. simpliciella*, Busck, Proc. U. S. Mus. Vol. 27, p. 761 (1904). Distr. Columbia.
 23. *T. inserrata*, Walsingham, Trans. Amer. Ent. Soc. Vol. 10, p. 184 (1882). E. United States.
 24. *T. leucocosma*, Meyrick, Exot. Microlep. Vol. 1, p. 584 (1916). Nyassaland.
 25. *T. flavocostella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 162 (1860). — E. United States, Canada.
- PI. 4, FIG. 78; PI. 5, FIG. 124.**
26. *T. manella*, Möschler, Ab. Senck. Ges. Vol. 15, p. 344 (1890). Portorico.
 27. *T. eupatoriella*, Chambers, Canad. Ent. Vol. 4, p. 221 (1872). E. United States.
 dolabella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 288, pl. 4, f. 30 (1873).
 28. *T. setosella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 166 (1860). E. United States, Canada.
 costolutella, Chambers, Canad. Ent. Vol. 4, p. 209 (1872).
 bilobella, Zeller, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 280, pl. 4, f. 28 (1873).
 29. *T. renascens*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 2, p. 96, pl. 3, f. 14 (1911). Mexico.
 30. *T. retracta*, Meyrick, Trans. Ent. Soc. Lond. p. 109 (1922). Brazil.
 31. *T. thrypsandra*, Meyrick, Exot. Microlep. Vol. 3, p. 5 (1923). Ecuador.
 32. *T. nessica*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 95 (1911). Panama.
 33. *T. varronia*, Busck, Insec. Inscit. Menstr. Vol. 1, p. 89 (1913). Panama, Guiana.
 34. *T. arotrosema*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 95 (1911). Mexico.
 35. *T. costirufella*, Chambers, Canad. Ent. Vol. 6, p. 240 (1874) (*costarufoella*). Texas.
 36. *T. ardesiella*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 96 (1911). Mexico.
 37. *T. washingtoniella*, Busck, Canad. Ent. Vol. 38, p. 121 (1906). Canada, Pennsylvania.
 38. *T. condaliivorella*, Busck, Proc. U. S. Mus. Vol. 23, p. 232 (1900) (-*iav-*). Florida.
 39. *T. melantherella*, Busck, ibidem, Vol. 23, p. 232 (1900). Florida.
 40. *T. trinotella*, Busck, Canad. Ent. Vol. 38, p. 122 (1906). Pennsylvania.
 41. *T. leuconotella*, Busck, Proc. U. S. Mus. Vol. 27, p. 762 (1904). Distr. Columbia.
 42. *T. juncidella*, Clemens, Proc. Acad. Nat. Sc. Philad. p. 166 (1860). E. United States, Canada.
 pallipalpis, Walker, List Lep. Brit. Mus. Vol. 29, p. 596 (1864).
 dubitella, Chambers, Canad. Ent. Vol. 4, p. 92 (1872).
 43. *T. levitella*, Fyles, ibidem, Vol. 36, p. 211 (1904). Canada.
 44. *T. carinella*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 99, pl. 3, f. 20 (1911). Mexico.
 45. *T. caerulescens*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 180 (1913). Assam.
 46. *T. plumosa*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 302 (1913). Transvaal, Comoro Isl*.
 47. *T. seminata*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 275 (1911). Aldabra.
 48. *T. rubidula*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 303 (1913). Transvaal.
 49. *T. asteropis*, Meyrick, ibidem, Vol. 8, p. 83 (1921). Rhodesia.
 50. *T. stromatias*, Meyrick, ibidem, Vol. 6, p. 23 (1918). Zululand.
 51. *T. cocta*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 179 (1913). Assam.
 52. *T. amphichlora*, Meyrick, Exot. Microlep. Vol. 3, p. 4 (1923). Assam.
 53. *T. cellaria*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 180 (1913). Assam.

54. <i>T. contentella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 638 (1864).	Borneo.
55. <i>T. illucescens</i> , Meyrick, Exot. Microlep. Vol. 2, p. 151 (1918).	Assam.
56. <i>T. siranta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 179 (1913).	Assam.
57. <i>T. immerita</i> , Meyrick, ibidem, Vol. 22, p. 178 (1913).	Ceylon.
58. <i>T. fungifera</i> , Meyrick, ibidem, Vol. 22, p. 177 (1913).	Assam.
59. <i>T. procrossa</i> , Meyrick, ibidem, Vol. 22, p. 177 (1913).	S. India.
60. <i>T. lissota</i> , Meyrick, ibidem, Vol. 22, p. 177 (1913).	Assam.
61. <i>T. deceptella</i> , Snellen, Tijdschr. v. Ent. Vol. 46, p. 40, pl. 4, f. 9 (1903).	Java.
62. <i>T. corniculata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 177 (1913).	Assam.
63. <i>T. crambaleas</i> , Meyrick, ibidem, Vol. 22, p. 178 (1913).	Assam.

Group 8 (Lecithocera type)

An extensive group with much generic development, of which the most noticeable prevailing character is the length of the antennae, which are commonly as long as or longer than the forewings, but this is not constant, and in some genera (especially but not exclusively the more primitive), which cannot be advantageously separated, they are no longer than normal; there is also a not infrequent tendency to long antennal ciliations in the ♂, or sometimes to thickening of the stalk, in those forms in which the antennae are not especially long. The palpi are usually long and simple in structure, rarely tufted, but sometimes curiously modified in the ♂. In the forewings the termination of vein 7 may be costal, apical, or terminal, the last-named being the original type; 2 and 3 are usually stalked, but sometimes even very widely remote in the more advanced forms. The hindwings generally show little or no terminal sinuation; veins 3 and 4 are usually connate, stalked, or coincident, 6 and 7 connate or more generally stalked; the cubital pecten is always absent. The group is especially characteristic of the Indo-Malayan region, where it forms a very prominent feature; a certain number of forms derived from this fauna are found in Africa, Australia and the Palaearctic region; from America the group is almost wholly absent, and it is also not found in New Zealand. The larvae sometimes feed on leaves, but also often on dead or dry vegetable substances, which was the primitive habit of the family.

274. GENUS OECIA, WALSINGHAM

Oecia, Walsingham, Proc. Zool. Soc. Lond. p. 111 (1897). — Type : *O. oecophila*, Staudinger.

Macroceras, Staudinger, Stett. Ent. Zeit. Vol. 37, p. 150 (1876) (praeocc. -a, -os, -us). — Type : *O. oecophila*, Staudinger.

Characters. — Head with appressed scales; ocelli posterior; tongue absent. Antennae over 1, in ♂ stout, simple, basal joint moderate, without pecten. Labial palpi long, rather curved, diverging, second joint thickened with dense scales, terminal joint nearly as long as second, moderate, pointed. Maxillary palpi very short. Posterior tibiae and basal joint of tarsi with long loose hairs. Forewings with 1b furcate, 2 from 4/5, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings hardly 1, ovate-lanceolate, cilia 2; 3 and 4 remote, 5 somewhat approximated, 6 and 7 long-stalked.

Remarks. — A peculiarly modified form, which has puzzled authors, but can only be regarded as a member of this group, with some affinity to *Symmoca*.

Geographical distribution of species. — Generally widespread in hot countries, being artificially conveyed by man; origin apparently uncertain.

Larva unrecorded, but must feed on domestic refuse, the insect being found in houses on the walls.

1. *O. oecophila*, Staudinger, Stett. Ent. Zeit. Vol. 37, p. 150 (1876) — PI. 5, S. Europe, S. W. Asia, India, Java, Queensl., Hawaiian Islands, S. America, Antilles, S. Africa.
Fig 109.
maculata, Walsingham, Proc. Zool. Soc. Lond. p. 111 (1897).

275. GENUS AMBLOMA, WALSHAM

Ambloma, Walsingham, Proc. Zool. Soc. Lond. p. 946 (1907). — Type : *A. brachyptera*, Walsingham.

Characters. — Head smooth. Antennae over 1, in ♂ simple, basal joint without pecten. Labial palpi moderately long, curved, ascending, second joint with scales slightly projecting at apex beneath, terminal joint short, acute. Maxillary palpi short. Posterior tibiae hairy. Forewings with cell short, 2 separate, 6 and 7 out of 8, 7 to costa. Hindwings 2/3, ovate-lanceolate, cilia 1 1/2; 3 and 4 stalked, 6 and 7 stalked.

Remarks. — Apparently a derivative of *Symmoca*, with some resemblances to the preceding.

Geographical distribution of species. — Canary Islands.

Larva unknown.

1. *A. brachyptera*, Walsingham, Proc. Zool. Soc. Lond. p. 947, pl. 51, f. 18 (1907). Canary Islands.

276. GENUS APOTISTATUS, WALSHAM

Apotistatus, Walsingham, Ent. M. Mag. Vol. 40, p. 271 (1904). — Type : *A. leucostictus*, Walsingham.

Characters. — Head with appressed scales; tongue absent. Antennae 2/3, in ♂ simple, basal joint without pecten. Labial palpi rather short, subascending, second joint rather densely scaled beneath, terminal joint shorter, smooth, obtuse. Maxillary palpi obsolete. Posterior tibiae hairy. Forewings with 1b furcate, 2 separate, 6 and 7 out of 8, 7 to costa, 11 from middle. Hindwings 1, elongate-ovate, cilia 1; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Not known to me, but doubtless referable here.

Geographical distribution of species. — Mediterranean.

Larva mining in leaves.

Foodplant *Limoniastrum* (*Plumbaginaceae*).

1. *A. leucostictus*, Walsingham, Ent. M. Mag. Vol. 40, p. 271 (1904). Algeria.

277. GENUS EREMICA, WALSHAM

Eremica, Walsingham, Ent. M. Mag. Vol. 40, p. 270 (1904). — Type : *E. saharae*, Walsingham.

Characters. — Head smooth; tongue developed. Antennae 4/5, in ♂ rather stout, simple, serrulate. Labial palpi long, recurved, second joint with appressed scales, terminal joint shorter than

second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 connate or stalked or coincident, 4 approximated or connate with 2, 7 and 8 stalked, 7 to apex. Hindwings 1, elongate-ovate, cilia 2; 3 and 4 stalked, 5 slightly approximated, 6 and 7 stalked.

Remarks. — Related to the preceding.

Geographical distribution of species. — Mediterranean.

Larva unknown.

- | | |
|---|----------|
| 1. <i>E. lithochroma</i> , Walsingham, Ent. M. Mag. Vol. 40, p. 271 (1904). | Algeria. |
| 2. <i>E. saharae</i> , Walsingham, ibidem, Vol. 40, p. 270 (1904). | Algeria. |

278. GENUS HOLCOPOGON, STAUDINGER

Holcopogon, Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 330 (1880). — Type : *H. bubulcella*, Staudinger.

Cynia, Walsingham, Ent. M. Mag. Vol. 36, p. 218 (1900). — Type : *H. bubulcella*, Staudinger.

Epistomotis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 416 (1906). — Type : *H. robusta*, Butler.

Characters. — Head with appressed scales, sidetufts raised, connivent; ocelli small, posterior; tongue short. Antennae 2/3, in ♂ variably ciliated, basal joint moderately elongate, without pecten. Labial palpi long, ascending, second joint with long projecting apical tuft beneath, terminal joint as long as second, erect, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 short, from near angle, 7 and 8 stalked, 7 to apex, or 7 absent, 11 from middle. Hindwings 1, elongate-ovate, cilia 1-1 1/2; 3 and 4 connate or short-stalked, 5 nearly parallel, 6 and 7 long-stalked or seldom coincident (6 absent).

Remarks. — This genus has been found perplexing, but is really a derivative of *Symmoca*, with the exceptional feature of a strong palpal tuft.

Geographical distribution of species. — Properly Mediterranean, but extending into India and South Africa.

Larva unknown.

- | | |
|--|------------------------|
| 1. <i>H. scaeocentra</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 93 (1921). — PI. 5, | Tanganyika Territ. |
| Fig. 110. | |
| 2. <i>H. rhyodes</i> , Meyrick, ibidem, Vol. 2, p. 16, pl. 5, f. 8 (1909). | Transvaal. |
| <i>amphicentra</i> , Meyrick, ibidem, Vol. 3, p. 326 (1913). | |
| 3. <i>H. psameticella</i> , Rebel, Iris, Vol. 28, p. 268 (1914). | Egypt. |
| 4. <i>H. robusta</i> , Butler, Proc. Zool. Soc. Lond. p. 174 (1883). | Bombay, Ceylon. |
| <i>penessa</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 416 (1906). | |
| 5. <i>H. geminella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 330 (1915). | Algeria. |
| 6. <i>H. helveolella</i> , Staudinger, Hor. Soc. Ent. Ross. Vol. 15, p. 330 (1880). | S. E. Europe, W. Asia. |
| 7. <i>H. bubulcella</i> , Staudinger, Stett. Ent. Zeit. Vol. 20, p. 245 (1859). | S. Europe, Palestine. |
| <i>pulverellus</i> , Constant, Ann. Soc. Ent. Fr. p. 191, pl. 7, f. 5 (1865) | |
| <i>barbata</i> , Walsingham, Ent. M. Mag. Vol. 36, p. 219 (1900). | |

279. GENUS STRENOPHILA, MEYRICK

Strenophila, Meyrick, Ann. Transv. Mus. Vol. 3, p. 306 (1913). — Type : *S. hyptiota*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ ciliated, basal joint elongate, with slight pecten. Labial palpi very long, recurved, with appressed scales, terminal

joint longer than second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 2/3, lanceolate, cilia 3; 3 and 4 short-stalked, 5 parallel, 6 and 7 long-stalked.

Remarks. — Derived from *Oegoconia*.

Geographical distribution of species. — African.

Larva unknown.

1. *S. hyptiota*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 306 (1913). — PI. 5, Transvaal.
Fig. 108.

280. GENUS OEGOCONIA, STANTON

Oegoconia, Stainton, Ins. Brit. Tin. p. 163 (1854). — Type : *O. quadripuncta*, Haworth.

Apatema, Walsingham, Ent. M. Mag. Vol. 36, p. 219 (1900). — Type : *O. fasciata*, Stainton.

Clerogenes, Meyrick, Ann. Transv. Mus. Vol. 8, p. 93 (1921). — Type : *O. meledantis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 2/3, in ♂ simple or ciliated, basal joint moderately elongate, with slight very fugitive pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint as long as second or shorter, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired. Forewings with 1 b furcate, 2 from towards angle, 3 and 4 sometimes connate, 7 and 8 stalked, 7 to costa, 6 sometimes out of 8 near base, 11 from middle. Hindwings 1, elongate-ovate or ovate-lanceolate, cilia over 1; 3 and 4 connate or stalked, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — A development of *Symmoeca*.

Geographical distribution of species. — Perhaps originally Mediterranean, but with representatives in S. Africa, Ceylon and Australia, and typical *quadripuncta* has been artificially introduced into North America and New Zealand.

Larva (*quadripuncta*) feeding on dry vegetable refuse.

- | | |
|---|--|
| 1. <i>O. isthmodes</i> , Meyrick, Exot. Microlep. Vol. 2, p. 506 (1922). | Queensland. |
| 2. <i>O. praeramis</i> , Meyrick, ibidem, Vol. 2, p. 155 (1918). | Ceylon. |
| 3. <i>O. cyrota</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 94 (1921). | Cape Colony. |
| 4. <i>O. quadripuncta</i> , Haworth, Lep. Brit. p. 557 (1828). | C. & S. Europe, N. Africa,
N. America, New
Zealand. |
| <i>bifasciella</i> , Stephens, Ill. Brit. Ent. Haust. Vol. 4, p. 217 (1835). | |
| <i>kindermannella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, f. 418 (1855). | |
| <i>deauratella</i> , Herrich-Schäffer, ibidem, Vol. 5, p. 135 (1855). | |
| <i>novimundi</i> , Busck, Proc. Ent. Soc. Wash. Vol. 17, p. 84 (1915). | |
| 5. <i>O. fasciata</i> , Stainton, Ann. Mag. Nat. Hist. (3), Vol. 3, p. 213 (1859). | S. Europe, N. Africa,
W. C. Asia, Madeira,
Canaries. |
| <i>coarctella</i> , Rebel, Ann. Hofmus. Wien, Vol. 11, p. 129, pl. 3, f. 11 (1896). | |
| <i>mediopalidum</i> , Walsingham, Ent. M. Mag. Vol. 36, p. 220 (1900). | |
| 6. <i>O. lucida</i> , Walsingham, Proc. Zool. Soc. Lond. p. 945, pl. 52, f. 3 (1907). | Canaries. |
| 7. <i>O. meledantis</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 93 (1921). | Cape Colony. |

281. GENUS SYMMOCA, HÜBNER

Symmoeca, Hübner, Verz. bek. Schmett. p. 403 (1826). — Type : *S. signella*, Hübner.

Paradoria, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 740 (1907). — Type : *S. anaphracta*,
Meyrick.

Characters. — Head with loosely appressed scales, sidetufts rather raised; ocelli posterior; tongue developed. Antennae 2/3-3/4, in ♂ simple or minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi moderate or long, recurved, second joint thickened with dense scales, sometimes somewhat projecting at apex beneath, terminal joint as long as second or rather shorter, moderate or slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings sometimes with tufts of scales; 1b furcate, 2 from towards angle, 3 from angle, seldom 3 and 4 stalked, 7 and 8 stalked, 7 to apex or termen, 11 from middle. Hindwings 1, elongate-ovate or ovate lanceolate, cilia 1-1 1/4; 3 and 4 connate or stalked, 5 parallel, 6 and 7 stalked.

Remarks. — This genus, with its accessories, appears to form a subgroup or separate line of descent parallel with the rest of the *Lecithocera* group and derived with it from primitive forms of the *Autosticha* group, thus tending to exhibit some of the early Oecophorid-like features of its ancestors.

Geographical distribution of species. — Essentially Mediterranean, with a homogeneous colony in India.

Larva (4 known) feeding on dead bark and dry vegetable refuse.

1. <i>S. crocodesma</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 70 (1912).	Transvaal.
2. <i>S. amphicalyx</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 735 (1911). — Pl. 5, Fig. III.	S. India.
3. <i>S. dolabrata</i> , Meyrick, Exot. Microlep. Vol. 1, p. 589 (1916).	S. India.
4. <i>S. palacta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 736 (1911).	Bombay, Coorg.
5. <i>S. rhodota</i> , Meyrick, ibidem, Vol. 20, p. 735 (1911). — Pl. 5, Fig. II 2.	S. India.
6. <i>S. stesichora</i> , Meyrick, ibidem, Vol. 20, p. 735 (1911).	Bombay, Coorg, S. India.
7. <i>S. anaphracta</i> , Meyrick, ibidem, Vol. 17, p. 740 (1907).	India.
8. <i>S. alacris</i> , Meyrick, Exot. Microlep. Vol. 2, p. 156 (1918).	Kanara.
9. <i>S. indagata</i> , Meyrick, ibidem, Vol. 2, p. 155 (1918).	Kanara.
10. <i>S. acatharta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 736 (1911).	Kanara, Coorg.
11. <i>S. vitiosella</i> , Zeller, Stett. Ent. Zeit. Vol. 29, p. 139 (1868).	Greece, Asia Minor.
12. <i>S. minimella</i> , Caradja, Iris, Vol. 34, p. 120 (1920).	W. Siberia.
13. <i>S. mendosella</i> , Zeller, Stett. Ent. Zeit. Vol. 29, p. 137 (1868).	Tyrol, Carinthia.
14. <i>S. pyrrhella</i> , Ragonot, Bull. Soc. Ent. Fr. p. 107 (1895).	Asia Minor.
15. <i>S. muricella</i> , Chrétien, Naturaliste, p. 104 (1896).	Pyrenees.
16. <i>S. zeitunella</i> , Rebel, Iris, Vol. 15, p. 111 (1902).	Asia Minor.
17. <i>S. monochromella</i> , Rebel, ibidem, Vol. 15, p. 110, pl. 4, f. 5 (1902).	Asia Minor.
18. <i>S. uniformella</i> , Rebel, ibidem, Vol. 13, p. 166 (1900).	Spain.
19. <i>S. tristella</i> , Caradja, ibidem, Vol. 34, p. 119 (1920).	Spain.
20. <i>S. contrastella</i> , Caradja, ibidem, Vol. 34, p. 119 (1920).	Asia Minor.
21. <i>S. turana</i> , Caradja, ibidem, Vol. 34, p. 119 (1920).	Ferghana.
22. <i>S. oenophila</i> , Staudinger, Cat. Eur. Lep. p. 426 (1871).	S. France, Spain.
23. <i>S. pleostigmella</i> , Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 67, p. (51) (1917).	Spain.
24. <i>S. latiusculella</i> , Stanton, Tin. Syr. p. 55 (1867).	Asia Minor.
25. <i>S. caliginella</i> , Mann, Verh. Zool.-bot. Ges. Wien, Vol. 17, p. 842 (1867).	Tyrol.
26. <i>S. achrestella</i> , Rebel, ibidem, Vol. 39, p. 320, pl. 8, f. 4 (1889).	Tyrol, Carinthia.
27. <i>S. hispanella</i> , Rebel, ibidem, Vol. 67, p. (49) (1917).	Spain.
28. <i>S. syriacella</i> , Ragonot, Bull. Soc. Ent. Fr. p. 108 (1895).	S. E. Europe, Syria.
<i>pannosella</i> , Rebel, Berl. Ent. Zeitschr. Vol. 50, p. 309 (1905).	
29. <i>S. calidella</i> , Walsingham, Ent. M. Mag. Vol. 41, p. 37 (1905).	Algeria.
30. <i>S. atricanella</i> , Rebel, Berl. Ent. Zeitschr. Vol. 50, p. 308 (1905).	Greece.
31. <i>S. sparsella</i> , Joannis, Bull. Soc. Ent. Fr. p. 84 (1891).	Syria.
32. <i>S. tectaphella</i> , Rebel, Ann. Hofmus. Wien, Vol. 30, p. 161, pl. 4, f. 12 (1916).	Crete.
33. <i>S. designatella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 111, f. 384 (1855).	Dalmatia, Greece, Asia Minor.
<i>bifasciata</i> , Staudinger, Hor. Soc. Ent. Ross. Vol. 7, p. 248 (1870).	
34. <i>S. nigromaculella</i> , Ragonot, Bull. Soc. Ent. Fr. p. 194 (1875).	Portugal.

35. *S. Helleri*, Rebel, Ann. Hofmus. Wien, Vol. 24, p. 356, pl. 12, f. 5 (1910). Canaries.
36. *S. petrogenes*, Walsingham, Ent. M. Mag. Vol. 43, p. 213 (1907). Spain.
37. *S. longipalpella*, Rebel, Iris, Vol. 28, p. 269 (1914). Egypt.
38. *S. signatella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 111, f. 380 (1855). S. & W. C. Europe,
- scriptella*, Duponchel, Hist. Nat. Lép. Fr. Vol. 11, pl. 298, f. 3 (1838) Algeria.
- (nec Hübner).
39. *S. rosmarinella*, Walsingham, Ent. M. Mag. Vol. 37, p. 177 (1901). S. France.
40. *S. dodecatella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 239 (1859). Spain.
41. *S. Husadeli*, Rebel, Ann. Hofmus. Wien, Vol. 24, p. 353, pl. 12, f. 7 (1910). Canaries.
42. *S. oxybiella*, Millière, Pet. Nouv. Ent. Vol. 4, p. 172 (1872). S. France, Tyrol, Algeria.
43. *S. delicatella*, Walsingham, Ent. M. Mag. Vol. 37, p. 178 (1901). Corsica.
44. *S. albicanella*, Zeller, Stett. Ent. Zeit. Vol. 29, p. 136 (1868). S. E. Europe.
- signella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 111, f. 338 (1855).
45. *S. ratella*, Herrich-Schäffer, ibidem, Vol. 5, p. 211, f. 427, 428 (1855). S. Europe, Asia Minor,
- sericeella*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 67, p. 52 (1917). Algeria.
46. *S. signella*, Hübner, Samml. Eur. Schmett. Tin. f. 211 (1796). Alps.
47. *S. designella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 111, f. 461 (1855). Hungary.
48. *S. ponerias*, Walsingham, Ent. M. Mag. Vol. 41, p. 37 (1905). Algeria.
49. *S. alhambrella*, Walsingham, ibidem, Vol. 47, p. 212 (1911). Spain.
50. *S. tofosella*, Rebel, Stett. Ent. Zeit. Vol. 54, p. 53 (1893). Spain.
51. *S. cedestiella*, Zeller, ibidem, Vol. 29, p. 140 (1868). S. Russia, Asia Minor.
52. *S. perpygmaeella*, Walsingham, Ent. M. Mag. Vol. 37, p. 179 (1901). Corsica.
53. *S. undecimpunctella*, Mann, Wien. Ent. Monatsschr. Vol. 8, p. 185, pl. 4, f. 17 (1864). S.E. Europe, Asia Minor.
- ? *dissoluta*, Staudinger, Hor. Soc. Ent. Ross. Vol. 7, p. 250 (1870).
54. *S. virginella*, Rebel, Iris, Vol. 15, p. 112, pl. 4, f. 6 (1902). Asia Minor.
55. *S. aegrella*, Walsingham, Proc. Zool. Soc. Lond. p. 949, pl. 52, f. 2 (1907). Canaries.
56. *S. canariensis*, Rebel, Ann. Hofmus. Wien, Vol. 21, p. 38 (1906). Canaries.
57. *S. griseosericiella*, Ragonot, Bull. Soc. Ent. Fr. p. 140 (1879). Portugal.
58. *S. albidella*, Rebel, Iris, Vol. 13, p. 166 (1900). Algeria.
59. *S. sericiella*, Walsingham, Ent. M. Mag. Vol. 40, p. 273 (1904). Algeria.
60. *S. pallida*, Staudinger, Stett. Ent. Zeit. Vol. 37, p. 149 (1876). Sicily, Dalmatia.
61. *S. exigua*, Chrétien, Ann. Soc. Ent. Fr. p. 335 (1915). Algeria.
62. *S. oblitterata*, Walsingham, Ent. M. Mag. Vol. 41, p. 38 (1905). Algeria.
63. *S. molitor*, Walsingham, ibidem, Vol. 41, p. 38 (1905). Algeria.
64. *S. quadrifariella*, Mann, Verh. Zool.-bot. Ges. Wien, Vol. 5, p. 563 (1855). Corsica.
65. *S. saharae*, Oberthür, Et. Ent. Vol. 12, p. 44, pl. 6, f. 34k (1888). Algeria.
66. *S. ochreopicta*, Walsingham, Ent. M. Mag. Vol. 37, p. 178 (1901). Corsica.
67. *S. cryptogamarum*, Millière, Pet. Nouv. Ent. Vol. 4, p. 172 (1872). S. France.
68. *S. orphnella*, Rebel, Stett. Ent. Zeit. Vol. 54, p. 54 (1893). Italy.
69. *S. muscularia*, Staudinger, Hor. Soc. Ent. Ross. Vol. 7, p. 251, pl. 3, f. 9 (1870). Greece.

282. GENUS CHERSOGENES, WALSHAM

Chersogenes, Walsingham, Proc. Zool. Soc. Lond. p. 947 (1907). — Type : *C. victimella*, Walsingham.

Epanastasis, Walsingham, ibidem, p. 948 (1907). — Type : *C. sophroniella*, Rebel.

Characters. — Head with appressed scales; tongue developed. Antennae 1, in ♂ simple, basal joint without pecten. Labial palpi very long, curved, ascending, second joint with dense scales rather triangularly projecting beneath at apex, terminal joint shorter, slender, acute. Maxillary palpi short. Posterior tibiae clothed with hairs above. Forewings sometimes with tufts of scales; 2 separate, 7 and 8

stalked, 7 to termen. Hindwings 1, elongate, pointed, termen faintly sinuate, cilia 1-1 1/2; 3 and 4 stalked, 5 nearly parallel, 6 and 7 long-stalked.

Remarks. — Probably correlated with *Symmoca*.

Geographical distribution of species. — Canary Islands.

Larva unknown.

1. *C. sophroniella*, Rebel, Ann. Hofmus. Wien, Vol. 9, p. 89 (1895). Canaries.
2. *C. victimella*, Walsingham, Proc. Zool. Soc. Lond. p. 947, pl. 51, f. 17 (1907). Canaries.

283. GENUS CEUTHOMADARUS, MANN

Ceuthomadarus, Mann, Wien. Ent. Monatsschr. Vol. 8, p. 188 (1864). — Type : *C. tenebrionellus*, Mann.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli inferior; tongue absent. Antennae 4/5, in ♂ rather stout, simple, basal joint rather elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, rather rough beneath, terminal joint somewhat shorter, moderate, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings 1, elongate-ovate, cilia 3/4; 3 and 4 connate, 5 parallel, 6 and 7 long-stalked. Wings in *viduellus* ♀ abbreviated.

Remarks. — Nearly related to *Symmoca*.

Geographical distribution of species. — Mediterranean.

Larva unknown.

1. *C. viduellus*, Rebel, Ann. Hofmus. Wien, Vol. 18, p. 332, pl. 3, f. 15 (1903). Bulgaria.
2. *C. tenebrionellus*, Mann, Wien. Ent. Monatsschr. Vol. 8, p. 188, pl. 5, f. 1, 2 (1864). Asia Minor, N. Persia.

284. GENUS TOGIA, WALKER

Togia, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 791 (1864) — Type : *T. nemophorella*, Walker.

Characters. — Head smooth. Antennae over 1 (?), somewhat flattened, slightly roughened above. Labial palpi very long, recurved, slender, smooth, terminal joint as long as second, acute. Anterior coxae and middle tibiae clothed with long hairs. Hindwings trapezoidal; 3 and 4 stalked, 5 connate, 6 and 7 stalked.

Remarks. — Characters taken from original type, the only example known, but unset and too much damaged for further investigation; probably allied to *Tisis*, and recognisable superficially by its Adelid (not Nemophorid) appearance, and structurally by its leg-characters.

Geographical distribution of species. — Malayan.

Larva unknown.

1. *T. nemophorella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 792 (1864). Borneo.

285. GENUS TISIS, WALKER

Tisis, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 793 (1864). — Type : *T. bicolorella*, Walker.

Tonosa, Walker, ibidem, Vol. 29, p. 796 (1864). — Type : *T. seclusella*, Walker.

Tingentera, Walker, ibidem, Vol. 29, p. 798 (1864). — Type : *T. meliorella*, Walker.

Tipha, Walker, ibidem, Vol. 29, p. 798 (1864). — Type : *T. chalybaeella*, Walker.

Tirallis, Walker, ibidem, Vol. 29, p. 806 (1864). — Type : *T. chalybaeella*, Walker.

Cacogamia, Snellen, Tijdschr. v. Ent. Vol. 46, p. 48 (1903). — Type : *T. elegans*, Snellen.

Characters. — Head smooth-scaled; ocelli posterior; tongue developed. Antennae over 1, in ♂ with sinuation and thickening of dense scales near base, in ♀ more or less roughened with scales near base, basal joint long, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute, or in ♂ sometimes clothed with dense rough scales to apex. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs. Forewings with 1 b furcate, 2 from angle, 3 absent (seldom visible as a minute furcation at extreme apex of 2), 4 absent, 7 and 8 stalked, 7 to costa, 9 sometimes out of 7, 11 from middle. Hindwings 1 or over 1, elongate-trapezoidal, termen more or less rounded, cilia 1/2-3/4; 4 connate with or out of 3 or absent. 5 closely approximated or out of 3, 6 and 7 stalked, 6 sometimes to costa or absent.

Remarks. — Correlated with *Doxogenes* and *Timyra* as similar but distinct developments from a common origin. Elegant insects, often with ornamental orange and metallic colouring.

Geographical distribution of species. — Malayan, with centre of origin in Borneo.

Larva unknown.

- | | |
|--|----------|
| 1. <i>T. meliorella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 798 (1864). | Borneo. |
| 2. <i>T. hemixysta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 438 (1910). | Borneo. |
| 3. <i>T. mesozosta</i> , Meyrick, Suppl. Ent. p. 50 (1914). | Formosa. |
| 4. <i>T. hyacinthina</i> , Meyrick, Trans. Ent. Soc. Lond. p. 439 (1910). | Borneo. |
| 5. <i>T. argyrophaea</i> , Meyrick, ibidem, p. 439 (1910). | Borneo. |
| 6. <i>T. eurylampis</i> , Meyrick, ibidem, p. 438 (1910). | Borneo. |
| 7. <i>T. helioclina</i> , Meyrick, ibidem, p. 19 (1894). | Burma. |
| 8. <i>T. elegans</i> , Snellen, Tijdschr. v. Ent. Vol. 46, p. 49, pl. 5, f. 10 (1903). | Java. |
| 9. <i>T. chalybaeella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 799 (1864). | Borneo. |

Pl. 4, Fig. 80.

latifasciella, Walker, ibidem, Vol. 29, p. 806 (1864).

- | | |
|--|------------------------|
| 10. <i>T. charadraea</i> , Meyrick, Trans. Ent. Soc. Lond. p. 439 (1910). | Malay States. |
| 11. <i>T. luteella</i> , Snellen, Tijdschr. v. Ent. Vol. 46, p. 50, pl. 5, f. 13 (1903). | Java, Borneo, Sumatra. |
| 12. <i>T. bicolorella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 793 (1864). | Borneo. |
| 13. <i>T. imperatrix</i> , Meyrick, Trans. Ent. Soc. Lond. p. 440 (1910). | Borneo. |
| 14. <i>T. seclusella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 796 (1864). | Borneo. |

286. GENUS MNESTERIA, MEYRICK

Mnesteria, Meyrick, Trans. Ent. Soc. Lond. p. 438 (1910). — Type : *M. pharetrata*, Meyrick.

Characters. — Head smooth-scaled; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten, in ♂ with slight apical scale-projection. Labial palpi

in ♂ with second joint expanded with dense scales, oval, concave internally, terminal joint absent; in ♀ very long, recurved, second joint smooth-scaled, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae smooth, with somewhat expanded whorls at origin of spurs. Forewings with 1b short-furcate, 2 remote, 3 and 4 closely approximated from angle, 5 and 6 parallel, 7 absent, 11 from beyond middle. Hindwings 1, elongate-trapezoidal, cilia nearly 1; 3 and 4 stalked, 5 rather near and parallel, 6 and 7 approximated at base; in ♂ an expandible hair-pencil from base in submedian groove.

Remarks. — Apparently a development of *Doxogenes*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *M. sideraula*, Meyrick, Exot. Microlep. Vol. 1, p. 573 (1916). Ceylon.
2. *M. basanistis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 453 (1908). Ceylon.
3. *M. pharetrata*, Meyrick, ibidem, Vol. 16, p. 593 (1905). — **Pl. 4, Fig. 81.** Ceylon.

287. GENUS DINOCHARES, NOV. GEN.

Type : *D. conotoma*, Meyrick.

Characters. — Head smooth-scaled; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, slender, without pecten. Labial palpi very long, recurved, second joint smooth-scaled, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae smooth, with somewhat expanded whorls at origin of spurs. Forewings 1b short-furcate, 2 and 4 connate or closely approximated from angle, 3 absent, 5 and 6 parallel, 7 absent, 11 from 2/3. Hindwings 1, elongate-trapezoidal, cilia over 1; 3 and 4 connate, 5 rather near and parallel, 6 and 7 stalked; in ♂ with expandible hair-pencil from base in submedian groove.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *D. conotoma*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 453 (1908). Ceylon.

288. GENUS DOXOGENES, NOV. GEN.

Type : *D. brochias*, Meyrick.

Characters. — Head smooth-scaled; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi in ♂ with second joint expanded with dense scales, oval, concave internally, terminal joint absent; in ♀ very long, recurved, second joint smooth-scaled, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae smooth, with somewhat expanded whorls at origin of spurs, or seldom rough-scaled above (*spectralis*). Forewings with 1b short-furcate, 2 and 4 almost connate or stalked from angle, 3 absent, 5 and 6 parallel, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings about 1, elongate-trapezoidal, cilia 3/4-1; 3 and 4 connate or short-stalked, 5 rather near and parallel, 6 and 7 closely approximated or stalked; in ♂ with expandible hair-pencil from base in submedian groove.

Remarks. — Derivable from *Timyra*; handsome insects, with orange and metallic colouring as in the three preceding genera.

Geographical distribution of species. — Ceylon.

Larva unknown.

- | | |
|--|---------|
| 1. <i>D. phalaritis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 594 (1905). | Ceylon. |
| 2. <i>D. philodoxa</i> , Meyrick, ibidem, Vol. 18, p. 452 (1908). | Ceylon. |
| 3. <i>D. brochias</i> , Meyrick, ibidem, Vol. 16, p. 594 (1905). | Ceylon. |
| 4. <i>D. pyrophanes</i> , Meyrick, ibidem, Vol. 16, p. 594 (1905). | Ceylon. |
| 5. <i>D. ecliptica</i> , Meyrick, ibidem, Vol. 18, p. 451 (1908). | Ceylon. |
| 6. <i>D. thoracias</i> , Meyrick, ibidem, Vol. 18, p. 452 (1908). | Ceylon. |
| 7. <i>D. spectralis</i> , Meyrick, ibidem, Vol. 16, p. 593 (1905). | Ceylon. |

289. GENUS OXYGNOSTIS, NOV. GEN.

Type: *O. diacma*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi in ♂ with second joint expanded with dense scales, oval, concave internally, terminal joint absent; in ♀ very long, recurved, slender, second joint smooth, scales somewhat rough at apex beneath, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae and tarsi rough-scaled above. Forewings with 1 b long-furcate, 2 remote, 3 from angle, 4 and 5 stalked, 8 and 9 out of 7, 7 to costa, 11 from beyond middle. Hindwings 1, elongate-trapezoidal, cilia over 1; 3 and 4 connate, 5 obsolete, 6 and 7 stalked, 6 in ♂ to costa, in ♀ to apex; in ♂ with expansible hairpencil from base in submedian groove.

Remarks. — A development of *Teucrodoxa*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *O. diacma*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 142 (1906). Ceylon.

290. GENUS TEUCRODOXA, NOV. GEN.

Type: *T. spiculifera*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint moderate, without pecten, in ♂ with slight apical scale-projection. Labial palpi in ♂ with second joint expanded, with dense scales, oval, concave internally, terminal joint absent; in ♀ very long, recurved, second joint with appressed scales, roughly projecting at apex beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae and tarsi with rough hairscales above. Forewings with 1 b long-furcate, 2 from 3/4, 3-5 closely approximated or 3 and 4 connate, 8 and 9 out of 7, 7 to costa, 11 from near middle. Hindwings 1, elongate-trapezoidal, cilia 1; 3 and 4 connate, 5 obsolete, 6 and 7 stalked, 6 to apex or termen; in ♂ an expansible hairpencil from base in submedian groove.

Remarks. — A derivative from *Timyra*.

Geographical distribution of species. — Ceylon.

Larva unknown.

- | | |
|--|---------|
| 1. <i>T. spiculifera</i> , Meyrick, Exot. Microlep. Vol. 2, p. 152 (1918). | Ceylon. |
| 2. <i>T. monetella</i> , Felder, Reise Novara, Lep. pl. 139, f. 19 (1875). | Ceylon. |

291. GENUS ALCIPHANES, NOV. GEN.

Type : *A. molybdantha*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten, in ♂ with slight apical scale-projection. Labial palpi in ♂ with second joint expanded with dense rough projecting hair-scales beneath and also slightly above, terminal joint obsolete. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae densely scaled, with loosely expanded rough whorls at origin of spurs, basal joint of tarsi rough-scaled above. Forewings with 2 from 3/5, 3 and 4 short-stalked from angle, 5 approximated, 7 and 8 stalked, 7 to costa, 11 from beyond middle. Hindwings under 1, elongate-trapezoidal, cilia over 1; 3 and 4 connate, 5 obsolete, 6 and 7 stalked; in ♂ with expandible hair-pencil from base in submedian groove.

Remarks. — A development of *Phanoschista*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *A. molybdantha*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 454 Ceylon.
(1908).

292. GENUS PHANOSCHISTA, NOV. GEN.

Type : *P. meryntis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, with thickened scale-projection near base of stalk, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint somewhat thickened with smooth scales, terminal joint as long as second, slender, acute. Maxillary palpi obsolete. Posterior tibiae with loosely appressed hairs above. Forewings 1b furcate, 2 remote, 3-5 approximated, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings 1, elongate-trapezoidal, cilia 1; 3 and 4 connate, 5 near and parallel, 6 and 7 stalked; in ♂ with expandible fringe of long hairs in submedian groove.

Remarks. — Derived from *Timyra*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *P. meryntis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 455 (1908). S. India.

293. GENUS TECHNOGRAPHA, NOV. GEN.

Type : *T. ephestrus*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten, in ♂ with small apical scale-tooth. Labial

palpi in ♂ subascending, second joint expanded with rough scales towards apex above and beneath, terminal joint very short, slender, pointed. Maxillary palpi rudimentary. Posterior tibiae and tarsi rough-scaled above, with strong median tuft on tibiae. Forewings with 1b simple, 2 from 3/4, 3 from near angle, 4 and 5 stalked from angle, 8 and 9 out of 7, 8 to costa, 11 from beyond middle. Hindwings 1, elongate-trapezoidal, cilia 1; 3 and 4 connate, 5 obsolete, 6 and 7 stalked, 6 to termen; in ♂ an expandible hair-pencil from base in submedian groove.

Remarks. — A derivative of *Timyra*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *T. ephestris*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 454 (1908). Ceylon.

294. GENUS MAGONYMPHA, MEYRICK

Magonympha, Meyrick, Exot. Microlep. Vol. 1, p. 572 (1916). — Type : *M. chrysocosma*, Meyrick.

Characters. — Head smooth, sidetufts raised; ocelli posterior; tongue developed. Antennae 1, in ♂ simple, basal joint very long, slender, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi obsolete. Posterior tibiae with median tuft of rough projecting scales. Forewings with 1b furcate, 2 from 4/5, 3 from angle, 7 and 8 stalked, 7 to apex or hardly below, 11 from middle. Hindwings under 1, triangular-lanceolate, cilia nearly 2; 3 and 4 connate, transverse vein obsolete, 5 parallel to 4 but obsolete except towards extremity, 6 and 7 long-stalked.

Remarks. — This and the next three genera are correlatives of the *Timyra* group.

Geographical distribution of species. — Indian.

Larva unknown.

1. *M. chrysocosma*, Meyrick, Exot. Microlep. Vol. 1, p. 572 (1916). S. India.

295. GENUS MONERISTA, NOV. GEN.

Type : *M. hippastis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae and basal joint of tarsi with rough hairscales above, forming dense tufts at origin of spurs. Forewings with 1b short-furcate, 2 from towards angle, 4 absent, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1, ovate-lanceolate, cilia over 1; 4 absent, 5 tolerably parallel, 6 and 7 stalked, 6 to costa.

Remarks. — Nearly allied to the preceding.

Geographical distribution of species. — Indian; one of the two furthest emigrants of this group.

Larva unknown.

1. *M. hippastis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 451 (1908). Assam.

296. GENUS HETERALCIS, NOV. GEN.

Type : *H. tetricina*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint moderate or elongate, without pecten. Labial palpi in ♀ very long, recurved, second joint with appressed scales, sometimes roughly projecting at apex beneath, terminal joint as long as second, slender, acute; in ♂ sometimes similar, or with fine long expandible hairs above, or with second joint broadly expanded with dense projecting scales beneath, concave internally, and terminal joint much reduced or absent. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above, with dense or long rough tufts at origin of spurs. Forewings with 1b short-furcate, 2 remote, 3 from angle, 4 and 5 stalked, 7 and 8 stalked, 7 to costa, 9 sometimes out of 7, 11 from middle. Hindwings about 1, trapezoidal, termen slightly sinuate, cilia nearly 1; 3 and 4 connate, 5 partially or wholly obsolete, 6 and 7 stalked; in ♂ with submedian groove.

Remarks. — Naturally discriminated by neuration of forewings.

Geographical distribution of species. — Mainly Ceylonese, but extending into India.

Larva unknown.

- | | |
|--|-------------------|
| 1. <i>H. isochra</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 444 (1908). | Ceylon, S. India. |
| 2. <i>H. platycapna</i> , Meyrick, Exot. Microlep. Vol. 1, p. 569 (1916). | Ceylon. |
| 3. <i>H. palathodes</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 144 (1906). | Ceylon. |
| 4. <i>H. rhizophora</i> , Meyrick, Exot. Microlep. Vol. 2, p. 236 (1919). | Madras. |
| 5. <i>H. tetricina</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 143 (1906). | Ceylon. |
| 6. <i>H. holocoma</i> , Meyrick, ibidem, Vol. 18, p. 446 (1908). | Ceylon. |

297. GENUS OLBOTHREPTA, NOV. GEN.

Type : *O. hydrosema*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae rough-haired above, with median tuft of scales. Forewings with 1b short-furcate, 2 remote, 3 from angle, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, termen sinuate, cilia 1 or over 1; 2 remote, 3 and 4 connate, 5 absent, 6 and 7 stalked.

Remarks. — Distinguishable from *Timyra* by form of hindwings, as well as by ♂ neuration.

Geographical distribution of species. — Southern India.

Larva unknown.

- | | |
|---|-----------|
| 1. <i>O. hydrosema</i> , Meyrick, Exot. Microlep. Vol. 1, p. 572 (1916). | S. India. |
| 2. <i>O. corythista</i> , Meyrick, ibidem, Vol. 2, p. 98 (1918). | S. India. |
| 3. <i>O. sphaeristis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 450 (1908). | S. India. |

298. GENUS TIMYRA, WALKER

Timyra, Walker, List. Lep. Het. Brit. Mus. Vol. 29, p. 782 (1864). — Type : *T. phycidella*, Walker.

Decuaria, Walker, ibidem, Vol. 29, p. 797 (1864). — Type : *T. mendicella*, Walker.

Uipsa, Walker, ibidem, Vol. 29, p. 828 (1864). — Type : *T. phycidella*, Walker.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten, in ♀ sometimes densely tufted. Labial palpi long, recurved, second joint in ♂ sometimes clothed with long loose rough hairs or with fine expandible hairs internally, in ♀ with appressed scales, rough beneath towards apex, terminal joint in ♂ sometimes very short, thickened with scales, rough anteriorly, or obsolete, in ♀ (and sometimes ♂) nearly or quite as long as second, slender or loosely scaled, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae more or less rough-scaled, with large dense median tuft above; anterior tibiae with brush of scales beneath. Forewings with 1b furcate, 2 remote, 3 from angle, 7 and 8 stalked, 7 to apex or termen, 9 sometimes out of 7, 11 from middle. Hindwings 1 or over 1, trapezoidal, obtuse, cilia 1/2-1; in ♂ 3 absent, 2 and 4 nearly approximated or stalked, in ♀ 2 remote, 3 and 4 connate or stalked; 5 absent, 6 and 7 stalked; in ♂ with expandible hair-pencil in submedian groove.

Remarks. — Developed from a form approaching *Frisilia*. The tufted posterior legs of this and the allied genera are obviously intended for display (the tuft is often partly metallic), and in an example of *T. phorcis* the two posterior legs are projected beneath the wings and erected on each side of the head, which I suppose to be the natural attitude of display, similar to that assumed by *Thysostoma* and the *Heliodinidae*, but I am not aware of any record of the habit, and cannot explain its object. The species of this genus are rather large insects, more or less conspicuously decorated, sometimes with metallic markings; they evidently often occur plentifully, and their quaint appearance invites the attention of observers.

Geographical distribution of species. — Characteristic of Ceylon, with some stragglers in Southern India, one in Assam and one in Africa. This and the twelve preceding genera constitute a striking natural group which seems to have originated in Ceylon and spread thence to Southern India, only two species reaching Assam.

Larva unknown; it is not unlikely that the larval habits of the group may prove to be unusual and distinctive.

- | | |
|---|-------------------|
| 1. <i>T. extranea</i> , Walsingham, Trans. Ent. Soc. Lond. p. 105, pl. 5, f. 41 (1891). | Gambia. |
| 2. <i>T. phorcis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 444 (1908). | Ceylon. |
| 3. <i>T. orthadia</i> , Meyrick, ibidem, Vol. 17, p. 145 (1906). | Ceylon. |
| 4. <i>T. irrorella</i> , Walsingham, Moore's Lep. Ceyl. Vol. 3, p. 517, pl. 209, f. 9 (1886). | Ceylon, S. India. |
| 5. <i>T. pastas</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 444 (1908). | Coorg, Kanara. |
| 6. <i>T. autarcha</i> , Meyrick, ibidem, Vol. 18, p. 445 (1908). | S. India. |
| 7. <i>T. stachyophora</i> , Meyrick, ibidem, Vol. 18, p. 445 (1908). | Ceylon. |
| 8. <i>T. cincinnata</i> , Meyrick, Exot. Microlep. Vol. 1, p. 570 (1916). | Ceylon. |
| 9. <i>T. phycidella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 783 (1864) (- <i>cisella</i>).
<i>peronella</i> , Walker, ibidem, Vol. 29, p. 828 (1864). | Ceylon. |
| 10. <i>T. marmoritis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 144 (1906). | Ceylon. |
| 11. <i>T. selmatias</i> , Meyrick, ibidem, Vol. 18, p. 450 (1908). | Ceylon. |
| 12. <i>T. praeceptrix</i> , Meyrick, ibidem, Vol. 20, p. 461 (1910). | Ceylon. |
| 13. <i>T. mendicella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 797 (1864).
<i>torrentis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 449 (1908). | Ceylon. |
| 14. <i>T. aulonitis</i> , Meyrick, ibidem, Vol. 18, p. 447 (1908). | Ceylon. |
| 15. <i>T. xanthaula</i> , Meyrick, ibidem, Vol. 18, p. 448 (1908). | S. India. |
| 16. <i>T. toxastis</i> , Meyrick, ibidem, Vol. 18, p. 449 (1908). | Coorg, Kanara. |
| 17. <i>T. schoenola</i> , Meyrick, ibidem, Vol. 18, p. 448 (1908). | S. India. |
| 18. <i>T. machlas</i> , Meyrick, ibidem, Vol. 16, p. 595 (1905). — PI. 4, FIG. 82. | Ceylon. |
| 19. <i>T. temenodes</i> , Meyrick, Ent. Mitt. Vol. 11, p. 44 (1922). | Ceylon. |

20. *T. cingalensis*, Walsingham, Moore's Lep. Ceyl. Vol. 3, p. 522, pl. 209, Ceylon.
f. 12 (1886).
21. *T. parochra*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 146 Ceylon.
(1906).
22. *T. stasiotica*, Meyrick, ibidem, Vol. 18, p. 447 (1908). Ceylon.
23. *T. lecticaria*, Meyrick, Exot. Microlep. Vol. 1, p. 571 (1916). Ceylon.
24. *T. metallanthes*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 596 Ceylon.
(1905).
25. *T. peronetris*, Meyrick, ibidem, Vol. 17, p. 145 (1906). Ceylon.
26. *T. crassella*, Felder, Reise Novara, Lep. Vol. 2, pl. 139, f. 22 (1877).
sphenias, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 596 (1905). Ceylon.
27. *T. alloptila*, Meyrick, Exot. Microlep. Vol. 1, p. 570 (1916). Ceylon.
28. *T. tinctella*, Walsingham, Moore's Lep. Ceyl. Vol. 3, p. 517, pl. 209, Ceylon.
f. 11 (1886).
29. *T. dipsalea*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 446 (1908). Assam.
30. *T. pristica*, Meyrick, Exot. Microlep. Vol. 1, p. 571 (1916). Ceylon.

299. GENUS ANAXYRINA, MEYRICK

Anaxyrina, Meyrick, Exot. Microlep. Vol. 2, p. 98 (1918). — Type : *A. cyanopa*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli posterior; tongue developed. Antennae over 1, in ♂ rather stout, serrulate, simple, basal joint moderate, without pecten. Labial palpi moderately long, recurved, second joint with appressed scales, slightly rough beneath, terminal joint half second, evenly scaled and slightly roughened anteriorly, obtuse. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above towards base, with large expanded median and apical tufts of rough scales, basal joint of tarsi rough-scaled above. Forewings with 1b short-furcate, 2 from 4/5, 3 absent, 4 and 5 stalked from angle, 7 absent, 11 from beyond middle. Hindwings 1, trapezoidal, apex obtuse, termen sinuate, cilia 1; 2 remote, 3 from angle, 4 absent, 5 approximated, 6 and 7 stalked.

Remarks. — Closely related to *Canthonistis*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *A. cyanopa*, Meyrick, Exot. Microlep. Vol. 2, p. 99 (1918). Coorg.

300. GENUS HETERODELTIS, NOV. GEN.

Type : *H. trichroa*, Meyrick.

Characters. — Head smooth-scaled; ocelli posterior; tongue developed. Antennae 1 1/2, in ♂ simple, basal joint elongate, without pecten, in ♀ with small apical scale-projection. Labial palpi in ♂ with second joint expanded with dense scales, oval, concave internally, terminal joint absent; in ♀ very long, recurved, second joint smooth-scaled, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae and tarsi rough-scaled above, with large dense median tuft on tibiae. Forewings with 1b simple, 2 from angle, 3 absent, 4 and 5 stalked, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings 1, elongate-trapezoidal, cilia over 1; 3 and 4 connate, 5 obsolete, 6 and 7 long-stalked; in ♂ an expandible hairpencil from base in submedian groove.

Remarks. — Immediate affinity dubious, but perhaps nearer *Heteralcis*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *H. trichroa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 142 (1906). Ceylon.

301. GENUS CANTHONISTIS, MEYRICK

Canthonistis, Meyrick, Zool. Med. Leid. Vol. 7, p. 82 (1922). — Type : *C. amphicarpa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1 (?), in ♂ rather stout, simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, slender, pointed. Maxillary palpi rudimentary. Posterior tibiae with median tuft of long scales, apical half with appressed scales. Forewings with 1b simple, 2 from towards angle, 3 absent, 4 and 5 short-stalked, 7 absent, 8 and 9 stalked, 11 from before middle. Hindwings 1, elongate-trapezoidal-ovate, cilia 1; 3 and 4 coincident, 5 closely approximated at base, 6 and 7 stalked.

Remarks. — Probably derivable from *Habrogenes*.

Geographical distribution of species. — Malayan.

Larva unknown.

1. *C. amphicarpa*, Meyrick, Zool. Med. Leid. Vol. 7, p. 82 (1922). Java.

302. GENUS HABROGENES, MEYRICK

Habrogenes, Meyrick, Exot. Microlep. Vol. 2, p. 102 (1918). — Type : *H. eupatris*, Meyrick.

Characters. — Head with appressed scales, sidetufts somewhat raised; ocelli very small, posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely haired above, with rough expanded median tuft of scales. Forewings with 1b furcate, 2 and 3 long-stalked, 8 and 9 out of 7, 7 to apex, 11 from beyond middle. Hindwings 1, trapezoidal, termen hardly sinuate, cilia 3/4; 3 and 4 stalked, 5 slightly approximated, 6 and 7 stalked.

Remarks. — This and the preceding genera probably indicate a line of development collateral with the *Timyra* group.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

1. *H. leucochlora*, Meyrick, Trans. Ent. Soc. Lond. p. 448 (1910). Borneo.
2. *H. eupatris*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 443 (1910). Assam.
— Pl. 4, Fig. 87.

303. GENUS HYPTIASTIS, MEYRICK

Hyptiastis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 733 (1911). — Type : *H. clematis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1, somewhat thickened towards base, in ♂ with short very fine sparse cilia towards base, basal joint elongate, without pecten. Labial palpi very long, curved, ascending, second joint thickened with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae densely hairy above. Forewings with 2 and 3 short-stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, apex obtuse, termen hardly sinuate, cilia 1/2-3/5; 4 and 5 out of 3, transverse vein absent between 5 and 6, 6 and 7 stalked.

Remarks. — A peculiar form, which may be related to *Timyra*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *H. clematis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 734 (1911). S. India.

304. GENUS FRISILIA, WALKER

Frisilia, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 795 (1864). — Type : *F. nesciatella*, Walker.

Tipasa, Walker, ibidem, Vol. 29, p. 804 (1864) (praeocc.). — Type : *F. nesciatella*, Walker.

Macrernis, Meyrick, Trans. Ent. Soc. Lond. p. 275 (1887). — Type : *F. helipta*, Meyrick.

Characters. — Head with appressed scales, sidetufts in ♂ projecting; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, more or less thickened and roughened towards base, basal joint moderate, without pecten. Labial palpi in ♂ with second joint short, with dense projecting tuft of scales beneath, terminal joint wholly clothed with dense rough scales, bent over so as to form a longer tuft lying above former; in ♀ recurved, second joint rough-scaled above and tufted beneath, terminal joint longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings in ♂ with subdorsal groove edged with dense scales from base to tornus; 2 and 3 stalked or 3 usually absent in ♂ and sometimes in ♀, 4 out of 2 in ♂ and seldom in ♀, 7 to termen, 8 and 9 out of 7 or 8 absent, 11 from middle. Hindwings 1, trapezoidal, termen somewhat sinuate, cilia 2/3-4/5; 3 and 4 stalked, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — A well-marked and distinct genus; the species however are plainly and simply marked, and difficult of discrimination.

Geographical distribution of species. — Chiefly Indian, extending into the Malayan and African regions.

Larva unknown.

- | | |
|---|-------------------|
| 1. <i>F. melanardis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 446 (1910). | Borneo. |
| 2. <i>F. homochlora</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 437 (1910). | S. India. |
| 3. <i>F. verticosa</i> , Meyrick, ibidem, Vol. 22, p. 772 (1914). | S. India. |
| 4. <i>F. rostrata</i> , Meyrick, ibidem, Vol. 17, p. 147 (1906). — Pl. 4, Fig. 85. | Ceylon. |
| 5. <i>F. compstoma</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 87 (1921). | Rhodesia. |
| 6. <i>F. sulcata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 437 (1910). | Assam, Himalayas. |
| 7. <i>F. dipsia</i> , Meyrick, ibidem, Vol. 20, p. 437 (1910). | Ceylon. |
| 8. <i>F. procentra</i> , Meyrick, Exot. Microlep. Vol. 1, p. 573 (1916). | S. India. |
| 9. <i>F. nesciatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 796 (1864).
<i>basaliella</i> , Walker, ibidem, Vol. 29, p. 805 (1864). | Ceylon. |
| 10. <i>F. strepsiptila</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 436 (1910). | Coorg. |

11. *F. notifica*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 438 (1910). Ceylon.
 12. *F. triturata*, Meyrick, Exot. Microlep. Vol. 1, p. 277 (1914). Nyassaland.
 13. *F. senilis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 436 (1910). Coorg.
 14. *F. heliaptia*, Meyrick, Trans. Ent. Soc. Lond. p. 275 (1887). Ceylon.
 15. *F. indigens*, Meyrick, Suppl. Ent. p. 50 (1914). Formosa.

305. GENUS NARTHECOCEROS, MEYRICK

Narthecoceros, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 148 (1906). — Type : *N. platyconta*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1, thick, flatly compressed throughout, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint shorter than second, acute. Maxillary palpi rudimentary. Posterior tibiae rough-haired above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, termen faintly sinuate, cilia 1; 3 and 4 connate, 5 parallel, 6 and 7 long-stalked.

Remarks. — Allied to *Frisilia*; specially distinguished by the peculiar taeniate antennae.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *N. logica*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 435 (1910). Ceylon.
 2. *N. platyconta*, Meyrick, ibidem, Vol. 16, p. 597 (1905). — Pl. 4, Fig. 86. Ceylon.
 3. *N. xyloides*, Meyrick, ibidem, Vol. 17, p. 148 (1906). Ceylon.

306. GENUS TELEPHATA, MEYRICK

Telephata, Meyrick, Exot. Microlep. Vol. 1, p. 592 (1916). — Type : *T. cheramopis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1, in ♂ very short, compressed, simple, basal joint moderate, without pecten. Labial palpi moderately long, curved, ascending, second joint much thickened with dense scales, rough beneath, terminal joint somewhat shorter than second, moderate, scaled, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with 2 and 3 stalked from angle, 7 and 8 long-stalked, 7 to termen, 11 from beyond middle. Hindwings 1, elongate-trapezoidal, termen faintly sinuate, cilia 1; 2 and 3 slightly approximated towards base, 3 and 4 widely remote, parallel, 5 hardly approximated to 4, 6 and 7 long-stalked.

Remarks. — Possibly allied to the preceding.

Geographical distribution of species. — Papuan.

Larva unknown.

1. *T. cheramopis*, Meyrick, Exot. Microlep. Vol. 1, p. 593 (1916). New Guinea.

307. GENUS STEREMNIODES, MEYRICK

Steremniodes, Meyrick, Exot. Microlep. Vol. 3, p. 37 (1923). — Type : *S. sciactis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ stout, serrulate, simple, basal joint moderate, without pecten. Labial palpi moderately long,

curved, ascending, second joint much thickened with dense appressed scales, terminal joint shorter than second, stout, pointed. Maxillary palpi rudimentary. Posterior tibiae rough-haired above. Forewings with 2 and 3 stalked from angle, 4 approximated, 5 absent, 6 from near 7, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, termen faintly sinuate, cilia 3/5; 3 and 4 stalked, 5 rather approximated, 6 and 7 stalked.

Remarks. — Although the structural characters are not strongly pronounced, this genus seems naturally referable here, though geographically exceptional.

Geographical distribution of species. — South American.

Larva unknown.

1. *S. sciactis*, Meyrick, Exot. Microlep. Vol. 3, p. 38 (1923)

Guiana, Brazil.

308. GENUS SYNCATHEDRA, MEYRICK

Syncathedra, Meyrick, Exot. Microlep. Vol. 3, p. 37 (1923). — Type: *S. criminata*, Meyrick.

Characters. — Head with appressed scales, sidetufts rather loose; ocelli posterior; tongue developed. Antennae nearly 1, somewhat flattened, not tapering, in ♂ stout, simple, in ♀ rather slender, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint shorter than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with rough hairs above. Forewings with 1b furcate, 2 and 3 stalked from angle, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, elongate-ovate, rather pointed, cilia 1; 3 and 4 connate, 5 nearly parallel, 6 and 7 long-stalked.

Remarks. — Probably related to *Narthecoceros*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *S. criminata*, Meyrick, Exot. Microlep. Vol. 3, p. 37 (1923).

Assam.

309. GENUS PSAMMORIS, MEYRICK

Psammoris, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 149 (1906). — Type: *P. carpaea*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 1, in ♂ rather thick, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with scales, shortly projecting beneath towards apex, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with loosely appressed hairs. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings 1, trapezoidal, termen sinuate, cilia 1; 4 absent, 3 and 5 connate or short-stalked, 6 and 7 stalked.

Remarks. — A development of *Pseudodraces*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *P. carpaea*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 149 (1906). Ceylon.

310. GENUS PSEUDOCRATES, MEYRICK

Pseudocrates, Meyrick, Exot. Microlep. Vol. 2, p. 99 (1918). — Type : *P. antisphena*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli small, posterior; tongue developed. Antennae 4/5, in ♂ serrulate, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, loose beneath towards apex, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above. Forewings with 1b furcate, 2 from near angle, 7 and 8 stalked, 7 to termen, 9 connate with 7, 11 from middle. Hindwings 1, pointed-trapezoidal, termen slightly sinuate, cilia 1 1/4; 4 absent, 3 and 5 connate or short-stalked, 6 and 7 stalked.

Remarks. — Derivable from *Homaloxestis*.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|--|-----------|
| 1. <i>P. antisphena</i> , Meyrick, Exot. Microlep. Vol. 2, p. 99 (1918). | S. India. |
| 2. <i>P. soritica</i> , Meyrick, ibidem, Vol. 2, p. 100 (1918). | S. India. |

311. GENUS HELIANGARA, MEYRICK

Heliangara, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 147 (1906). — Type : *H. lampetis*, Meyrick.

Characters. — Head with appressed scales, face retreating; ocelli posterior; tongue developed. Antennae over 1, thick, compressed, in ♂ simple, basal joint moderate, without pecten. Labial palpi moderately long, curved, ascending, smooth-scaled, terminal joint shorter than second, acute. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 from angle, 3 absent, 7 and 8 stalked, 7 to costa, 9 and 10 from near 7, 11 from before middle. Hindwings 1, elongate-ovate, cilia 1 1/2; 3 and 4 stalked, 5 parallel, 6 and 7 long-stalked.

Remarks. — A development of *Mnesistega*, with interesting features; the unusual metallic coppery or purple colouring probably indicates the habit of flying in sunshine.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|--|------------------------|
| 1. <i>H. ericydes</i> , Meyrick, Exot. Microlep. Vol. 1, p. 573 (1916). | Ceylon. |
| 2. <i>H. macaritis</i> , Meyrick, Rec. Ind. Mus. Vol. 5, p. 221 (1910). | Bombay, Bengal, Assam. |
| 3. <i>H. lampetis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 147 (1906). — Pl. 4, Fig. 87. | Ceylon. |

312. GENUS MNESISTEGA, MEYRICK

Mnesistega, Meyrick, Exot. Microlep. Vol. 2, p. 101 (1918). — Type : *M. talantodes*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ somewhat stout, simple, basal joint moderate, with pecten. Labial palpi moderately long, recurved, second joint rather thickened with scales, somewhat rough towards apex beneath, terminal

joint nearly as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to termen, 9 from near 7, 11 from beyond middle. Hindwings 1, trapezoidal, narrowed posteriorly, apex pointed, termen sinuate, cilia 1 1/2; 3 and 4 connate, 5 nearly parallel, 6 and 7 closely approximated towards base.

Remarks. — Of doubtful immediate affinity; the sporadic appearance of an antennal pecten is again a curious feature.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|---|--------|
| 1. <i>M. talantodes</i> , Meyrick, Exot. Microlep. Vol. 2, p. 101 (1918). | Coorg. |
| 2. <i>M. convexa</i> , Meyrick, ibidem, Vol. 3, p. 36 (1923). | Assam. |

313. GENUS PARELLIPTIS, MEYRICK

Parelliptis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 439 (1910). — Type: *P. scytalias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1 or over 1, rather stout towards base, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense appressed scales, slightly rough beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 from towards angle, 4 and 5 sometimes stalked, 7 absent, 8 and 9 sometimes stalked, 11 from middle. Hindwings slightly over 1, elongate-trapezoidal, termen slightly sinuate, cilia about 1; 4 absent, 3 and 5 rather approximated at base or stalked, 6 and 7 stalked.

Remarks. — Probably derived from *Homaloxestis*.

Geographical distribution of species. — Indo-Malayan.

Larva unknown,

- | | |
|---|---------|
| 1. <i>P. librata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 445 (1910). | Java. |
| 2. <i>P. scytalias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 439 (1910). | Ceylon. |

314. GENUS DOLICHOTORNA, MEYRICK

Dolichotorna, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 438 (1910). — Type: *D. tholias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with scales, slightly rough towards apex beneath, terminal joint as long as second, in ♂ posteriorly with erect tuft from base and short median projection of scales, in ♀ slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long hairs above. Forewings with 2 from towards angle, 7 absent, 11 from middle. Hindwings 1, very elongate-trapezoidal, apex obtuse, termen rounded, cilia 1 2/3; 3 and 4 out of 5, cell apparently open, 6 and 7 long-stalked.

Remarks. — A derivative of *Homaloxestis*.

Geographical distribution of species. — Ceylon.

Larva unknown.

- | | |
|---|---------|
| 1. <i>D. tholias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 439 (1910). | Ceylon. |
|---|---------|

315. GENUS ACHORIA, MEYRICK

Achoria, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 405 (1904). — Type : *A. inopina*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened beneath with rough scales triangularly projecting at apex, terminal joint as long as second, slightly rough anteriorly, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 and 3 parallel, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings 1, elongate, apex and termen obtusely rounded, cilia 1 1/4; 3 separate, 4 and 5 stalked, transverse vein absent, 6 and 7 stalked, 6 to costa; in ♂ with hair-pencil from base in discal groove.

Remarks. — A development of *Neocorodes*.

Geographical distribution of species. — Australian.

Larva unknown.

1. *A. inopina*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 405 (1904). Queensland, New South Wales.

316. GENUS NEOCORODES, MEYRICK

Neocorodes, Meyrick, Exot. Microlep. Vol. 3, p. 36 (1923). — Type : *N. amnesta*, Meyrick.

Characters. — Head loosely scaled, sidetufts raised, connivent; ocelli posterior; tongue developed. Antennae 1, in ♂ rather stout, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint slightly longer, slender, acute. Maxillary palpi rudimentary. Forewings with 2 from towards angle, 3 and 4 approximated from angle, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings 1, elongate-trapezoidal-ovate, cilia over 1; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Derived from *Homaloxestis*.

Geographical distribution of species. — Mediterranean.

Larva unknown.

1. *N. amnesta*, Meyrick, Exot. Microlep. Vol. 3, p. 36 (1923). Cyprus.

317. GENUS ENTHETICA, MEYRICK

Enthetica, Meyrick, Exot. Microlep. Vol. 1, p. 574 (1916). — Type : *E. picryntis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint tufted with long rough projecting hairs towards apex beneath, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long rough hairs above. Forewings with 2 from towards angle, angularly bent and connected by bar with 1c. 3 from angle, 7 and 8 stalked, 7 to

costa, 11 from before middle. Hindwings 1, elongate-trapezoidal, termen somewhat sinuate, cilia 1 1/4; 3 and 4 connate or stalked, 5 nearly parallel, 6 and 7 long-stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|--|-----------|
| 1. <i>E. picryntis</i> , Meyrick, Exot. Microlep. Vol. 1, p. 574 (1916). | S. India. |
| 2. <i>E. tribrachia</i> , Meyrick, ibidem, Vol. 3, p. 38 (1923). | S. India. |

318. GENUS PHARANGITIS, MEYRICK

Pharangitis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 597 (1905). — Type : *P. spathias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ thick, simple, basal joint moderate, without pecten. Labial palpi long, curved, ascending, second joint thickened with scales, roughly projecting towards apex beneath, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from near angle, 3 absent, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, elongate-trapezoidal, termen faintly sinuate, cilia 1; 3 and 4 remote, 5 parallel, 6 and 7 long-stalked.

Remarks. — A distinct form, of uncertain affinity.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *P. spathias*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 597 Ceylon.
(1905). — Pl. 4, Fig. 93.

319. GENUS CROCOGMA, MEYRICK

Crocogma, Meyrick, Exot. Microlep. Vol. 2, p. 100 (1918). — Type : *C. isocola*, Meyrick.

Demopractis, Meyrick, ibidem, Vol. 2, p. 154 (1918). — Type : *C. isocola*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli small, posterior; tongue developed. Antennae 1, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint smooth-scaled, terminal joint shorter than second, moderate, pointed, in ♂ with rough spreading hairs posteriorly. Maxillary palpi rudimentary. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 3 and 4 connate from angle, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia nearly 1, 3 and 4 coincident, 5 nearly parallel, 6 and 7 stalked.

Remarks. — Derivable from *Homaloxestis*.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|--|--------|
| 1. <i>C. isocola</i> , Meyrick, Exot. Microlep. Vol. 2, p. 100 (1918) (♂). | Assam. |
| <i>tonaea</i> , Meyrick, ibidem, Vol. 2, p. 154 (1918) (♀). | |

320. GENUS ERIDACHTHA, MEYRICK

Eridachtha, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 440 (1910). — Type : *E. prolocha*, Meyrick.

Corthyntis, Meyrick, Exot. Microlep. Vol. 1, p. 574 (1916). — Type : *E. prolocha*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1, in ♂ simple, sometimes with small basal notch, basal joint moderate, without pecten, in ♂ sometimes with apicat tuft. Labial palpi long, recurved, second joint thickened with appressed scales, in ♂ sometimes with large expandible tuft of hairs above, terminal joint as long as second or in ♂ rather shorter, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1^b furcate, 2 from near angle, 3 normally separate, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, termen slightly sinuate, cilia nearly 1; 3 and 5 stalked, 4 absent, 6 and 7 stalked.

Remarks. — A development of *Homaloxestis*.

Geographical distribution of species. — Chiefly African, with one species in India.

Larva unknown.

- | | |
|---|---------------|
| 1. <i>E. prolocha</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 440 S. India.
(1910) (♀). | |
| <i>chlorotricha</i> , Meyrick, Exot. Microlep. Vol. 1, p. 575 (1916) (♂). | |
| 2. <i>E. crossogramma</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 87 (1921). | Rhodesia. |
| 3. <i>E. longicornella</i> , Chrétien, Ann. Soc. Ent. Fr. p. 335 (1915). | Algeria. |
| 4. <i>E. parvella</i> , Chrétien, ibidem, p. 335 (1915). | Algeria. |
| 5. <i>E. phaeochlora</i> , Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 76 (1920). | Kenya Colony. |
| 6. <i>E. calamopsis</i> , Meyrick, ibidem, Vol. 2, p. 76 (1920). | Kenya Colony. |
| 7. <i>E. cosymbota</i> , Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 285 (1920). | Cape Colony. |

321. GENUS CELETODES, MEYRICK

Celetodes, Meyrick, Zool. Med. Leid. Vol. 6, p. 166 (1921). — Type : *C. dracopis*, Meyrick.

Characters. — Head with appressed hairs, sidetufts slightly raised; ocelli posterior; tongue developed. Antennae 1, in ♂ stout, serrulate, simple, basal joint moderate, without pecten. Labial palpi moderate & long, curved, ascending, second joint thickened with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae loosely haired above. Forewings with 2 from towards angle, 3 and 4 long-stalked or coincident, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings 1, trapezoidal, apex tolerably pointed, termen slightly sinuate, cilia over 1; 3 and 4 connate, 5 rather approximated, 6 and 7 long-stalked

Remarks. — Probably derived from *Homaloxestis*.

Geographical distribution of species. — Malayan.

Larva unknown.

- | | |
|--|-------|
| 1. <i>C. dracopis</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 166 (1921). | Java. |
|--|-------|

322. GENUS ATHRINACIA, WALSINGHAM

Athrinacia, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 104 (1911). — Type : *A. xanthographa*, Walsingham.

Characters. — Head with appressed scales, sidetufts spreading; ocelli posterior; tongue developed. Antennae 4/5 to almost 1, in ♂ rather stout, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint smooth-scaled, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b slightly furcate or simple, 2 from towards angle, 3-5 approximated, 7 absent, 11 from middle or 4/5. Hindwings over 1, trapezoidal, termen somewhat sinuate, cilia 1/2; 3 and 4 coincident, 5 somewhat approximated, 6 and 7 approximated towards base.

Characters. — A genus of uncertain affinity, but possibly allied to the following.

Geographical distribution of species. — Tropical American.

Larva unknown

- | | |
|--|---------|
| 1. <i>A. cosmophragma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 116 (1922). | Brazil. |
| 2. <i>A. trifasciata</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 105 (1911). | Mexico. |
| 3. <i>A. xanthographa</i> , Walsingham, ibidem, Vol. 4, p. 105, pl. 3, fig. 27 (1911). | Mexico. |
| 4. <i>A. leucographa</i> , Walsingham, ibidem, Vol. 4, p. 106 (1911). | Mexico. |

323. GENUS THRYPSIGENES, MEYRICK

Thrypsigenes, Meyrick, Trans. Ent. Soc. Lond. p. 272 (1914). — Type : *T. colluta*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ rather stout, simple, basal joint moderate, without pecten. Labial palpi moderately long, curved, ascending, second joint thickened with scales angularly at apex beneath, terminal joint 2/3 of second, moderate, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 3 from angle, 7 absent, 8 and 9 short-stalked or connate, 11 from beyond middle. Hindwings under 1, elongate-trapezoidal, apex tolerably pointed, termen slightly sinuate, cilia 1 1/2; 3 and 4 connate, 5 absent, 6 and 7 long stalked.

Remarks. — A development of *Lioclepta*.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|---|-----------------|
| 1. <i>T. colluta</i> , Meyrick, Trans. Ent. Soc. Lond. p. 272 (1914). | Guiana, Brazil. |
| 2. <i>T. furvescens</i> , Meyrick, ibidem, p. 272 (1914). | Guiana, Brazil. |

324. GENUS LIOCLEPTA, MEYRICK

Lioclepta, Meyrick, Trans. Ent. Soc. Lond. p. 115 (1922). — Type : *L. complanata*, Meyrick.

Characters. — Head smooth, sidetufts slightly raised; ocelli posterior; tongue developed. Antennae 5/6, in ♂ stout, simple, basal joint elongate, without pecten. Labial palpi moderately long.

recurved, second joint with scales roughly tufted at apex beneath, terminal joint somewhat shorter than second, slender, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long rough hairs above. Forewings with 1b short-furcate, 2 from towards angle, 7 to costa, 8 and 9 out of 7, 11 from beyond middle. Hindwings somewhat under 1, trapezoidal, termen slightly sinuate, cilia 3/4; 3 and 4 stalked, 5 absent, 6 and 7 stalked.

Remarks. — Probably derivable from *Homaloxestis*.

Geographical distribution of species. — South American.

Larva unknown.

1. *L. complanata*, Meyrick, Trans. Ent. Soc. Lond. p. 116 (1922). Peru.

325. GENUS EUPRAGIA, WALSINGHAM

Eupragia, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 106 (1911). — Type: *E. solida*, Walsingham.

Characters. — Head loosely scaled, sidetufts raised; tongue developed. Antennae 3/4, in ♂ stout, simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint longer than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b short-furcate, 2 from 3/4, 3 and 4 closely approximated from angle, 7 absent, 8 and 9 approximated at base, 11 from middle, 12 sinuate and closely approximated to 11 near beyond origin. Hindwings over 1, trapezoidal, termen faintly sinuate; 3 and 4 in ♂ coincident, in ♀ stalked, 5 rather approximated, 6 and 7 almost connate.

Remarks. — Of doubtful immediate affinity.

Geographical distribution of species. — North American.

Larva unknown.

1. *E. solida*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 107, pl. 3, Mexico.
f. 28 (1911).

326. GENUS PHTHORACMA, MEYRICK

Pthoracma, Meyrick, Ann. Transv. Mus. Vol. 8, p. 87 (1921). — Type: *P. blanda*, Meyrick.

Characters. — Head with appressed scales, sidetufts somewhat raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ rather stout, serrulate, minutely ciliated, basal joint elongate, without pecten. Labial palpi with second joint long, curved, ascending, rather slender, with appressed scales, terminal joint (♂) minute, rudimentary, filiform. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long fine hairs above. Forewings with cell 3/4 of wing, 2 from near angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal, termen sinuate beneath apex, cilia 1; 3 and 4 connate, 5 absent, 6 and 7 stalked.

Remarks. — Also of uncertain location.

Geographical distribution of species. — African.

Larva unknown.

1. *P. blanda*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 87 (1921). Transvaal.

327. GENUS APHNOGENES, MEYRICK

Aphnogenes, Meyrick, Ann. Transv. Mus. Vol. 8, p. 88 (1921). — Type : *A. zonata*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint somewhat thickened with appressed scales, terminal joint nearly as long as second, moderate, pointed. Maxillary palpi rudimentary. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 short-stalked from angle, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings somewhat under 1, elongate-trapezoidal, apex obtuse, termen hardly sinuate, cilia 1; 2 and 3 stalked, 4 absent, 5 rising out of stalk of 2 and 3, transverse vein absent, 6 free, 7 wholly absent.

Remarks. — An abnormally modified form, of which the ♀ would probably possess more regular characters.

Geographical distribution of species. — African.

Larva unknown.

1. *A. zonata*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 88 (1921). Rhodesia.

328. GENUS CARTERICA, NOV. GEN.

Type : *C. phthoneropa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6 (?), in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 7 absent, 8 and 9 connate, 11 from middle. Hindwings 1, trapezoidal; 3 and 4 connate, 5 rather approximated, 6 and 7 connate.

Remarks. — Perhaps allied to the following.

Geographical distribution of species. — Chinese.

Larva unknown.

1. *C. phthoneropa*, Meyrick, Exot. Microlep. Vol. 2, p. 505 (1922). China.

329. GENUS CYMATOPLEX, NOV. GEN.

Type : *C. aestuosa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint 3/5 of second, thick, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 3-5 approximated, 7 absent, 8 and 9 connate, 11 from middle. Hindwings 1, elongate-trapezoidal, termen sinuate, cilia 1; 3 and 4 connate, 5 rather approximated, 6 and 7 connate.

Remarks. — An inconspicuous form of doubtful affinity.

Geographical distribution of species. — African.

Larva unknown.

1. *C. aestuosa*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 295 (1913).

Transv., Natal, Comoro Is.

330. GENUS CARODISTA, NOV. GEN.

Type : *C. flagitiosa*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5/6, in ♂ stout, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 2 from towards angle, 7 and 8 stalked, 7 to apex, 9 and 10 stalked. Hindwings 1, trapezoidal; 3 and 4 connate, 5 rather approximated, 6 and 7 stalked.

Remarks. — Probably a derivative of *Homaloxestis*.

Geographical distribution of species. — African.

Larva unknown.

1. *C. flagitiosa*, Meyrick, Exot. Microlep. Vol. 1, p. 198 (1914).

Nyassaland.

331. GENUS HOMALOXESTIS, MEYRICK

Homaloxestis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 440 (1910). — Type : *H. endocoma*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1 or over 1, rather stout towards base, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 from towards angle, 3 separate, 4 and 5 sometimes stalked, 7 and 8 stalked, 7 to apex or termen, 9 often out of 7, 11 from middle. Hindwings 1, elongate-trapezoidal, apex more or less pointed, termen faintly sinuate or rounded, cilia 2/3-2; 3 and 4 connate or stalked, 5 rather approximated, 6 and 7 stalked.

Remarks. — A development from *Lecithocera*, and with similar neural variation. The males are often variously furnished with expandible tufts of hairs.

Geographical distribution of species. — Indo-Malayan; the common and widely distributed *H. cholopis* is probably spread artificially, and reaches Africa.

Larva (*cholopis*) feeding in spun leaves.

Foodplant *Mallotus* (*Euphorbiaceae*).

- | | |
|---|---------|
| 1. <i>H. lochitis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 101 (1918). | Coorg. |
| 2. <i>H. turbinata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 442 (1910). | Assam. |
| 3. <i>H. melicrata</i> , Meyrick, ibidem, Vol. 20, p. 442 (1910). | Assam. |
| 4. <i>H. galeodes</i> , Meyrick, ibidem, Vol. 20, p. 442 (1910). | Assam. |
| 5. <i>H. cribanota</i> , Meyrick, ibidem, Vol. 20, p. 442 (1910). | Assam. |
| 6. <i>H. antibathra</i> , Meyrick, Exot. Microlep. Vol. 1, p. 575 (1916). | Kanara. |

7. <i>H. callitricha</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 440 (1910).	Assam.
8. <i>H. periseucta</i> , Meyrick, ibidem, Vol. 20, p. 441 (1910).	Assam.
9. <i>H. perichlora</i> , Meyrick, Trans Ent. Soc. Lond. p. 446 (1910).	Borneo.
10. <i>H. queribunda</i> , Meyrick, Zool. Med. Leid. Vol. 7, p. 83 (1922).	Java.
11. <i>H. liciata</i> , Meyrick, ibidem, Vol. 7, p. 83 (1922).	Java.
12. <i>H. ceroxesta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 102 (1918).	Bombay, Java.
13. <i>H. chiloptila</i> , Meyrick, ibidem, Vol. 2, p. 435 (1921).	Assam.
14. <i>H. endocoma</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 441 (1910).	S. India.
15. <i>H. plocamandra</i> , Meyrick, ibidem, Vol. 17, p. 737 (1907).	Sikkim, Bhotan, Assam.
16. <i>H. ochroscles</i> , Meyrick, ibidem, vol. 20, p. 441 (1910).	Ceylon.
17. <i>H. xylotripta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 102 (1918).	Himalayas.
18. <i>H. tenuipalpella</i> , Snellen, Tijdschr. v. Ent. Vol. 46, p. 35 (1903).	Java. [mosa, S. Africa.]
19. <i>H. cholopis</i> , Meyrick. Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 149 (1906).	India, Burma, Java, For-

332. GENUS PROCHARISTA, MEYRICK

Procharista, Meyrick, Zool. Med. Leid. Vol. 7, p. 82 (1922). — Type : *P. sardonias*, Meyrick.

Characters. — Head smooth. sidetufts loosely raised. Antennae over 1, in ♂ simple, basal joint elongate. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Forewings with 1b short-furcate, 2 and 4 connate from angle, 3 absent, 6 absent, 8 and 9 out of 7, 7 to costa, 11 from beyond middle. Hindwings 1, elongate-trapezoidal-ovate, cilia nearly 1; 3 and 4 stalked, 5 absent, 6 and 7 long-stalked.

Remarks. — A development of *Lecithocera*.

Geographical distribution of species. — Malayan.

Larva unknown.

1. *P. sardonias*, Meyrick, Zool. Med. Leid. Vol. 7, p. 83 (1922). Java.

333. GENUS PHATNOTIS, MEYRICK

Phatnotis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 180 (1913). — Type : *P. factiosa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae almost 1, in ♂ ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with scales roughly expanded towards apex above and dense rather rough scales beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above and beneath. Forewings with 1b furcate, 2 and 3 long-stalked, 7 to termen, 8-10 out of 7, 11 from beyond middle. Hindwings somewhat over 1, trapezoidal, termen somewhat sinuate, cilia 1/2; 3 and 4 connate or stalked, 5 approximated, 6 and 7 stalked.

Remarks. — Correlated with *Epharmonia*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *P. factiosa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 181 (1913). S. India.
2. *P. legata*, Meyrick, ibidem, Vol. 22, p. 181 (1913). S. India.

334. GENUS EPHARMONIA, NOV. GEN.

Type *E. ardua*, Meyrick.

Characters. — Head with appressed scales, sidetufts slightly raised; ocelli posterior; tongue developed. Antenne 5/6, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 4 and 5 separate, 7 to termen, 8 and 9 out of 7, 11 from middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia 1/2; 3 and 5 stalked, 4 absent, 6 and 7 stalked.

Remarks. — Related to *Deltoplastis*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *E. pselaphistis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 456 Assam.
(1910).
2. *E. ardua*, Meyrick, ibidem. Vol. 20, p. 458 (1910). Assam.

335. GENUS SISYRODONTA, MEYRICK

Sisyrodonta, Meyrick, Ark. f. Zool. Vol. 14, n° 15, p. 5 (1922). — Type : *S. ochrosidera*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue imperceptible (?). Antennae under 1 (?), in ♂ stout, strongly lamellate, stalk and lamellae wholly clothed with dense scales, basal joint stout, without pecten. Labial palpi very long, recurved, second joint thickened with dense appressed scales, terminal joint longer than second, rather slender, scaled, acute. Maxillary palpi imperceptible. Forewings with 2 and 4 stalked from angle, 3 absent, 5 connate with 2, 8 and 9 out of 7, 7 to apex (indefinite), 11 from middle. Hindwings 1, trapezoidal, termen faintly sinuate, cilia 1/2; 3 and 5 stalked, 4 absent, 6 and 7 stalked.

Remarks. — Correlated with *Ptilothyris*, an interesting connection.

Geographical distribution of species. — Australian.

Larva unknown.

1. *S. ochrosidera*, Meyrick, Ark. f. Zool. Vol. 14, n° 15, p. 6 (1922). N. W. Australia.

336. GENUS PTILOTHYRIS, WALSHAM

Ptilothyris, Walsingham, Trans. Ent. Soc. Lond. p. 37 (1897). — Type : *P. purpurea*, Walsingham.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 4/5, in ♂ stalk thickened with scales except towards apex, bipectinated, basal joint elongate, without pecten. Labial palpi very long, recurved, smooth, second joint thickened with appressed scales, compressed, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with appressed scales. Forewings with 1b furcate, 3 and 4 out of 2, 5 approximated,

7 absent, 8 and 9 stalked, 11 from middle. Hindwings over 1, trapezoidal-ovate, termen hardly sinuate, cilia 1/6; 2 remote, 3 and 5 connate, 4 absent, transverse vein obsolete, 6 and 7 stalked.

Remarks. — A derivative of *Deltoplastis*. The neuration as given by Walsingham is incorrect.

Geographical distribution of species. — African.

Larva unknown.

1. *P. purpurea*, Walsingham, Trans. Ent. Soc. Lond. p. 38, pl. 2, f. 2 (1897). S. Nigeria, Fr. Congo.

337. GENUS IDIOPTERYX, WALSINGHAM

Idiopteryx, Walsingham, Trans. Ent. Soc. Lond. p. 104 (1891). — Type : *I. obliquella*, Walsingham.

Dragmatucha, Meyrick, Proc. Zool. Soc. Lond. p. 726 (1908). — Type : *I. proaula*, Meyrick.

Characters. — Head with appressed hairs; ocelli posterior; tongue small. Antennae 4/5, in ♂ ciliated or stout, simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, sometimes with loose tuft beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform. Posterior tibiae clothed with very long rough spreading hairs. Forewings with 1b furcate, 3 and 4 out of 2 or sometimes 3 absent, 5 approximated, 7 to costa, 8 and 9 out of 7, 11 from beyond middle. Hindwings over 1, trapezoidal, apex obtuse, termen hardly sinuate, cilia 1/2; 4 absent, 5 nearly parallel to 3, 6 and 7 stalked.

Remarks. — Correlated with the preceding. In this genus also the neuration is given wrongly by Walsingham.

Geographical distribution of species. — African.

Larva unknown.

1. *I. proaula*, Meyrick, Proc. Zool. Soc. Lond. p. 726 (1908). Pl. 4, Fig. 94. Transvaal, Rhodesia.

2. *I. obliquella*, Walsingham, Trans. Ent. Soc. Lond. p. 254, pl. 11, f. 22 (1881). Natal.
obsepta, Meyrick, Ann. Transv. Mus. Vol. 6, p. 24 (1918).

3. *I. haeresiella*, Wallengren, Oef. Af. Kon. Vet. Akad. För. p. 128 (1875). Transvaal.
4. *I. bivia*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 25 (1918). Natal.

338. GENUS HYPEROCHTHA, NOV. GEN.

Type : *H. butyropa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ moderately ciliated or simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 4 connate with 2 or 5 or absent, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 3/4; 4 absent, 3 and 5 stalked, 6 and 7 stalked.

Remarks. — A derivative of *Deltoplastis*.

Geographical distribution of species. — Indian and African.

Larva unknown.

- | | |
|--|------------|
| 1. <i>H. dischema</i> , Meyrick, Exot. Microlep. Vol. 1, p. 576 (1916). | Nyassaland |
| 2. <i>H. butyropa</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 458 (1910). | Ceylon. |
| 3. <i>H. justa</i> , Meyrick, ibidem, Vol. 20, p. 458 (1910). | Ceylon. |
| 4. <i>H. metriodes</i> , Meyrick, Exot. Microlep. Vol. 2, p. 112 (1918). | Bombay. |

339. GENUS DELTOPLASTIS, NOV. GEN.

Type : *D. ocreata*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked or coincident, 4 and 5 connate or stalked or coincident, 8 and 9 out of 7, 7 to apex, 11 from beyond middle. Hindwings 1 or over 1, trapezoidal, termen sinuate, cilia 1/2-4/5; 4 absent, 3 and 5 stalked, 6 and 7 stalked.

Remarks — Probably derived from *Gasmara*.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

- | | |
|--|-----------------------|
| 1. <i>D. coercita</i> , Meyrick, Exot. Microlep. Vol. 3, p. 42 (1923). | Coorg. |
| 2. <i>D. scopulosa</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 450 (1910). | Coorg. |
| 3. <i>D. clerodotis</i> , Meyrick, ibidem, Vol. 20, p. 450 (1910). | Ceylon. |
| 4. <i>D. amicella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 634 (1864).
<i>obligatella</i> , Walker, ibidem, Vol. 30, p. 1023 (1864). | Assam, Burma, Ceylon. |
| 5. <i>D. tetradelta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 150 (1906). | Ceylon. |
| 6. <i>D. causidica</i> , Meyrick, ibidem, Vol. 20, p. 451 (1910). | Assam. |
| 7. <i>D. ocreata</i> , Meyrick, ibidem, Vol. 20, p. 451 (1910). | Bombay, Madras. |
| 8. <i>D. similella</i> , Snellen, Tijdschr. v. Ent. Vol. 46, p. 44, pl. 5, f. 5 (1903). | Java. |
| 9. <i>D. horistis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 452 (1910). | Assam. |
| 10. <i>D. balanitis</i> , Meyrick, ibidem, Vol. 20, p. 452 (1910). | S. India. |
| 11. <i>D. cremuaspis</i> , Meyrick, ibidem, Vol. 16, p. 598 (1905). — Pl. 4, Fig. 92. | Ceylon. |
| 12. <i>D. straminicornis</i> , Meyrick, ibidem, Vol. 20, p. 453 (1910). | Ceylon. |
| 13. <i>D. caduca</i> , Meyrick, ibidem, Vol. 20, p. 454 (1910). | Assam. |
| 14. <i>D. figurata</i> , Meyrick, ibidem, Vol. 20, p. 453 (1910). | Ceylon. |
| 15. <i>D. propensa</i> , Meyrick, ibidem, Vol. 20, p. 456 (1910). | Ceylon. |
| 16. <i>D. byssina</i> , Meyrick, ibidem, Vol. 20, p. 457 (1910). | Ceylon. |
| 17. <i>D. acrophanes</i> , Meyrick, ibidem, Vol. 20, p. 457 (1910). | Ceylon. |
| 18. <i>D. commodata</i> , Meyrick, Exot. Microlep. Vol. 3, p. 44 (1923). | Ceylon. |
| 19. <i>D. leptobrocha</i> , Meyrick, ibidem, Vol. 3, p. 44 (1923). | Coorg. |

340. GENUS COYDALLA, WALKER

Coydalla, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1037 (1864). — Type : *C. interguttella*, Walker.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae over 1, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint smooth, terminal joint much

longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 from near angle, 3 absent, 4 and 5 connate from angle, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 3/5; 4 absent, 3 and 5 stalked, 6 and 7 stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Malayan

Larva unknown.

1. *C. interguttella*, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1038 (1864). Borneo.

341. GENUS HARMATITIS, MEYRICK

Harmatitis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 460 (1910). — Type : *H. sphecopa*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue strong. Antennae 5/6, in ♂ stout, scaled, strongly ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, rather rough beneath, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae smooth-scaled, with apical expanded whorl, basal joint of tarsi rough-scaled above. Forewings with 1b furcate, 2 and 3 coincident, 4 and 5 connate from angle, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings somewhat over 1, trapezoidal, termen faintly sinuate, cilia 1/2; 4 absent, 5 rather approximated to 3, 6 and 7 stalked.

Remarks. — A development of *Gasmara*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *H. sphecopa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 460 Ceylon. (1910) (♂).

? *eucerella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 636 (1864).

hippotarcha, Meyrick, Exot. Microlep. Vol. 3, p. 41 (1923) (♀).

342. GENUS GASMARA, WALKER

Gasmara, Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1040 (1864). — Type : *G. coelatella*, Walker.

Antiochtha, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 598 (1905). — Type : *G. balbidota*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked or coincident, 4 usually out of 2, 5 in a straight line with lower margin of cell, connate or nearly with 4 or 3, 7 to termen, 8 and 9 out of 7, 11 from beyond middle. Hindwings 1, trapezoidal, termen somewhat sinuate, cilia 3/4; beneath with scattered blackish scales on costal half; 4 absent, 3 and 5 connate or seldom separate, 6 and 7 stalked.

Remarks. — Derived from *Lecithocera*. The curious persistence of the scattered black scales on undersurface of hindwings, an unusual character, makes it worth noticing as a generic feature.

Geographical distribution of species. — Characteristically Ceylonese, but extending into Southern India.

Larva unknown.

- | | |
|--|-----------|
| 1. <i>G. stellulata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 149 (1906). | Ceylon. |
| 2. <i>G. periastra</i> , Meyrick, ibidem, Vol. 20, p. 454 (1910). | Ceylon. |
| 3. <i>G. oxyzona</i> , Meyrick, ibidem, Vol. 20, p. 455 (1910). | Ceylon. |
| 4. <i>G. coelatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 30, p. 1040 (1864). | Ceylon. |
| 5. <i>G. achnastis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 150 (1906). | Ceylon. |
| 6. <i>G. balbidota</i> , Meyrick, ibidem, Vol. 16, p. 598 (1905). | Ceylon. |
| 7. <i>G. foederalis</i> , Meyrick, Exot. Microlep. Vol. 3, p. 43 (1923). | S. India. |
| 8. <i>G. leucograpta</i> , Meyrick, ibidem, Vol. 3, p. 45 (1923). | S. India. |
| 9. <i>G. vigilax</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 455 (1910). | Ceylon. |
| 10. <i>G. cataclina</i> , Meyrick, Exot. Microlep. Vol. 3, p. 44 (1923). | Ceylon. |

343. GENUS CYNICOSTOLA, NOV. GEN.

Type : *C. pogonias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint beneath tufted with long rough projecting scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1^b furcate, 2 and 4 stalked, 3 absent, 5 connate with 2, in a line with lower margin of cell, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings 1, elongate-trapezoidal, termen slightly sinuate, cilia 2/3; 4 absent, 3 and 5 stalked, 6 and 7 stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|--|-----------|
| 1. <i>C. pogonias</i> , Meyrick, Exot. Microlep. Vol. 3, p. 43 (1923). | S. India. |
|--|-----------|

344. GENUS THYMBRITIS, NOV. GEN.

Type : *T. molybdias*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6, in ♂ simple, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1^b furcate, 2 and 3 stalked, 4 and 5 separate, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 1; 4 absent, 3 and 5 stalked, 6 and 7 stalked.

Remarks. — Probably derived from *Lecithocera*.

Geographical distribution of species. — Ceylon.

Larva unknown.

- | | |
|---|---------|
| 1. <i>T. molybdias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 456 (1910). | Ceylon. |
|---|---------|

345. GENUS CROCANTHES, MEYRICK

Crocantes, Meyrick, Trans. Ent. Soc. Lond. p. 277 (1886). — Type : *C. prasinopis*, Meyrick.

Aprosoesta, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 151 (1919). — Type : *C. pancala*, Turner.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 1 or over 1, in ♂ simple or ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, smooth, second joint slender or somewhat thickened, terminal joint as long as second or longer, acute, in ♂ sometimes much modified. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with 1b furcate, 2 and 4 connate or stalked, 3 absent. 7 absent, 9 and 10 sometimes out of 8, or 9 sometimes absent, 11 from beyond middle. Hindwings 1, trapezoidal, termen somewhat sinuate, cilia 1/2-1; 4 absent, 3 and 5 stalked, transverse vein more or less obsolete, 6 and 7 stalked.

Remarks. — Probably a development from *Sarisophora*. The species are elegant insects, more or less adorned with yellow or sometimes rosy colouring; the typical species is remarkable in life for its emerald-green eyes. The antennae are frequently porrected in repose.

Geographical distribution of species. — Australian and Papuan; originating in New Guinea.

Larva unknown, not improbably of exceptional habit.

- | | |
|---|--|
| 1. <i>C. sidonia</i> , Meyrick, Trans. Ent. Soc. Lond. p. 442 (1910). — Pl. 4, Fig. 83. | Queensl., New Guinea. |
| 2. <i>C. temeraria</i> , Meyrick, ibidem, p. 443 (1910). | New Guinea. |
| 3. <i>C. rhodantha</i> , Meyrick, Exot. Microlep. Vol. 2, p. 97 (1918). | New Guinea. |
| 4. <i>C. pancala</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 151 (1919). | Queensland. |
| 5. <i>C. scioxantha</i> , Meyrick, Trans. Ent. Soc. Lond. p. 443 (1910). | New Guinea. |
| 6. <i>C. halurga</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 399 (1904). | Queensland. |
| 7. <i>C. zonodesma</i> , Lower, ibidem, Vol. 25, p. 50 (1900). | Queensl., N. Australia. |
| 8. <i>C. thermobapta</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 44, p. 65 (1920). | Queensland. |
| 9. <i>C. prasinopis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 277 (1886). — Pl. 5,
Fig. 119 a, b. | E. Australia, Tasmania,
New Guinea. |
| 10. <i>C. glycina</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 400 (1904). | Victoria, Tasmania. |
| 11. <i>C. acroxantha</i> , Lower, Trans. Roy. Soc. S. Austr. Vol. 20, p. 170 (1896). | Queensland. [Wales. |
| 12. <i>C. perigrapta</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 402 (1904). | Queensland, New South
N. Australia. |
| 13. <i>C. characotis</i> , Meyrick, Exot. Microlep. Vol. 1, p. 592 (1916). | Queensland. |
| 14. <i>C. chordotona</i> , Meyrick, ibidem, Vol. 1, p. 592 (1916). | Queensland. |
| 15. <i>C. trizona</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 40, p. 543 (1916). | Queensland. |
| 16. <i>C. epitherma</i> , Lower, ibidem, Vol. 20, p. 170 (1896). | Queensland. [Wales. |
| 17. <i>C. micradelpha</i> , Lower, ibidem, Vol. 21, p. 56 (1897). | Queensland, New South
New Guinea. |
| 18. <i>C. fallax</i> , Durrant, Lep. Wollaston Exped. p. 164 (1915). | New Guinea. |
| 19. <i>C. crypsichola</i> , Durrant, ibidem, p. 165 (1915). | New Guinea. |
| 20. <i>C. celema</i> , Durrant, ibidem, p. 152 (1915). | New Guinea. |
| 21. <i>C. diula</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 398 (1904). | Queensland. |
| 22. <i>C. doliope</i> , Meyrick, Exot. Microlep. Vol. 2, p. 434 (1921). | Queensland. |
| 23. <i>C. zonias</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 403 (1904). | Queensland. |
| 24. <i>C. gelastis</i> , Meyrick, Exot. Microlep. Vol. 2, p. 97 (1918). | New Guinea. |
| 25. <i>C. thrasydora</i> , Meyrick, Trans. Ent. Soc. Lond. p. 444 (1910). | New Guinea. |
| 26. <i>C. millina</i> , Durrant, Lep. Wollaston Exped. p. 151 (1915). | New Guinea. |
| 27. <i>C. carcharias</i> , Meyrick, Trans. Ent. Soc. Lond. p. 444 (1910). — Pl. 4,
Fig. 99. | New Guinea. |
| 28. <i>C. eurypyra</i> , Meyrick, Exot. Microlep. Vol. 2, p. 97 (1918). | New Guinea. |
| 29. <i>C. sceletopa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 445 (1910). | Moluccas. |
| 30. <i>C. pyrochorda</i> , Meyrick, ibidem, p. 442 (1910). — Pl. 4, Fig. 84. | New Guinea. |

346. GENUS GONAEPA, WALKER

Gonaepa, Walker, List Lep. Het. Brit. Mus. Vol. 35, p. 1840 (1866). — Type : *G. josianella*, Walker.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae nearly 1, in ♂ strongly biciliated, basal joint elongate, without pecten. Labial palpi very long, recurved slender, smooth, terminal joint as long as second, acute. Maxillary palpi rudimentary. Posterior tibiae somewhat rough-scaled above. Forewings with 1b simple, 2 and 4 stalked, 3 absent, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 1/3; 4 absent, 3 and 5 connate or stalked, 6 and 7 connate or stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Papuan.

Larva unknown.

- | | |
|--|-------------|
| 1. <i>G. actinis</i> , Walsingham, Lep. Wollaston Exped. p. 149 (1915). | New Guinea. |
| 2. <i>G. heliarcha</i> , Meyrick, Trans. Ent. Soc. Lond. p. 278 (1886). | New Guinea. |
| 3. <i>G. josianella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 35, p. 1840 (1866). | New Guinea. |

347. GENUS PLACANTHES, MEYRICK

Placanthes, Meyrick, Exot. Microlep. Vol. 3, p. 42 (1923). — Type : *P. xanthomorpha*, Meyrick.

Characters. — Head smooth; ocelli very small, posterior; tongue developed. Antennae 1 (?), basal joint elongate, without pecten. Labial palpi long, recurved, second joint somewhat thickened, smooth, terminal joint longer than second, slender, acute. Maxillary palpi very short, loosely scaled. Posterior tibiae shortly rough-scaled above. Forewings with 1b furcate, 2 and 4 stalked from angle, 3 absent, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, costa with some rough projecting scales anteriorly, apex forming a long pointed projection (nearly 1/3 of wing), termen rectangularly emarginate beneath this, cilia 1/5; 4 absent, 3 and 5 connate, 6 and 7 stalked.

Remarks. — An interesting and undoubtedly connecting link between the preceding and following genera.

Geographical distribution of species. — Malayan.

Larva unknown.

- | | |
|--|--------------|
| 1. <i>P. xanthomorpha</i> , Meyrick, Exot. Microlep. Vol. 3, p. 42 (1923). | Philippines. |
|--|--------------|

348. GENUS NOSPHISTICA, MEYRICK

Nosphistica, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 733 (1911). — Type : *N. erraticata*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 5.6, in ♂ with long fine ciliations, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint smooth-scaled, terminal joint as long as second, slender, acute. Maxillary palpi short, filiform, appressed to tongue. Posterior tibiae rough-scaled above, basal joint of tarsi tufted at apex.

Forewings with 3 and 4 out of 2, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings over 1, trapezoidal, termen irregularly sinuate, cilia 2/5, costa with projecting scaleteeth; 4 absent, 3 and 5 connate, 6 and 7 stalked.

Remarks. — Nearly related to the following.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *N. erratica*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 733 (1911). Ceylon.

Pl. 4, Fig. 89.

349. GENUS PHILOPTILA, MEYRICK

Philoptila, Meyrick, Exot. Microlep. Vol. 2, p. 111 (1918). — Type : *P. effrenata*, Meyrick.

Characters. — Head smooth, face retreating; ocelli small, posterior; tongue developed. Antennae over 1, in ♂ strongly ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Anterior tibiae very short, tarsi long; posterior legs with rough scaletufts above on origin of tibial spurs and apex of basal joint of tarsi. Forewings with 1b furcate, 3 and 4 out of 2, 5 approximated, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, costa with strong rough scale-projections before and beyond middle, termen bisinuate, cilia 1/3; 4 absent, 3 and 5 connate, 6 and 7 connate.

Remarks. — Probably a development of *Thubana*. The bisinuation of hindwings is exaggerated by the cilia, and with the costal tufts produces a singular ragged effect, probably displayed in repose.

Geographical distribution of species. — Indian.

Larva unknown.

1. *P. effrenata*, Meyrick, Exot. Microlep. Vol. 2, p. 111 (1918).

Kanara.

350. GENUS COPROPTILIA, SNELLEN

Coproptilia, Snellen, Tijdschr. v. Ent. Vol. 46, p. 32 (1903). — Type : *C. glebicolorella*, Snellen.

Characters. — Head with appressed scales, sidetufts somewhat raised; tongue developed. Antennae nearly 1, in ♂ ciliated. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint as long as second, slender, acute. Posterior tibiae rough-scaled above. Forewings with 3 and 4 out of 2, 5 connate, 8 and 9 out of 7, 7 to termen, 10 connate with 7, 11 absent. Hindwings over 1, trapezoidal, cilia 1/4; 4 absent, 3 and 5 connate, 6 and 7 connate.

Remarks. — Not known to me, but obviously referable here; apparently derived from *Thubana*. The absence of vein 11 of forewings seems improbable, but is explicitly stated by Snellen.

Geographical distribution of species. — Malayan.

Larva unknown.

1. *C. glebicolorella*, Snellen, Tijdschr. v. Ent. Vol. 46, p. 34, pl. 5, f. 4 (1903). Java, Sumatra.

351. GENUS ADELOMORPHA, SNELLEN

Adeломорфа., Snellen, Tijdschr. v. Ent. Vol. 28, p. 31 (1885). — Type : *A. Ritsemae*, Snellen.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 1 in ♂ fasciculate-ciliated towards base, basal joint elongate, without pecten. Labial palpi very long, recurved, smooth, second joint somewhat thickened, terminal joint as long as second, slender, acute. Maxillary palpi very short, loosely scaled, appressed to tongue. Posterior tibiae loosely scaled above. Forewings with 1^b furcate, 3 and 4 out of 2, 5 connate with 2 from angle, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings over 1, trapezoidal, termen sinuate, cilia 1/4; 3 and 4 short-stalked, 5 equidistant, 6 and 7 short-stalked.

Remarks. — A strikingly conspicuous form, presumably derived from *Thubana*.

Geographical distribution of species. — Malayan.

Larva unknown.

1. *A. Ritsemae*, Snellen, Tijdschr. v. Ent. Vol. 28, p. 32, pl. 3, f. 1-3 (1885). Celebes, Ceram.

352. GENUS SPHENOCRATES, NOV. GEN.

Type : *S. aulodocha*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 1, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint smooth-scaled, terminal joint longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae shortly rough-scaled above. Forewings with 2 and 4 stalked from angle, 3 absent, 5 connate with 2, in a line with lower margin of cell, 8 and 9 out of 7, 7 to apex, 10 absent, 11 from beyond middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 1/2; 4 absent, 3 and 5 connate, 6 and 7 stalked.

Remarks. — Probably a derivative of *Thubana*.

Geographical distribution of species. — Papuan.

Larva unknown.

1. *S. aulodocha*, Meyrick, Exot. Microlep. Vol. 2, p. 98 (1918).

New Guinea.

353. GENUS THUBANA, WALKER

Thubana, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 814 (1864). — Type : *T. bisignatella*, Walker.

Tiva, Walker, ibidem, Vol. 29, p. 821 (1864). — Type : *T. bisignatella*, Walker.

Inapha, Walker, ibidem, Vol. 30, p. 999 (1864). — Type : *T. bisignatella*, Walker.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1 or over 1, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1^b furcate, 3 and 4 out of 2, or 3 absent, 5 connate with 2 from angle of cell, 8 and 9 out of 7, 7 to apex or

termen, or sometimes 7 absent, 8 and 9 stalked. Hindwings over 1, trapezoidal, termen more or less sinuate, cilia 2/5-3/5; 3 and 4 stalked, 5 nearer 4, somewhat approximated at base, 6 and 7 stalked.

Remarks. — A development of *Lecithocera*.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

1. <i>T. bisignatella</i> , Walker, List. Lep. Het Brit. Mus. Vol. 29, p. 814 (1864).	Borneo, Formosa.
<i>binotella</i> , Walker, ibidem, Vol. 29, p. 822 (1864).	
<i>lampronalis</i> , Walker, ibidem, Vol. 30, p. 1000 (1864).	
2. <i>T. albisignis</i> , Meyrick, Suppl. Ent. p. 50 (1914).	Formosa.
3. <i>T. costimaculella</i> , Snellen, Tijdschr. v. Ent. Vol. 46, p. 37, pl. 4, f. 7 (1903).	Java.
4. <i>T. bullulata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 443 (1910).	Assam.
5. <i>T. nodosa</i> , Meyrick, Trans. Ent. Soc. Lond. p. 447 (1910).	Malay States.
6. <i>T. Heylaertsi</i> , Snellen, Tijdschr. v. Ent. Vol. 46, p. 38, pl. 4, f. 8 (1903).	Java.
7. <i>T. laxata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 713 (1911).	Assam.
8. <i>T. nardinopa</i> , Meyrick, Exot. Microlep. Vol. 2, p. 103 (1918).	Coorg.
9. <i>T. isocrypta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 709 (1911).	Ceylon.
10. <i>T. xylogramma</i> , Meyrick, Zool. Med. Leid. Vol. 7, p. 86 (1922).	Java.
11. <i>T. xanthoteles</i> , Meyrick, Exot. Microlep. Vol. 3, p. 38 (1923).	Assam, Burma.
12. <i>T. melitopyga</i> , Meyrick, ibidem, Vol. 3, p. 41 (1923).	Ceylon.
13. <i>T. residua</i> , Meyrick, ibidem, Vol. 3, p. 41 (1923).	Assam.

354. GENUS BRACHYERGA, NOV. GEN.

Type : *B. hemiacma*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1, in ♂ stout, simple, basal joint elongate, without pecten. Labial palpi long, curved, ascending, second joint with appressed scales, terminal joint half second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 4 and 5 separate, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, termen faintly sinuate, cilia 3/4; 3 and 4 stalked, 5 nearer 4, somewhat approximated at base, 6 and 7 stalked.

Remarks. — A derivative of *Lecithocera*.

Geographical distribution of species. — Malayan.

Larva unknown.

1. <i>B. hemiacma</i> , Meyrick, Trans. Ent. Soc. Lond. p. 448 (1910).	Borneo.
--	---------

355. GENUS PERIPHORECTIS, NOV. GEN.

Type : *P. ichorodes*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint somewhat thickened with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae with appressed scales. Fore-

wings with 1b furcate, 2 and 3 stalked from angle, 4 and 5 separate, 8 and 9 out of 7, 7 to costa, 11 from middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia $\frac{3}{4}$; 3 and 4 stalked, 5 rather approximated at base, much nearer 4, 6 and 7 stalked.

Remarks. — Derived from *Lecithocera*, from which it differs essentially only by costal termination of vein 7 of forewings.

Geographical distribution of species. — Indian.

Larva unknown.

1. *P. ichorodes*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 445 (1910). S. India.

356. GENUS SARISOPHORA, MEYRICK.

Sarisophora. Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 403 (1904). — Type : *S. leptoglypta*, Meyrick.

Styloceros. Meyrick, ibidem, Vol. 29, p. 408 (1904). — Type : *S. cyclonitis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae over 1, in ♂ more or less stout, simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, sometimes somewhat rough towards apex beneath, terminal joint as long as second, moderate or slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to apex or termen, 9 seldom out of 7, 11 from middle. Hindwings 1, trapezoidal, termen not or slightly sinuate, cilia $\frac{3}{4}$ -1; 4 absent, 3 and 5 connate or stalked, 6 and 7 stalked.

Remarks. — A development of *Lecithocera*. In some of the species the antennae are porrected in repose.

Geographical distribution of species. — Australian and European; but it is perhaps still uncertain whether connecting forms may not be found in the intermediate region.

Larva (2 known) feeding on dead leaves.

- | | |
|--|------------------------------|
| 1. <i>S. leptoglypta</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 404 (1904). | Queensland. |
| Pl. 4, Fig. 98. [Wales.] | |
| 2. <i>S. chlaenota</i> , Meyrick, ibidem, Vol. 29, p. 404 (1904). | Queensland, New South Wales. |
| 3. <i>S. nyctiphylax</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 154 (1919). | Queensland. |
| 4. <i>S. ptochomorpha</i> , Meyrick, Exot. Microlep. Vol. 3, p. 36 (1923). | Cyprus. |
| 5. <i>S. sicculella</i> , Wocke, Nat. Sicil. Vol. 9, p. 2 (1889). | Sicily. |
| 6. <i>S. briantiella</i> , Turati, Bull. Soc. Ent. Ital. p. 202, pl. 8, f. 15 (1879). | S. Europe, Asia Minor. |
| 7. <i>S. pallicornella</i> , Staudinger, Stett. Ent. Zeit. Vol. 20, p. 239 (1859). | Spain, S. France. |
| 8. <i>S. flavissimella</i> , Mann, Wien. Ent. Monatsschr. Vol. 6, p. 402, pl. 3, f. 14 (1862). | Macedonia, Asia Minor. |
| 9. <i>S. tamides</i> , Meyrick, Trans. Ent. Soc. Lond. p. 445 (1910). | New Guinea. |
| 10. <i>S. terrena</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 153 (1919). | Queensland. |
| 11. <i>S. tenella</i> , Turner, ibidem, Vol. 31, p. 153 (1919). | Queensland. |
| 12. <i>S. leucoscia</i> , Turner, ibidem, Vol. 31, p. 154 (1919). | Queensland. |
| 13. <i>S. pycnospila</i> , Turner, ibidem, Vol. 31, p. 155 (1919). | Queensland. |
| 14. <i>S. cyclonitis</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 409 (1904). | Queensland. |
| 15. <i>S. brachymita</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 157 (1919). | Queensland. |
| 16. <i>S. dispila</i> , Turner, ibidem, Vol. 31, p. 157 (1919). | North Australia. |

357. GENUS LECITHOCERA, HERRICH-SCHÄFFER

Lecithocera, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 45 (1853). — Type : *L. luteicornella*, Zeller.

Tiriza, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 790 (1864). — Type : *L. leucotella*, Walker.

Titana, Walker, ibidem, Vol. 29, p. 813 (1864). — Type : *L. adelella*, Walker.

Tirasia, Walker, ibidem, Vol. 29, p. 817 (1864) (praeocc.). — Type : *L. punctigeneralis*, Walker.

Patouissa, Walker, ibidem, Vol. 29, p. 820 (1864). — Type : *L. dissonella*, Walker.

? **Andusia**, Walker, ibidem, Vol. 35, p. 1836 (1866). — Type : *L. alternella*, Walker.

Slovata, Walker, ibidem, Vol. 35, p. 1837 (1866). — Type : *L. pulcherrimella*, Walker.

Macrotona, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 405 (1904). — Type : *L. sobria*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennæ 1 or over 1, more or less thickened towards base, in ♂ simple or rarely ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, often somewhat rough towards apex beneath, rarely with rough projecting scales beneath, terminal joint as long as second or longer, slender, acute, or seldom in ♂ flat, obtuse. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with hairs above. Forewings with 1b furcate, 2 and 3 stalked or seldom coincident, 4 seldom stalked with 2 or sometimes stalked with 5, 5 from above angle, 7 to apex or termen, 8 and sometimes 9 out of 7, or seldom 8 absent, or 8 and 9 stalked, 7 absent, 11 from middle. Hindwings 1 or somewhat over 1, trapezoidal, termen hardly sinuate, cilia 2/3-4/5; 3 and 4 connate or stalked or sometimes coincident, 5 rather approximated or nearly parallel, nearer 4 than 6, 6 and 7 stalked.

Remarks. — A development of *Brachmia*. The neuration varies sometimes within the limits of the species, but usually furnishes good specific distinctions, which repay attention especially in the unicolorous forms. The numerous species are often very similar and obscurely coloured, but their differences, though small, are usually not difficult of apprehension; doubtless however a large number of species remain to be discovered. Even the typical European *luteicornella* has been the subject of much confusion among authors who have neglected to examine the neuration (correctly given by Herrich-Schäffer).

Geographical distribution of species. — Properly Indo-Malayan, in which region the development is extensive, but with considerable colonies in Africa and Australia, barely represented in Europe, and entirely absent from America (except one species apparently artificially introduced into South America from the Philippines) and from New Zealand.

Larva entirely unknown; not improbably the habit observed in *Sarisophora* of feeding on dead leaves may furnish a clue; also at least three species seem to have been spread artificially.

- | | |
|--|--------------|
| 1. <i>L. compsoiphila</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 709 (1911). | Ceylon. |
| 2. <i>L. innotatella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 807 (1864). | Borneo. |
| 3. <i>L. delestra</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 711 (1911). | Ceylon. |
| 4. <i>L. ptochas</i> , Meyrick, Exot. Microlep. Vol. 2, p. 104 (1918). | Bengal. |
| 5. <i>L. ochrocapna</i> , Meyrick, ibidem, Vol. 3, p. 40 (1923). | Philippines. |
| 6. <i>L. orbata</i> , Meyrick, Trans. Ent. Soc. Lond. p. 450 (1910). | Borneo. |
| 7. <i>L. adelella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 814 (1864). | Borneo. |
| 8. <i>L. recurvata</i> , Meyrick, Exot. Microlep. Vol. 3, p. 39 (1923). | Philippines. |

- | | | |
|-----|--|--|
| 9. | <i>L. picrodora</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 294 (1913). | Transvaal. |
| 10. | <i>L. protoma</i> , Meyrick, Exot. Microlep. Vol. 1, p. 198 (1914). | Gold Coast. [saland |
| 11. | <i>L. iresia</i> , Meyrick, Journ. Bombay. Nat. Hist. Soc. Vol. 20, p. 709 (1911). | S. India, Ceylon, Nyas- |
| 12. | <i>L. alcestis</i> , Meyrick, Exot. Microlep. Vol. 3, p. 40 (1923). | Coorg, Kanara. |
| 13. | <i>L. bracculata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 710
(1911). | Assam. |
| 14. | <i>L. argocrossa</i> , Meyrick, ibidem, Vol. 20, p. 710 (1911). | Ceylon. |
| 15. | <i>L. philosopha</i> , Meyrick, ibidem, Vol. 20, p. 708 (1911). | Assam. |
| 16. | <i>L. barbifera</i> , Meyrick, Zool. Med. Leid. Vol. 7, p. 84 (1922). | Java. |
| 17. | <i>L. invariella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 641 (1864).
<i>thranewta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 712 (1911).
<i>caligula</i> , Meyrick, Exot. Micr. Vol. 2, p. 104 (1918). | India, Ceylon, Borneo,
New Guinea. |
| 18. | <i>L. pellax</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 713 (1911). | Assam. |
| 19. | <i>L. metasaris</i> , Meyrick, ibidem, Vol. 20, p. 712 (1911). | Assam. |
| 20. | <i>L. opportuna</i> , Meyrick, Exot. Microlep. Vol. 3, p. 38 (1923). | Assam. |
| 21. | <i>L. corsota</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 712 (1911). | Assam. |
| 22. | <i>L. digna</i> , Meyrick, Exot. Microlep. Vol. 2, p. 105 (1918). | Assam. |
| 23. | <i>L. chlorogastra</i> , Meyrick, Zool. Med. Leid. Vol. 7, p. 84 (1922). | Java. |
| 24. | <i>L. diligens</i> , Meyrick, ibidem, Vol. 7, p. 84 (1922). | Java. |
| 25. | <i>L. omphacias</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 445
(1910). | Ceylon. |
| 26. | <i>L. contracta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 107 (1918). | Kanara. |
| 27. | <i>L. integrata</i> , Meyrick, ibidem, Vol. 2, p. 107 (1918). | Kanara. |
| 28. | <i>L. semirupta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 445
(1910). | Assam. |
| 29. | <i>L. cornutella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 632 (1864). | Ceylon. |
| 30. | <i>L. elutens</i> , Meyrick, Exot. Microlep. Vol. 2, p. 108 (1918). | S. India. |
| 31. | <i>L. insidians</i> , Meyrick, ibidem, Vol. 2, p. 108 (1918). | Coorg. |
| 32. | <i>L. syntropha</i> , Meyrick, ibidem, Vol. 2, p. 109 (1918). | W. Himalayas. |
| 33. | <i>L. responsa</i> , Meyrick, ibidem, Vol. 2, p. 108 (1918). | Bombay. |
| 34. | <i>L. linocoma</i> , Meyrick, ibidem, Vol. 1, p. 593 (1916). | North Australia. |
| 35. | <i>L. noseropa</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 158 (1919). | Queensland. |
| 36. | <i>L. poliocoma</i> , Meyrick, Exot. Microlep. Vol. 1, p. 593 (1916). | North Australia. |
| 37. | <i>L. anympha</i> , Meyrick, ibidem, Vol. 1, p. 593 (1916). | North Australia. |
| 38. | <i>L. micromela</i> , Lower, Trans. Roy. Soc. S. Australia, Vol. 21, p. 55 (1897). | Queensland, New South
Wales, Victoria. |
| 39. | <i>L. terrigena</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 406 (1904). | New South Wales. |
| 40. | <i>L. imprudens</i> , Meyrick, Exot. Microlep. Vol. 1, p. 201 (1914). | New South Wales. |
| 41. | <i>L. alampes</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 156 (1919). | Queensland. |
| 42. | <i>L. distigmatella</i> , Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 366 (1877). | ? Queensland. |
| 43. | <i>L. chamela</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 155 (1919). | Queensland. |
| 44. | <i>L. cyamitis</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 407 (1904). | Queensland. |
| 45. | <i>L. sobria</i> , Meyrick, ibidem, Vol. 29, p. 407 (1904). | Queensland, New South
Queensland. [Wales. |
| 46. | <i>L. concinna</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 157 (1919). | Queensland. |
| 47. | <i>L. isophanes</i> , Turner, ibidem, Vol. 31, p. 158 (1919). | Queensland. |
| 48. | <i>L. neosticta</i> , Meyrick, Exot. Microlep. Vol. 2, p. 107 (1918). | Coorg. |
| 49. | <i>L. masina</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 445 (1910). | Punjab. |
| 50. | <i>L. proclivis</i> , Meyrick, ibidem, Vol. 20, p. 446 (1910). | S. India. |
| 51. | <i>L. itinea</i> , Meyrick, ibidem, Vol. 20, p. 444 (1910). | India, Ceylon. |
| 52. | <i>L. pachyntis</i> , Meyrick, Trans. Ent. Soc. Lond. p. 17 (1894). | Burma. |
| 53. | <i>L. perspensa</i> , Meyrick, Exot. Microlep. Vol. 2, p. 153 (1918). | Assam. |
| 54. | <i>L. chersitis</i> , Meyrick, ibidem, Vol. 2, p. 106 (1918). | Korea. |
| 55. | <i>L. desolata</i> , Meyrick, ibidem, Vol. 2, p. 105 (1918). | Bombay, S. India. |
| 56. | <i>L. glaphyritis</i> , Meyrick, ibidem, Vol. 2, p. 106 (1918). | Ceylon. |

57. *L. sinuosa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 444 (1910). Ceylon.
 58. *L. macella*, Meyrick, Exot. Microlep. Vol. 2, p. 105 (1918). S. India.
 59. *L. perfida*, Meyrick, ibidem, Vol. 2, p. 105 (1918). S. India.
 60. *L. acolasta*, Meyrick, ibidem, Vol. 2, p. 236 (1919). Bombay.
 61. *L. caustospila*, Meyrick, ibidem, Vol. 2, p. 109 (1918). Assam.
 62. *L. metacausta*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 446 (1910) Assam.
 63. *L. antiphractis*, Meyrick, Exot. Microlep. Vol. 2, p. 435 (1921). Assam.
 64. *L. aulias*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 447 (1910). Assam.
 65. *L. jugalis*, Meyrick, Exot. Microlep. Vol. 2, p. 109 (1918). Bombay.
 66. *L. strangalistis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 722 (1911). Assam.
 67. *L. citrostrota*, Meyrick, ibidem, Vol. 20, p. 721 (1911). Assam.
 68. *L. hemichrysa*, Meyrick, ibidem, Vol. 20, p. 447 (1910). **Pl. 4, Fig. 90.** Assam.
 69. *L. megalopis*, Meyrick, Exot. Microlep. Vol. 1, p. 575 (1916). Philippines.
 70. *L. choritis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 448 (1910). S. India.
 71. *L. affusa*, Meyrick, Exot. Microlep. Vol. 3, p. 40 (1923). Assam. [gentina].
 72. *L. fausta*, Meyrick, Trans. Ent. Soc. Lond. p. 449 (1910). Philippines, Brazil, Ar-
 73. *L. bipunctella*, Snellen, Tijdschr. v. Ent. Vol. 46, p. 36, pl. 4, f. 6 (1903). Java.
 74. *L. dissonella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 821 (1864). Borneo.
 75. *L. querula*, Meyrick, Trans. Ent. Soc. Lond. p. 449 (1910). Java.
 76. *L. subservitella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 639 (1864). Borneo.
 77. *L. homocentra*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 449 (1910). Ceylon.
 78. *L. diplosticta*, Meyrick, Zool. Med. Leid. Vol. 7, p. 84 (1922). Java.
 79. *L. combusta*, Meyrick, Exot. Microlep. Vol. 2, p. 110 (1918). Ceylon.
 80. *L. caecilia*, Meyrick, ibidem, Vol. 2, p. 110 (1918). Ceylon.
 81. *L. autologa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 446 (1910). Ceylon.
 82. *L. pauperella*, Rebel, Denk. Akad. Wien, Vol. 93, p. 443 (1916). Egypt.
 83. *L. luridella*, Christoph, Bull. Soc. Nat. Mosc. p. 33 (1882). E. Siberia.
 84. *L. orsoviella*, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 361 (1870). Roumania.
 85. *L. luticornella*, Zeller, Isis, p. 197 (1839). C. & S. Europe.
 86. *L. xanthochalca*, Meyrick, Exot. Microlep. Vol. 1, p. 199 (1914). Nyassaland, Natal,
 flavipalpis, Walsingham, Trans. Ent. Soc. Lond. p. 105, pl. 5, fig. 40 (1891)
 (praeocc.).
 87. *L. acrosphales*, Meyrick, Exot. Microlep. Vol. 2, p. 108 (1918). Madagascar.
 88. *L. sceptrarcha*, Meyrick, Voyage Alluaud Jean. Lep. Vol. 2, p. 77 (1920). Kenya Colony.
 89. *L. isomitra*, Meyrick, Exot. Microlep. Vol. 1, p. 277 (1914). Nyassaland.
 90. *L. officialis*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 67 (1912). Transvaal.
 91. *L. frustrata*, Meyrick, Exot. Microlep. Vol. 2, p. 107 (1918). French Congo.
 92. *L. myopa*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 294 (1913). Transvaal.
 93. *L. propitia*, Meyrick, ibidem, Vol. 3, p. 293 (1913). Transvaal.
 94. *L. xanthocosma*, Meyrick, Exot. Microlep. Vol. 3, p. 46 (1923). Uganda.
 95. *L. flavipalpella*, Walsingham, Trans. Ent. Soc. Lond. p. 262, pl. 12, f. 31 (1881). Natal.
 96. *L. hybrista*, Meyrick, Exot. Microlep. Vol. 2, p. 505 (1922). Upper Senegal.
 97. *L. lucernata*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 294 (1913). Transvaal.
 98. *L. craniota*, Meyrick, ibidem, Vol. 3, p. 293 (1913). Transvaal.
 99. *L. anthologella*, Wallengren, Oef. Af. Kon. Vet. Akad. Foer, p. 129 (1875). Transvaal.
 100. *L. binotata*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 24 (1918). Natal.
 101. *L. obsignata*, Meyrick, Exot. Microlep. Vol. 1, p. 277 (1914). Nyassaland.
 102. *L. malacta*, Meyrick, ibidem, Vol. 2, p. 110 (1918). Comoro Islands.
 103. *L. pyxinodes*, Meyrick, ibidem, Vol. 2, p. 109 (1918). Madagascar.
 104. *L. spiladias*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 88 (1921). Port. E. Africa.

LEPIDOPTERA HETEROPTERA

105. *L. marginata*, Walsingham, Trans. Ent. Soc. Lond. p. 104, pl. 5, f. 39 Gambia, Egypt (1891).
 106. *L. coleasta*, Meyrick, Exot. Microlep. Vol. 2, p. 103 (1918). New Guinea.
 107. *L. oxyconia*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 444 (1910). India.
 108. *L. deloma*, Durrant, Lep. Wollaston Exped. p. 165 (1915). New Guinea.
 109. *L. thiodora*, Meyrick, Suppl. Ent. p. 51 (1914). Formosa.
 110. *L. perigypsa*, Meyrick, Zool. Med. Leid. Vol. 7, p. 85 (1922). Celebes.
 111. *L. storestis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 711 (1911). Ceylon.
 112. *L. nomaditis*, Meyrick, Exot. Microlep. Vol. 1, p. 594 (1916). Solomon Islands.
 113. *L. prudens*, Meyrick, ibidem, Vol. 2, p. 106 (1918). New Guinea.
 114. *L. nefasta*, Meyrick, ibidem, Vol. 1, p. 575 (1916). Kanara.
 115. *L. crebrata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 447 (1910). Coorg.
 116. *L. immobilis*, Meyrick, Exot. Microlep. Vol. 2, p. 103 (1918). S. India.
 117. *L. geraea*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 717 (1911). Ceylon.

Pl. 5, Fig. 118a, b.

118. *L. percobela*, Meyrick, ibidem, Vol. 20, p. 723 (1911). S. India.
 119. *L. puleolata*, Meyrick, ibidem, Vol. 20, p. 716 (1911). S. India.
 120. *L. elephantopa*, Meyrick, Rec. Ind. Mus. Vol. 5, p. 222 (1910). Bombay, S. India.
 121. *L. lycopis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 717 (1911). Ceylon.
 122. *L. haemylopis*, Meyrick, ibidem, Vol. 20, p. 716 (1911). Ceylon.
 123. *L. octonias*, Meyrick, Trans. Ent. Soc. Lond. p. 447 (1910). Borneo.
 124. *L. biferrinella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 642 (1864). Borneo.
 125. *L. cassiterota*, Meyrick, Exot. Microlep. Vol. 3, p. 40 (1923). Philippines.
 126. *L. lamprodesma*, Meyrick, Zool. Med. Leid. Vol. 7, p. 85 (1922). Celebes.
 127. *L. bitinctella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 675 (1864). Borneo.
 128. *L. leucotella*, Walker, ibidem, Vol. 29, p. 791 (1864). Borneo.
 129. *L. fuscedinella*, Snellen, Tijdschr. v. Ent. Vol. 44, p. 88, pl. 5, f. 8 (1901). Java.
 130. *L. pulcherrimella*, Walker, List Lep. Het. Brit. Mus. Vol. 35, p. 1838 (1866). Java.
 131. *L. clastrata*, Meyrick, Trans. Ent. Soc. Lond. p. 448 (1910). — **Pl. 4,** Borneo.

Fig. 91.

132. *L. ancylola*, Meyrick, ibidem, p. 17 (1894). Assam, Burma.
 133. *L. arcifera*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 738 (1907). Bhotan, S. India.
 134. *L. parallactis*, Meyrick, Trans. Ent. Soc. Lond. p. 17 (1894). Burma.
 135. *L. octavana*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 714 (1911). Assam.
 136. *L. sortilega*, Meyrick, ibidem, Vol. 20, p. 714 (1911). Assam.
 137. *L. protrocha*, Meyrick, Exot. Microlep. Vol. 1, p. 576 (1916). S. India.
 138. *L. paroena*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 148 (1906). Ceylon.
 139. *L. epomia*, Meyrick, ibidem, Vol. 16, p. 599 (1905). Ceylon.
 140. *L. paroristis*, Meyrick, ibidem, Vol. 20, p. 718 (1911). Ceylon.
 141. *L. praeses*, Meyrick, Exot. Microlep. Vol. 2, p. 236 (1919). Himalayas, Celebes.
 142. *L. isomila*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 718 (1911). Ceylon.
 143. *L. strophopa*, Meyrick, Exot. Microlep. Vol. 3, p. 46 (1923). S. India.
 144. *L. exophthalma*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 720 (1911). Ceylon.
 145. *L. rusticana*, Meyrick, Exot. Microlep. Vol. 2, p. 113 (1918). S. India.
 146. *L. parasema*, Meyrick, ibidem, Vol. 1, p. 66 (1913). Bengal, Kanara.
 147. *L. oxalea*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 449 (1910). Coorg.
 148. *L. fortis*, Meyrick, Exot. Microlep. Vol. 2, p. 111 (1918). S. India.
 149. *L. exaeca*, Meyrick, Zool. Med. Leid. Vol. 7, p. 86 (1922). Java.
 150. *L. carcerata*, Meyrick, Exot. Microlep. Vol. 3, p. 39 (1923). Assam.
 151. *L. punctigeneralis*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 818 (1864). Borneo.
 152. *L. trigonopsis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 737 (1907). Himalayas.

153. <i>L. strigosa</i> , Durrant, Lep. Wollaston Exped. p. 165 (1915).	New Guinea.
154. <i>L. dirupta</i> , Meyrick, Exot. Microlep. Vol. 3, p. 39 (1923).	S. India.
155. <i>L. deltospila</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 706 (1911).	Assam.
156. <i>L. signifera</i> , Felder, Reis. Novar. Lep. Vol. 2, pl. 139, f. 23 (1875).	Ceylon.
157. <i>L. cordata</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 720 (1911).	S. India.
158. <i>L. fornacalis</i> , Meyrick, ibidem, Vol. 20, p. 719 (1911).	Ceylon.
159. <i>L. nubigena</i> , Meyrick, ibidem, Vol. 20, p. 720 (1911).	Ceylon.
160. <i>L. epigompha</i> , Meyrick, ibidem, Vol. 20, p. 448 (1910).	Ceylon.
161. <i>L. capnaula</i> , Meyrick, ibidem, Vol. 20, p. 719 (1911).	Ceylon.
162. ? <i>L. alternella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 35, p. 1836 (1866).	Java.

358. GENUS ASMENISTIS, NOV. GEN.

Type : *A. cucullata*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 1 (?), basal joint elongate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 2 and 3 stalked, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal; 4 absent, 3 and 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Of rather doubtful affinity.

Geographical distribution of species. — African.

Larva unknown.

1. *A. cucullata*, Meyrick, Exot. Microlep. Vol. 1, p. 199 (1914). Nyassaland.

359. GENUS LARCOPHORA, NOV. GEN.

Type : *L. sophronistis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, basal joint elongate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 from near angle, 3 absent, 4 from angle, 5 remote, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings 1, trapezoidal, apex pointed, termen sinuate, cilia nearly 1; 4 absent, 3 and 5 connate, 6 and 7 stalked.

Remarks. — Probably correlated with *Lecithocera*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *L. sophronistis*, Meyrick, Exot. Microlep. Vol. 2, p. 112 (1918). Kanara.

360. GENUS CHLOROLYCHNIS, NOV. GEN.

Type : *C. agnatella*, Walker.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 5/6 in ♂ rather stout, simple, basal joint moderate, without pecten. Labial palpi long, recurved, second

joint thickened with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1 b furcate, 3 and 4 out of 2, 5 connate, 8 and 9 out of 7, 7 to apex, 11 from beyond middle. Hindwings slightly over 1, trapezoidal, termen hardly sinuate, cilia 2/3; 4 absent, 3 and 5 stalked, 6 and 7 stalked.

Remarks. — A derivative of *Protolychnis*.

Geographical distribution of species. — Indo-Malayan.

Larva unknown.

1. *C. agnatella*, Walker, List. Lep. Het. Brit. Mus. Vol. 29, p. 633 (1864). India, Ceylon, Java.

361. GENUS PROTOLYCHNIS, NOV. GEN.

Type: *P. maculata*, Walsingham.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ thick, simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1 b furcate, 2 and 3 stalked, 4 and 5 separate, 8 and 9 out of 7, 7 to apex, 11 from beyond middle. Hindwings over 1, trapezoidal, termen hardly sinuate, cilia 1/2; 4 absent, 3 and 5 stalked, 6 and 7 stalked.

Remarks. — Correlated with *Cophomantis*.

Geographical distribution of species. — African and Australian.

Larva unknown.

1. *P. trigonias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 408 (1904). Queensland.
 2. *P. chlorotoma*, Meyrick, Exot. Microlep. Vol. 1, p. 200 (1914). Nyassaland.
 3. *P. maculata*, Walsingham, Trans. Ent. Soc. Lond. 276, pl. 11, f. 18 (1881). C. et S. Africa.

362. GENUS COPHOMANTIS, NOV. GEN.

Type: *C. elaphopis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ moderately or strongly ciliated or simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second or shorter, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1 b furcate, 2 and 3 stalked, 4 out of 2 or absent, 7 absent, 8 and 9 stalked, 11 from beyond middle. Hindwings over 1, trapezoidal, termen slightly sinuate, cilia 1/4; 4 absent, 3 and 5 connate or stalked, 6 and 7 stalked.

Remarks. — A derivative of *Organitis*.

Geographical distribution of species. — Indo-Malayan and African, extending into North Australia; probably African in origin.

Larva unknown.

1. *C. elaphopis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 459 (1910). Assam, Borneo.
 2. *C. myadelpha*, Meyrick, ibidem, Vol. 20, p. 459 (1910). India.

3. *C. eremota*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 706 (1911). Ceylon.
 4. *C. lychnocentra*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 409 North Australia.
 (1904).
 5. *C. crypsizyga*, Meyrick, Exot. Microlep. Vol. 1, p. 278 (1914). Nyassaland. [Terr.
 6. *C. bifrenata*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 89 (1921). Transvaal, Tanganyika
 7. *C. bythota*, Meyrick, Exot. Microlep. Vol. 1, p. 576 (1916). Gold Coast.
 8. *C. homogramma*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 25 (1918). Natal.
 9. *C. furnaria*, Meyrick, ibidem, Vol. 3, p. 294 (1913). Transvaal.
 10. *C. cubiculata*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 273 (1911). Seychelles.

363. GENUS ORGANITIS, MEYRICK

Organitis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 151 (1906) — Type : *O. characopa*,
 Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ shortly ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 4 absent, 8 and 9 out of 7, 7 to apex, 11 from beyond middle. Hindwings 1, trapezoidal, termen hardly sinuate, cilia 3/5; 4 absent, 3 and 5 connate or stalked, 6 and 7 stalked.

Remarks. — Correlated with *Prosodarma*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *O. characopa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 151 (1906). Ceylon.
 2. *O. lubrica*, Meyrick, ibidem, Vol. 20, p. 460 (1910). India.

364. GENUS STELECHORIS, NOV. GEN.

Type : *S. exaema*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple or pubescent, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second or longer, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 3 and 4 out of 2, 5 approximated, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 3/5-4/5, 3 and 5 stalked, 5 nearly approximated at base, 6 and 7 stalked.

Remarks. — Derived from *Hygroplasta*.

Geographical distribution of species. — Indian, probably originating in Ceylon.

Larva unknown.

1. *S. cherandra*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 151 (1905). Ceylon.
 2. *S. exaema*, Meyrick, ibidem, Vol. 20, p. 707 (1911). Ceylon, S. India.

365. GENUS PROSODARMA, NOV. GEN.

Type : *P. fibularis*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ thick, simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint longer than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 4 and 5 connate, 8 and 9 out of 7. 7 to apex, 11 from beyond middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 1/2; 5 absent, 3 and 4 stalked, 6 and 7 stalked.

Remarks. — Correlated with the following.

Geographical distribution of species. — Malayan.

Larva unknown.

1. *P. fibularis*, Meyrick, Zool. Med. Leid. Vol. 6, p. 167 (1921).

Java, Celebes.

366. GENUS HYGROPLASTA, NOV. GEN.

Type : *H. spoliatella*, Walker.

Characters. — Head with appressed scales, sidetufts somewhat raised; ocelli posterior; tongue developed. Antennae 5/6, in ♂ fasciculate-ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint somewhat longer or shorter than second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 4 and 5 rather approximated at base, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings over 1, trapezoidal, termen faintly sinuate, cilia 2/5; 3 and 4 connate or very short-stalked, 5 much nearer 4 than 6, nearly parallel or slightly approximated at base, 6 and 7 stalked.

Remarks. — Allied to *Torodora*.

Geographical distribution of species. — Indian.

Larva (*spoliatella*) densely clothed with hairs so that the segments are not distinguishable, and larva looks like a mass of hairs (Fletcher); feeding on dry dead leaves.

- 1. *H. continetella*, Walker, List Lep. Het Brit. Mus. Vol. 29, p. 565 (1864). S. India.
- 2. *H. spoliatella*, Walker, ibidem, Vol. 29, p. 659 (1864). Ceylon, Kanara.
diluticornis, Walsingham, Moore's Lep. Ceyl. Vol. 3, p. 519, pl. 209, f. 7 (1886).
- 3. *A. monodryas*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 773 S. India.
(1914).
- 4. *H. lygaea*, Meyrick, ibidem, Vol. 20, p. 707 (1911). Kashmir.

367. GENUS TORODORA, MEYRICK

Torodora, Meyrick, Trans. Ent. Soc. Lond. p. 16 (1894). — Type : *T. characteris*, Meyrick.

Characters. — Head with appressed scales, sidetufts somewhat spreading; ocelli posterior; tongue developed. Antennae 5/6, in ♂ serrate, moderately ciliated, basal joint elongate, without pecten.

Labial palpi very long, recurved, second joint much thickened with appressed scales; terminal joint rather longer than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 8 and 9 out of 7, 7 to apex, 11 from middle. Hindwings over 1, trapezoidal, termen sinuate, cilia 1/2-2/3; 3 and 4 connate, 5 much nearer 4 than 6, slightly approximated at base, 6 and 7 stalked, 7 to costa.

Remarks. — Perhaps a development of *Brachmia*.

Geographical distribution of species. — Indian.

Larva unknown.

- | | |
|--|-----------|
| 1. <i>T. characteris</i> , Meyrick, Trans. Ent. Soc. Lond. p. 16 (1894). | Burma. |
| 2. <i>T. spilotella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 746 (1864). | Ceylon. |
| <i>tripustulata</i> , Walsingham, Moore's Lep. Ceyl. Vol. 3, p. 520, pl. 209, f. 8 (1886). | |
| 3. <i>T. typhlopis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 715 (1911). — Pl. 4, Fig. 100. | S. India. |
| 4. <i>T. artiasta</i> , Meyrick, ibidem, Vol. 20, p. 715 (1911). | S. India. |
| 5. <i>T. syrphetodes</i> , Meyrick, ibidem, Vol. 17, p. 152 (1906). | Ceylon. |

368. GENUS ERYTHRIASTIS, NOV. GEN.

Type : *E. rubentula*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint nearly as long, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings under 1, trapezoidal, termen sinuate, cilia 1 1/2; 3 and 4 connate, 5 curved, connate or closely approximated at base to 4, 6 and 7 stalked.

Remarks. — Derived from *Pachnistis*.

Geographical distribution of species. — South American.

Larva unknown.

- | | |
|---|-----------------|
| 1. <i>E. rubentula</i> , Meyrick, Trans. Ent. Soc. Lond. p. 273 (1914). | Guiana. |
| 2. <i>E. rhodocrossa</i> , Meyrick, ibidem, p. 273 (1914). | Guiana, Brazil. |

369. GENUS PACHNISTIS, MEYRICK

Pachnistis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 737 (1907). — Type : *P. cephalochra*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second or nearly, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings 1, trapezoidal, termen somewhat sinuate, cilia 1/3-3/5; 3 and 4 connate or stalked, 5 nearly parallel, not much nearer 4 than 6, 6 and 7 connate or stalked.

Remarks. — Presumably derived from *Brachmia*.

Geographical distribution of species. — Indo-Malayan and African.

Larva unknown.

1. <i>P. autophanta</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 89 (1921).	Rhodesia.
2. <i>P. microphanta</i> , Meyrick, ibidem, Vol. 8, p. 89 (1921).	Rhodesia.
3. <i>P. consors</i> , Meyrick, ibidem, Vol. 8, p. 89 (1921).	Cape Colony.
4. <i>P. cephalochra</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 737 (1907).	Punjab.
5. <i>P. cremnobathra</i> , Meyrick, Zool. Med. Leid. Vol. 7, p. 86 (1922).	Java.
6. <i>P. inhomesta</i> , Meyrick, Exot. Microlep. Vol. 1, p. 579 (1916).	Kanara.
7. <i>P. morologa</i> , Meyrick, Bull. Mus. Hist. Nat. Paris, no 8, p. 564 (1923).	Angola.
8. <i>P. nubivaga</i> , Meyrick, Zool. Med. Leid. Vol. 6, p. 167 (1921).	Celebes.
9. <i>P. arenis</i> , Meyrick, Exot. Microlep. Vol. 1, p. 66 (1913).	Bengal.

370. GENUS PARALLACTIS, NOV. GEN.

Type : *P. plaeziodes*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with dense appressed scales, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 and 9 stalked, 7 to apex, 8 absent, 11 from middle. Hindwings 1, trapezoidal-ovate, cilia 3/5; 3 and 4 connate, 5 nearly parallel, 6 and 7 connate.

Remarks. — A derivative of *Brachmia*. In this family the termination of vein 7 of forewings may be costal, apical, or terminal, but in no instance (except *Amphigenes*) can the termination of vein 8 be shown to be anything but costal; hence we must accept the view that the absent vein in the present genus is 8.

Geographical distribution of species. — African.

Larva unknown.

1. <i>P. panchlora</i> , Meyrick, Ann. Transv. Mus. Vol. 2, p. 232 (1911).	Transvaal.
2. <i>P. mitigata</i> , Meyrick, Exot. Microlep. Vol. 1, p. 278 (1914).	Nyassaland, Kenya
3. <i>P. periochra</i> , Meyrick, ibidem, Vol. 1, p. 577 (1916).	Nyassaland. [Colony.]
4. <i>P. plaeziodes</i> , Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 78 (1920). derogata, Meyrick, Ann. Transv. Mus. Vol. 8, p. 91 (1921).	Tanganyika Terr., Port. E. Africa.
5. <i>P. fuitima</i> , Meyrick, ibidem, Vol. 8, p. 88 (1921).	Rhodesia.

371. GENUS DEROXENA, MEYRICK

Deroxena, Meyrick, Exot. Microlep. Vol. 1, p. 153 (1913). — Type : *D. venosulella*, Möschler.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ moderately ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, terminal joint as long as second, somewhat thickened with scales and slightly roughened anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 11 from middle. Hindwings somewhat over 1, oblong-ovate, cilia 1/2; 3 and 4 connate, 5 slightly approximated, 6 and 7 closely approximated towards base.

Remarks. — Of doubtful affinity; European authors used to include the species in *Depressaria*, which is wholly out of the question; it is certainly a rather primitive form of the present family, but somewhat peculiar.

Geographical distribution of species. — European.

Larva unknown.

1. *D. venosulella*, Möschler, Wien. Ent. Monatsschr. Vol. 6, p. 142, pl. 1, f. 15 S.E. Europe, Asia Minor. (1862).
neglectella, Lederer, ibidem, Vol. 7, p. 46, pl. 1, f. 12 (1863).

372. GENUS PHILARACHNIS, NOV. GEN.

Type : *P. xerophaga*, Meyrick.

Characters. — Head with loosely appressed scales, sidetufts rather raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, minutely ciliated, basal joint moderate, without pecten. Labial palpi long, recurved, second joint with appressed scales, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-haired above and beneath. Forewings with 1b furcate, 2 and 3 stalked, 8 and 9 out of 7, 7 to termen, 11 from middle. Hindwings hardly 1, trapezoidal, termen sinuate, cilia 4/5; 3 and 4 connate, 5 slightly approximated, 6 and 7 short-stalked.

Remarks. — Probably derived from *Apethistis*.

Geographical distribution of species. — Indian.

Larva living in the large web-nests of a gregarious spider (*Stegodyphus*), and feeding on the dry remains of insects captured, thus helping the spider by cleansing the fabric.

1. *P. xerophaga*, Meyrick, Ent. M. Mag. Vol. 50, p. 219 (1914). Orissa, Madras, Ceylon.

373. GENUS SYNDESMICA, TURNER

Syndesmica, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 150 (1919). — Type : *S. homogenes*, Turner.

Characters. — Head with appressed scales; tongue developed. Antennae 3/4, in ♂ simple, basal joint without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, slightly rough anteriorly, somewhat expanded at apex, terminal joint stout, acute. Forewings with 2 and 3 stalked, 8 and 9 out of 7, 7 to apex. Hindwings over 1, trapezoidal, apex pointed, termen sinuate; 3 and 4 connate, 5 parallel, 6 and 7 connate.

Remarks. — Not known to me.

Geographical distribution of species. — Australian.

Larva unknown.

1. *S. homogenes*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 150 (1919). Queensland.

374. GENUS BRACHMIA, HÜBNER

Brachmia, Hübner, Verz. bek. Schmett. p. 419 (1826). — Type : *B. dimidiella*, Schiffmüller.

Ceratophora, Heinemann, Schmett. Deutschl. (2), Vol. 2, p. 325 (1870) (praeocc.). — Type : *B. rufescens*, Haworth.

Cladodes, Heinemann, ibidem (2), Vol. 2, p. 330 (1870) (praeocc.). — Type : *B. dimidiella*, Schiffmüller.

Eudodaces, Snellen, Tijdschr. v. Ent. Vol. 32, p. 204 (1889). — Type : *B. dimidiella*, Schiffmüller.

Aulacomima, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 395 (1904). — Type : *B. trinervis*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ simple or ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second or somewhat longer or shorter, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b fuscate, 2 and 3 stalked, 7 and 8 stalked, 7 to apex, 9 sometimes out of 7, 11 from middle. Hindwings 1 or over 1, trapezoidal, termen sinuate, cilia 1/3-1: 3 and 4 connate or stalked, 5 somewhat approximated, 6 and 7 stalked, 7 to apex.

Remarks. — Probably derived from *Apethistis*. The stalking of vein 9 with 7 in the forewings cannot be used generically, as it occurs or is absent in closely allied species, but is apparently reliable specifically.

Geographical distribution of species. — Indo-Malayan, African, and European, with stragglers in America and Australia; absent from New Zealand.

Larva (11 known) feeding in rolled or spun leaves.

Foodplants Gramineae (5), Convolvulaceae (4), Compositae, Solanaceae.

- | | |
|--|-----------------------------|
| 1. ? <i>B. biareatella</i> , Erschoff, Fedtsch. Lep. Turkest. p. 102, pl. 6, f. 116 (1874). | Turkestan. |
| 2. <i>B. lineolella</i> , Zeller, Isis, p. 197 (1839). | E. Europe. |
| 3. <i>B. graphicodes</i> , Meyrick, Ann. Transv. Mus. Vol. 4, p. 194 (1914). | Natal. |
| 4. <i>B. malacogramma</i> , Meyrick, ibidem, Vol. 2, p. 14, pl. 5, f. 2 (1909). | Transvaal. |
| 5. <i>B. nesidias</i> , Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 273 (1911). | Seychelles. |
| 6. <i>B. arotraea</i> , Meyrick, Trans. Ent. Soc. Lond. p. 15 (1894). | India, Ceylon, Burma. |
| 7. <i>B. phryganitis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 722 (1911). — Pl. 5, Fig. 101. | Ceylon. [Tonkin, Java.] |
| 8. <i>B. radiosella</i> , Erschoff, Fedtsch. Lep. Turkest. p. 102, pl. 6, f. 115 (1874). | Turkestan. [Australia.] |
| 9. <i>B. trinervis</i> , Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 395 (1904). | New South Wales, W. |
| 10. <i>B. modicella</i> , Christoph, Bull. Soc. Nat. Mosc. p. 28 (1882). | E. Siberia, Bengal. |
| 11. <i>B. impunctella</i> , Caradja, Iris, Vol. 34, p. 111 (1920). | E. Siberia. |
| 12. <i>B. triannulella</i> , Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 201, f. 458 (1855). | C. & S. Europe, Asia Minor. |
| sepiella, Steudel, Stett. Ent. Zeit. Vol. 27, p. 312 (1866). | |
| 13. ? <i>B. Hedemanni</i> , Caradja, Iris, Vol. 34, p. 112 (1920). | Punjab. |
| 14. <i>B. xerastis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 599 (1905). | Punjab. |
| 15. <i>B. triophthalma</i> , Meyrick, Rec. Ind. Mus. Vol. 5, p. 220 (1910). | Travancore. |
| 16. <i>B. hystricella</i> , Braun, Ent. News Philad. Vol. 32, p. 11 (1921). | Ohio. |

17. *B. subalbella*, Chambers, Canad. Ent. Vol. 6, p. 242 (1894) (-*albusella*). United States.
parvipulvella, Chambers, ibidem, Vol. 6, p. 242 (1894).
inaequipulvella, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 239 (1875).
chambersella, Murtfeldt, Canad. Ent. Vol. 13, p. 242 (1881).
18. *B. lyrella*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 101, pl. 3, f. 19 (1911). Guatemala.
19. *B. lutatella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 201, f. 467 (1855). C. Europe.
20. *B. engrapta*, Meyrick, Exot. Microlep. Vol. 2, p. 114 (1918). India.
21. *B. inornatella*, Douglas, Trans. Ent. Soc. Lond. (2), Vol. 1, p. 65 (1850). England, Holland, Ger-
22. *B. badia*, Braun, Ent. News Philad. Vol. 32, p. 12 (1921). California. [many.]
23. *B. robustella*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 60, p. (28), (1910). Herzegovina.
24. *B. rufescens*, Haworth, Lep. Brit. p. 555 (1828). C. & S. Europe.
- terrella*, Fischer von Rösterstamm, Abbild. Schmett. pl. 96, f. a-k (1844).
diaphanella, Zeller, Isis, p. 285 (1846).
isabella, Zeller, Stett. Ent. Zeit. Vol. 11, p. 151 (1850).
25. *B. sigillatrix*, Meyrick, Rec. Ind. Mus. Vol. 5, p. 222 (1910). Kanara.
26. *B. dimidiella*, Schiffermüller, Syst. Verz. Schmett. Wien, p. 141 (1776). Europe, Turkestan.
- costiguttella*, Zeller, Isis, p. 290 (1846).
Kneri, Nowicki, Enum. Lep. Hal. p. 191 (1860).
27. *B. procursella*, Rebel, Verh. Zool.-bot. Ges. Wien, Vol. 53, p. 97 (1903) Austria.
28. *B. gerronella*, Zeller, Stett. Ent. Zeit. Vol. 11, p. 155 (1850). C. & S. Europe, Palestine.
29. *B. hapalyntis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 724 (1911). India, Ceylon, Queensl.
30. *B. idiastris*, Meyrick, Exot. Microlep. Vol. 1, p. 577 (1916). India
31. *B. simplex*, Walsingham, Bull. Liverp. Mus. Vol. 3, p. 2 (1900). Sokotra.
32. *B. obfuscata*, Meyrick, Exot. Microlep. Vol. 2, p. 436 (1921). Queensland.
33. *B. tristella*, Snellen, Tijdschr. v. Ent. Vol. 44, p. 85, pl. 6, f. 2 (1901). Java.
34. *B. aruritis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 723 (1911). Ceylon.
35. *B. albicincta*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 90 (1921). Cape Colony.
36. *B. fiscinata*, Meyrick, ibidem, Vol. 6, p. 26 (1918). Zululand.
37. *B. craticula*, Meyrick, ibidem, Vol. 8, p. 90 (1921). Port. E. Africa.
38. *B. hemiopa*, Meyrick, ibidem, Vol. 8, p. 90 (1921). Rhodesia. [tia.]
39. *B. rasilella*, Herrich-Schäffer, Schmett. Eur. Vol. 5, p. 202, f. 459 (1855). Austria, Hungary, Croa-
40. *B. verberata*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 68 (1912). Transvaal, Natal.
41. *B. velitaris*, Meyrick, ibidem, Vol. 3, p. 295 (1913). Transvaal.
42. *B. lochistis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 723 (1911). Ceylon, Coorg.
43. *B. apricata*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 295 (1913). Transvaal.
44. *B. ochrobyrsa*, Meyrick, ibidem, Vol. 8, p. 92 (1921). Rhodesia.
45. *B. ochrostoma*, Meyrick, ibidem, Vol. 3, p. 296 (1913). Transvaal.
46. *B. neurograpta*, Meyrick, ibidem, Vol. 8, p. 91 (1921). Rhodesia.
47. *B. musicopa*, Meyrick, Proc. Zool. Soc. Lond. p. 727 (1908). Transvaal.
48. *B. philomusa*, Meyrick, Exot. Microlep. Vol. 2, p. 114 (1918). Ceylon, Bengal.
49. *B. ceramochroa*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 150 (1919). Queensland.
50. *B. octophora*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 25 (1918). Natal.
51. *B. torva*, Meyrick, Exot. Microlep. Vol. 1, p. 278 (1914). Nyassaland.
52. *B. pantheropa*, Meyrick, Ann. Transv. Mus. Vol. 3, p. 296 (1913). Transvaal, Natal.
53. *B. cricopa*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 274 (1911). Seychelles.
54. *B. discoannulella*, Chambers, Cinc. Quart. Journ. Sc. Vol. 2, p. 254 (1875) Texas.
(-*anu*-).
55. *B. emigrans*, Meyrick, Exot. Microlep. Vol. 2, p. 435 (1921). Barbados.
56. *B. trichocyma*, Meyrick, ibidem, Vol. 3, p. 47 (1923). Guiana, Brazil.
57. *B. anisopa*, Meyrick, ibidem, Vol. 2, p. 140 (1918). Colombia.
58. *B. cerinura*, Meyrick, ibidem, Vol. 3, p. 47 (1923). Brazil.
59. *B. convolvuli*, Walsingham, Proc. Zool. Soc. Lond. p. 944, pl. 51, f. 16 (1907). India, Java, S. Africa,
crypsilychna, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 773 (1914). Canaries, Comoros.
dryadopa, Meyrick, Ann. Transv. Mus. Vol. 6, p. 25 (1918).
efferia, Meyrick, Exot. Microlep. Vol. 2, p. 104 (1918).

60. *B. microsema*, Meyrick, Trans. Linn. Soc. Lond. Vol. 14, p. 274 (1911). Seychelles.
 61. *B. pullifimbriella*, Clemens, Proc. Ent. Soc. Philad. Vol. 2, p. 120 (1863). Canada, Maine.
 62. *B. legalis*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 91 (1911). Port. E. Africa.
 63. *B. amphisticta*, Meyrick, Exot. Microlep. Vol. 1, p. 279 (1914). Port. E. Africa.
 64. *B. ternatella*, Staudinger, Stett. Ent. Zeit. Vol. 20, p. 240 (1859). Spain.
 65. *B. strigosa*, Meyrick, Trans. Ent. Soc. Lond. p. 450 (1910). Borneo.
 66. *B. craterospila*, Meyrick, Exot. Microlep. Vol. 3, p. 46 (1923). Assam.
 67. *B. autonoma*, Meyrick, Trans. Ent. Soc. Lond. p. 369 (1910). Bengal, Chagos Islands.
 68. *B. ptochodryas*, Meyrick, Exot. Microlep. Vol. 3, p. 46 (1923). Assam.
 69. *B. resoluta*, Meyrick, ibidem, Vol. 2, p. 113 (1918). Bengal.
 70. *B. melicephala*, Meyrick, ibidem, Vol. 2, p. 114 (1918). Burma.
 71. *B. insuavis*, Meyrick, Suppl. Ent. p. 51 (1914). Formosa.
 72. *B. nephelopis*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 91 (1911). Rhodesia.
 73. *B. circumfusa*, Meyrick, Exot. Microlep. Vol. 2, p. 506 (1922). French Guinea.
 74. *B. sterictis*, Meyrick, Proc. Zool. Soc. Lond. p. 727 (1908). Transvaal, Natal, Port.
 75. *B. ochyrota*, Meyrick, Ann. S. Afr. Mus. Vol. 17, p. 285 (1920). Natal. [E. Africa.
 76. *B. liberta*, Meyrick, Exot. Microlep. Vol. 3, p. 1 (1925). Madagascar.
 77. *B. crateropsis*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 91 (1911). Rhodesia.
 78. *B. serialis*, Meyrick, Proc. Zool. Soc. Lond. p. 727 (1908). Transvaal.
 79. *B. inconspicua*, Walsingham, Trans. Ent. Soc. Lond. p. 103, pl. 5, f. 38 (1891). Gambia.
 80. *B. marginata*, Walsingham, ibidem, p. 99, pl. 4, f. 35 (1891). Gambia.
 81. *B. tepidata*, Meyrick, Exot. Microlep. Vol. 2, p. 505 (1922). China.
 82. *B. deltopis*, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 79 (1920). Kenya Colony.
 83. *B. insulsa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 22, p. 774 (1914). India.
 84. *B. perumbrata*, Meyrick, Exot. Microlep. Vol. 2, p. 113 (1918). Bengal.
 85. *B. juridiva*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 716 (1911). Ceylon.
 86. *B. obtrectata*, Meyrick, Exot. Microlep. Vol. 2, p. 506 (1922). China.
 87. *B. vecors*, Meyrick, ibidem, Vol. 2, p. 112 (1918). S. India, Tonkin, China.
 88. *B. japonicella*, Zeller, Hor. Soc. Ent. Ross. Vol. 13, p. 365, pl. 5, f. 124 (1877). Japan.
 89. *B. virescens*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 84, pl. 3, f. 5 (1911). Mexico.
 90. *B. carpoides*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 459 (1908). Assam.
 91. *B. syntonopis*, Meyrick, Exot. Microlep. Vol. 3, p. 48 (1923). Bombay.
 92. *B. antichroa*, Meyrick, ibidem, Vol. 2, p. 156 (1918). Ceylon.
 93. *B. custos*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 725 (1911). S. India.

375. GENUS APETHISTIS, MEYRICK

Apethistis, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 459 (1908). — Type : *A. metocca*, Meyrick.

Characters. — Head with appressed scales, sidetuffs somewhat raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, shortly ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint as long as second or nearly, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 and 8 stalked, 7 to termen, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 3/4-1; 3 and 4 connate, 5 rather approximated, 6 and 7 stalked.

Remarks. — Derivable from a form approaching *Protobathra*. Obscure insects of similar appearance.

Geographical distribution of species. — Indian.

Larva unknown.

1. *A. dolosa*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 724 (1911). Ceylon.
2. *A. alienella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 634 (1864). Ceylon.
3. *A. metoeca*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 460 (1908). Ceylon.
4. *A. officiosa*, Meyrick, Exot. Microlep. Vol. 2, p. 113 (1918). Madras.
5. *A. sitiens*, Meyrick, ibidem, Vol. 2, p. 114 (1918). S. India.
6. *A. cenchritis*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 721 (1911). Assam.
7. *A. inspersa*, Meyrick, Exot. Microlep. Vol. 2, p. 436 (1921). Assam.
8. *A. consummata*, Meyrick, ibidem, Vol. 3, p. 47 (1923). Assam.
9. *A. epistica*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 16, p. 599 (1905). Ceylon.

Pl. 5, Fig. 102.

376. GENUS LACISTODES, MEYRICK

Lacistodes, Meyrick, Ann. Transv. Mus. Vol. 8, p. 92 (1921). — Type : *L. tauropis*, Meyrick.

Characters. — Head smooth, sidetufts loosely raised; ocelli posterior; tongue developed. Antennae 3/4, in ♂ minutely ciliated, basal joint elongate, without pecten. Labial palpi long, recurved, second joint much thickened with appressed scales, laterally compressed, somewhat rough beneath, terminal joint as long as second, moderate, acute. Maxillary palpi short, filiform, porrected. Posterior tibiae clothed with long rough hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings slightly over 1, trapezoidal, apex obtuse, termen faintly sinuate, cilia 1/2; 3 and 4 connate, 5 rather approximated to 4, 6 and 7 closely approximated at base.

Remarks. — Apparently allied to *Brachmia*.

Geographical distribution of species. — African.

Larva unknown.

1. *L. tauropis*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 92 (1921). Rhodesia.

377. GENUS ENCOLPOTIS, MEYRICK

Encolpotis, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 352 (1909). — Type : *E. xanthoria*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderately elongate, without pecten. Labial palpi long, recurved, thickened with appressed scales, terminal joint as long as second, slender or rather stout, acute. Maxillary palpi very short, drooping, filiform. Posterior tibiae clothed with long hairs above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 7 to termen, 9 sometimes out of 7, 11 from middle. Hindwings 1, trapezoidal, apex produced, termen sinuate, cilia 1; 3 and 4 connate or short-stalked, 5 rather approximated, 6 and 7 stalked.

Remarks. — An early form of this group.

Geographical distribution of species. — African and Indian.

Larva (*xanthoria*) feeding on colonies of a scale-insect, *Icerya* (*Coccidae*).

1. *E. heliopepta*, Meyrick, Exot. Microlep. Vol. 2, p. 153 (1918). Assam.
2. *E. scioplasta*, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 79 (1920). Kenya Colony.
3. *E. xanthoria*, Meyrick, Ann. S. Afr. Mus. Vol. 5, p. 352 (1909). — **Pl. 5**, Nyassaland, Transvaal, Natal, Cape Colony
Fig. 114.

Group 9 (Autosticha type)

In this group are included a small number of early genera characterised by the absence of vein 7 and stalking of veins 2 and 3 of forewings, together with the remnants of those ancestral forms from which they and the rest of the family appear to have been derived, approaching the *Oecophorid* type. The cubital pecten is absent in the hindwings. The larvae appear to have the primitive habit of feeding on vegetable refuse, lichens, etc. Geographically the group appears to be primarily Indian, yet with a representative branch in America, and small colonies in Africa and Australia.

378. GENUS ISCHNODORIS, MEYRICK

Ischnodoris, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 726 (1911). — Type : *I. sigalota*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely spreading; ocelli posterior; tongue developed. Antennae 4/5, in ♂ shortly ciliated, basal joint moderately elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked, 7 absent, 6 and 8 stalked, 11 from middle. Hindwings 1, elongate-trapezoidal, apex pointed, termen hardly sinuate, cilia 1 1/4; 3 and 4 connate, 5 somewhat approximated, 6 and 7 stalked.

Remarks. — Derivable from *Autosticha*.

Geographical distribution of species. — Indian.

Larva unknown.

1. *I. sigalota*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 726 (1911). Ceylon.

379. GENUS SCEPTEA, WALSINGHAM

Sceptea, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 108 (1911). — Type : *S. decedens*, Walsingham.

Characters. — Head with appressed scales; tongue developed. Antennae 3/4, in ♂ simple, basal joint without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint nearly as long as second, moderate, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae rough-scaled above. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 6 and 8 stalked, 11 from middle. Hindwings over 1, trapezoidal-ovate, termen slightly sinuate; 3 and 4 connate or short-stalked, 5 parallel, 6 and 7 stalked.

Remarks. — Probably derived from *Glyphidocera*.

Geographical distribution of species. — North American.

Larva unknown.

1. *S. aberratella*, Busck, Journ. N. York Ent. Soc. Vol. 15, p. 138 (1907). Maryland.

2. *S. decedens*, Walsingham, Biol. Centr. Amer. Lep. Het. Vol. 4, p. 109, Mexico.
pl. 3, f. 30 (1911).

380. GENUS PTILOSTONYCHIA, WALSINGHAM

Ptilostonychia, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 109 (1911). — Type: *P. plicata*, Walsingham.

Characters. — Head with appressed scales; tongue developed. Antennae 4/5, in ♂ with notch above basal joint. Labial palpi very long, recurved, second joint tufted with long rough projecting hair-scales, terminal joint shorter than second, also with rough projecting scales, less marked in ♀. Maxillary palpi very short. Posterior tibiae somewhat rough-scaled above. Forewings with 2 and 3 stalked, 7 absent. Hindwings over 1, trapezoidal-ovate; 3 and 4 stalked, 5 approximated, 6 and 7 stalked.

Remarks. — A development of *Glyphidocera*.

Geographical distribution of species. — Central American.

Larva unknown.

1. *P. plicata*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 109, pl. 3, Panama.
f. 31 (1911).

381. GENUS GLYPHIDOCERA, WALSINGHAM

Glyphidocera, Walsingham, Proc. Zool. Soc. Lond. p. 531 (1891). — Type: *G. audax*, Walsingham.

Harpagandra, Meyrick, Exot. Microlep. Vol. 2, p. 210 (1918). — Type: *H. cryphiodes*, Meyrick.

Characters. — Head with dense appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ with notch and scale-projection at base of stalk, basal joint elongate, sometimes thickened, without pecten. Labial palpi very long, recurved, second joint thickened with appressed scales, terminal joint somewhat shorter, moderate or stout, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough scales. Forewings with 1b furcate, 2 and 3 stalked, 4 sometimes out of 2, 7 absent, 8 and 9 sometimes stalked, 11 from middle. Hindwings over 1, trapezoidal-ovate, cilia 1/4-1/3; 3 and 4 connate or short-stalked, 5 nearly parallel, 6 and 7 stalked.

Remarks. — This may be regarded as the American representative of *Autosticha*.

Geographical distribution of species. — American.

Larva unknown, but probably refuse-feeders.

- | | |
|--|--|
| 1. <i>G. lactiflorella</i> , Chambers, Bull. U. S. Geol. Surv. Vol. 4, p. 89 (1878). Texas.
(-osella). | |
| 2. <i>G. hypochloea</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 102, Mexico.
pl. 3, f. 23 (1911). | |
| 3. <i>G. dimorphella</i> , Busck, Journ. N. York Ent. Soc. Vol. 15, p. 136 (1907). Maryland. | |
| 4. <i>G. meyrickella</i> , Busck, ibidem, Vol. 15, p. 138 (1907). Maryland. | |
| 5. <i>G. speratella</i> , Busck, Proc. Ent. Soc. Wash. Vol. 9, p. 88 (1908). Pennsylvania. | |
| 6. <i>G. septentrionella</i> , Busck, Proc. U. S. Mus. Vol. 28, p. 762 (1904). Brit. Columbia. | |
| 7. <i>G. aequipulvella</i> , Chambers, Canad. Ent. Vol. 4, p. 192 (1872) (<i>aeque-</i>). Texas, Kentucky, California. | |
| 8. <i>G. rhypara</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 111 (1911). Mexico. [nia, Colorado. | |
| 9. <i>G. elpista</i> , Walsingham, ibidem, Vol. 4, p. 110 (1911). Panama. | |
| 10. <i>G. cinctalis</i> , Meyrick, Exot. Microlep. Vol. 3, p. 49 (1923). Ecuador. | |
| 11. <i>G. percnoleuca</i> , Meyrick, ibidem, Vol. 3, p. 48 (1923). Brazil. | |
| 12. <i>G. reparabilis</i> , Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 112 (1911). Panama. | |
| 13. <i>G. psammolitha</i> , Meyrick, Exot. Microlep. Vol. 3, p. 48 (1923). Brazil. | |

14. *G. vestita*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 112 (1911). Panama.
 15. *G. orthocleis*, Meyrick, Exot. Microlep. Vol. 3, p. 49 (1923). Brazil.
 16. *G. crocogramma*, Meyrick, ibidem, Vol. 3, p. 49 (1923). Brazil.
 17. *G. umbrata*, Walsingham, Biol.-Centr. Amer. Lep. Het. Vol. 4, p. 111, pl. 3, f. 34 (1911). Guatemala.
 18. *G. exsiccata*, Meyrick, Trans. Ent. Soc. Lond. p. 273 (1914). Guiana.
 19. *G. stenomorpha*, Meyrick, Exot. Microlep. Vol. 3, p. 49 (1923). Guiana.
 20. *G. floridanella*, Busck, Proc. Ent. Soc. Wash. Vol. 2, p. 474 (1901). Florida.
 21. *G. perobscura*, Walsingham, Biol. Centr.-Amer. Lep. Het. Vol. 4, p. 112 pl. 3, f. 35 (1911). Mexico.
 22. *G. saline*, Walsingham, ibidem, Vol. 4, p. 110, pl. 3, f. 33 (1911). Mexico.
 23. *G. recticostella*, Walsingham, Proc. Zool. Soc. Lond. p. 97 (1897). Grenada.
 24. *G. dominicella*, Walsingham, ibidem, p. 97 (1897). Dominica.
 25. *G. audax*, Walsingham, ibidem, p. 531 (1891). St Vincent.
 26. *G. caribea*, Busck, Bull. Trinid. Dep. Agric. Vol. 9, p. 244 (1910) (*caribea*). Trinidad.
 27. *G. inurbana*, Meyrick, Trans. Ent. Soc. Lond. p. 273 (1914). Guiana, Brazil.
 28. *G. cryphoides*, Meyrick, Exot. Microlep. Vol. 2, p. 210 (1918). Guiana, Brazil.

382. GENUS ANAPTILORA, MEYRICK

Anaptilora, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 390 (1904). — Type: *A. isocosma*, Meyrick

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ serrulate, simple, basal joint elongate, without pecten. Labial palpi long, recurved, second joint thickened with dense appressed scales, projecting towards apex above, terminal joint in ♂ rather shorter, thickened with dense scales, with long loosely expanded hair-scales posteriorly, in ♀ as long as second, with appressed scales, pointed. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough scales above. Forewings with 1b furcate, 2 and 3 connate or stalked, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings 1, oblong-trapezoidal, termen slightly sinuate, cilia 2/3; 3 and 4 connate, 5 tolerably parallel, 6 and 7 stalked.

Remarks. — A development of *Autosticha*. Although the other members of the genus all share the usual sombre and very commonplace colouring of the group, the type-species affords a remarkable instance of presumable mimicry, exactly imitating a conspicuously marked *Oecophorid* of totally different appearance, which occurs with it.

Geographical distribution of species. — Australian.

Larva unknown.

1. *A. isocosma*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 390 (1904). Queensland.
Pl. 4, Fig. 97.
 2. *A. basiphaea*, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 149 (1919). Queensland.
 3. *A. haplospila*, Turner, ibidem, Vol. 31, p. 149 (1919). Queensland.
 4. *A. parasira*, Meyrick, Exot. Microlep. Vol. 1, p. 591 (1916). North Australia.
 5. *A. ephelotis*, Meyrick, ibidem, Vol. 1, p. 591 (1916). North Australia.
 6. *A. eremias*, Meyrick, Proc. Linn. Soc. N. S. Wales, Vol. 29, p. 391 (1904). Queensl., New Guinea.
 7. *A. homoclera*, Meyrick, Exot. Microlep. Vol. 1, p. 591 (1916). North Australia.

383. GENUS STOEBERHINUS, BUTLER

Stoeberhinus, Butler, Ann. Mag. Nat. Hist. (5), Vol. 7, p. 402 (1881). — Type: *S. testacea*, Butler.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint elongate, without pecten. Labial palpi in ♂ very long, strongly recurved, basal

joint very long, nearly reaching crown, second joint strongly reflexed, smooth, terminal joint shorter, with posterior brush of very long expansible hairscales; in ♀ long, smooth, recurved, basal joint normal, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, drooping. Posterior tibiae with appressed scales. Forewings with 1^b furcate, 2 and 3 stalked, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 2/3; 3 and 4 connate or stalked, 5 nearly parallel, 6 and 7 closely approximated or stalked.

Remarks. — Correlated with the preceding.

Geographical distribution of species. — Pacific Ocean, probably spread artificially.

Larva feeding on dry vegetable refuse.

1. *S. testacea*, Butler, Ann. Mag. Nat. Hist. Vol. (5), 7, p. 402 (1881). Fiji, Marquesas, Hawaii.
demias, Meyrick, Trans. Ent. Soc. Lond. p. 281 (1886) (♀).

384. GENUS SYRMADAULA, MEYRICK

Syrmadaula, Meyrick, Ann. Transv. Mus. Vol. 6, p. 26 (1918). — Type: *S. automorpha*, Meyrick.

Characters. — Head with appressed scales; ocelli small, posterior; tongue developed. Antennae 3/4, in ♂ very shortly ciliated, basal joint rather elongate, without pecten. Labial palpi long, recurved, second joint thickened with appressed scales, somewhat rough beneath, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairscales above. Forewings with 2 and 3 short-stalked from angle, 4 and 5 somewhat approximated, 7 absent, 11 from middle. Hindwings somewhat under 1, elongate-trapezoidal, termen beneath apex very oblique, slightly sinuate, cilia 1; 3-5 equidistant, remote, 6 and 7 stalked.

Remarks. — A development of *Autosticha*.

Geographical distribution of species. — African.

Larva unknown.

1. *S. automorpha*, Meyrick, Ann. Transv. Mus. Vol. 6, p. 26 (1918). Transvaal, Rhodesia.

385. GENUS ULIARIA, DUMONT

Uliaria, Dumont, Bull. Soc. Ent. Fr. p. 329 (1920). — Type: *U. insulella*, Dumont.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ minutely ciliated. Labial palpi long, recurved, second joint thickened with appressed scales beneath, with rough projecting scales above, terminal joint nearly as long as second, moderate, acute. Posterior tibiae rough-haired above. Forewings with 1^b furcate, 2 and 3 stalked from angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate; 3 and 4 nearly approximated at base, 5 parallel, 6 and 7 nearly approximated at base.

Remarks. — Apparently a derivative of *Autosticha*.

Geographical distribution of species. — Western Europe.

Larva unknown.

1. *U. insulella*, Dumont, Bull. Soc. Ent. Fr. p. 330 (1920). France.

386. GENUS AUTOSTICHA, MEYRICK

Autosticha, Meyrick, Trans. Ent. Soc. Lond. p. 281 (1886). — Type : *A. pelodes*, Meyrick.

Automola, Meyrick, Ent. M. Mag. Vol. 20, p. 34 (1883) (praeocc.). — Type : *A. pelodes*, Meyrick.

Epicharma, Walsingham, Trans. Ent. Soc. Lond. p. 38 (1897). — Type : *A. nothriiformis*, Walsingham.

Epicoenia, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 140 (1906). — Type : *A. chernetis*, Meyrick.

Prosomura, Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 147 (1919). — Type : *A. symmetra*, Turner.

Characters. — Head with appressed scales, sidetufts somewhat raised; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, simple or shortly ciliated or rarely with whorls of long cilia, basal joint moderate, without pecten. Labial palpi long, recurved, second joint rather thickened with appressed scales, sometimes somewhat dilated towards apex, terminal joint as long as second or nearly, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with rough scales. Forewings with 1b furcate, 2 and 3 stalked from angle, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen sinuate, cilia 4/5-1; 3 and 4 connate, 5 approximated, 6 and 7 stalked or approximated.

Remarks. — Correlated with *Apethistis*. The species are often obscure and similar, and require careful attention.

Geographical distribution of species. — Primarily Indo-Malayan, and chiefly characteristic of Ceylon (which should be its place of origin) but extending into Africa, and with stragglers in Australia and Fiji; the occurrence of the type-species in Hawaii is probably due to artificial introduction, and this may be the case with the Fijian species also.

Larva (6 species known) feeding on moss, lichens, and vegetable refuse.

- | | |
|---|------------------|
| 1. <i>A. chlorodelta</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 140 (1906). | Ceylon. |
| 2. <i>A. auxodelta</i> , Meyrick, Exot. Microlep. Vol. 1, p. 585 (1916). | Assam. |
| 3. <i>A. exemplaris</i> , Meyrick, ibidem, Vol. 1, p. 586 (1916). | S. India. |
| 4. <i>A. iterata</i> , Meyrick, ibidem, Vol. 1, p. 585 (1916). | Ceylon. |
| 5. <i>A. vicularis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 20, p. 725 (1911). | Ceylon. |
| 6. <i>A. conciliata</i> , Meyrick, Exot. Microlep. Vol. 2, p. 154 (1918). | Madras. |
| 7. <i>A. relaxata</i> , Meyrick, ibidem, Vol. 1, p. 586 (1916). | Ceylon. |
| 8. <i>A. authaema</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 141 (1906). | Ceylon. |
| 9. <i>A. flavescens</i> , Meyrick, Exot. Microlep. Vol. 1, p. 557 (1916). | Ceylon. |
| 10. <i>A. petrotoma</i> , Meyrick, ibidem, Vol. 1, p. 587 (1916). | Ceylon. |
| 11. <i>A. acharacta</i> , Meyrick, ibidem, Vol. 2, p. 153 (1918). | N. W. India. |
| 12. <i>A. encycota</i> , Meyrick, ibidem, Vol. 2, p. 506 (1922). | Madras. |
| 13. <i>A. cheruetis</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 141 (1906). | Ceylon. |
| 14. <i>A. phaulodes</i> , Meyrick, ibidem, Vol. 18, p. 459 (1908). | Ceylon. |
| 15. <i>A. emmetra</i> , Meyrick, Ann. Transv. Mus. Vol. 8, p. 93 (1921). | Rhodesia. |
| 16. <i>A. symmetra</i> , Turner, Proc. Roy. Soc. Queensl. Vol. 31, p. 148 (1919). | Queensland. |
| 17. <i>A. solita</i> , Meyrick, Exot. Microlep. Vol. 3, p. 45 (1923). | Fiji. |
| 18. <i>A. strenuella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 632 (1864). | Ceylon. |
| 19. <i>A. pelodes</i> , Meyrick, Ent. M. Mag. Vol. 20, p. 34 (1883). | Celebes, Hawaii. |
| 20. <i>A. demotica</i> , Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 458 (1908). | Ceylon, Celebes. |
| 21. <i>A. nothriiformis</i> , Walsingham, Trans. Ent. Soc. Lond. p. 39, pl. 2, f. 3 (1897). | French Congo. |
| 22. <i>A. stagmatopis</i> , Meyrick, Exot. Microlep. Vol. 3, p. 50 (1923). | Madras. |

23. *A. tetrapeda*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 458 S. India.
(1908).
 24. *A. calceata*, Meyrick, ibidem, Vol. 18, p. 456 (1908). Ceylon.
 25. *A. pelaea*, Meyrick, ibidem, Vol. 18, p. 456 (1908). Ceylon.
 26. *A. nothropis*, Meyrick, Ann. Transv. Mus. Vol. 8, p. 92 (1921). Rhodesia.
 27. *A. academica*, Meyrick, Zool. Med. Leid. Vol. 7, p. 87 (1922). Java.
 28. *A. demetrias*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 457 (1908). Ceylon.
 29. *A. xanthographa*, Meyrick, Exot. Microlep. Vol. 1, p. 588 (1916). Ceylon.
 30. *A. aspasta*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 457 (1908). Ceylon.

Pl. 5, Fig. 103.

31. *A. protypa*, Meyrick, ibidem, Vol. 18, p. 457 (1908). Ceylon.
 32. *A. thermopsis*, Meyrick, Exot. Microlep. Vol. 3, p. 50 (1923). Ceylon.
 33. *A. crocothicta*, Meyrick, ibidem, Vol. 1, p. 588 (1916). Ceylon.
 34. *A. affixella*, Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 632 (1864). Ceylon.
 35. *A. deductella*, Walker, ibidem, Vol. 29, p. 631 (1864). Ceylon.
 36. *A. euryterma*, Meyrick, Voyage Alluaud Jean. Lép. Vol. 2, p. 80 (1920). Kenya Colony.
 37. *A. spilochorda*, Meyrick, Exot. Microlep. Vol. 1, p. 588 (1916). S. India.
 38. *A. perixantha*, Meyrick, ibidem, Vol. 1, p. 202 (1914). Port. E. Africa.
 39. *A. aureolata*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 18, p. 455 (1908). Ceylon.
 40. *A. binaria*, Meyrick, ibidem, Vol. 18, p. 456 (1908). Ceylon.
 41. *A. nauylchna*, Meyrick, ibidem, Vol. 18, p. 456 (1908). — **Pl. 5, Fig. 104.** Ceylon.

387. GENUS DEMIOPHILA, MEYRICK

Demiophila, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 152 (1906). — Type : *D. psaphara*, Meyrick.

Characters. — Head with appressed scales; ocelli posterior; tongue developed. Antennae 4/5, in ♂ serrulate, minutely ciliated, basal joint moderately elongate, without pecten. Labial palpi long, recurved, second joint thickened with dense appressed scales, slightly rough beneath at apex, terminal joint as long as second, moderate, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with long rough hairs above. Forewings with 1b furcate, 2 from near angle, 3 and 4 stalked, 5 approximated, 7 absent, 9 approximated, 11 from middle. Hindwings over 1, trapezoidal-ovate, cilia 1/2; 3 and 4 connate or short-stalked, 5 tolerably parallel, 6 and 7 connate.

Remarks. — Probably derived from *Encrasima*.

Geographical distribution of species. — Ceylon.

Larva unknown.

1. *D. psaphara*, Meyrick, Journ. Bombay Nat. Hist. Soc. Vol. 17, p. 152 Ceylon.
(1906). — **Pl. 5, Fig. 105.**

388. GENUS PANICOTRICA, MEYRICK

Panicotricha, Meyrick, Ann. Transv. Mus. Vol. 3, p. 296 (1923). — Type : *P. prographa*, Meyrick.

Characters. — Head above with long rough spreading hairs, face with appressed scales; ocelli posterior; tongue short. Antennae 4/5, basal joint moderate, without pecten. Labial palpi very long, curved, ascending, slender, second joint with appressed scales, terminal joint almost as long as second, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough

hairs above. Forewings with 2 from near angle, curved, 2-5 approximated at base, 6 and 8 approximated at base, 6 and 8 approximated, 7 absent, 11 from middle. Hindwings 1, trapezoidal, termen slightly sinuate, cilia 1; 3 and 4 connate or stalked, 5 approximated, transverse vein very oblique, 6 and 7 stalked.

Remarks. — Also apparently a derivative of *Encrasima*; but the curious rough head is unique in the family.

Geographical distribution of species. — African.

Larva unknown.

- | | |
|---|------------|
| 1. <i>P. pographa</i> , Meyrick, Ann. Transv. Mus. Vol. 3, p. 296 (1913). | Transvaal. |
|---|------------|

389. GENUS ENCRASIMA, MEYRICK

Encrasima, Meyrick, Exot. Microlep. Vol. 1, p. 594 (1916). — Type : *E. reversa*, Meyrick.

Characters. — Head smooth; ocelli posterior; tongue developed. Antennae 3/4, in ♂ simple, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough hairs above. Forewings with 2 from towards angle, 3-5 approximated, 7 absent, 11 from middle. Hindwings somewhat over 1, trapezoidal-ovate, termen slightly sinuate, cilia 2/5; 3 and 4 connate, 5 slightly approximated, 6 and 7 connate.

Remarks. — Derived from *Protobathra*.

Geographical distribution of species. — Indian, extending to China and Madagascar.

Larva unknown.

- | | |
|--|-------------|
| 1. <i>E. insularis</i> , Butler, Ann. Mag. Nat. Hist. (5), Vol. 5, p. 394 (1880). | Madagascar. |
| 2. <i>E. xanthoclista</i> , Meyrick, Exot. Microlep. Vol. 3, p. 50 (1923). | Ceylon. |
| 3. <i>E. reversa</i> , Meyrick, ibidem, Vol. 1, p. 594 (1916). | Ceylon. |
| 4. <i>E. retractella</i> , Walker, List Lep. Het. Brit. Mus. Vol. 29, p. 630 (1864). | China. |
| 5. <i>E. elacopis</i> , Meyrick, Exot. Microlep. Vol. 1, p. 594 (1916). — Pl. 5, | Ceylon. |
- Fig. 106.**
6. *E. simpliciella*, Stainton, Trans. Ent. Soc. Lond. (2), Vol. 5, p. 118 (1858). — *E. communicata*, Meyrick, Exot. Microlep. Vol. 2, p. 154 (1918). — Bengal.

390. GENUS PROTOBATHRA, MEYRICK

Protobathra, Meyrick, Exot. Microlep. Vol. 1, p. 595 (1916). — Type : *P. crista*, Meyrick.

Characters. — Head with appressed scales, side tufts raised; ocelli posterior; tongue developed. Antennae 2/3, in ♂ simple or ciliated, basal joint elongate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint rather shorter than second, slender, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiae clothed with rough scales above. Forewings with 2 from towards angle, 3-5 approximated, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings somewhat over 1, trapezoidal-ovate, termen slightly sinuate, cilia 1/2; 3 and 4 short stalked, 5 rather approximated, 6 and 7 connate.

Remarks. — This genus is assumed to approach closely the primitive type of the family, being such that all the preceding genera can be with probability derived from it, and differing so little in all respects from *Cryptolechia* in the *Oecophoridae*, that it may naturally indicate the connection with one of the earliest forms of that family.

Geographical distribution of species. — Indian, extending to Japan.

Larva unknown.

1. *P. leucostola*, Meyrick, Exot. Microlep. Vol. 2; p. 436 (1921). — **Pl. 5**, Japan.

Fig. 107.

2. *P. coenotypa*, Meyrick, ibidem, Vol. 2, p. 155 (1918). Ceylon.

3. *P. erista*, Meyrick, ibidem, Vol. 1, p. 595 (1916). S. India.

391. GENUS AMPHIGENES, MEYRICK

Amphigenes, Meyrick, Exot. Microlep. Vol. 2, p. 436 (1921). — Type : *A. tartarea*, Meyrick.

Characters. — Head with appressed scales, sidetufts loosely spreading; ocelli posterior; tongue short, slender. Antennae $\frac{3}{4}$, in ♂ strongly ciliated, basal joint moderate, without pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint almost as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with rather rough scales above. Forewings with 1b furcate, 2 from towards angle, 7 and 8 stalked, 8 to apex, 11 from middle. Hindwings 1, trapezoidal-ovate, cilia $\frac{1}{4}$; 3 and 4 short-stalked, 5 hardly approximated, 6 and 7 closely approximated at base.

Remarks. — Closely allied to the preceding genus, with which it is apparently correlated, but differing from it and every other genus of the family by the apical termination of vein 8 of the forewings.

Geographical distribution of species. — Papuan.

Larva unknown.

1. *A. tartarea*, Meyrick, Exot. Microlep. Vol. 2, p. 437 (1921). New Guinea.



INDEX

(The names of species in *italics* are synonyms)

Pages	Pages	Pages			
abdominella, Busck	77	acrogypsa, Turn.	176	agathopa, Meyr.	179
abella, Busck	80	acroleuca, Meyr.	148	Agellarchis (genus), Meyr.	172
aberratella, Busck	252	acrolychna, Meyr.	175	agnatella, Walk.	242
abhaustella, Chrét.	141	acromelas, Turn.	171	Agnippe (genus), Chamb.	54
abietisella, Pack.	58	acronipha, Turn.	102	agraphella, Rag.	30
abjunctella, Walk.	81	acropasta, Turn.	24	Agriastis, Meyr.	125
abnormella, Herr.-Sch.	42	acropelta, Meyr.	115	agricola, Wals.	153
abolitella, Walk.	122	acrophanes, Meyr.	228	agrifolia, Braun	70
abortiva, Wals.	189	acrophylla, Meyr.	72	agrimoniella, Clem.	123
abradescens, Braun	84	acrosphales, Meyr.	239	ainella, Chrét.	44
abruptella, Walk.	121, 152	acrotoma, Meyr.	153	alacella, Clem.	130
abscessella, Walk.	178	acroxantha, Low.	2	alacella, Dup.	124
absconditella, Walk.	44	acroxantha, Low.	231	alaecilla, Meyr.	201
absoluta, Meyr.	91	actias, Meyr.	32	alaicella, Car.	107
acaciae, Wals.	154	actiella, Barnes	35	alampes, Turn.	238
acaciella, Busck	72, 105	actinis, Wals.	232	alaopis, Meyr.	113
academica, Meyr.	257	actinota, Meyr.	148	alaricella, Busck	90
Acanthephila (genus), Hein.	123	acuminata, Staud.	175	albicanelia, Zell.	202
acanthyllidis, Wals.	112	acuminata, Wals.	116	albicans, Hein.	81
acatharta, Meyr.	201	acuminatella, Sirc.	95	albiceps, Zell.	52
acaudella, Reb.	107	acupediella, Frey	81	albicincta, Meyr.	249
accensa, Meyr.	71	acutangulella, Hein.	42	albidella, Chrét.	21
accincta, Meyr.	24	acutella, Chrét.	23	albidella, Reb.	40, 202
accinctella, Walk.	122	adactella, Walk.	177	albidella, Snell.	164
acerata, Meyr.	153	adaequata, Meyr.	138	albidorsella, Snell.	58
acerella, Herr.-Sch.	96	adamantina, Meyr.	104	albifaciella, Hein.	82
acharacta, Meyr.	256	adelella, Walk.	237	albifemorella, Hofm.	77
acherusia, Meyr.	163	adelocentra, Meyr.	174	albiflora, Meyr.	77
achlyodes, Meyr.	176	Adeleomorpha (genus) Snell.	234	albifrons, Turn.	149
achnastis, Meyr.	230	Adrasteia, Chamb.	69	albifrontella, Hein.	96, 112
achnias, Meyr.	24	adumbratella, Snell.	63	albilingua, Wals.	67
Achoria (genus), Meyr.	218	aduncella, Zell.	123	albilocrella, Zell.	76
achrestella, Reb.	201	adustipennis, Wals.	169	albimarginella, Chamb.	90
achroa, Low.	35	aegrella, Wals.	202	albinervella, Strand	89
achyrota, Meyr.	99	aenictodes, Turn.	34	albipalpella, Herr.-Sch.	112
acicularis, Meyr.	116	aeolopis, Meyr.	149	albipectus, Wals.	83
aclera, Meyr.	148	Aeolotrocha (genus), Meyr.	140	albiramis, Meyr.	23
acolasta, Meyr.	239	aequata, Meyr.	177	albiramosella, Christ.	30
Acompsia (genus), Hübn.	141	aequipulvella, Chamb.	253	albascripta, Meyr.	110
acosmeta, Wals.	122	aequorea, Meyr.	58	albisignis, Meyr.	235
Acoraeologa (genus), Meyr.	38	aerobatis, Meyr.	80	albisparsella, Chamb.	77
acrantha, Meyr.	102	aestivella, Herr.-Sch.	30	albistrigella, Chamb.	116
acrema, Meyr.	30	aestuosa, Meyr.	224	albocinctella, Dup.	58
Acribolega (genus), Meyr.	171	aethiops, Westw.	69	albogrisea, Wals.	156
acricula, Meyr.	149	affinis, Dougl.	74	albomaculella, Chamb.	78
acrocelea, Turn.	101	affixella, Walk.	257	alburnella, Dup.	71
acrochlora, Meyr.	110	affusa, Meyr.	239	alcestis, Meyr.	238
acrodactyla, Meyr.	99	agatha, Meyr.	43	alcimacha, Meyr.	105

	Pages		Pages		Pages
Alophanes (genus), Meyr.	207	amphichroma, Meyr.	71	anthracura, Meyr.	121
alella, Fabr.	70	amphicoma, Meyr	178	anthyllidella, Hübn.	111
alella, Wood	52	Amphigenes (genus), Meyr.	259	antibathra, Meyr.	224
aletris, Wals.	47	amphiptila, Meyr.	31	anticentra, Meyr.	103
aleurodes, Meyr.	154	amphiscolia, Meyr.	118	antichroa, Meyr.	250
alexandriella, Chamb.	72	amphisticta, Meyr.	250	antileuca, Meyr.	88
algeriella, Bak.	43	amphizeucta, Meyr.	171	Antiochtha, Meyr.	229
alhambrella, Wals.	202	ampliata, Meyr.	179	antipala, Meyr.	42
alienella, Walk.	251	ampycota, Meyr.	192	antiphractis, Meyr.	239
allactis, Meyr.	87	Anacampsis (genus), Curt.	125	antiplaca, Meyr.	121
Allocota, Meyr.	155	anagramma, Meyr	81	antirrhinella, Mill.	142
alloea, Wals.	160	analoxa, Meyr.	187	antisaris, Meyr.	156
alloptila, Meyr.	211	ananeura, Meyr.	149	antisphena, Meyr.	216
allotria, Meyr.	81	Anaphaula, Wals.	41	antizyga, Meyr.	176
alnifructella, Busck	58	anaphracta, Meyr.	201	anxia, Meyr.	66
alphitodes, Meyr.	23	Anaptilara (genus), Meyr.	254	anympha, Meyr.	238
alpicola, Frey	79	Anarsia (genus), Zell.	153	aoropis, Meyr.	64
alpicolella, Hein.	75	anarsiella, Chamb.	98	Apatema, Wals.	200
alsinella, Zell.	96	anarsiella, Chrét.	81	Apatetris (genus), Staud.	22
Alsodryas (genus), Meyr.	124	Anasphealtis (genus), Meyr.	107	apentheta, Turn.	170
altercata, Meyr.	153	Anaxyrina (genus), Meyr.	211	Apethistis (genus), Meyr.	250
alternatella, Kearn.	79	ancillula, Wals.	45	Aphanaula, Meyr.	57
alternella, Hübn.	70	ancylosticha, Turn.	192	aphanopa, Meyr.	177
alternella, Walk.	241	ancylota, Meyr.	240	aphileta, Meyr.	34
alternella, Walk.	179	andropogonis, Braun	37	aphiltra, Meyr.	45
alticolans, Wals.	116	Andusia, Walk.	237	Aphnogenes (genus), Meyr.	223
altisona, Meyr.	95	anguinea, Meyr.	157	aphroditella, Chamb.	130
altithermella, Wals.	23	anguinella, Herr.-Sch.	70	aphromorpha, Meyr.	46
amalleuta, Meyr.	154	angulatella, Chrét.	79	apicilinella, Clem.	117
amauropis, Meyr.	189	angulifera, Wals.	141	apicipuncta, Busck	156
amaurota, Meyr.	121	angustella, Hein.	83	apicipunctella, Staint.	31
amazonica, Meyr.	121	angustipennella, Clem.	43	apicistrigella, Chamb.	117
ambitiosa, Meyr.	153	animosella, Walk.	101	apicistrigella, Herr.-Sch.	72
Ambloma (genus) Wals.	198	anisaula, Meyr.	23	apicitripunctella, Clem.	58
amblopis, Meyr.	149	anisectis, Meyr.	78	apludella, Led.	176
Amblypalpis (genus), Rag.	21	anisopa, Meyr.	249	Apocritica (genus), Meyr.	64
ambrosiella, Chamb.	90	Anisoplaca (genus), Meyr.	98	Apodia, Hein.	41
ambusta, Wals.	121	anisosticta, Meyr.	108	apolectella, Wals.	77
amesella, Chrét.	42	annularia, Turn.	150	Aponoea (genus), Wals.	165
amethystias, Meyr.	131	annulicornis, Steph.	60	Apopira, Wals.	115
amethystina, Meyr.	138	annulicornis, Wals.	57	Apotactis (genus), Meyr.	159
ametris, Meyr.	81	anoma, Wals.	131	Apothetoea (genus), Meyr.	73
amicella, Walk.	228	Anomoxena (genus), Meyr.	39	Apotistatus (genus), Wals.	198
ammonura, Meyr.	157	Anorthosia, Clem.	174	apparitrix, Meyr.	157
ammoxantha, Meyr.	175	anteliella, Busck	35	apraca, Meyr.	169
amnesta, Meyr.	218	Anterethista (genus), Meyr.	114	aprlicata, Meyr.	249
amoenella, Frey	63	anthemidella, Wocke	35	aprilella, Herr.-Sch.	30
amoenella, Joan.	46	Anthinora (genus), Meyr.	126	Aproaerema, Durr.	111
amorphella, Chamb.	84	Anthistarcta (genus), Meyr.	67	Aprosoesta, Turn.	231
amphicalyx, Meyr.	201	anthochra, Low.	78	aptatella, Walk.	94
amphicarpa, Meyr.	212	anthologella, Wall.	239	aquilina, Meyr.	90
amphicentra, Meyr.	199	anthracina, Meyr.	148	aquosa, Meyr.	46
amphichlora, Meyr.	196	anthracopa, Meyr.	81	archaea, Wals.	67

LEPIDOPTERA HETEROCHERA

Pages	Pages	Pages			
arcifera, Meyr.	240	arundinella, Staint.	42	auritogata, Wals.	181
arcicella, Walk.	38	aruritis, Meyr.	249	auvorella, Frey.	98
arcuata, Wals.	44	asinella, Hübn.	98	austerodes, Meyr.	157
ardeliella, Hübn.	142	Asmenistis (genus), Meyr.	241	autarcha, Meyr.	210
ardesiella, Wals.	196	aspasta, Meyr.	257	authaema, Meyr.	256
ardiophora, Meyr.	118	aspera, Haw.	71	autodesma, Meyr.	131
ardua, Meyr.	226	aspetodes, Meyr.	150	autoleuca, Meyr.	23
arens, Meyr.	246	asphaltitis, Meyr.	66	autologa, Meyr.	239
Arogha (genus), Chrét.	141	aspretella, Led.	30	Automola, Meyr.	256
arganthes, Meyr.	150	astemphella, Meyr.	34	automorpha, Meyr.	255
argentaria, Meyr.	176	asterella, Treitsch.	46	Autoneda, Busck	33
argentialbella, Chamb.	58	asterisci, Wals.	42	autonoma, Meyr.	250
argentifera, Busck	47	asteropis, Meyr.	196	autophanta, Meyr.	246
argentifrons, Wals.	130	asthenodes, Meyr.	43	autopis, Meyr.	148
argentinella, Berg	175	astragali, Staud.	84	Autosticha (genus), Meyr	256
argentipunctella, Ely	75	astrapaea, Meyr.	109	auturga, Meyr.	33
argillosella, Herr.-Sch.	85	astroconis, Meyr.	117	auxiliaris, Meyr.	131
argocentra, Meyr.	148	atalopis, Meyr.	92	auxodelta, Meyr.	256
argocrossa, Meyr.	238	Atasthalitis (genus), Meyr.	136	aversella, Walk.	149
argodecta, Meyr.	45	atelesta, Meyr.	118	axenopsis, Meyr.	91
argogastra, Wals.	195	atterimella, Reb.	80	Axyrostola (genus), Meyr.	163
argonota, Low.	34	Athrinacia (genus), Wals.	221	azosterella, Herr.-Sch.	112
argosema, Meyr.	75	attractodes, Meyr.	102	baccata, Meyr.	177
argoxantha, Meyr.	102	atrella, Haw.	42	bacchariella, Busck	90
argyracma, Meyr.	120	atricanella, Reb.	201	Bactrolopha, Low.	36
argyracta, Meyr.	116	atrifascis, Meyr.	91	badia, Braun.	249
argyractis, Meyr.	46	triplex, Busck	90	bagriotella, Dup.	77
argyreia, Turn.	102	triplicella, Fisch. v. Rösl.	93	balanitis, Meyr.	228
argyromitra, Meyr.	166	triplicella, Strand	90	balanocentra, Meyr.	44
argyroneurella, Staud.	34	atropictella, Dietz	59	balanopa, Meyr.	102
argyrophaea, Meyr.	204	attenta, Meyr.	174	balbidota, Meyr.	230
argyrothamniella, Busck	123	attenuata, Meyr.	157	balcanica, Reb.	112
atidella, Walk	157	attonita, Meyr.	179	baldiana, Barnes	72
arignota, Meyr.	156	attributella, Walk.	63	balia, Wals.	122
aristella, Busck	77	attritella, Staud.	34	baliodes, Low.	157
Aristotelia (genus), Hübn.	40	attritella, Walk.	58	balioneura, Meyr.	152
aristotelis, Mill.	84	audax, Wals.	254	balneariella, Chrét.	28
arizonella, Busck	76	augusta, Meyr.	131	balsamopa, Meyr.	62
armata, Meyr.	138	Aulacomima, Meyr.	245	balteata, Meyr.	138
arminiella, Frey	47	aulacopis, Meyr.	44	banausodes, Meyr.	150
arnoldiella, Reb.	47	aulias, Meyr.	230	banksiella, Busck	90
Arogalea (genus), Wals.	67	Aulidiotis (genus), Meyr.	182	barathrodes, Meyr.	187
arotraea, Meyr.	248	aulodocha, Meyr.	234	barathrota, Meyr.	103
Arotria (genus), Meyr.	160	aulonitis, Meyr.	210	barbata, Wals.	199
arotrias, Meyr.	81	aulonota, Meyr.	45	barbella, Hübn.	179
arotrosema, Wals.	106	aulotoma, Meyr.	176	barberella, Busck.	25
arsenopa, Meyr.	154	aurantiaca, Wals.	131	barbifera, Meyr.	238
artemisiella, Kearf.	91	aurantiella, Chrét.	81	barcinonella, Mill.	26
artemisiella, Treitsch.	94	aurantiella, Reb.	45	barnesiella, Busck.	80, 106
arthrodes, Meyr.	103	aureolata, Meyr.	257	barydelta, Meyr.	62
artista, Meyr.	245	aurimaculella, Chamb.	94	barydesma, Meyr.	131
articulata, Meyr.	40	aurimaculella, Hofn.	42	barysphena, Meyr.	168
artificella, Herr.-Sch.	43	aurisulcata, Meyr.	195		

Pages		Pages		Pages	
<i>basaliella</i> , Walk.	213	<i>bilobella</i> , Zell.	196	<i>branella</i> , Busck	75
<i>basalis</i> , Staud.	113	<i>bimaculella</i> , Chamb.	76	<i>brevipalpella</i> , Reb.	75
<i>basaltinella</i> , Zell.	74	<i>biminimaculella</i> , Chamb.	83	<i>briantiella</i> , Tur.	236
<i>basanistis</i> , Meyr.	205	<i>binaria</i> , Meyr.	257	<i>brizella</i> , Treitsch.	46
<i>basifasciella</i> , Zell.	70	<i>biniveipunctata</i> , Wals.	87	<i>brochias</i> , Meyr.	206
<i>basiguttella</i> , Hein.	81	<i>binotata</i> , Meyr.	239	<i>brochodesma</i> , Meyr.	44
<i>basiphæa</i> , Turn	254	<i>binotatella</i> , Walk.	121	Brochometis (genus), Meyr.	180
<i>basipunctella</i> , Herr.-Sch.	81	<i>binotella</i> , Fisch. v. Rösl.	34	<i>brochospila</i> , Meyr.	121
<i>basistriata</i> , Wals.	179	<i>binotella</i> , Walk.	235	<i>brontophora</i> , Meyr.	93
<i>basistrigella</i> , Zell.	71	<i>bipunctella</i> , Snell.	239	<i>brucinella</i> , Mann	47
<i>batanella</i> , Busck	90	<i>bipunctella</i> , Wals.	177	<i>brumella</i> , Clem.	83
Batenia (genns), Chrét.	60	<i>bisignatella</i> , Walk.	235	<i>Brunhildae</i> , Schaw.	92
<i>bathrarcha</i> , Meyr.	112	<i>bisignella</i> , Snell.	179	<i>Brunicki</i> , Reb.	42
<i>bathropis</i> , Meyr.	81	<i>bispiculata</i> , Meyr.	80	<i>brunnea</i> , Schöy.	80
Battaristis (genus), Meyr.	117	<i>bistellella</i> , Walk.	150	<i>brunneella</i> , Herr.-Sch.	122
<i>Baueri</i> , Reb.	96	<i>bistrigata</i> , Meyr.	149	<i>brunneus</i> , Busck.	140
<i>Begoe</i> , Chamb.	195	<i>bistrigella</i> , Busck	117	<i>brunniceps</i> , Feld.	117
<i>behenella</i> , Const.	96	<i>bistrigella</i> , Chamb.	80	<i>bryochlora</i> , Meyr.	124
<i>belangerella</i> , Chamb.	70	<i>bitinctella</i> , Walk.	240	<i>bryophilella</i> , Wals.	164
<i>belfragesella</i> , Chamb.	49	<i>bivia</i> , Meyr.	227	<i>Bryotropha</i> , Hein.	73
<i>bellella</i> , Walk.	46	<i>bivittella</i> , Chrét.	113	<i>bubulcella</i> , Staud.	199
<i>belonodes</i> , Meyr.	24	<i>blacica</i> , Turn.	149	<i>bucolica</i> , Meyr.	93
<i>Benedeni</i> , Wey.	42	<i>blanda</i> , Meyr.	222	<i>buffonella</i> , Mill.	42
<i>beneficentella</i> , Murtf.	82	<i>blandella</i> , Dougl.	95	<i>bufo</i> , Wals.	75
<i>benitella</i> , Barnes	78	<i>blandiella</i> , Walk.	138	<i>bullata</i> , Meyr.	103
<i>bergiella</i> , Teich	80	<i>blandulella</i> , Tutt	95	<i>bullulata</i> , Meyr.	235
<i>betulea</i> , Haw.	79	<i>blandulella</i> , Walk.	76	<i>busckella</i> , Ely	38
<i>betulella</i> , Meyr.	70	<i>blapsigona</i> , Meyr.	94	<i>busckiella</i> , Kearf.	90
<i>biangulata</i> , Meyr.	112	<i>blattariella</i> , Hübn.	126	<i>butyraula</i> , Meyr.	108
<i>biannulella</i> , Chamb.	98	<i>blepharopa</i> , Meyr.	120	<i>butyropa</i> , Meyr.	228
<i>biareatella</i> , Ersch.	248	<i>bleptodes</i> , Turn.	147	<i>byrsoxantha</i> , Meyr.	189
<i>bicolor</i> , Wals.	24	<i>bletrias</i> , Meyr.	81	<i>byssina</i> , Meyr.	228
<i>bicolorella</i> , Chamb.	54	<i>bohemiella</i> , Nick.	32	<i>bythota</i> , Meyr.	243
<i>bicolorella</i> , Reb.	42	<i>bolshevikella</i> , Car.	46		
<i>bicolorella</i> , Treitsch.	70	<i>boreella</i> , Dougl.	83	<i>Cacelice</i> , Busck	51
<i>bicolorella</i> , Walk.	204	<i>boreae</i> , Wals.	47	<i>cachrydias</i> , Meyr.	175
<i>bicostimaculella</i> , Chamb.	63, 72	<i>bosquella</i> , Chamb.	87	<i>cacoderma</i> , Wals.	80
<i>bicristata</i> , Meyr.	50	<i>brabylitis</i> , Meyr.	138	<i>Cacogamia</i> , Snell.	204
<i>bicristatella</i> , Chamb.	71	<i>bracculata</i> , Meyr.	238	<i>cacomictra</i> , Wals.	42
<i>bicunea</i> , Meyr.	137	Brachmia (genus), Hübn.	248	<i>cacuminum</i> , Frey	95
<i>bidiscomaculella</i> , Chamb.	196	Brachyacma (genus), Meyr.	168	<i>caduca</i> , Meyr.	228
<i>bifasciata</i> , Staud.	201	<i>Brachycrossata</i> , Hein.	141	<i>caeca</i> , Meyr.	186
<i>bifasciella</i> , Busck.	46	Brachyerga (genus), Meyr.	235	<i>caecella</i> , Zell.	83
<i>bifasciella</i> , Chamb.	51	<i>brachygrapha</i> , Meyr.	174	<i>caecigena</i> , Meyr.	70
<i>bifasciella</i> , Steph.	200	<i>brachymetra</i> , Meyr.	177	<i>caecilia</i> , Meyr.	239
<i>biferrinella</i> , Walk.	240	<i>brachymita</i> , Turn.	236	<i>caelata</i> , Meyr.	70
<i>bifidella</i> , Dietz	24	<i>brachyntha</i> , Meyr.	152	<i>caeligena</i> , Meyr.	133
<i>biforella</i> , Busck	78	<i>brachyptera</i> , Wals.	198	<i>caerulescens</i> , Meyr.	196
<i>bifractella</i> , Dougl.	42	<i>brachyptila</i> , Meyr.	174	<i>caespitella</i> , Zell.	79
<i>bifrenata</i> , Meyr.	243	<i>brachyrhiza</i> , Meyr.	156	<i>calamogona</i> , Meyr.	34
<i>bigella</i> , Busck	80	<i>brachysticha</i> , Turn.	109	<i>calamopis</i> , Meyr.	220
<i>biguttella</i> , Herr.-Sch.	112	<i>brackenridgella</i> , Busck	90	<i>calastomella</i> , Christ.	46
<i>bilbainella</i> , Rössl.	98	<i>brahamiella</i> , Heyd.	95	<i>calathaea</i> , Meyr.	71

LEPIDOPTERA HETEROCHERA

Pages	Pages	Pages			
calceata, Meyr.	257	carpotoma, Meyr.	108	centrosema, Meyr.	34
calculatrix, Meyr.	45	Carterica (genus), Meyr.	223	centrospila, Turn.	170
calens, Meyr.	45	carycastis, Meyr.	138	centrothetis, Meyr.	170
calidella, Wals.	201	carycina, Meyr.	180	centrotypa, Meyr.	113
californica, Busck	60	caryfoliella, Chamb.	176	cephalella, Car.	87
caliginella, Mann	201	caryivorella, Pack.	71	cephalochra, Meyr.	246
caliginosa, Turn.	147	caryodora, Meyr.	156	cephalota, Meyr.	149
<i>caligula</i> , Meyr.	238	caryophragma, Meyr.	180	ceponoma, Meyr.	176
callichroma, Meyr.	133	caryoterra, Meyr.	121	ceramica, Meyr.	149
Calliprora (genus), Meyr.	106	<i>cassella</i> , Walk.	45	ceramochroa, Turn.	249
callirrhoda, Meyr.	45	cassidata, Meyr.	120	Ceratophora, Hein.	248
callitechna, Meyr.	70	cassiterota, Meyr.	240	cercerella, Chamb.	76
callitricha, Meyr.	225	castiliella, Möschl.	30	cerealella, Ol.	38
callyntrophora, Reb.	46	castrigera, Meyr.	71	cerinura, Meyr.	249
calumnians, Meyr.	118	cataclina, Meyr.	230	cerostomella, Wals.	79
Calyptrotis, Meyr.	22	catagrapha, Meyr.	27	ceroxesta, Meyr.	225
caminariella, Fuchs	79	Catalexis (genus), Wals.	123	cerussata, Wals.	76
caminopsis, Meyr.	148	catalinella, Busck	77	cervinella, Eversm.	47
campalea, Wals.	121	Catameces (genus), Turn.	32	Cerycangela (genus), Meyr.	134
campicolella, Mann	44	catarrhacta, Meyr.	149	Ceuthomadarus (genus), Mann	203
canariensis, Reb.	202	catectis, Meyr.	253	chalazodes, Turn.	149
canariensis, Wals.	70	Cataphrys (genus), Meyr.	182	chalcopera, Wals.	137
cancellata, Meyr.	159	catharodes, Meyr.	77	chalcophaea, Meyr.	176
candida, Staint.	164	catharosema, Meyr.	114	chalcoscia, Meyr.	34
candidella, Chrét.	65	catharurga, Meyr.	107	chalcotora, Turn.	81
caneodes, Meyr.	125	Cathégesis, Wals	142	chalcura, Meyr.	108
canicostella, Wals.	35	catopta, Turn.	144	Chaliniaestis (genus), Meyr.	109
canofusella, Walk.	120	Catoptristis (genus), Meyr.	134	chalybaeella, Walk.	204
<i>canopulvella</i> , Chamb.	82	cauligenella, Schmid	96	chalybitis, Meyr.	190
Canthoniastis (genus), Meyr.	212	causidica, Meyr.	228	chalybochroa, Wals.	45
capillata, Wals.	175	caustonota, Meyr.	193	chalyburga, Meyr.	138
capitella, Fabr.	87	caustospila, Meyr.	239	chambersella, Dyar	80
capitochrella, Chamb.	81	<i>cautella</i> , Zell.	83	chambersella, Murtf.	249
capnaula, Meyr.	241	ceanothiella, Braun	58	chamelia, Turn.	238
<i>capnella</i> , Const.	75	ceanothiella, Busck	75	characias, Meyr.	102
capnites, Meyr.	179	cecidiella, Chrét.	92	characopa, Meyr.	243
capsophilella, Chrét.	96	Cecidophaga, Wals.	22	characotis, Meyr.	231
captivella, Herr.-Sch.	112	cedestiella, Zell.	202	characteris, Meyr.	245
capucinella, Hübn.	177	celema, Durr.	231	charadraea, Meyr.	204
capyrodes, Meyr.	126	<i>colerella</i> , Staint	82	chariphila, Meyr.	69
Carbatina (genus), Meyr.	184	celeris, Meyr.	33	Charistis (genus), Meyr.	133
carbonaria, Meyr.	153	Celestodes (genus), Meyr.	220	charonea, Meyr.	110
carcerata, Meyr.	240	celidophora, Turn.	150	chartaria, Meyr.	185
carcharias, Meyr.	231	cellaria, Meyr.	196	Cheloria (genus), Haw.	155
carchariella, Zell.	43	<i>celphodes</i> , Meyr.	33	chelidonia, Meyr.	120
caribbea, Busck	254	cemiomostomella, Zell.	105	chelophora, Meyr.	100
carinata, Meyr.	63	cenchrritis, Meyr.	251	chenias, Meyr.	149
carinella, Wals.	196	cenelpis, Wals.	125	chenopodiella, Busck	90
carlinella, Staint.	30	centracma, Meyr.	189	cheradias, Meyr.	95
Carna, Walk.	174	centritis, Meyr.	102	cheramopis, Meyr.	214
Carodista (genus), Meyr.	224	centrocrossa, Meyr.	106	cherandra, Meyr.	243
carpaea, Meyr.	215	centrophora, Meyr.	104	chernetis, Meyr.	256
carphodes, Meyr.	250	centrosema, Low.	43	cherregella, Chrét.	88

Pages	Pages	Pages			
chersaea, Meyr.	61	cinerella, Clerck	142	coelatella, Walk.	230
chersitis, Meyr.	238	cinerella, Murtf.	93	coenotypa, Meyr.	259
chersochlora, Meyr.	93	cinerosella, Tengst.	75	coenulentella, Herr.-Sch.	36
Chersogenes (genus), Wals.	202	cingalensis, Wals.	211	coercita, Meyr.	228
chersophila, Meyr.	92	cinnabarina, Meyr.	194	coeruleopictella, Car.	46
chilonella, Treitsch.	107	cinnamicostella, Zell.	188	cognatella, Hein.	82
Chilopselaphus (genus), Mann	28	circaeа, Meyr.	112	cognatella, Walk.	120
chiloptila, Meyr.	225	circumfusa, Meyr.	250	colasta, Meyr.	92
chinganella, Christ.	98	Cirrha, Chamb.	73	coleasta, Meyr.	240
chionarcha, Meyr.	106	cirrhaea, Meyr.	23	coleella, Const.	102
chionitis, Meyr	164	cirrhocoma, Meyr.	113	Coleostoma (genus), Meyr.	119
chionocephala, Low.	24	cirrhospila, Meyr.	157	collecta, Meyr.	24
Chionodes, Hübn.	73	cirrhostola, Turn.	176	colleta, Wals	105
chionostigma, Wals.	85	cisti, Staint.	72	collina, Meyr.	186
chionura, Meyr.	115	cisti, Staud.	176	collinella, Chamb.	90
chiquitella, Busck	90	cistiflorella, Const.	94	collocatella, Walk.	121
chiradia, Meyr.	150	cistulata, Meyr.	120	colluta, Meyr.	221
chiridota, Meyr	108	citharista, Meyr.	171	Colobodes, Meyr.	108
chlauenota, Meyr.	236	citanthes, Meyr.	62	Colonanthes (genus), Meyr.	55
chloranthes, Low.	170	citrifoliella, Chamb.	175	Colopteryx (genus) Hofm.	26
chloristis, Meyr.	170	citrinella, Barnes	62	coloradensis, Busck	76
chlorodelta, Meyr.	256	citrocosma, Meyr.	45	colubrinae, Busck	58
chlorogastra, Meyr.	238	citromitra, Meyr.	153	combusta, Meyr	239
chlorographa, Meyr.	46	citrostrota, Meyr.	239	comedonella, Staud.	72
Chlorolychnis (genus) Meyr.	241	cladara, Meyr.	148	comis, Meyr.	47
chloroneura, Meyr.	43	Cladodes, Hein.	248	comissata, Meyr.	87
chlorophracta, Meyr.	176	clandestina, Meyr.	75	commaculata, Meyr.	70
chlorophthalma, Meyr.	160	clarella, Treitsch.	176	Commatica (genus), Meyr.	115
chloroschema, Meyr.	78	clarescens, Meyr.	177	commodata, Meyr.	228
chlorotoma, Meyr.	68, 242	clastrata, Meyr.	240	communicata, Meyr.	258
chlorotricha, Meyr.	220	clavata, Meyr.	47	complanata, Meyr.	222
cholopis, Meyr.	225	claviculata, Meyr.	189	complicata, Meyr.	102
chordotona, Meyr.	231	clematis, Meyr.	213	composita, Meyr.	167
choristis, Meyr.	138	Cleodora, Steph.	29	compositella, Chrét.	65
choritis, Meyr.	239	cleodorella, Chrét.	36	compositella, Walk.	76
chromatica, Meyr.	64	cleodorella, Wals.	105	compsochroa, Meyr.	149
chrysanthema, Meyr.	102	cleodorella, Zell.	102	Compsalechia (genus), Meyr.	119
chrysanthemella, Hofm.	95	clerica, Ros.	73	compsophila, Meyr.	237
chrysocosma, Meyr.	208	clerodotis, Meyr.	228	Compsosaria (genus), Meyr.	56
Chrysoesthia, Hübn.	40	Clerogenes, Meyr.	200	compsostoma, Meyr.	213
chrysopa, Meyr.	102	clidias, Meyr.	102	compsotoma, Meyr.	115
chrysoplaca, Meyr.	122	clinopeda, Meyr.	102	conchylidella, Hofm.	26
Chrysopora, Clem.	41	Clistothrypis (genus), Zell.	61	conciliata, Meyr.	256
chrysostoma, Meyr.	127	cistrodoma, Meyr.	78	concinna, Turn.	238
cicinnota, Meyr.	210	clitella, Reb.	30	concinna, Wals.	78
ciliatella, Herr.-Sch.	75	clitellaria, Meyr.	49	concinnalis, Feisth.	136
ciliolineella, Chamb.	35	Cnaphostola (genus), Meyr.	104	concinnusella, Chamb.	117
cinclidias, Meyr.	193	coarctata, Wals.	188	conclusa, Meyr.	184
cinctella, Hübn.	112	coarctatella, Zell.	38	concreta, Walk.	125
cinctella, Walk.	188	coarctella, Reb.	200	concreta, Meyr.	94
cincticulella, Herr.-Sch.	112	cockerella, Busck	47	condalivorella, Busck	196
cinctipunctella, Ersch.	77	Cockerelli, Busck	81	condylodes, Meyr.	189
cinerella, Chrét.	58	cocta, Meyr.	196	confinis, Staint.	74

LEPIDOPTERA HETEROCHERA

Pages		Pages		Pages
confirmata, Meyr.	53	cornifera, Wals.	125	crepitatrix, Meyr.
confixa, Meyr.	69	cornuta, Busck	136	crepusculella, Teich.
conficta, Meyr.	105	cornutella, Walk.	238	crescentifasciella, Chamb.
<i>confusella</i> , Chamb.	84	coronilla, Treitsch.	112	cretigena, Meyr.
confusella, Hein.	82	corsicella, Herr.-Sch.	168	cribanota, Meyr.
congressariella, Bruand	98	corsota, Meyr.	238	cricopa, Meyr.
conifera, Meyr.	90	Corthyntis, Meyr.	220	criminata, Meyr.
coniferella, Kearn.	58	corymbas, Meyr.	122	criodes, Meyr.
coniodes, Meyr.	35	Corynaea (genus), Turn.	146	cristatella, Chamb.
coniogramma, Meyr.	34	corynetis, Meyr.	156	cristifasciella, Chamb.
Coniogrya (genus), Meyr.	64	corythista, Meyr.	209	critica, Wals.
coniombra, Meyr.	24	cosmella, Const.	107	Crocanthes (genus), Meyr.
coniosema, Meyr.	118	cosmia, Meyr.	126	crocina, Meyr.
conistica, Wals.	126	cosmodes, Low.	87	crocipunctella, Wals.
conjugella, Car.	87	cosmographa, Meyr.	46	crocodesma, Meyr.
conotoma, Meyr.	205	cosmophragma, Meyr.	221	crocodilopa, Meyr.
consanguinella, Herr.-Sch.	107	cosmotis, Meyr.	148	crocodora, Meyr.
conscriptella, Hübn.	156	costalis, Busck	178	Crocegma (genus), Meyr.
consertella, Christ.	175	costella, Westw.	94	crocogramma, Meyr.
considerata, Meyr.	125	<i>costigittella</i> , Zell.	249	crocoleuca, Meyr.
consociata, Meyr.	62	costimaculella, Snell.	235	croconota, Meyr.
consona, Meyr.	75	<i>costipunctella</i> , Möschl.	87	crocothicta, Meyr.
consonella, Zell.	123	costirufella, Chamb.	196	Croesopola, Meyr.
consors, Meyr.	246	<i>costolutella</i> , Chamb.	196	Crossobela (genus), Meyr.
consortiella, Car.	34	cosymbota, Meyr.	220	crossogramma, Meyr.
conspersa, Braun	82	coticola, Busck	77	crotalariella, Busck
conspersella, Herr.-Sch.	42	cotifera, Meyr.	176	<i>crotalariella</i> , Busck
conspurcatella, Hein.	82	Cotyloscia (genus), Meyr.	193	crotalodes, Meyr.
constantina, Bak.	35	Coudia (genus), Chrét.	65	crudescens, Meyr.
constrictella, Zell.	51	coussonella, Chrét.	96	cruenta, Meyr.
consummata, Meyr.	251	coverdalella, Kearn.	120	crustaria, Meyr.
contentella, Walk.	197	Coydalia (genus), Walk.	228	cryolopha, Meyr.
<i>contigua</i> , Haw.	96	crambaleas, Meyr.	197	cryphias, Meyr.
continctella, Walk.	244	crambinella, Zell.	105	cryphoides, Meyr.
continella, Zell.	78	Crambedoxa (genus), Meyr.	104	crysibatis, Meyr.
contracta, Meyr.	238	craniota, Meyr.	239	crypsichola, Durr.
contraria, Braun	90	Crasimorpha (genus), Meyr.	160	<i>crysilychna</i> , Meyr.
contristella, Car.	201	Craspedotis (genus), Meyr.	144	crypsixantha, Turn.
contrita, Meyr.	186	cramella, Feld.	211	crypsizygia, Meyr.
<i>contubernatellus</i> , Fitch	175	crassicornis, Wals.	44	cryptina, Wals.
<i>contuberniella</i> , Staud.	122	<i>crataegella</i> , Busck	58	cryptogamarum, Mill.
convergens, Wals.	105	crateraula, Meyr.	33	cryptosperma, Meyr.
convexa, Meyr.	217	crateropis, Meyr.	250	crystallista, Meyr.
convolvuli, Wals.	249	craterospila, Meyr.	250	cubiculata, Meyr.
cophias, Meyr.	84	craterota, Meyr.	71	cucullata, Meyr.
Cephomantis (genus), Meyr.	242	craticula, Meyr.	249	cuencella, Car.
Copocercia, Zell.	105	crauropa, Meyr.	62	<i>culminicolella</i> , Staud.
Cepreptilia (genus), Snell.	233	creberima, Wals.	78	cuneatella, Dougl.
coquillettella, Busck	90	crebrata, Meyr.	240	cuneifera, Wals.
coracina, Meyr.	111	cremnaspis, Meyr.	228	cuneiformis, Meyr.
corallina, Wals.	45	cremnobathra, Meyr.	246	cuprimarginella, Chrét.
cordata, Meyr.	241	crenoides, Meyr.	165	curonella, Standf.
corniculata, Meyr.	197	creperrima, Turn.	148	curtella, Busck

Pages		Pages		Pages
<i>curvatella</i> , Busck	86	Delmnestra (genus), Meyr.	189	diacma, Meyr.
<i>curvistrigella</i> , Chamb.	72	<i>dejectella</i> , Staud.	40	diacnista, Meyr.
<i>custos</i> , Meyr.	250	<i>delacma</i> , Meyr.	102	<i>diaphanella</i> , Zell.
<i>cyamitis</i> , Meyr.	238	<i>delatrix</i> , Meyr.	70	Diastalтика (genus), Wals.
<i>cyanoneura</i> , Meyr.	195	<i>deleastræ</i> , Meyr.	237	diasticha, Turn.
<i>cyanopæ</i> , Meyr.	211	<i>delicatella</i> , Wals.	202	<i>diazeucta</i> , Meyr.
<i>cyanorrhœa</i> , Meyr.	115	<i>delminiella</i> , Reb.	63	<i>dicausta</i> , Meyr.
<i>cyanozona</i> , Meyr.	131	<i>delochorda</i> , Low.	23	<i>dicax</i> , Meyr.
<i>cyclælla</i> , Busck	117	<i>deloma</i> , Durr.	240	Dichomeris (genus), Hübn.
<i>cyclonitis</i> , Meyr.	236	<i>delotella</i> , Busck	174	<i>dichroa</i> , Wals.
<i>cyclopila</i> , Meyr.	188	<i>delphinatella</i> , Const.	96	<i>dictyomorpha</i> , Low.
<i>cycnobathra</i> , Low.	171	<i>deltaspis</i> , Meyr.	179	<i>Didactylota</i> , Wals.
<i>cycota</i> , Meyr.	71	<i>deltochlora</i> , Meyr.	124	<i>didymopa</i> , Meyr.
<i>cymatodes</i> , Meyr.	188	<i>deltodes</i> , Low.	149	<i>dietziella</i> , Busck
<i>Cymatomorpha</i> , Meyr.	155	<i>deltopis</i> , Meyr.	250	<i>difficilisella</i> , Chamb.
Cymatoplex (genus), Meyr.	223	Deltoplastis (genus), Meyr.	228	<i>diffinis</i> , Haw.
Cymotricha (genus), Meyr	188	<i>deltospila</i> , Meyr.	241	<i>diffuella</i> , Hein.
<i>cymotrocha</i> , Meyr.	171	<i>demetrias</i> , Meyr.	257	<i>digitata</i> , Meyr.
Cynicostola (genus), Meyr.	230	<i>demias</i> , Meyr.	255	<i>digna</i> , Meyr.
<i>cynthia</i> , Meyr.	45	Demiophila (genus), Meyr.	257	<i>dignella</i> , Wals.
<i>Cyrnia</i> , Wals.	199	<i>Demopractis</i> , Meyr.	219	<i>dilechria</i> , Turn.
<i>cyrota</i> , Meyr.	200	<i>demonstrata</i> , Meyr.	156	<i>diligens</i> , Meyr.
<i>cyrtopleura</i> , Turn.	157	<i>demotica</i> , Meyr.	256	<i>dilucescens</i> , Meyr.
<i>cytheraea</i> , Meyr.	45	<i>densata</i> , Meyr.	91	<i>diluticornis</i> , Wals.
<i>cytisella</i> , Curt.	36	<i>dentata</i> , Meyr.	171	<i>dimidiella</i> , Schiff.
<i>cytisella</i> , Treitsch.	78	<i>dentella</i> , Busck	77	<i>diminutella</i> , Zell.
Dactylethora (genus), Meyr.	164	Decclona (genus), Busck	183	<i>dimorpha</i> , Petr.
<i>Dactylota</i> , Snell.	22	<i>depressostrigella</i> , Chamb.	81	<i>dimorphella</i> , Busck
<i>daedalea</i> , Wals.	138	<i>depuratella</i> , Busck	77	Dinochares (genus), Meyr.
<i>damonella</i> , Mill.	43	<i>dermatica</i> , Meyr.	156	<i>diortha</i> , Meyr.
<i>dasyopoda</i> , Wals.	45	<i>derogata</i> , Meyr.	246	<i>diplanetis</i> , Meyr.
<i>daturae</i> , Zell.	94	Deroxena (genus), Meyr.	246	<i>diplobathra</i> , Meyr.
<i>deauratella</i> , Herr.-Sch.	200	<i>desectella</i> , Zell.	122	<i>diplodelta</i> , Meyr.
<i>decaspila</i> , Low.	149	<i>desertella</i> , Dougl.	75	<i>diplolychna</i> , Meyr.
Decatopseustis (gen.), Meyr.	140	<i>desertella</i> , Reb.	92	<i>diplonesa</i> , Meyr.
<i>decedens</i> , Wals.	252	<i>deserticolella</i> , Staud.	95	<i>diplosticta</i> , Meyr.
<i>decemmaculella</i> , Chamb.	81	<i>designatella</i> , Herr.-Sch.	201	<i>dipsalea</i> , Meyr.
<i>deceptella</i> , Snell.	197	<i>designatella</i> , Walk.	189	<i>dipsia</i> , Meyr.
<i>declaratella</i> , Staud.	98	<i>designella</i> , Herr.-Sch.	202	<i>directa</i> , Meyr.
<i>decolorella</i> , Hein.	84	<i>desiliens</i> , Meyr.	80	<i>dirupta</i> , Meyr.
<i>decolorella</i> , Herr.-Sch.	43	<i>desmanthes</i> , Low.	81	<i>dischema</i> , Meyr.
<i>decoratella</i> , Staud.	46	<i>desmatra</i> , Low.	148	<i>discissa</i> , Meyr.
<i>decorella</i> , Haw.	69	Desmaucha (genus), Meyr.	127	<i>disclosa</i> , Meyr.
<i>decrepidella</i> , Herr.-Sch	75	<i>desolata</i> , Meyr.	238	<i>discoannulella</i> , Chamb.
<i>Dectobathra</i> , Meyr.	137	<i>despectella</i> , Walk.	61	<i>discocellella</i> , Chamb.
<i>Decuria</i> , Walk.	209	<i>destillans</i> , Meyr.	70	<i>discomaculella</i> , Chamb.
<i>decuriella</i> , Mann	70	<i>detersella</i> , Clem.	90	<i>disconotella</i> , Chamb.
<i>decurtella</i> , Hübn.	46	<i>detersella</i> , Zell.	113	<i>discontinuella</i> , Reb.
<i>decusella</i> , Walk.	179	<i>deuteraula</i> , Meyr.	105	<i>discostrigella</i> , Chamb.
<i>deductella</i> , Walk.	257	<i>Deuteroptila</i> , Meyr.	155	<i>discretella</i> , Reb.
<i>deflecta</i> , Busck	192	<i>deverrae</i> , Wals.	113	<i>discriminata</i> , Meyr.
<i>deflectivella</i> , Herr.-Sch.	177	<i>devia</i> , Meyr.	76	<i>discura</i> , Meyr.
		<i>deviella</i> , Walk.	163	<i>disjectella</i> , Staud.
				94

LEPIDOPTERA HETEROCHERA

Pages	Pages	Pages			
<i>disparata</i> , Meyr.	108	<i>dryozona</i> , Meyr.	58	<i>Encoentrotis</i> (genus) , Meyr.	27
<i>dispensata</i> , Meyr.	92	<i>dubitella</i> , Chamb.	196	<i>enchotypa</i> , Turn.	149
<i>dispila</i> , Turn.	236	<i>dudiella</i> , Busck	89	<i>Enchrysa</i> , Zell.	41
<i>displacitella</i> , Walk.	122	<i>durdhamella</i> , Staint.	173	<i>Encelapta</i> (genus) , Meyr.	158
<i>Disquei</i> , Meess	126	<i>Duvita</i> , Busck	117	<i>Encelpotis</i> (genus) , Meyr.	251
<i>disrupta</i> , Meyr.	66	<i>dyariella</i> , Busck	77	<i>Encrasima</i> (genus) , Meyr.	258
<i>dissectella</i> , Zell.	47	<i>dysorata</i> , Turn.	176	<i>encycota</i> , Meyr.	256
<i>dissidens</i> , Meyr.	156	<i>Dzieduszycki</i> , Now.	77	<i>endocoma</i> , Meyr.	225
<i>dissimilella</i> , Treitsch.	77			<i>englypta</i> , Meyr.	149
<i>dissoluta</i> , Staud.	202	<i>Eatoni</i> , Wals.	30	<i>engrapta</i> , Meyr.	249
<i>dissonella</i> , Herr.-Sch.	102	<i>eburata</i> , Meyr.	84	<i>enoptrias</i> , Meyr.	132
<i>dissonella</i> , Walk.	239	<i>eburnella</i> , Christ.	154	<i>Enthetia</i> (genus) , Meyr.	218
<i>Dissoptila</i> (genus) , Meyr.	66	<i>echiochilonella</i> , Chrét.	33	<i>entryphopa</i> , Meyr.	119
<i>distigmatella</i> , Zell.	238	<i>ecliptica</i> , Meyr.	206	<i>eodryas</i> , Meyr.	71
<i>distinctella</i> , Zell.	82	<i>effera</i> , Meyr.	249	<i>epactaea</i> , Meyr.	81
<i>diula</i> , Meyr.	231	<i>effrenata</i> , Meyr.	233	<i>Epanastis</i> , Wals.	202
<i>diversella</i> , Busck	80	<i>egenella</i> , Herr.-Sch.	113	<i>epermeniella</i> , Wals.	44
<i>dividua</i> , Philp.	80	<i>Eidothea</i> , Chamb.	57	<i>Epharmonia</i> (genus) , Meyr.	226
<i>divisella</i> , Dougl.	43	<i>elaboratella</i> , Braun	78	<i>Ephelictis</i> (genus) , Meyr.	151
<i>dodecatella</i> , Staud.	202	<i>elachistella</i> , Busck	53	<i>ephelotis</i> , Meyr.	254
<i>dodecella</i> , Linn.	60	<i>elachistella</i> , Staint.	111	<i>ephestris</i> , Meyr.	208
<i>dodicella</i> , Wood.	71	<i>elachistella</i> , Zell.	42	<i>ephialtes</i> , Wals.	86
<i>dolabella</i> , Zell.	196	<i>elaeocoma</i> , Meyr.	112	<i>ephippias</i> , Meyr.	153
<i>dolabrata</i> , Meyr.	201	<i>elaeopis</i> , Meyr.	258	<i>ephoria</i> , Meyr.	45
Dolbyi, Wals.	76	<i>elaphopis</i> , Meyr.	242	<i>Ephyteris</i> (genus) , Meyr.	60
<i>Dolerotricha</i> (genus) , Meyr.	151	<i>Elasipora</i> (genus) , Meyr.	55	<i>epibola</i> , Wals.	123
<i>Dolichoterna</i> (genus) , Meyr.	217	<i>elassopis</i> , Turn.	148	<i>Epibrontis</i> (genus) , Meyr.	143
<i>doliodes</i> , Meyr.	43	<i>elatella</i> , Herr.-Sch.	77	<i>epicentra</i> , Meyr.	94, 137
<i>doliopa</i> , Meyr.	231	<i>electella</i> , Zell.	78	<i>Epicharma</i> , Wals.	256
<i>dolosa</i> , Meyr.	251	<i>elegans</i> , Snell.	204	<i>epicharta</i> , Turn.	45
<i>dolosella</i> , Zell.	34	<i>elegantella</i> , Chamb.	47	<i>epichorda</i> , Meyr.	29
<i>domestica</i> , Haw.	74	<i>elegantior</i> , Wals.	44	<i>epichorda</i> , Turn.	169
<i>dominicella</i> , Wals.	254	<i>elephantopa</i> , Meyr.	240	<i>epiclista</i> , Meyr.	102
<i>donatella</i> , Walk	87	<i>elephas</i> , Wals.	123	<i>epicyclosta</i> , Meyr.	145
<i>dorsalis</i> , Busck	117	<i>ellipsias</i> , Meyr.	189	<i>Epicoenia</i> , Meyr.	256
<i>dorsivittella</i> , Zell.	58	<i>elongella</i> , Hein.	44	<i>Epicorthyllis</i> (genus) , Zeller	192
<i>Doryonepa</i> (genus) , Low.	36	<i>elpista</i> , Wals.	253	<i>Epidola</i> (genus) , Staud.	26
<i>Doryphora</i> , Hein.	41	<i>elpistis</i> , Meyr.	148	<i>epidolella</i> , Chrét.	27
<i>doxarcha</i> , Meyr.	132	<i>elucidella</i> , Barnes	59	<i>spigatella</i> , Chamb.	75
<i>Doxogenes</i> (genus) , Meyr.	205	<i>eludens</i> , Meyr.	238	<i>epigompha</i> , Meyr.	241
<i>drachmaea</i> , Meyr.	121	<i>emancipata</i> , Meyr.	94	<i>Epimesophleps</i> (genus) , Reb.	169
<i>dracopis</i> , Meyr.	220	<i>emblematica</i> , Meyr.	144	<i>epimetalla</i> , Meyr.	47
<i>Dragmatucha</i> , Meyr.	227	<i>emblemella</i> , Clem.	131	<i>Epimimaetis</i> (genus) , Meyr.	144
<i>Drepaneterma</i> (genus) , Wals.	129	<i>embrocha</i> , Meyr.	112	<i>epiochra</i> , Meyr.	169
<i>drimylota</i> , Meyr.	150	<i>emerita</i> , Meyr.	63	<i>Epiparaesia</i> (genus) , Reb.	29
<i>dromicella</i> , Busck	77	<i>emicans</i> , Meyr.	86	<i>epiphanta</i> , Meyr.	119
<i>drosias</i> , Meyr.	24	<i>emigrans</i> , Meyr.	249	<i>epiphloea</i> , Meyr.	87
<i>dryadella</i> , Zell.	74	<i>emissurella</i> , Walk.	117	<i>Epiphthora</i> , Meyr.	22
<i>drvadopa</i> , Meyr	249	<i>emmetra</i> , Meyr.	256	<i>Episacta</i> , Turn.	155
<i>dryinodes</i> , Low.	179	<i>Empalactis</i> (genus) , Meyr.	170	<i>episema</i> , Wals.	121
<i>dryinopa</i> , Low.	164	<i>Empedaula</i> (genus) , Meyr.	47	<i>epispila</i> , Low.	40
<i>dryobathra</i> , Meyr.	80	<i>emphanista</i> , Meyr.	70	<i>episticta</i> , Meyr.	251
<i>dryocrossa</i> , Meyr.	122	<i>emplasta</i> , Meyr.	115	<i>Epistomotis</i> , Meyr.	199

	Pages		Pages		Pages
Epitheatis (genus), Meyr.	62	euchorda, Meyr.	157	excoriata, Meyr.	177
epitherma, Low.	131	euchroa, Low	136	exemplaris, Meyr.	256
epithymella, Staud.	93	Eucordylea (genus), Dietz	59	exiguella, Chrét.	202
epitricha, Meyr.	95	Eudactylota, Wals.	25	exilis, Wals.	34
epiula, Meyr.	153	<i>eudela</i> , Turn.	138	exoena, Meyr.	79
epomia, Meyr.	240	Eudodacles, Snell.	248	exoleteella, Ersch.	34
epomidella, Tengst.	70	eumela, Low.	150	exophthalma, Meyr.	240
Eporastis (genus), Meyr.	184	eumeris, Meyr.	46	Exoteleia (genus), Wall.	59
epotias, Meyr.	154	Eunebristis (genus), Meyr.	193	exsiccata, Meyr.	254
Eppelshimi, Staud.	47	Eunomarcha (genus), Meyr.	128	extincta, Meyr.	63
erebaula, Meyr.	39	euparypha, Meyr.	195	exsulata, Meyr.	112
erebenna, Meyr.	54	eupatoriella, Chamb.	196	exteriorella, Walk.	133
erebodelta, Meyr.	122	eupatris, Meyr.	212	externella, Zell.	189
eremaula, Meyr.	92	eupecta, Meyr.	120	extoris, Meyr.	92
eremella, Chrét.	80	euphorodes, Meyr.	153	extranea, Wals.	72, 210
eremias, Meyr.	254	<i>euphracta</i> , Meyr.	134	extremella, Walk.	115
Eremica (genus), Wals.	198	euplecta, Meyr.	157	exustella, Zell.	107
eremna, Meyr.	115	Eupolis (genus), Meyr.	180		
eremopis, Meyr.	68	Eupragia (genus), Wals.	222	<i>facteta</i> , Meyr.	112
eremota, Meyr.	243	<i>euratella</i> , Herr.-Sch.	70	factiosa, Meyr.	225
erethistis, Meyr.	106	euryanthes, Meyr.	81	fagella, Busck	61
ergasima, Meyr.	94	euryarga, Turn.	148	fagoniae, Meyr.	88
Ergatis, Hein.	41	eurybatis, Meyr.	73	fagoniae, Wals.	141
ericetella, Hübn.	79	eurydelta, Meyr.	106	falcatella, Walk.	115
ericinella, Dup.	46	eurydoxa, Meyr.	106	fallax, Durr.	231
ericnista, Meyr.	92	eurygypsa, Meyr.	121	fallax, Mann	28
ericta, Meyr.	156	eurylampis, Meyr.	204	famosa, Meyr.	188
ericydes, Meyr.	216	eurynota, Wals.	172	famulata, Meyr.	175
Eridachtha (genus), Meyr.	220	eurypsola, Turn.	45	fanatica, Meyr.	92
eridantis, Meyr.	177	eurypyra, Meyr.	231	Fapua, Strand	89
eridora, Meyr.	97	Eurysara, Turn.	174	farinata, Meyr.	63
erigeronella, Braun	90	eurysema, Meyr.	145	farinosa, Teich	82
eriozona, Meyr.	156	curyterma, Meyr.	257	farinosae, Staint.	42
Eripnura (genus), Meyr.	129	Euryzancla, Turn.	174	fasciata, Staint.	200
erista, Meyr.	259	euryzeucta, Meyr.	70	fasciella, Chamb.	71
erixantha, Meyr.	177	eustacta, Meyr.	176	fasciella, Chrét.	60
eromene, Wals.	58	eustephana, Turn.	150	fasciella, Feld.	120
erosella, Snell.	163	eutacta, Meyr.	153	fasciella, Hübn.	177
erratica, Meyr.	233	euxena, Meyr.	34	fastidiosa, Meyr.	66
Erschoffii, Frey	69	Euzonomacha (genus), Meyr.	133	fausta, Meyr.	239
erubescens, Wals.	79	Evagora, Clem.	57	faustella, Busck	90
erudita, Meyr.	149	Eville (genus), Chamb.	53	<i>favillaticella</i> , Zell.	65
erycina, Meyr.	45	evippella, Busck	54	febriculella, Zell.	58
Erythriastis (genus), Meyr.	245	evitata, Wals	195	fecunda, Meyr.	71
erythrogramma, Meyr.	100	exacta, Meyr.	95	femoralis, Staud.	70
escharitis, Meyr.	144	exaema, Meyr.	243	ferulella, Zell.	76
eschatopsis, Meyr.	93	exarista, Meyr.	148	ferax, Meyr.	79
ethicodes, Meyr.	28	excaecata, Meyr.	240	ferella, Berg	93
Ethirostoma (genus) Meyr.	117	excavata, Busck	195	fernaldella, Busck	196
euargyra, Turn.	138	excelsa, Frey	95	ferrata, Meyr.	177
Eucatoptus, Wals.	41	excepta, Meyr.	189	<i>ferrea</i> , Frey	43
eucerella, Walk.	229	<i>excisrella</i> , Walk.	175	ferreata, Meyr.	120
eucharacta, Meyr.	127	exclarella, Möschl.	77	ferritincta, Turn.	45

LEPIDOPTERA · HETEROCHERA

Pages		Pages		Pages
ferruginosa, Meyr.	175	fortis, Meyr.	240	galactopa, Meyr.
ferulata, Meyr.	66	fracticostella, Wals.	188	galbanella, Zell.
fervidella, Mann	47	fragariae, Busck	43	galenaea, Meyr.
festa, Meyr.	184	fragariella, Busck	122	galeodes, Meyr.
fibularis, Meyr.	244	fragmentella, Zell.	71	galeotis, Meyr.
Ficulea, Walk.	73	frankeniae, Wals.	46	gallaesteris, Kell.
fida, Meyr.	175	frankeniivorella, Chrét.	78	gallaediplopappi, Fyles
figulella, Staud.	75	fraternella, Dougl.	95	gallaesolidaginis, Ril.
figurata, Meyr.	228	frenigera, Meyr.	110	gallicella, Const.
figurella, Busck	80	frequens, Meyr.	93	gallicola, Busck
filicornis, Zell.	58	Frisilia (genus), Meyr.	213	galligenitella, Clem.
finitima, Meyr	246	frugalis, Meyr.	148	gallinolella, Mann.
finitimella, Reb	107	frustrata, Meyr.	239	<i>gallinella</i> , Treitsch.
fischerella, Treitsch.	96	frustulenta, Meyr.	52	Gasmara (genus), Walk.
fiscinata, Meyr.	249	fruticella, Wals.	48	gecko, Wals.
fissilis, Meyr.	100	fugacea, Zell.	71	gelastis, Meyr.
fiumella, Krone	94	fugitivella, Zell.	71	Gelechia (genus), Hüb.
flabellifera, Reb.	154	fuliginosa, Feld.	117	geminella, Chrét.
flagellifera, Wals.	58	fuliginosella, Snell.	74	geminella, Ril.
flagitiosa, Meyr.	224	fullonella, Zell.	122	gemmella, Linn.
flammella, Treitsch.	142	fulminella, Busck	76	generosa, Meyr.
flammulella, Wals.	78	fulminella, Mill.	83	Geniadophora, Wals.
flavella, Dup.	142	fulvicilia, Meyr.	196	geniatella, Busck
flavescens, Meyr.	256	fulvidella, Wals.	192	genistae, Staint.
flavicapitella, Chrét.	43	fulvistillella, Reb.	112	genistae, Wals.
flavicomella, Zell.	82	fumatella, Dougl.	82	geochrota, Meyr.
flavicorporella, Wals.	80	funestella, Hüb.	76	geomicta, Meyr.
flavidella, Busck	56	fungifera, Meyr.	197	georgiella, Walk.
flavipalpella, Tengst.	75	fungivorella, Clem.	46	geraca, Meyr.
flavipalpella, Wals.	239	furcifera, Turn.	149	germarella, Hüb.
<i>flavipalpis</i> , Wals.	239	furfurella, Staud.	94	germarella, Nolck.
flavissimella, Mann.	236	furnaria, Meyr.	243	gerronella, Zell.
flaviterminella, Wals.	127	furtiva, Meyr.	44	geryella, Chrét.
<i>flavivittellus</i> , Clem.	175	furvella, Zell.	188	gibbosella, Zell.
flavocostella, Clem.	196	furvescens, Meyr.	221	gibsonella, Kearf.
flavonigrella, Chrét.	79	fuscantella, Hein.	79	gibsonella, Busck
flexilis, Meyr.	148	fuscata, Meyr.	29	gigantea, Swaz.
flexiloqua, Meyr.	126	fuscedinella, Snell.	240	<i>gilviscopella</i> , Zell.
flexurella, Clem.	83	<i>fuscipennis</i> , Westw.	36	gilvolinella, Clem.
florella, Busck	90	fuscocristatella, Chamb.	49	gilvomaculella, Clem.
floridanella, Busck	254	fuscoluteella, Chamb.	82	glabrella, Hein.
fluctuans, Meyr.	188	fuscomaculella, Chamb.	83	gladiata, Meyr.
fluidescens, Meyr.	44	<i>fuscoochrella</i> , Chamb.	78	glagera, Turn.
fluitans, Meyr.	176	<i>fuscopallidella</i> , Chamb.	57	glandiferella, Zell.
fluvialella, Busck	84	<i>fuscopulvella</i> , Chamb.	54, 83	glaphyra, Wals.
foederalis, Meyr.	230	<i>fuscopunctella</i> , Clem.	72	Glaphyrurga (genus), Meyr.
fondella, Busck	78	<i>fuscostrigella</i> , Chamb.	105	glaphyritis, Meyr.
formidolosa, Meyr.	156	<i>fuscotaeniella</i> , Chamb.	80	Glauce (genus), Chamb.
formosella, Hüb.	142			glaucitis, Meyr.
formosella, Murtf.	83			glaucodes, Meyr.
formulata, Meyr.	170			glaucoterna, Meyr.
fornacalis, Meyr.	241			glebicolarella, Ersch.
fornacaria, Meyr.	94			glebicolarella, Snell.

Pages	Pages	Pages			
<i>gleditschiaeella</i> , Chamb.	51	<i>halonella</i> , Herr.-Sch.	95	<i>Herbichi</i> , Now.	94
<i>glenias</i> , Meyr.	101	<i>halurga</i> , Meyr.	231	<i>hercogramma</i> , Meyr.	188
<i>globulata</i> , Meyr.	164	<i>halymella</i> , Mill.	92	<i>Heringia</i> , Hed.	59
<i>glochinella</i> , Zell.	93	<i>hamulata</i> , Meyr.	102	<i>hermannella</i> , Fabr.	47
<i>glycina</i> , Meyr.	231	Hapalonoma (genus), Meyr.	116	<i>hessleriellus</i> , Rössl.	34
<i>glynopis</i> , Meyr.	128	Hapalosaris (genus), Meyr.	55	<i>hetaera</i> , Meyr.	101
<i>glycyrrhizella</i> , Chamb.	82	<i>hapalyntis</i> , Meyr.	249	<i>hetaeria</i> , Wals.	79
Glyphidocera (genus), Wals.	253	Haplochela (genus), Meyr.	157	<i>heteracma</i> , Meyr.	177
<i>gnophrina</i> , Feld.	136	<i>hapospila</i> , Turner	254	<i>heteractis</i> , Meyr.	114
Gnorimoschema (genus), Busck	89	Harmatitis (genus), Meyr.	220	Heteralcis (genus), Meyr.	209
<i>Godmani</i> , Wals.	102	<i>harmonias</i> , Meyr.	176	Heterodeltis (genus), Meyr.	211
Gonaepa (genus), Walk.	232	<i>Harpagandra</i> , Meyr.	253	<i>heterospora</i> , Meyr.	95
<i>goodellella</i> , Chamb.	139	Harpagidia (genus), Rag.	134	Heterozanca (genus), Turn	150
<i>gorgonias</i> , Meyr.	150	<i>harpastis</i> , Meyr.	24	<i>hexacentra</i> , Meyr.	115
<i>gorgopa</i> , Meyr.	136	Harpograptis (genus), Meyr.	126	<i>hexagramma</i> , Meyr.	23
<i>gossypiella</i> , Saund.	85	<i>harpophora</i> , Meyr.	157	<i>hexasticta</i> , Wals.	176
<i>gracilella</i> , Chamb.	177	<i>harrisonella</i> , Busck	43	<i>Heylaertsi</i> , Snell.	235
<i>gracilella</i> , Chrét.	44	<i>hastata</i> , Meyr.	98	<i>hiberna</i> , Meyr.	33
<i>gracilella</i> , Staint.	95	<i>hastella</i> , Chrét.	29	<i>hibiscella</i> , Busck.	81
<i>gradata</i> , Meyr.	137	<i>hedana</i> , Turn.	150	<i>hibisci</i> , Staint.	138
<i>grammitis</i> , Meyr.	102	<i>Hedemannia</i> , Car.	248	<i>hieracitis</i> , Meyr.	105
<i>Granti</i> , Wals.	179	<i>Helcystogramma</i> , Zell.	137	Hierangela (genus), Meyr.	100
<i>graphicella</i> , Busck	58	<i>heliacella</i> , Herr.-Sch.	46	<i>hieroglyphica</i> , Wals.	45
<i>graphicodes</i> , Meyr.	248	Heliangara (genus), Meyr.	216	<i>hieropla</i> , Meyr.	136
<i>gravata</i> , Meyr.	153	<i>helianthemi</i> , Wals.	177	<i>hilara</i> , Turn.	148
<i>gregalis</i> , Meyr.	91	<i>helipta</i> , Meyr.	214	<i>hilarella</i> , Car.	30
<i>gregariella</i> , Zell.	95	<i>heliarcha</i> , Meyr.	232	<i>hippastis</i> , Meyr.	208
<i>griseella</i> , Car.	83	<i>helicaula</i> , Meyr.	33	<i>hippeis</i> , Meyr.	95
<i>griseella</i> , Chamb.	196	Helice (genus), Chamb.	51	<i>hippica</i> , Meyr.	155
<i>griseella</i> , Hein.	43	<i>helicopis</i> , Meyr.	130	<i>hippocoma</i> , Meyr.	153
<i>grisella</i> , Chamb.	95	<i>heligmatodes</i> , Wals.	78	<i>hippophaella</i> , Schranck	81
<i>griseosericiella</i> , Rag.	202	<i>heliochares</i> , Low.	37	<i>hippotarcha</i> , Meyr.	229
<i>grisefasciella</i> , Chamb.	125	<i>heliocrina</i> , Meyr.	204	<i>hirculella</i> , Busck	175
<i>gudmannella</i> , Wals.	93	<i>heliopa</i> , Low.	94	<i>hirsutella</i> , Const.	122
Guebla (genus), Chrét.	65	<i>heliopepta</i> , Meyr.	251	<i>hispanella</i> , Reb.	201
<i>Guerini</i> , Staint.	100	<i>Helleria</i> , Reb.	202	<i>histrionella</i> , Hübn.	79
<i>guimarensis</i> , Wals.	42	<i>helotella</i> , Staud.	43	<i>Hoefneria</i> , Reb.	94
<i>guyonella</i> , Guen.	22	<i>heluanensis</i> , Debsk.	84	<i>hoffmanniella</i> , Strand.	98
<i>gypsaspis</i> , Meyr.	138	<i>helveolella</i> , Staud.	199	Holaxyra (genus), Meyr.	191
<i>gypsella</i> , Const.	94	<i>hemeropa</i> , Meyr.	191	Holcophora (genus) Staud.	166
<i>gypsocrana</i> , Turn.	149	<i>hemiacma</i> , Meyr.	235	Holcopogon (genus), Staud.	199
<i>gypsophilae</i> , Staint.	96	Hemarcha (genus), Meyr.	146	<i>holocyna</i> , Meyr.	209
<i>gyralea</i> , Meyr.	193	<i>hemichlaena</i> , Low.	144	<i>holomelas</i> , Low.	176
<i>habrias</i> , Meyr.	108	<i>hemichrysa</i> , Meyr.	239	Holophysa (genus), Wals.	130
<i>habrochitona</i> , Wals.	195	<i>hemileucas</i> , Meyr.	175	<i>holosericella</i> , Herr.-Sch.	82
Habrogenes (genus), Meyr.	212	<i>hemiopa</i> , Meyr.	249	<i>homaloxesta</i> , Meyr.	189
<i>haemaspila</i> , Low.	149	<i>hemiphaea</i> , Meyr.	108	Homaloxestis (genus), Meyr.	224
<i>haemylopis</i> , Meyr.	240	<i>hemiphaea</i> , Turn.	102	<i>homocentra</i> , Meyr.	239
<i>haeresiella</i> , Wall.	227	<i>hemisarca</i> , Low.	46	<i>homochlora</i> , Meyr.	213
<i>halimilignella</i> , Wals.	23	<i>hemixysta</i> , Meyr.	204	<i>homoclera</i> , Meyr.	254
<i>halimodendri</i> , Christ.	154	<i>henshawiella</i> , Busck	90	<i>homogenes</i> , Turn.	247
<i>halmyra</i> , Meyr.	121	<i>heptacentra</i> , Meyr.	30	<i>homogramma</i> , Meyr.	243
				<i>hoplitis</i> , Meyr.	34

LEPIDOPTERA HETEROCHERA

Pages		Pages		Pages
hoplomacha, Meyr.	102	Idiophantis (genus), Meyr.	108	inferialis, Meyr.
hoplophora, Meyr.	138	Idiopteryx (genus), Wals.	227	infernalis, Herr.-Sch.
hora, Busck	157	idioptila, Meyr.	153	infestella, Reb.
hordei, Kirb.	38	idiospila, Meyr.	147	infibulata, Meyr.
horiaula, Meyr.	76	ignavella, Zell.	63	inficeta, Meyr.
horides, Meyr.	175	igneella, Tengst.	30	infirma, Meyr.
horista, Wals.	116	ignobilieilla, Hein.	112	inflammatella, Christ.
horistis, Meyr.	228	ignorantella, Herr.-Sch.	84	inflatella, Chrét.
hormodes, Meyr.	149	ignorata, Meyr.	176	infracta, Wals.
Hornigi, Staud.	42	Illogiotis (genus), Meyr.	190	inficta, Meyr.
horogramma, Meyr.	59	illucescens, Meyr.	197	ingloria, Meyr.
horticolella, Rössl.	93	illuminata, Meyr.	33	ingloriella, Hein.
horticolla, Pey.	96	illustrella, Hübn.	107	ingravata, Meyr.
hortuella, Wood	83	ilmatarrella, Hoffm.	83	inhonesta, Meyr.
hortulana, Meyr.	189	ilyella, Zell.	90	innocuella, Zell.
hostilis, Meyr.	106	imbricata, Meyr.	177	innotatella, Chrét.
Howardi, Wals.	46	immaculatella, Dougl.	42	innotatella, Walk.
huebnerella, Don.	156	immerita, Meyr.	197	inopella, Zell.
Huebneri, Haw.	95	immeritella, Walk.	137	inopina, Meyr.
humeralis, Zell.	69	immobilis, Meyr.	240	inornatella, Dougl.
humerala, Schiff.	107	imparella, Fisch. v. Rösl.	34	Inotica (genus), Meyr.
Hungariae, Staud.	82	imperatrix, Meyr.	204	inquieta, Meyr.
Husadeli, Reb.	202	imperitella, Staud.	75	inquilinella, Busck
hutchinsonella, Wals.	79	impigra, Meyr.	177	inscripta, Wals.
hyacinthina, Meyr.	204	improba, Meyr.	156	inscriptella, Christ.
hyalombra, Meyr.	175	imprudens, Meyr.	238	inserrata, Wals.
hybrista, Meyr.	239	impudica, Wals.	93	insidians, Meyr.
hydrosema, Meyr.	209	impunctella, Car.	248	insignifcans, Wals.
Hygreplasta (genus), Meyr.	244	impurgata, Wals.	78	insipiens, Meyr.
hylias, Meyr.	140	inaequalis, Busck.	78	insomnis, Meyr.
Hylograpta (genus) Meyr.	177	inaequalis, Wals.	78	inspersa, Meyr.
hylurga, Meyr.	177	inaequipulvella, Chamb.	249	instabilella, Dougl.
Hydectis (genus), Meyr.	164	Inapha, Walk.	234	instabilella, Hein.
hyoscyamella, Reb.	78	incertella, Herr.-Sch.	29	instans, Meyr.
hyoscyamella, Staint	94	incitata, Meyr.	45	instaurata, Meyr.
Hypelictis (genus), Meyr.	100	incognita, Wals.	30	insuavis, Meyr.
Hyperecta (genus), Meyr.	132	incognitella, Car.	70	insularis, Butl.
Hyperochtha (genus), Meyr	227	incola, Meyr.	24	insularis, Staud.
hyperythra, Meyr.	113	incomptella, Herr.-Sch.	82	insularis, Wals.
hypochloa, Wals.	253	incondita, Meyr.	164	insulella, Dum.
hypocneca, Turn.	148	inconspicua, Wals.	250	insulella, Hein.
hypoleuca, Meyr.	148	inconspicuella, Murtf.	93	insulsa, Meyr.
Hyptiastis (genus), Meyr.	212	inculta, Wals.	153	intaminatella, Staint.
hypotiota, Meyr.	200	incurva, Meyr.	122	integrata, Meyr.
hystricella, Braun	248	indagata, Meyr.	201	intensa, Meyr.
		indica, Wals.	156	intentella, Walk.
ianthes, Meyr.	174	indigens, Meyr.	214	interalbicella, Herr.-Sch.
ichnota, Meyr.	118	indigna, Wals.	177	interguttella, Walk.
ichrodes, Meyr.	236	indignella, Staud.	75	intermedia, Braun
ichthyochroa, Wals.	44	indistinctella, Reb.	82	intermediella, Chamb.
idiastis, Meyr.	214	inermis, Meyr.	166	intermissella, Zell.
Idiobela, Turn.	30	inxperta, Meyr.	93	internella, Zell.
idiocentra, Meyr.	126	infelix, Wals.	30	interpolata, Meyr.

Pages	Pages	Pages			
interruptella, Hübn.	79	isomila, Meyr.	240	Klosi, Reb.	77
interruptella, Steph.	79	isomitra, Meyr.	239	knaggsiella, Staint.	95
interstratella, Christ.	46	isonira, Meyr	24	Kneri, Now.	249
intestina, Meyr.	92	isophanes, Turn.	238	Korbi, Car.	47
intestinella, Mann	30	Isophrictis (genus), Meyr.	35	kroesmanniella, Herr.-Sch.	95
intorta, Meyr.	105	isoptila, Meyr.	156		
inulella, Curt.	42	<i>isoscelixantha</i> , Low.	111	labradorica, Möschl.	142
inulella, Hübn.	75	isosema, Meyr	157	labradoriella, Clem.	76
inumbrata, Meyr.	112	isotis, Meyr.	34	laceratella, Ztl.	95
inurbana, Meyr.	254	isotricha, Meyr	156	laceratella, Wals.	84
inusta, Meyr.	121	isthmodes, Meyr.	200	Lachnostola (genus), Meyr.	171
inustella, Herr.-Sch.	94	istrella, Mann	78	lachtensis, Ersch.	112
invalida, Meyr.	148	Istrianis (genus), Meyr.	62	Lacistodes (genus), Meyr.	251
invariabilis, Kearn.	80	iterata, Meyr.	256	lacrimosa, Meyr.	63
invariella, Walk.	238	ithycosma, Meyr.	130	lactaria, Meyr.	124
invenustella, Berg	79	ithyxyla, Meyr.	40	lacteochrella, Chamb.	123
aversella, Zell.	192	itrinea, Meyr.	238	lacticaudella, Wals.	129
invictella, Busck	58	Iulota (genus), Meyr.	40	lacticeps, Wals.	76
invida, Meyr.	33	ivae, Busck	46	lacticoma, Meyr.	78
invisella, Const.	35			lactifera, Meyr.	156
involuta, Meyr.	95	Jakovlevi, Krul.	82	lactiflora, Meyr.	77
involuta, Turn.	149	jamaicensis, Wals.	92	lactiflorella, Chamb	253
Iochares (genus), Meyr.	184	janitrix, Meyr.	102	lacunosa, Meyr.	63
iochlaena, Meyr.	148	japonicella, Zell	250	lagopella, Herr.-Sch.	34
iodora, Meyr.	176	jaspidella, Chrét.	96	laguna, Busck	90
ioloncha, Meyr.	169	joannisiella, Rag.	25	lagunculariella, Busck	122
iomarmara, Meyr.	45	jordanella, Reb.	142	lagunella, Chrét.	93
iophaea, Meyr.	169	josianella, Walk.	232	lakatensis, Reb.	66
iophana, Meyr.	156	jugalis, Meyr.	239	lampetis, Meyr.	216
ioploca, Meyr.	134	jugata, Wals	189	lamprodesma, Meyr.	240
iospila, Meyr.	33	juncidella, Clem.	196	lampronialis, Walk.	235
iresia, Meyr.	238	junctella, Dougl.	96	lamprostoma, Zell.	138
iriantha, Meyr.	133	junctella, Herr.-Sch.	95	Lamprotes, Hein.	41
iridipennella, Clem.	130	junctivittella, Christ.	26	lanaiensis, Wals.	44
iriditis, Meyr.	71	juniperella, Kearn.	57	lanceolella, Zell.	34
iridosoma, Meyr.	138	juniperella, Linn.	177	languens, Meyr.	125
ironica, Meyr.	188	juridica, Meyr.	250	lapidella, Wals.	125
irrigata, Meyr.	39	justa, Meyr.	228	lapidescens, Meyr.	80
irrorella, Wals.	210	juventella, Wals.	116	lapillosa, Meyr.	80
isabella, Zell.	249			lappella, Linn.	30
Isohnoderis (genus), Meyr.	252	kearfottella, Busck	123	Larcophora (genus), Meyr.	241
ischnoptera, Meyr.	120	kearfottella, Busck	43	lasciva, Wals.	62
ischnoptila, Turn.	31	kefersteiniella, Zell.	35	Lata, Strand	89
ischnora, Turn.	40	kermella, Chrét.	113	latella, Herr.-Sch.	75
ischnota, Meyr.	35	Kiefferi, Strand	89	laterestriata, Wals.	102
Isochasta, Meyr.	41	<i>Kiesenwetteri</i> , Heuäck.	82	latescens, Wals.	192
isochorda, Meyr.	76	Killiasi, Frey	69	Lathontogenus, Wals.	168
isochra, Meyr.	209	kincaidella, Busck	77	lathridia, Meyr.	71
isoclera, Meyr.	192	<i>kindermannella</i> , Herr.-Sch.	200	lathyri, Staint.	63
isocola, Meyr.	219	kingerella, Herr.-Sch.	96	laticinctella, Wood.	126
isocosma, Meyr.	254	kinkerella, Snell.	23	latifasciella, Chamb.	72
isocrypta, Meyr.	235	kinzelella, Busck	24	latifasciella, Walk.	204
isogona, Meyr.	153	kittella, Wals.	58	latipalpis, Wals.	192

LEPIDOPTERA HETEROCHERA

Pages		Pages		Pages	
latistriga, Wals.	116	leucostictus, Wals.	198	livoniella, Teich	96
latiuscula, Hein.	43	leucostola, Meyr.	259	Locharcha (genus), Meyr.	86
latiusculella, Staint.	201	leucotella, Walk.	240	lochistis, Meyr.	249
Latrologa (genus), Meyr.	63	leucothicta, Meyr.	179	lochitis, Meyr.	224
laudatella, Wals.	95	leucothona, Meyr.	102	loemias, Meyr.	150
laurocistella, Chrét.	68	leucoxosta, Meyr.	24	logica, Meyr.	214
lavernella, Chamb.	90	leucura, Wals.	130	longicornella, Chrét.	220
laxata, Meyr.	235	Leuronema (genus), Meyr.	68	longicornis, Curt.	79
lechriosema, Turn.	150	levata, Meyr.	157	longifasciella, Clem.	72
Lecithocera (gen.), Herr.-Sch.	237	levigata, Meyr.	185	longinqua, Meyr.	110
lecticaria, Meyr.	211	levipedella, Clem.	120	longipalpella, Reb.	202
legalis, Meyr.	250	levisella, Fyles	196	longipalpella, Teich	82
legata, Meyr.	225	Lexiarcha (genus), Meyr.	145	longivitella, Reb.	29
lemniscella, Zell.	98	liberata, Meyr.	81	lophella, Busck	71
lemurella, Meyr.	23	liberta, Meyr.	250	loquax, Meyr.	91
lenta, Meyr.	91	libertinella, Zell.	47	lotella, Const.	174
lentiginosella, Zell.	84	libidinosa, Staud.	82	loxodesma, Meyr.	146
Leobatus (genus), Wals.	141	librata, Meyr.	217	loxogramma, Meyr.	122
Leonhardi, Krone	46	liciata, Meyr.	225	loxosaris, Meyr.	157
lepidella, Zell.	52	ligulella, Hübn.	175	lubrica, Meyr.	243
leptobrocha, Meyr.	228	ligulella, Zell.	112	lucernata, Meyr.	239
leptocentra, Meyr.	43	ligyra, Meyr.	189	lucida, Wals.	200
leptoconia, Turn.	23	limata, Turn.	110	lucidella, Steph.	44
Leptogenesia (genus), Meyr.	50	limbipunctella, Mill.	177	lucrifuga, Meyr.	175
leptoglypta, Meyr.	236	limbipunctella, Staud.	176	luctiferella, Herr.-Sch.	76
leptoicta, Turn.	148	limenaea, Meyr.	72	lucticella, Hübn.	76
leucaniella, Busck	75	limitanella, Reb.	82	luctuella, Dup.	76
leucatella, Clerck	58	limosella, Schläg.	177	luctuella, Hübn.	76
Leuo (genus), Chamb.	49	lindanella, Barnes	46	luctuella, Wood	71
leucocephala, Low.	93	lindenella, Busck.	78	luculella, Hübn.	71
leucocephala, Wals.	92	lineatella, Zell.	35, 154	lugens, Car.	126
leucoachalca, Meyr.	103	linella, Chrét.	112	lugubrella, Fabr.	76
leucohloria, Meyr.	212	lineolella, Zell.	248	luminosa, Busck	136
leucocosma, Meyr.	196	lingulacella, Clem.	47	lunariella, Wals.	77
leucocras, Wals.	105	lingulata, Meyr.	120	lunatella, Zett.	76
leucodoxa, Meyr.	77	linocoma, Meyr.	238	lupata, Meyr.	110, 115
leucogaea, Meyr.	23	Liolepta (genus), Meyr.	221	lupinella, Busck	122
leucoglypta, Meyr.	23	liota, Meyr.	148	luridella, Christ.	239
leucographa, Wals.	221	Lipatia, Busck	168	luridella, Teich	95
leucograpta, Meyr.	230	lipatiella, Busck	76	lusoria, Meyr.	183
leucomelanella, Zell.	96	lissota, Meyr.	197	lutatella, Herr.-Sch.	249
leucomella, Quens.	76	Lita, Treitsch.	73	luteella, Hein.	43
leucomichla, Meyr.	24	lithina, Low.	148	luteella, Snell.	204
leuconota, Zell.	54	lithochroma, Wals.	199	luteostrigella, Chamb.	105
leuconotella, Busck.	196	lithodelta, Meyr.	125	Intescens, Const.	75
leucopalpella, Herr.-Sch.	112	lithodes, Meyr.	80	luticarella, Zell.	239
leucophanta, Meyr.	45	lithodes, Wals.	80	luticostella, Chrét.	154
leucophora, Meyr.	153	lithomorpha, Meyr.	120	lutilabrella, Mann	80
leucoplecta, Meyr.	134	lithostrota, Meyr.	138	lutivittata, Meyr.	176
leucopleura, Meyr.	136	litigiosa, Meyr.	75	lutraula, Meyr.	75
leucorrhapta, Meyr.	121	litigiosella, Mill.	30	lutulentella, Zell.	44
leucoscia, Turn.	236	littorella, Dougl.	30	lynchnocentra, Meyr.	243
leucostena, Wals.	178	liturosella, Zell.	78	lyciella, Wals.	94

Pages	Pages	Pages			
lycopersicella, Wals.	45	manella, Möschl.	196	melanostictella , Rag.	71
lycopis, Meyr.	240	mangelivora, Wals.	122	melanostictella, Zell.	58
<i>lyellella</i> , Westw.	69	mappigera, Meyr.	66	melanota, Wals.	189
lygaea, Meyr.	244	maraschella, Car.	113	melanotephrella, Ersch.	96
lygropa, Low.	176	margarodes, Meyr.	102	melanozona, Meyr.	70
lynceella, Zell.	83	marginata, Wals.	240, 250	melantherella, Busck	196
lypetica, Wals.	175	marginella, Fabr.	176	melantypella, Mann.	77
lyrella, Wals.	249	marginipunctella, Staint.	52	meledantis, Meyr.	200
macaritis, Meyr.	216	mariae, Frey	77	melicentra, Meyr.	72
macella, Meyr.	239	marina, Meyr.	93	melicephala, Meyr.	250
machlas, Meyr.	210	mariona, Heinr.	72	melichroa, Meyr.	176
Machiotricha (genus), Meyr.	186	marmaritis, Meyr.	210	melicrata, Meyr.	224
Macrenches (genus), Meyr.	72	marmorata, Wals.	179	melicrata, Turn.	148
Macrernis, Meyr.	213	marmorea, Haw.	96	meliorella, Walk.	204
Macroceras, Staud.	197	marmorea, Low.	37	meliphanes, Low.	170
macrodelta, Meyr.	100	<i>marmorella</i> , Chamb.	94	melissia, Wals.	189
macroplaca, Low.	147	Martini, Petr.	42	melitopis, Meyr.	34
macroptera, Meyr.	175	maturata, Meyr.	184	melitoptila, Meyr.	67
macrosema, Low.	170	matutina, Meyr.	79	melitopyga, Meyr.	235
macrosphena, Meyr.	178	mauricaudella, Oberth.	113	Melitoxestis (genus), Meyr.	113
macrothecta, Meyr.	45	mazina, Meyr.	238	melitura, Meyr.	179
Macrotona, Meyr.	237	mechanistis, Meyr.	149	memnonia, Meyr.	189
macroxyla, Meyr.	188	meconitis, Meyr.	139	mendax, Wals.	44
Macrozancla, Turn.	174	mediatrix, Zell.	107	mendica, Turn.	176
macrynta, Meyr.	152	mediofuscella, Clem.	78	mendicella, Walk.	210
maculata, Staud.	47	<i>mediopallidum</i> , Wals.	200	mendosella, Zell.	201
maculata, Wals.	242	medjella, Chrét.	63	mendozella, Strand	89
maculata, Wals.	198	medullella, Busck	72	mercedella, Wals.	138
maculatella, Hübn.	78	Megaeraspedus (genus), Zell.	33	meridionella, Herr.-Sch.	35
maculaticornis, Wals.	44	megalarthra, Meyr.	151	meridionella, Wals.	177
maculea, Haw.	95	megalommata, Meyr.	150	Meridorma (genus), Meyr.	31
maculiferella, Dougl.	96	megalopis, Meyr.	239	Merimnetria (genus), Wals.	127
maculiferella, Staint.	96	megalornis, Meyr.	24	merismatella, Zell.	57
maculimarginella, Chamb.	83	megalosticta, Turn.	149	merista, Meyr.	32
maculosella, Herr.-Sch.	142	melagonella, Const.	112	mersinella, Staud.	71
magnanima, Meyr.	33	<i>melaleuella</i> , Const.	77	meryntis, Meyr.	207
magnella, Busck	35	melanamba, Meyr.	118	mesacta, Meyr.	79
magnetella, Staud.	75	melanaphra, Meyr.	43	mesochra, Low.	149
Magonympha (genus), Meyr.	208	melanardis, Meyr.	213	mesoctenis, Meyr.	176
majorella, Reb.	102	<i>melanarthra</i> , Low.	38	mesodelta, Meyr.	120
malachias, Meyr.	191	melanattracta, Meyr.	33	mesoglena, Meyr.	177
malacodes, Meyr.	171	melanecta, Meyr.	157	mesoleuca, Low.	63
malacogramma, Meyr.	248	melanella, Hein.	95	Mesophleps (genus), Hübn.	168
Malacotricha, Zell.	195	melanobathra, Meyr.	116	mesophracta, Turn.	171
malacta, Meyr.	239	melanocampta, Meyr.	93	mesopsamma, Turn.	150
maledicta, Meyr.	112	melanoleuca, Wals.	72	mesoxysta, Meyr.	43
<i>malifoliellus</i> , Fitch	175	melanombra, Meyr.	24	mesozosta, Meyr.	204
malindella, Busck	78	melanophylla, Turn.	176	Metabolaea (genus), Meyr.	160
malthacopa, Meyr.	195	melanoplecta, Meyr.	154	metacausta, Meyr.	239
malvella, Hübn.	85	melanoptila, Low.	83	metachalca, Meyr.	167
mancipata, Meyr.	156	melanosacta, Meyr.	108	metadupa, Wals.	120
mandella, Busck	77	melanosoma, Meyr.	188	metagramma, Meyr.	123
				metallanthes, Meyr.	211

LEPIDOPTERA HETEROCEA

Pages		Pages		Pages
metallica, Braun	75	Mnesteria (genus), Meyr.	204	muscosella, Zell.
metallica, Wals.	131	mniocosma, Meyr.	121	musculina, Staud.
Metanarsia (genus), Staud.	25	mochlopis, Meyr.	188	musicopa, Meyr.
metaphorica, Meyr.	157	modesta, Staud.	26	Musurga (genus), Meyr.
metasaris, Meyr.	238	modesta, Wals.	35	mutabilis, Meyr.
Meteorisitis (genus), Meyr.	143	modicella, Christ.	248	myadelpha, Meyr.
metochra, Meyr.	115	modicella, Dev.	111	Myconita (genus), Meyr.
metoeca, Meyr.	251	molestella, Zell.	46	mylicotis, Meyr.
Metopleura (genus), Busck	162	molitor, Wals.	92, 202	myopa, Meyr.
metorcha, Meyr.	158	mollis, Barnes	177	myricariella, Frey
metriodes, Meyr.	228	Melapostola (genus), Meyr.	118	Myrophila (genus), Meyr.
metrodes, Meyr.	176	molybdantha, Meyr.	207	myrrhina, Turn.
Metzneria (genus), Zell.	29	molybdias, Meyr.	230	Mythographa (genus), Meyr.
metzneriella, Staint.	30	molybdina, Wals	122	
mexicana, Wals.	175	molybdota, Meyr.	153	Naera, Chamb.
meyrickella, Busck	253	molyntis, Meyr.	150	<i>naevifera</i> , Dup.
miarodes, Meyr.	23	Mometa (genus), Durr.	85	<i>nana</i> , Low.
micella, Hüb.	46	monactis, Meyr.	43	nanella, Hüb.
micella, Schiff.	46	monella, Busck.	98	Nannodia, Hein
micradelpha, Low.	231	Monerista (genus), Meyr.	268	nanodella, Busck
micradelpha, Wals.	94	monetella, Feld.	207	nardinopa, Meyr.
microdora, Meyr.	148	monilella, Barnes	47	Narthecoceras (genus), Meyr.
microgramma, Meyr.	157	Monochroa, Hein.	41	<i>natarella</i> , Busck
micromela, Low.	238	monochroa, Wals.	30	<i>natalis</i> , Heinr.
micropa, Meyr.	149	monochromella, Const.	80	naufraga, Meyr.
microphanta, Meyr.	246	monochromella, Reb.	201	naulychna, Meyr.
microptila, Meyr.	143	monochromella, Walk.	122	nea, Wals.
microsema, Meyr.	250	monococca, Meyr.	188	Nealyda (genus), Dietz
microsperma, Meyr.	71	monodryas, Meyr.	244	neatodes, Meyr.
microsphena, Meyr	179	<i>monoleuca</i> , Low.	150	<i>nebulosa</i> , Steph.
microtima, Meyr.	24	monolorella, Reb.	34	<i>nebulosella</i> , Hein.
Milleri, Busck	58	monophragma, Meyr.	80	Neda, Chamb.
milleriella, Chamb.	93	montanella, Chrét.	95	nefasta, Meyr.
<i>millerellus</i> , Staint.	176	montivaga, Wals.	82	neglecta, Philp.
miltina, Durr.	231	monumentella, Chamb.	82	<i>neglectella</i> , Led.
miltophragma, Meyr.	188	morenella, Busck	77	negundella, Heinr.
mimella, Clem.	81	moreonella, Heinr.	58	nemophorella, Walk.
minimaculella, Chamb.	75	moritzella, Treitsch	95	neochalca, Meyr.
minimella, Car.	201	morologa, Meyr.	246	Neochrista (genus), Meyr.
minimella, Chamb.	43	morosa, Mühl.	42	Neocorodoe (genus), Meyr.
minor, Busck	93	morphochroma, Wals.	44	Neodactyota (genus), Busck
minorella, Hein.	75	<i>mouffetella</i> , Linn.	70	neograpta, Meyr.
minorella, Reb.	142	mouffetella, Schiff.	63	neosticta, Meyr.
mirabella, Staud.	23	mucronatella, Chrét.	96	neotrophella, Heinr.
mirabilis, Christ.	46	mulinella, Zell.	79	nephelombra, Meyr.
mirandella, Chrét.	46	multifasciella, Chamb.	57	nephelopis, Meyr.
mistipalpis, Wals.	105	multifida, Meyr.	106	nephelota, Meyr.
mitescens, Meyr.	154	multinotata, Meyr.	125	nephodesma, Meyr.
mitigata, Meyr.	246	multistrigella, Rag.	42	neptica, Wals.
mitophora, Turn.	148	mundana, Meyr.	157	<i>nerteria</i> , Meyr.
mitrella, Wals.	112	mundella, Dougl.	75	<i>nerterodes</i> , Meyr.
mixolitha, Meyr.	62	muricella, Chrét.	201	nesciatella, Walk.
Mnesistega (genus), Meyr.	216	murinella, Herr.-Sch.	95	nesidias, Meyr.

	Pages		Pages		Pages
Nesolechia (genus), Meyr.	59	noseropa, Turn.	238	ocelligera, Butl.	122
nessica, Wals.	196	Nesophistica (genus), Meyr.	232	ocellinella, Chrét.	81
neurograpta, Meyr.	249	notandella, Busck	98	ochnias, Meyr.	79
neurophora, Meyr.	122	notata, Wals.	44	ochraceella, Chrét.	94
neuropterella, Zell.	30	notatella, Hübn.	70	ochraula, Meyr.	33
Nevadia (genus), Car.	27	nothostigma, Meyr.	58	<i>ochricostella</i> , Chamb.	123
nigra, Busck	58	nothriformalis, Wals.	256	<i>ochreofuscella</i> , Chamb.	81
nigra, Haw.	83	Nothris (genus), Hübn.	97	<i>ochreopicta</i> , Wals.	202
<i>nigralbella</i> , Herr.-Sch.	52	nothrodes, Meyr.	150	<i>ochreostrella</i> , Chamb.	81
nigratomella, Clem.	117	nothropis, Meyr.	257	<i>ochreostrigella</i> , Chamb.	90
nigrella, Chamb.	112	notifica, Meyr.	214	<i>ochreosuffusella</i> , Chamb.	81
<i>nigricans</i> , Hein.	82	notospila, Meyr.	62	<i>ochreoviridella</i> , Pag.	136
nigricella, Chrét.	74	<i>novimundi</i> , Busck	200	<i>ochricapilla</i> , Reb.	47
nigriciliella, Wals.	44	nubigena, Meyr.	241	<i>ochrifoliata</i> , Wals.	70
<i>nigricostella</i> , Dougl.	63	nubivaga, Meyr.	246	<i>ochripalpella</i> , Frey	82
<i>nigricostella</i> , Dup.	63	nucifera, Wals.	80	<i>ochripalpella</i> , Zell.	139
nigridorsis, Meyr.	68	nugatricella, Reb.	113	<i>ochrisignella</i> , Nolck.	84
nigrimaculata, Braun	71	numidella, Chrét.	28	<i>ochrobyrsa</i> , Meyr.	249
nigrimaculella, Busck	83	nundinella, Zell.	82	<i>ochrocapna</i> , Meyr.	237
nigrinotella, Zell.	52	nyctias, Meyr.	148	<i>ochrocosma</i> , Meyr.	63
<i>nigripalpella</i> , Hein.	95	nyctiphylax, Turn.	236	<i>ochrodetata</i> , Meyr.	92
nigripectus, Wals.	83	obeliscota, Meyr.	149	<i>ochrophanes</i> , Meyr.	174
nigristrigella, Wocke	82	obfuscata, Meyr.	249	<i>ochropis</i> , Meyr.	86
<i>nigritella</i> , Staint.	111	objecta, Meyr.	71	<i>ochropyga</i> , Wals.	189
nigritella, Zell.	112	obligata, Busck	72	<i>ochrosceles</i> , Meyr.	225
nigromaculella, Mill.	94	<i>obligatella</i> , Walk.	228	<i>ochrosidera</i> , Meyr.	226
nigromaculella, Rag.	201	<i>obliquella</i> , Rag.	112	<i>ochrostoma</i> , Meyr.	249
nigrorosea, Wals.	77	<i>obliquella</i> , Wals.	227	<i>ochrotoma</i> , Meyr.	72
<i>nigrovittella</i> , Dup.	52	<i>obliquifasciella</i> , Chamb.	72	<i>ochroxesta</i> , Meyr.	189
nimbosa, Meyr.	154	<i>obliquistrigella</i> , Chamb.	57	<i>ochyrota</i> , Meyr.	250
niphastis, Meyr.	103	<i>obliterata</i> , Wals.	202	<i>ocreata</i> , Meyr.	228
niphaula, Meyr.	11	obruta, Meyr.	81	<i>octavana</i> , Meyr.	240
niphocentra, Meyr.	121	<i>obscurecinerea</i> , Nolck.	74	<i>octomaculella</i> , Chamb.	90
niphodes, Low.	34	<i>obscurella</i> , Chamb.	83	<i>octonias</i> , Meyr.	240
nitentella, Fuchs	93	<i>obscurella</i> , Fisch. v. Rösl.	122	<i>octophora</i> , Meyr.	249
nitidula, Staint	47	<i>obscurella</i> , Hein.	75	<i>ocymoidella</i> , Wals.	97
nitrariella, Chrét.	63	<i>obscuripennis</i> , Frey	98	<i>odorifera</i> , Meyr.	149
<i>nivea</i> , Haw.	52	<i>obscurocellella</i> , Chamb.	83	Oecla (genus), Wals.	197
niveisignella, Zell.	85	<i>obscurosuffusella</i> , Chamb.	82	Oecocecis (genus), Guen.	23
niveopulvella, Chamb.	123	<i>obsepta</i> , Meyr.	227	<i>oeconomica</i> , Meyr.	33
<i>nivella</i> , Wood	95	<i>obseratella</i> , Zell.	138	<i>oecophilala</i> , Staud.	198
nocturna, Meyr.	125	<i>obsignata</i> , Meyr.	239	Oegoconia (genus), Staint.	200
nocturnella, Staud.	95	<i>obsoleta</i> , Christ.	30	<i>oenochyta</i> , Meyr.	142
nodosa, Meyr.	235	<i>obsoletella</i> , Fisch. v. Rösl.	93	<i>oenombra</i> , Meyr.	188
Noeza (genus), Walk.	190	<i>obtrectata</i> , Meyr.	250	<i>oenophila</i> , Staud.	201
nomadella, Zell.	44	<i>obtruncata</i> , Meyr.	156	<i>Oeseis</i> , Chamb.	97
nomaditis, Meyr.	240	<i>obtusipalpis</i> , Wals.	165	Oestomorpha (genus), Wals.	160
Nomia, Clem.	41	<i>obvoluta</i> , Meyr.	103	<i>officialis</i> , Meyr.	239
nomias, Meyr.	92	<i>occidentella</i> , Chamb.	83	<i>officiosa</i> , Meyr.	251
nonstrigella, Busck	123	<i>oceania</i> , Meyr.	177	<i>olbiaella</i> , Mill.	79
nonstrigella, Chamb.	139	<i>ocellatella</i> , Boyd.	93	Olbothrepta (genus), Meyr.	209
nonyma, Turn.	93	<i>ocellella</i> , Chamb.	98	<i>oleata</i> , Meyr.	177
Norvegiae, Strand.	82				

LEPIDOPTERA HETEROCHERA

Pages	Pages	Pages			
olga, Meyr.	135	Oxybelia, Hübn.	174	palustrella, Dougl.	43
olivescens, Meyr.	179	oxybiella, Mill.	202	pammacha, Meyr.	33
olivierella, Rag.	21	oxycedrella, Mill.	169	pamphaea, Meyr.	42
olympiadella, Zell.	76	oxycona, Meyr.	240	panaula, Meyr.	79
omachella, Oberth.	92	Oxycriptis (genus), Meyr.	179	pancalala, Turn.	231
omoptila, Meyr.	154	Oxygnostis (genus), Meyr.	206	pancaliella, Staud.	46
omphacias, Meyr.	238	oxygramma, Meyr.	102	panchlora, Meyr.	246
omphalopa, Meyr.	88	oxygrapha, Meyr.	188	Pancoenia (genus), Meyr.	151
oncodes, Meyr.	112	Oxylechia (genus), Meyr.	52	pancratiastis, Meyr.	103
onctera, Wals.	193	oxyphanes, Meyr.	34	panella, Busck	77
Onebala (genus), Walk.	137	oxyprora, Meyr.	100	panglycera, Turn.	102
onzella, Christ.	26	Oxypteryx, Reb.	142	Panicotricha (genus), Meyr.	257
opaula, Meyr.	79	Oxysactis, Meyr.	188	<i>pannosella</i> , Reb.	201
operaria, Meyr.	103	oxyspila, Meyr.	111	panormitella, Car.	122
operculella, Zell.	93	oxystola, Meyr.	105	pantalaena, Wals.	43
operosa, Meyr.	33	oxytheces, Meyr.	102	pantheropa, Meyr.	249
ophitis, Meyr.	76	oxytypa, Meyr.	158	paphia, Meyr.	45
ophitis, Wals.	115	oxyzona, Meyr.	230	paphlactis, Meyr.	83
opificella, Mann	95	Pachniatia (genus), Meyr.	245	Pappophorus (genus), Wals.	172
opisthopa, Meyr.	29	Pachygenia (genus), Meyr.	49	Parachronistis (genus), Meyr.	52
oppletella, Herr.-Sch.	82	pachynitis, Meyr.	234	paracma, Meyr.	106
opportuna, Meyr.	238	Pachysaris (genus), Meyr.	186	paraconta, Meyr.	102
opsonoma, Meyr.	106	pacifica, Meyr.	148	paradesma, Meyr.	44
orbata, Meyr.	237	packardella, Chamb.	77	Paradoris, Meyr.	200
oreocyrniella, Petr.	94	paenitens, Meyr.	186	Paralechia, Busck	59
Organitis (genus), Meyr.	243	palacta, Meyr.	201	Parallactis (genus), Meyr.	246
orgilopis, Meyr.	72	palathodes, Meyr.	209	parallactis, Meyr.	240
oribatis, Meyr.	45	<i>palaris</i> , Meyr.	38	paralogella, Busck	75
ornatifimbriella, Clem.	84	palermitta, Lah.	94	Paranarsia (genus), Rag.	25
ornatipalpella, Wals.	58	palinorsa, Meyr.	105	Paranoea (genus), Wals.	192
oronella, Wals.	70	Pallintropa (genus), Meyr.	155	parapleura, Meyr.	80
orophota, Meyr.	106	palirrhoa, Meyr.	115	paraplutella, Busck	75
orphnella, Reb.	202	pallacopa, Meyr.	144	Parapodia, Joan.	41
orsicoma, Meyr.	52	pallicornella, Staud.	236	Parapsectria (genus), Meyr.	65
orsoviella, Hein.	230	pallida, Staud.	202	paraptila, Meyr.	108
orthadia, Meyr.	210	pallidella, Chamb.	35	parasema, Meyr.	240
orthanotis, Low.	140	pallidella, Hein.	92	Parasia, Dup.	29
orthiastis, Meyr.	103	palliderosacella, Chamb.	70	parasira, Meyr.	254
orthocampta, Meyr.	118	pallidibasella, Rag.	135	Paraspistes, Meyr.	168
orthostenis, Meyr.	254	pallidigriseella, Chamb.	82	Parastega (genus), Meyr.	86
orthodesma, Low.	37	pallidipulchra, Wals.	142	parasticta, Meyr.	145
orthogonella, Staud.	42	pallidistrigella, Chamb.	35	paraula, Meyr.	70
orthophracta, Meyr.	120	pallidochrella, Chamb.	51-60	Parelectra (genus), Meyr.	120
Orthoptila (genus), Meyr.	151	pallipalpella, Snell.	82	Parelliptis (genus), Meyr.	217
orthostathma, Meyr.	157	<i>pallipalpis</i> , Walk.	196	<i>parenthesella</i> , Haw.	107
oschophora, Meyr.	61	palpialbella, Chamb.	44	<i>paripunctella</i> , Thunb.	71
osseella, Staint.	40	<i>palpiannulella</i> , Chamb.	44	Paristhomia (genus), Meyr.	187
ostariella, Wals.	57	palpigera, Wals.	169	parmata, Meyr.	122
ostensella, Walk.	189	palpilineella, Chamb.	112	parmulata, Meyr.	115
ostracodes, Meyr.	170	paltobola, Meyr.	102	parochra, Meyr.	211
ouedella, Chrét.	43	Paltedora (genus), Meyr.	36	paroetas, Meyr.	157
ovulata, Meyr.	163	paltodoriella, Busck.	120	paroena, Meyr.	240
oxalea, Meyr.	240			paroristis, Meyr.	240

Pages	Pages	Pages			
parthenica, Meyr.	103	percnacma, Meyr.	178	petrinodes, Meyr.	93
particulata, Meyr.	157	percnobela, Meyr	240	petrogenes, Wals.	202
partitella, Christ.	70	percnoleuca, Meyr.	253	petrographa, Meyr.	125
parvella, Chrét.	220	percnopholis, Wals.	175	petromorpha, Meyr.	120
parvipulex, Wals.	92	percnospila, Meyr.	122	petrotoma, Meyr.	256
<i>parvipulvella</i> , Chamb.	249	perdita, Low.	93	Petryi, Hofm.	96
parvula, Staud.	43	perfida, Meyr.	239	petulans, Meyr.	55
pascuicola, Staud.	80	perfidiosa, Meyr.	93	phacelota, Meyr.	157
pastas, Meyr.	210	perfossa, Meyr.	45	phaeochlora, Meyr.	220
paterata, Meyr.	45	pergrandella, Reb.	80	phaeopa, Meyr.	111
Patouissa, Walk.	237	periastra, Meyr.	230	phaeostrota, Meyr.	172
patruella, Mann	112	peribapta, Low.	43	phaeotoxa, Meyr.	120
patulella, Walk.	153	perichlora, Meyr.	225	phaetus, Chamb.	53
<i>pauciguttellus</i> , Clem.	175	periculella, Busck	77	phagnalella, Const.	94
<i>paucipunctella</i> , Dougl.	30	perigrapta, Meyr.	231	phalacra, Wals.	87
paucipunctella, Zell.	30	perigypsa, Meyr.	240	phalacodes, Meyr.	94
paulella, Busck	77	perinaeta, Wals.	118	phalaritis, Meyr.	206
<i>paupella</i> , Zell.	42	periochra, Meyr.	246	phanerozona, Meyr.	48
<i>pauperella</i> , Hein.	95	<i>perionella</i> , Walk.	210	phanocrossa, Meyr.	115
pauperella, Reb.	239	Perloristica (genus), Wals.	136	Phanoschista (genus), Meyr.	207
Pauroneura (genus), Turn.	108	periphora, Meyr.	151	phantasta, Meyr.	24
pausimacha, Meyr.	106	Peripherectis (genus), Meyr.	235	Pharangitis (genus) Meyr.	219
Pazsiezkyi, Reb.	94	peristylis, Meyr.	176	pharetrata, Meyr.	205
pectinalella, Chamb.	49	peritella, Const.	70	pharetria, Meyr.	80
Pectinophora, Busck	84	peritura, Meyr.	106	phasianis, Meyr.	150
peculella, Busck	120	perixantha, Meyr.	257	Phatnotis (genus), Meyr.	225
<i>pedisequella</i> , Hüb.	63	perizeucta, Meyr.	225	phaulodes, Meyr.	256
pedmontella, Chamb.	90	perlatella, Walk.	120	phauloptila, Turn.	40
pelaea, Meyr.	257	<i>perlevis</i> , Turn.	176	phaulosema, Meyr.	69
pelieilla, Treitsch.	84	perlustrata, Meyr.	33	phelotris, Meyr.	92
pelitis, Meyr.	189	permagna, Meyr.	122	phepsalitis, Meyr.	121
pellax, Meyr.	238	<i>permolestella</i> , Busck	51	Philarachnis (genus), Meyr.	247
pelodes, Meyr.	256	permundella, Walk.	189	philodoxa, Meyr.	206
pelogramma, Meyr.	149	pernigrella, Dougl.	122	philamus, Meyr.	249
peloptila, Meyr.	125	perobscura, Wals.	254	Philoptila (genus), Meyr.	233
pelota, Meyr.	151	<i>perobscurella</i> , Wals.	83	philosopha, Meyr.	238
peltosema, Low.	47	peronectis, Meyr.	79	Phloeocecis, Chrét.	88
pendens, Meyr.	92	peronetris, Meyr.	211	phloeodes, Meyr.	148
<i>penessa</i> , Meyr.	199	perpensa, Meyr.	238	Phloeographis (genus), Meyr.	152
penetrans, Meyr.	57	perpetuella, Herr.-Sch.	77	phloecropa, Turn.	149
penicillata, Wals.	45	perpygmaella, Wals.	202	<i>phoebusella</i> , Mill.	74
pennella, Busck	35	perquisita, Meyr.	125	phoenaula, Meyr.	79
pennyanica, Dietz	79	persicella, Murtf.	84	phorcis, Meyr.	210
pensilis, Meyr.	153	perspersella, Wocke	77	phorticata, Meyr.	153
pentacentra, Meyr.	33	perspicua, Wals.	70	phosphoropa, Meyr.	114
<i>pentadora</i> , Meyr.	67	perterrita, Meyr.	43	photinopa, Meyr.	34
pentagramma, Meyr.	106	perumbrata, Meyr.	250	Photodotis (genus), Meyr.	50
pentasta, Meyr.	120	Pessograpta (genus), Meyr.	158	phoxopterella, Snell.	183
pentheres, Wals.	34	petasis, Pfaff.	80	phryganitis, Meyr.	248
penthicles, Meyr.	149	Peterseni, Teich	75	phthoneropa, Meyr.	223
peracuta, Meyr.	34	petiginella, Mann	63	Phthoracma (genus), Meyr.	222
peragrata, Meyr.	161	petraea, Wals.	80	Phthorimaea (genus), Meyr.	91
perceptella, Busck	136	petrella, Busck	90	phycidella, Walk.	210

LEPIDOPTERA HETEROCHERA

	Pages		Pages		Pages
Phylopatria (genus), Meyr.	56	pleuroleuca, Turn.	176	praecipitata, Meyr.	112
physaliella, Chamb.	44	pleuropa, Meyr.	189	praedicata, Meyr.	71
Physalivorella , Chamb.	90	pleurophaea, Turn.	175	praefixa, Braun	72
physeta, Meyr.	188	pleurosaris, Meyr.	57	praenivea, Meyr.	120
phytomiella, Busck	125	plexigramma, Meyr.	180	praeramis, Meyr.	200
piceella, Kearf.	58	plicata, Wals.	253	praeses, Meyr.	240
picrocarpa, Meyr.	185	plinthactis, Meyr.	149	praesignis, Meyr.	99
picrodora, Meyr.	238	plinthodes, Low.	81	praevacua, Meyr.	176
picrophanes, Meyr.	192	plocamandra, Meyr.	225	pragmatica, Meyr.	145
picyntis, Meyr.	219	plumata, Meyr.	45	Pragmatodes (genus), Wals.	48
pictella, Zell.	47	plumbella, Hein.	42	prasina, Meyr.	125
picticornis, Wals.	121	plumbeolata, Wals.	122	prasinantha, Meyr.	161
picula, Wals.	58	plumbosa, Meyr.	196	prasinoleuca, Meyr.	71
Pilocrates (genus), Meyr.	159	plutella, Chamb.	34	prasinopsis, Meyr.	231
pilosella, Walk.	156	<i>plutella</i> , Chamb.	54, 196	prasinoptila, Meyr.	124
pinella, Busck	58	plutelliformis, Snell.	139	praticolella, Christ.	94
pinguinella, Treitsch.	83	plutelliformis, Staud.	79	pravinominella, Chamb.	81
pinifoliella, Chamb.	60	plutonella, Heinr.	53	prensans, Meyr.	175
piperata, Wals.	175	poecilopa, Meyr.	103	Pribitzeri , Reb.	83
piscipellis, Zell.	35	poecilosoma, Wals.	135	primella, Busck	107
pisoniae, Busck	24	Pegochaetia (genus), Staud.	97	primigenia, Meyr.	126
Pithanurga (genus), Meyr.	68	pogonias, Meyr.	230	primipilana, Meyr.	47
Pityeonra (genus), Meyr.	31	poliocoma, Meyr.	238	princeps, Busck	90
pityritis, Meyr.	34	polioleuca, Turn.	147	prioleuca, Meyr.	76
Placanthes (genus), Meyr.	232	poliombra, Meyr.	125	prisca, Meyr.	148
placidella, Zell.	43	poliopasta, Turn.	24	prismatopa, Meyr.	118
placoterra, Meyr.	115	politella, Staint.	75	pristica, Meyr.	211
pladarota, Meyr.	177	polostella, Busck	53	Proactica , Wals.	22
plaesiodes, Meyr.	246	polyaema, Meyr.	194	proaula, Meyr.	227
plaesiosema, Turn.	93	polyaula, Meyr.	103	probata, Meyr.	72
Plagiferella , Walk.	164	polychromella, Reb.	112	probolaea, Meyr.	156
planata, Meyr.	191	polygramma, Meyr.	81	probolopis, Meyr.	45
planodes, Meyr.	83	Polyhymno (genus), Chamb.	105	procax, Meyr.	156
plantaginella, Staint.	92	polyommata, Turn.	176	procentra, Meyr.	213
plantariella, Tengst.	75	pomaceella, Walk.	125	prochalina, Meyr.	50
plastica, Meyr.	174	<i>pometella</i> , Fitch.	175	Procharista (genus), Meyr.	225
Platanilla , Chamb.	77	ponerias, Wals.	202	Proclesis, Wals.	183
platiastis, Meyr.	120	popularis, Meyr.	35	proclivella, Fuchs	94
platyaula, Meyr.	104	populella, Clerck	126	proclivis, Meyr.	238
platycapna, Meyr.	209	<i>populella</i> , Hüb.	83	procrossa, Meyr.	197
platyconta, Meyr.	214	porcella, Hein.	92	procursella, Reb.	249
platydoxa, Meyr.	75	porphyraspis, Meyr.	134	procypoides, Meyr.	188
Platyedra (genus), Meyr.	84	porphyrogamma, Meyr.	195	Prodesiaroha (genus), Meyr.	146
platyleuca, Low.	145	porphyroloma, Low.	144	profusa, Meyr.	63
platyleuca, Meyr.	34	Porpodryas (genus), Meyr.	161	prographa, Meyr.	258
platysoma, Wals.	63	<i>portlandicella</i> , Rich.	75	prograpta, Meyr.	159
platyxypha, Meyr.	106	<i>portosanctana</i> , Staint.	93	<i>prohaskeella</i> , Reb.	46
plebeiella, Zell.	75	postpallescens, Wals.	87	prolapsa, Meyr.	112
plectanopa, Meyr.	56	potosi, Busck	162	prolocha, Meyr.	220
Plectrocossa (genus), Meyr.	103	praealbescens, Meyr.	194	prominula, Meyr.	45
plemochoa, Meyr.	95	praeceptrix, Meyr.	210	Promelezia (genus), Meyr.	118
pleostigmella, Reb.	201	<i>praecarella</i> , Hein.	82	promptella, Staud.	92
pleroma, Wals.	81	<i>praecarella</i> , Herr.-Sch.	82	propensa, Meyr.	228

	Pages		Pages		Pages
Prophoraula (genus), Meyr.	181	pteropis, Meyr.	102	pyretodes, Meyr.	182
propitia, Meyr.	239	ptilastis, Meyr.	58	pyrochorda, Meyr.	231
prorepta, Meyr.	76	ptilocompa, Meyr.	189	pyrocosma, Meyr.	136
proscripta, Meyr.	150			pyrodercia, Wals.	45
Proselotis (genus), Meyr.	30			Ptilostonychia (genus), Wals.	253
proserpinella, Frey	130	ptochas, Meyr.	237	Ptilothrys (genus), Wals.	226
Prosodarma (genus), Meyr.	244	Ptocheuusa, Hein.	41	<i>pyropella</i> , Hübñ.	168
prosoestea, Turn.	102	ptochodryas, Meyr.	250	pyrophanes, Meyr.	206
Prosmura, Turn.	256	ptochogramma, Meyr.	121	<i>pyropis</i> , Meyr.	181
Prostomeus (genus), Busck	140	ptochomorpha, Meyr.	236	pyrphora, Meyr.	102
Protobathra (genus), Meyr.	258			<i>pyrrhanthes</i> , Meyr.	93
Protolechia (genus), Meyr.	147			pyrrhella, Rag.	201
Protolychnis (genus), Meyr.	242	Ptycerata (genus), Ely	37	pyrrhica, Turn.	149
protoma, Meyr.	238	<i>ptychophora</i> , Meyr.	58	pyrrhitis, Meyr.	188
protrocha, Meyr.	240	ptychosema, Meyr.	179	pyrrhopis, Meyr.	181
protypa, Meyr.	257	ptyoptera, Meyr.	99	pyxinodes, Meyr.	239
provinciella, Staint.	96	<i>pudibundella</i> , Chamb.	45		
<i>proxima</i> , Haw.	96	<i>pudibundella</i> , Zell.	46	<i>quadrella</i> , Fabr.	76
proximella, Hübñ.	70	pudicella, Mann	189	<i>quadrifariella</i> , Mann	202
prozona, Meyr.	66	puertella, Busck	80	<i>quadrifascia</i> , Walk.	120
prudens, Clem.	61	pulcherrimella, Walk.	240	<i>quadrimaculata</i> , Wals.	131
prudens, Meyr.	240	pulchra, Wcll.	96	<i>quadrimaculella</i> , Chamb.	81, 123
pruinosa, Chrét.	165	pulicella, Wals.	44	<i>quadrinella</i> , Herr.-Sch.	173
pruinosa, Zell	63	<i>pullatella</i> , Hübñ.	154	<i>quadripuncta</i> , Haw.	200
<i>pruniella</i> , Clem.	154	pullatella, Tengst.	96	<i>quadripunctella</i> , Chrét.	113
<i>pruniella</i> , Hübñ.	58	pullella, Chamb.	43	<i>quaestionella</i> , Herr.-Sch.	42
prunifoliella, Chamb.	54	pullifimbriella, Clem.	250	querella, Lah.	126
psacasta, Meyr.	33	pulveratella, Herr.-Sch.	43	quercicola, Meyr.	176
psamathias, Meyr.	44	pulvherea, Braun	46	querciella, Chamb.	71
Psamathocrita (genus), Meyr.	40	<i>pulverellus</i> , Const.	199	<i>querciella</i> , Chamb.	176
psameticella, Reb.	199	pulverosella, Chrét.	22	<i>quercifoliella</i> , Chamb.	72
psammitella, Snell.	82	punctata, Staud.	94	<i>quercinigrella</i> , Chamb.	71
psammolitha, Meyr.	253	punctatella, Staud.	44	<i>quercipomonella</i> , Chamb.	175
Psammoris (genus), Meyr.	215	punctatella, Walk.	175	<i>quercivorella</i> , Chamb.	58
psaphara, Meyr.	257	punctidiscella, Clem.	177	queribunda, Meyr.	225
pselaphistis, Meyr.	226	punctigeneralis, Walk.	240	querula, Meyr.	239
psephias, Meyr.	148	punctipennella, Clem.	177	quinella, Zell.	76
pseudacaciella, Chamb	83	pungens, Meyr.	122	<i>quinqueannulella</i> , Chamb.	61
Pseudochelaria, Dietz	73	purpurea, Wals.	227	<i>quinquecristatella</i> , Chamb.	69
Pseudocrates (genus), Meyr.	216	<i>purpureofusca</i> , Wals.	139	quinquedentata, Wals.	70
pseudofondella, Busck	78	pusilla, Wals.	34	quinquepunctella, Busck	44
pseudolella, Christ.	96	pusillella, Reb.	92	<i>quinquepunctella</i> , Herr.-Sch.	30
pseudometra, Meyr.	189	putella, Busck	57	quinquepunctella, Wals.	126
psilella, Herr.-Sch.	94	puteolata, Meyr.	240		
psilodoxa, Meyr.	161	Pycnobathra , Low.	33	<i>radiatella</i> , Busck	90
psiloptera, Barnes	84	pycnoda, Low.	81	<i>radicata</i> , Meyr.	45
psimythota, Meyr.	76	pycnode, Meyr.	99	<i>radiella</i> , Krul.	71
<i>psolosticta</i> , Turn.	23	Pycnodytis (genus), Meyr.	39	<i>radiosella</i> , Ersch.	248
psoralella, Mill.	112	pycnospila, Turn.	236	<i>rancidella</i> Herr.-Sch.	95
psoraliella, Barnes.	122	Pycnostola (genus), Meyr.	32	<i>rasilella</i> , Herr.-Sch.	249
Psoricoptera, Staint.	155	pygmaea, Turn.	151	<i>rastrifera</i> , Meyr.	112
psoricopterella, Wals.	142	pygmaeella, Hein.	95	<i>ratella</i> , Herr.-Sch.	202
psychrodes, Meyr.	23	pylartis, Meyr.	100	<i>rebeliella</i> , Haud.	80
		<i>pyramidophora</i> , Turn.	47	<i>reciproca</i> , Meyr.	154
		pyrenaica, Petr.	77	<i>recta</i> , Meyr.	120

LEPIDOPTERA HETEROCHERA

	Pages		Pages		Pages
<i>recticostella</i> , Wals.	254	<i>rhodopa</i> , Meyr.	102	<i>ruderella</i> , Chamb.	175
<i>rectifasciella</i> , Fuchs	42	<i>rhodopetala</i> , Meyr.	133	<i>rufescens</i> , Haw.	249
<i>rectistrigella</i> , Barnes	80	<i>rhodophaea</i> , Meyr.	179	<i>rufitecta</i> , Meyr.	118
Recurvaria (genus), Haw.	57	<i>rhodoptera</i> , Mann	78	<i>rufulella</i> , Snell.	43
<i>recurvata</i> , Meyr.	237	<i>rhodota</i> , Meyr.	201	<i>rufusella</i> , Chamb.	122
<i>rsedella</i> , Chamb.	175	<i>rhoifructella</i> , Clem.	123	<i>rumicetella</i> , Hofm.	42
<i>refracta</i> , Meyr.	120	<i>rhombella</i> , Schiff.	81	<i>rumicivorella</i> , Mill.	80
<i>regia</i> , Meyr.	133	<i>rhombelliformis</i> , Staud.	81	<i>ruptella</i> , Const.	82
<i>relaxata</i> , Meyr.	256	<i>rhombica</i> , Meyr.	121	<i>rurigena</i> , Meyr.	186
<i>religata</i> , Meyr.	120	<i>rhomboidella</i> , Curt.	156	<i>rustica</i> , Wals.	175
<i>religiosa</i> , Meyr.	143	<i>rhombophorella</i> , Zell.	57	<i>rusticana</i> , Meyr.	240
<i>remissa</i> , Zell.	44	Rhynchosopacha (gen.), Staud.	165	<i>rusticella</i> , Walk.	138
<i>renascens</i> , Wals.	196	Rhynchotona (genus), Meyr.	172	<i>rutilella</i> , Snell.	47
<i>renigerella</i> , Zell.	107	<i>rhyodes</i> , Meyr.	199	<i>sabinella</i> , Zell.	98
<i>repandella</i> , Walk.	121	<i>rhypara</i> , Wals.	253	<i>sabulella</i> , Wals.	35
<i>reparabilis</i> , Wals.	253	<i>rhyptodes</i> , Wals.	79	<i>sabulosella</i> , Reb.	95
<i>repentella</i> , Chrét.	96	<i>ribbeella</i> , Car.	28	<i>sacricola</i> , Meyr.	134
<i>repentina</i> , Wals.	78	<i>ribbeella</i> , Zell.	138	Sagaritis , Chamb.	174
<i>repudiata</i> , Meyr.	42	<i>ribeella</i> , Chamb.	77	<i>saginella</i> , Zell.	96
<i>rescissella</i> , Zell.	76	<i>rileyella</i> , Chamb.	181	<i>sagittans</i> , Meyr.	107
<i>resecta</i> , Meyr.	71	<i>ripula</i> , Wals.	72	<i>sagittaria</i> , Meyr.	154
<i>residua</i> , Meyr.	235	<i>Ritsemae</i> , Snell.	234	<i>sagittella</i> , Car.	52
<i>resinosa</i> , Meyr.	45	<i>rivalis</i> , Meyr.	126	<i>sagittifera</i> , Low.	34
<i>resoluta</i> , Meyr.	250	<i>ritsella</i> , Möschl.	87	<i>sagmatica</i> , Meyr.	153
<i>responsa</i> , Meyr.	238	<i>robertsonella</i> , Curt.	154	<i>saharae</i> , Oberth.	202
<i>retamella</i> , Chrét.	154	<i>robinella</i> , Chrét.	35	<i>saharae</i> , Wals.	199
<i>retecta</i> , Meyr.	71	<i>robiniacfoliella</i> , Chamb.	57	<i>salebrosa</i> , Meyr.	120
<i>retiniella</i> , Barnes	81	<i>robiniella</i> , Fitch	57	<i>salicifungiella</i> , Clem.	45
<i>retracta</i> , Meyr.	196	<i>robusta</i> , Braun	43	<i>salicorniae</i> , Her.	92
<i>retractella</i> , Walk.	258	<i>robusta</i> , Butl.	199	<i>salinae</i> , Wals.	254
<i>retusella</i> , Reb.	43	<i>robustella</i> , Reb.	249	<i>salinaris</i> , Busck	90
<i>reussella</i> , Ratz.	60	<i>robustella</i> , Staud.	43	<i>salinella</i> , Zell.	94
<i>Reutitia</i> , Hofm.	101	<i>robustella</i> , Walk.	87	<i>salmonis</i> , Wals.	74
<i>reuttiella</i> , Hein.	83	<i>Rogenhoferi</i> , Staud.	46	<i>altenella</i> , Schöy.	92
<i>reversa</i> , Meyr.	258	<i>rosalbella</i> , Fol.	77	<i>saltuum</i> , Zell.	70
<i>revoluta</i> , Meyr.	71	<i>roseella</i> , Zett.	95	<i>salva</i> , Meyr.	92
<i>rhabdodes</i> , Wals.	126	<i>roseicrinella</i> , Busck	107	<i>samadensis</i> , Pfaff.	93
<i>rhabducba</i> , Meyr.	137	<i>roseocostellus</i> , Wals.	177	<i>sandaracota</i> , Meyr.	133
Rhadinophylla (genus), Turn.	173	<i>roseosuffusella</i> , Clem.	46	<i>sandycitis</i> , Meyr.	194
<i>rhedaria</i> , Meyr.	78	<i>rosmarinella</i> , Wals.	202	<i>sangiella</i> , Staint.	112
<i>rhenanella</i> , Heyd.	42	<i>rostella</i> , Feld.	115	<i>sanguinolentella</i> , Joan.	30
<i>rhicnota</i> , Meyr.	156	<i>rostrata</i> , Meyr.	213	<i>santolinae</i> , Staud.	107
<i>rhicnota</i> , Wals.	57	<i>rostrifera</i> , Meyr.	55	<i>saphirinella</i> , Chamb.	90
<i>Rhinosia</i> , Treitsch.	174	<i>rotundata</i> , Wals.	116	<i>sapphiritis</i> , Meyr.	130
<i>rhizogramma</i> , Meyr.	173	<i>Rougemonti</i> , Reb.	96	<i>sarcochroma</i> , Wals.	85
<i>rhizophora</i> , Meyr.	209	<i>rubensella</i> , Chamb.	45	<i>sarcodes</i> , Wals.	47
<i>Rhobonda</i> , Walk.	174	<i>rubentula</i> , Meyr.	245	<i>sarcographa</i> , Meyr.	79
<i>rhodantha</i> , Meyr.	231	<i>rubescens</i> , Wals.	122	<i>sardonias</i> , Meyr.	225
<i>rhodochra</i> , Meyr.	79	<i>rubida</i> , Turn.	150	<i>sarisias</i> , Meyr.	148
<i>rhodocosma</i> , Meyr.	48	<i>rubidella</i> , Chrét.	96	Sarlosphera (genus), Meyr.	236
<i>rhodocrossa</i> , Meyr.	245	<i>rubidella</i> , Clem.	45	<i>sarothamnella</i> , Zell.	112
<i>rhodogramma</i> , Meyr.	106	<i>rubidula</i> , Meyr.	196	Sarotorna (genus), Meyr.	97
<i>rhodomicta</i> , Meyr.	103	<i>rubiginosella</i> , Walk.	189		

Pages	Pages	Pages			
sartor, Wals.	58	scopulata, Meyr.	121	senectella, Zell.	75
Sathrogenes (genus), Meyr.	191	scopulosa, Meyr.	156, 228	senicula, Meyr.	35
Satrapodexa (genus), Meyr.	132	scordiscella, Reb.	44	senilis, Meyr.	214
saturata, Meyr.	175	scotaea, Meyr.	103	senticetella, Staud.	98
saturnina, Meyr.	45	scotia, Turn.	157	separabilis, Wals.	128
saulotis, Meyr.	102	scotinella, Herr.-Sch.	82	separatella, Fisch. v. Rösl.	34
saundersella, Chamb.	63	scotodes, Wals.	84	sepcolella, Herr.-Sch.	42
sauteriella, Zell.	76	scriniata, Meyr.	156	sepiella, Staud.	248
saxeia, Meyr.	57	scriptella, Dup.	202	septella, Zell.	138
scabidella, Zell.	77	scriptella, Hübn.	71	septentrionella, Busck	253
scabra, Staud.	69	scutata, Meyr.	138	septentrionella, Fyles	90
scabrella, Busck	84	scutella, Zell.	121	sequax, Haw.	72
scaeocentra, Meyr.	199	scutellariella, Chamb.	90	serena, Meyr.	189
scalata, Meyr.	125	scytalias, Meyr.	217	serialis, Meyr.	250
scalella, Scop.	70	scytina, Meyr.	149	serica, Meyr.	34
scatebrosa, Meyr.	103	seclusella, Walk.	204	sericella, Reb.	202
sceletodes, Meyr.	31	secretella, Walk.	120	sericiella, Wals.	202
sceletopa, Meyr.	231	secta, Meyr.	149	serotinella, Busck	83
scenites, Meyr.	177	sectella, Walk.	121	serratipalpella, Chamb.	89
Sceptea (genus), Wals.	252	secundella, Walk.	120	serrativittella, Zell.	196
sceptrarcha, Meyr.	239	sedata, Butl.	93	serratilella, Tengst.	75
Schemataspis (genus), Meyr.	137	seductella, Walk.	121	sertigera, Meyr.	138
schematias, Meyr.	47	sedulitella, Busck	71	servella, Zell.	43
schematica, Meyr.	80	segetella, Zell.	142	servula, Wals.	139
Schematistis (genus), Meyr.	187	segnis, Meyr.	29	servula, Meyr.	115
Schistophila (genus), Chrét.	68	selaginella, Mann.	30	sesamodes, Meyr.	121
schizogynae, Wals.	71	selectella, Car.	78	sesostrella, Reb.	79
Schleichi, Christ	96	selectella, Walk.	130	sestertiella, Herr.-Sch.	96
schmidella, Heyd.	173	selenia, Meyr.	150	setosella, Clem.	196
schoenota, Meyr.	210	sella, Chamb.	70	sevectella, Walk.	191
scholastica, Wals.	42	selmatias, Meyr.	210	severella, Walk.	117
scholias, Meyr.	121	sematacma, Meyr.	34	sexstrigella, Chamb.	105
sciactis, Meyr.	215	sematica, Meyr.	71	sibila, Meyr.	92
sciastes, Wals.	175	semiacma, Meyr.	117	sicariella, Zell.	107
Scindalmota (genus), Turn.	110	semicostella, Hübn.	107	siccifolii, Wals.	164
scintillella, Fisch. v. Rösl.	122	semicostella, Staud.	71	Sicera (genus), Chrét	21
scintillula, Wals.	130	semicuprata, Meyr.	195	siculella, Wocke.	236
sciodes, Meyr.	148	semicyclicella, Busck	90	sideraula, Meyr.	205
sciadora, Meyr.	176	semidecandrella, Threlf.	96	siderophaea, Wals.	120
sciograpta, Meyr.	157	seminata, Meyr.	196	siderosema, Turn.	173
sciomima, Meyr.	120	seminivora, Wals.	42	sidonia, Meyr.	231
scioplasta, Meyr.	251	semiographa, Turn.	150	Sieversi, Staud.	79
scioplecta, Meyr.	103	semiophanes, Meyr.	46	Semimeris (genus), Meyr.	182
sciopola, Meyr.	32	semirupta, Meyr.	238	siewersiellus, Christ	79
scioxantha, Meyr.	231	semiusta, Meyr.	70	sigalota, Meyr.	252
sciritis, Meyr.	189	semnochroa, Meyr.	33	sigillatrix, Meyr.	249
scissella, Chrét.	78	semnopha, Meyr.	153	signatella, Herr.-Sch.	202
scitella, Walk.	120	semnopa, Meyr.	103	signella, Herr.-Sch.	202
sciurella, Wals	96, 102	Semnostoma (genus), Meyr.	103	signifera, Feld.	241
Sclerocecis (genus), Chrét.	21	Semocharista (genus), Meyr.	147	silacea, Haw.	30
Sclerograptis (genus), Meyr.	158	Semodictis, Meyr.	155	silacella, Hübn.	168
sclerotricha, Meyr.	34	senariella, Zell.	57	silignitis, Turn.	93
scopolella, Hübn.	76	senecta, Wals.	67	silvestris, Meyr.	156

LEPIDOPTERA HETEROPTERA

Pages		Pages		Pages
similella, Chamb.	35	sparsella, Joan.	201	stasiarcha, Meyr.
similella, Snell.	228	<i>sparsiciliella</i> , Barr.	111	stasigastra, Meyr.
similis, Staint.	74	spartiella, Schrank	154	stasiotica, Meyr.
Simoneura (genus), Wals.	114	spathias, Meyr.	219	staticella, Mill.
simplex, Wals.	249	spathota, Meyr.	156	statices, Staud.
<i>simplicella</i> , Hein.	79	<i>speciosella</i> , Teich.	157	Stegasta (genus), Meyr.
simplicella, Walk.	111	<i>speciosella</i> , Walk.	120	Stelecharis (genus), Meyr.
simpliciella, Busck	196	spectralis, Meyr.	206	stellatella, Busck
<i>simpliciella</i> , Chamb.	93	spectrata, Meyr.	108	stelliferella, Walk.
simpliciella, Staint.	258	spectrella, Meyr.	24	stellulata, Meyr.
simulacrella, Meyr.	156	specularis, Meyr.	125	Stenolechia (genus), Meyr.
sinaica, Frauenf.	44	speratella, Busck	253	stenomorpha, Meyr.
singula, Staud.	94	spergulariella, Chrét.	92	Stenopherna, Low.
sinistra, Meyr.	45	sphaeristis, Meyr.	209	Stereomnides (genus), Meyr.
Sinoe, Chamb.	57	Sphagiocrates (genus), Meyr.	183	stereogramma, Meyr.
sinuatella, Wals.	79	Sphaleractis (genus), Meyr.	145	Stereomita (genus), Braun
sinuosa, Meyr.	239	sphecopa, Meyr.	229	sterictis, Meyr.
Siovata, Walk.	237	sphenias, Meyr.	211	stesichora, Meyr.
siranta, Meyr.	197	Sphenocrates (genus), Meyr.	234	steudeliella, Frey
Sirogenes, Meyr.	190	sphenodelta, Meyr.	118	sticheris, Turn.
sirota, Meyr.	43	Sphenegrypa (genus), Meyr.	128	sticta, Wals.
sistrella, Busck	77	sphenophora, Meyr.	156	stictocosma, Meyr.
sisyraea, Meyr.	149	sphenopis, Meyr.	142	stigma, Staud.
Sisyredonta (genus), Meyr.	226	sphyrocopa, Meyr.	189	stillata, Meyr.
sitiens, Meyr.	251	spicata, Meyr.	154	stipella, Hüb.
Sitotroga (genus), Hein.	38	spiculifera, Meyr.	207	Stiphrestola (genus), Meyr.
<i>smaragdella</i> , Walk.	120	spiladias, Meyr.	239	Stoeberhinus (genus), Bull.
Smenodeca (genus), Meyr.	54	spilochorda, Meyr.	257	stoica, Meyr.
snellenella, Wals.	25	spilodoma, Meyr.	50	Stomopteryx (genus), Hein.
sobria, Meyr.	238	spilotella, Barnes	81	stomylia, Snell.
sodalella, Walk.	150	spilotella, Walk.	245	storestis, Meyr.
<i>solanella</i> , Boisd.	93	spinigera, Meyr.	39	<i>stramineellus</i> , Chamb.
<i>solanella</i> , Chamb.	93	spintheropis, Meyr.	130	straminicornis, Meyr.
<i>solemnella</i> , Christ.	123	spiraeae, Staud.	166	straminis, Wals.
solida, Wals.	222	splendens, Staud.	112	strangalistis, Meyr.
solidella, Walk.	121	splendoriferella, Busck	90	stratella, Wals.
solita, Meyr.	256	<i>spodiella</i> , Treitsch	142	stratifera, Meyr.
solitaria, Staud	97	spoliatella, Walk.	244	stratigera, Meyr.
soloea, Meyr.	145	sporodeta, Turn.	150	stratimera, Low.
solutella, Zell.	83	sporogramma, Meyr.	170	streliciella, Herr -Sch.
solutrix, Meyr.	157	sporozona, Meyr.	120	strelitziella, Hein.
sonorensis, Wals.	80	spurcella, Herr -Sch.	82	Streniastis (genus), Meyr.
Sophronia (genus), Hüb.	106	squalens, Meyr	177	Strenopnilia (genus), Meyr.
sophroniella, Reb.	203	<i>squamigera</i> , Wals.	46	strenuella, Walk.
sophronistis, Meyr.	241	<i>squamulella</i> , Pey.	71	strepsiptila, Meyr.
sordidula, Meyr.	63	stachyophora, Meyr.	210	striatella, Busck.
soreuta, Meyr.	108	Stachycostoma (genus), Meyr.	161	striatella, Herr -Sch.
soritica, Meyr.	216	stagmatophoria, Wals.	131	striatella, Hüb.
soronella, Busck	67	stagmatopis, Meyr.	256	striatella, Murtf.
soroculella, Hüb.	83	Stagmaturgis (genus), Meyr.	114	striatopunctella, Reb.
Sorotacta (genus), Meyr.	124	stalactis, Meyr.	32	strictella, Chrét.
sortilega, Meyr.	240	<i>Stangei</i> , Her.	92	<i>strigiplenella</i> , Walk.
sparsella, Christ.	170			

Pages	Pages	Pages			
strigosa, Durr.	241	<i>suffusella</i> , Chamb.	46, 177	tabulata, Meyr.	149
strigosa, Meyr.	250	<i>suffusella</i> , Dougl.	44	tachyptilella, Reb.	96
<i>striolatella</i> , Hein.	82	<i>suffusella</i> , Walk.	121	Tachyptilia, Hein.	125
Strobisia (genus), Clem.	130	<i>suffusella</i> , Wals.	33	<i>tactella</i> , Walk.	189
stromatias, Meyr.	196	<i>sulcata</i> , Meyr.	213	<i>tactica</i> , Meyr.	177
strophopa, Meyr.	240	<i>sulcella</i> , Staud.	98	<i>taeniolella</i> , Zell.	112
Stryphnocopa (genus), Meyr.	185	<i>summata</i> , Meyr.	175	<i>talantodes</i> , Meyr.	217
studiosa, Meyr.	63	<i>sumptella</i> , Walk.	189	<i>tamaricicola</i> , Joan.	44
stygia, Meyr.	115	<i>superbella</i> , Chamb.	47	<i>tamaricicola</i> , Wals.	23
stygnota, Wals.	181	<i>superbella</i> , Zell.	47	<i>tamariciella</i> , Zell.	78
Styloceros, Meyr.	236	<i>superrella</i> , Walk.	120	<i>tamiodes</i> , Meyr.	236
stylota, Meyr.	154	<i>superfetella</i> , Pey.	82	<i>tanacetella</i> , Schranck	35
suaedella, Rich.	92	<i>superfusella</i> , Walk.	121	<i>tannuolella</i> , Reb.	79
suaedivorella, Chrét.	92	<i>suppleiella</i> , Wals.	84	<i>tanyrrhina</i> , Meyr.	100
suasella, Const.	95	<i>surda</i> , Meyr.	44	Taphrosaris (genus), Meyr.	194
suasoria, Meyr.	92	<i>susceptella</i> , Walk.	120	<i>tapinota</i> , Wals.	123
subactella, Walk.	125	<i>suspectella</i> , Hein.	82	<i>taracta</i> , Turn.	150
subaequalis, Wals.	105	<i>suspectella</i> , Walk.	121	<i>tarandella</i> , Wocke	77
subalbella, Chamb.	249	<i>suspensa</i> , Meyr.	76	<i>tardella</i> , Walk.	121
subapicalis, Walk.	121	<i>sylvaecolella</i> , Chamb.	76	<i>tarquiniella</i> , Staint.	47
subcaerulea, Meyr.	92	<i>sylvicolella</i> , Busck	63	<i>tartarea</i> , Meyr.	259
subdecurtella, Staint.	46	Symbatica (genus), Meyr.	66	<i>tauropis</i> , Meyr.	251
subdentata, Meyr.	189	<i>symbolica</i> , Meyr.	138	Taygete, Chamb.	62
subdiminutella, Staint.	92	Symbolitis (genus), Meyr.	166	Technographa (genus), Meyr.	207
subdolella, Staud.	33	<i>symmetra</i> , Turn.	256	Tecia (genus), Strand	89
subericinella, Herr.-Sch.	46	Symmoca (genus), Hübn.	200	<i>tectaphella</i> , Reb.	201
suberiella, Car.	126	<i>symmocella</i> , Reb.	169	<i>tectella</i> , Herr.-Sch.	75
subfulvescens, Meyr.	154	Sympanactis (genus), Meyr.	101	<i>tegminata</i> , Meyr.	144
subjectella, Walk.	133	<i>symphora</i> , Wals.	118	<i>tegulella</i> , Herr.-Sch.	74
sublatella, Walk.	121	<i>synacma</i> , Meyr.	103	<i>tegulella</i> , Wals.	139
sublustricella, Walk.	116	<i>synaphrista</i> , Meyr.	73	<i>tehuacana</i> , Busck	84
sublutella, Christ.	42	Syncathedra (genus), Meyr.	215	<i>telegraphella</i> , Walk.	190
submissella, Frey	112	Syncopacula (genus), Meyr.	72	Teleia, Hein.	69
submissella, Staint.	92	<i>syncosma</i> , Meyr.	129	Telephata (genus), Meyr.	214
subnexella, Walk.	150	<i>syncrypta</i> , Meyr.	156	Telephila (genus), Meyr.	173
ubocellea, Steph.	102	<i>syndelta</i> , Meyr.	70	<i>telifera</i> , Meyr.	63
subrosea, Meyr.	45	Syndesmica (genus), Turn.	247	<i>telopis</i> , Meyr.	148
subrosea, Wood	71	<i>synecta</i> , Meyr.	92	Telphusa (genus), Chamb.	69
subrubella, Chamb.	122	<i>syngraphopa</i> , Meyr.	118	<i>temenitis</i> , Meyr.	148
subutila, Meyr.	188	<i>syngrapta</i> , Meyr.	189	<i>temenodes</i> , Meyr.	210
subscriptella, Walk.	121	<i>synocha</i> , Meyr.	118	<i>temeraria</i> , Meyr.	231
subsecivella, Zell.	111	<i>synthetica</i> , Wals.	75	<i>temerella</i> , Zell.	122
subsequella, Hübn	122	<i>syntonopis</i> , Meyr.	250	<i>tenebrella</i> , Hübn.	42
subsequella, Treitsch.	80	<i>syntropha</i> , Meyr.	238	<i>tenebricosa</i> , Turn.	87
subservitella, Walk.	239	<i>syriacella</i> , Rag.	201	<i>tenebrionellus</i> , Mann.	203
subsimella, Clem.	63	Syrmadaula (genus), Meyr.	255	<i>tenebrosa</i> , Meyr.	157
substratella, Wals.	175	<i>syrphetodes</i> , Meyr.	245	<i>tenebrosella</i> , Teich.	82
subterranea, Busck	90	<i>syrticola</i> , Staud.	83	<i>tenebrosella</i> , Zell.	42
subtractella, Walk.	93	<i>Systasiota</i> , Wals.	130	<i>tenella</i> , Turn.	236
subvectella, Walk.	130	<i>tabacella</i> , Rag.	93	<i>tengstroemiella</i> , Joan.	94
succincta, Wals.	121	<i>tabellata</i> , Meyr.	169	<i>tentatella</i> , Walk.	131
succinctella, Zell.	60	Tabernillaea (genus), Wals.	85	<i>tenuiella</i> , Mann.	43
suffectella, Walk.	120			<i>tenuipalpella</i> , Snell.	225

LEPIDOPTERA. HETEROCHERA

Pages		Pages		Pages	
tenuis, Wals.	102	thanatodes, Low.	146	thysanora, Meyr.	121
tepens, Meyr.	189	thanatopsis, Low.	179	thysanota, Wals.	57
tephrastis, Meyr.	171	thaumalea, Wals.	138	tigrina, Christ.	70
tephriasella, Chamb.	125	Theisoa, Chamb.	51	timidella, Wocke	122
tephriditella, Dup.	82	Thelyasceta (genus), Meyr.	139	Timyra (genus), Walk.	209
tephrodes, Meyr.	191	themelia, Meyr.	189	tinctella, Wals.	211
tephroplintha, Meyr.	157	thererastis, Turn.	44	Tingentera, Walk.	204
tepidata, Meyr.	250	thermaea, Low.	167	Tipasa, Walk.	213
terebinthinella, Herr. Sch.	77	thermobapta, Low.	231	Tipha, Walk.	204
terminimaculella, Kearf.	78	thermochroa, Low.	147	Tirallis, Walk	204
ternariella, Zell.	76	thermodryas, Meyr.	175	Tiranimia (genus), Chrét.	27
ternatella, Staud.	250	thermophaea, Meyr.	191	Tirasia, Walk	237
terpnodes, Meyr.	56	thermopis, Meyr.	257	Tiriza, Walk.	237
terracotta, Wals.	193	thesmiopa, Meyr.	189	tischeriella, Zell.	96
terracocella, Busck	90	thetica, Meyr.	44	Tisia (genus), Walk.	204
terrella, Fisch. v. Rösl.	249	thinodes, Meyr.	145	Titana, Walk.	237
terrella, Hübn.	75	thiodes, Meyr.	57	titanota, Wals.	121
terrella, Walk.	93	thiodora, Meyr.	240	Titacula (genus), Walk.	162
terrena, Turn.	236	Thlegnatha (genus), Meyr.	167	Tiva, Walk.	234
terrenella, Busck.	122	thiophara, Turn.	32	Toomia (genus), Walk.	186
terrestrella, Zell.	82	Thlotricha (genus), Meyr.	101	tofosella, Reb.	202
terrigena, Meyr.	238	tholaea, Meyr.	66	Togia (genus), Walk.	203
tessella, Hübn.	76	Tholerestola (genus), Meyr.	88	tonaea, Meyr.	219
tessulata, Meyr.	157	tholias, Meyr.	217	Tonosa, Walk.	204
testacea, Butl.	255	tholochorda, Meyr.	162	tonsa, Meyr.	156
testacea, Meyr.	56	thomeriella, Chret.	71	tophella, Wals.	35
tetrachroa, Low.	170	thoracibella, Chamb.	75	toreuta, Turn.	156
tetraclina, Meyr.	209	<i>thoracella</i> , Wals.	63, 179	<i>toreutis</i> , Meyr.	210
tetracosma, Meyr.	44	thoracias, Meyr.	206	Tornodoxa (genus), Meyr.	162
tetradelta, Meyr.	228	thoracica, Meyr.	113	tornoptila, Meyr.	122
tedradymiella, Busck	90	thoracistrigella, Chamb.	83	Torodora (genus), Meyr.	244
tetragonella, Staint.	44	thoracochrella, Chamb.	83	torosulella, Reb.	30
tetraphaga, Meyr.	105	thorybodes, Meyr.	103	torrefacta, Meyr.	182
tetraleuca, Meyr.	76	<i>thrancita</i> , Meyr.	238	torrescens, Meyr.	184
tetrametra, Meyr.	164	thrasydora, Meyr.	231	torridella, Mann	30
tetrapeda, Meyr.	257	thrasynta, Meyr.	191	tortuosa, Meyr.	156
tetraphala, Meyr.	103	Thriophora (genus) Meyr.	163	torva, Meyr.	249
tetraplecta, Meyr.	106	thrombodes, Meyr.	31	Tosca (genus), Heinr.	53
tetraploa, Meyr.	148	thrysandra, Meyr.	196	<i>tonceyellus</i> , Busck	175
tetraptila, Meyr.	156	Thrypsigenes (genus) Meyr.	221	toxastis, Meyr.	210
tetrasticta, Turn.	147	thryptica, Meyr.	178	Toxoceras, Chrét.	33
tetraxoa, Meyr.	39	Thubana (genus), Walk.	234	trachycnemis, Meyr.	121
<i>tetroctas</i> , Meyr.	164	thujella, Kearf.	58	trachycosma, Meyr.	78
tetrortha, Meyr.	121	thulella, Staud.	74	trachydyta, Meyr.	156
Teuchophanes (genus), Meyr.	135	thyellias, Meyr.	23	trachyphanes, Meyr.	148
Teucrodexa (genus), Meyr.	206	Thymbritis (genus), Meyr.	230	traducella, Busck	76
texanella, Chamb.	80	thymifoliella, Const.	94	traganella, Chrét.	92
textifera, Meyr.	92	Thymosophia (genus), Meyr.	87	tragicella, Heyd.	82
textorella, Chrét.	82	thyraula, Meyr.	93	<i>trajectella</i> , Walk.	121
thalamias, Meyr	150	thyridota, Meyr.	148	transjectella, Walk.	122
thalamitis, Meyr.	45	thrysicola, Meyr.	190	translucida, Wals.	70
thalamopa, Meyr.	189	thyroptera, Meyr.	150	trapezias, Meyr.	121
thalpodes, Meyr.	188	Thrysostoma (genus), Meyr.	99	traumatias, Meyr.	180

Pages	Pages	Pages			
trauniella, Zell.	76	trinotella, Herr.-Sch.	45	undecimpunctella, Mann	202
trematias, Meyr.	44	triocellella, Chamb.	90	undina, Meyr.	23
tremella, Wood	71	triophthalma, Meyr.	248	unicipunctellus, Clem.	176
tremulella, Dup.	126	triparella, Zell.	71	unicorella, Dup.	42
triaenota, Meyr.	154	triphora, Low.	37	unifasciella, Busck	77
trialbimaculella, Chamb	75	triplacopis, Meyr.	77	uniformella, Reb.	201
triangularis, Braun	125	triplagella, Walk.	193	unistrigella, Busck	117
triangulella, Busck	77	tripunctella, Schiff.	142	unistrigella, Chamb.	78
triannulata, Meyr.	157	tripunctella, Snell.	43	Untomia (genus), Busck	116
triannulella, Herr.-Sch.	248	tripustulata, Wals.	245	untomiella, Busck	116
triatomea, Mühl.	63	triscelis, Meyr.	76	uranopis, Meyr.	132
tribrachia, Meyr.	219	trisignis, Meyr.	76	urosema, Meyr.	93
trichalina, Meyr.	149	trissoxantha, Meyr.	134	ursula, Wals.	126
trichaspis, Meyr.	52	tristella, Car.	201	ussuriella, Car.	112
trichella, Busck	86	tristella, Hein.	96	ustulatella, Staud.	94
Trichembola (genus), Meyr	28	tristella, Reb.	30	ustulella, Fabr.	177
trichinaspis, Meyr.	91	tristella, Snell.	249		
trichocyma, Meyr.	249	tristella, Teich.	82	vacatella, Walk.	149
trichodeta, Meyr.	154	tristicta, Busck	189	vacciniella, Busck	174
trichombra, Low.	171	tristicta, Wals.	34	vagatioella, Chamb	58
trichosema, Meyr.	148	tristis, Staud.	77	vagella, Walk.	78
trichostola, Meyr.	83	tristrigella, Wals.	123	valesiella, Staud.	94
Trichotaphe (genus), Clem.	195	Tritadelpha (genus), Meyr.	143	vallicola, Meyr.	44
trichroa, Meyr.	212	triturata, Meyr.	214	variabilis, Busck	80
tricolor, Feld.	136	trizona, Low.	231	variana, Meyr.	87
tricolorella, Haw.	96	trochias, Meyr.	149	variella, Chamb.	58
tricornis, Meyr.	153	trochilea, Wals.	120	varronia, Busck	196
Tricyanaula (genus), Meyr	131	trochilella, Hein.	95	vasconiella, Rössl.	93
tridecta, Low.	149	tropaea, Meyr.	105	Vazugada (genus), Walk.	178
tridentella, Wals.	126	trophella, Busck	78	vecors, Meyr.	259
trifasciata, Wals.	221	trossulella, Wals.	46	vectaria, Meyr.	153
trifasciella, Chamb.	76	Trypanisma (genus), Clem.	61	velatella, Busck	70
trifasciella, Reb.	71	turana, Car.	201	velitaris, Meyr.	249
trifoliella, Const.	122	turbatella, Treitsch.	46	velocella, Dup.	80
triforella, Zell.	152	turbida, Turn.	45	veneranda, Wals.	83
triglossa, Meyr	40	turbidella, Nolck.	82	venosulella, Möschl.	247
trigonella, Wals.	189	turbanata, Meyr.	224	ventosa, Meyr.	176
trigonias, Meyr.	242	turgida, Meyr.	194	ventralella, Zell.	91
trigonophorella, Zell.	57	turpella, Herr.-Sch.	83	ventrella, Fitch	176
trigonopis, Meyr.	240	turrita, Meyr.	189	venustella, Chamb.	131
trigonota, Wals.	179	tussilaginella, Hein.	94	vepretella, Zell.	82
trigramma, Meyr.	106	Tuta, Strand	89	verbascella, Hübn.	98
trijugella, Ersch.	60	Tutti, Wals.	34	verberata, Meyr.	249
trilineella, Chamb.	83	typhlopis, Meyr.	245	vernella, Murtf.	83
trimaculella, Chamb.	175, 196	Uipsa, Walk.	209	versatella, Walk.	121
trimaculella, Pack.	78	Uiliaria (genus), Dum.	255	versicolorella, Chamb.	90
trimetropis, Meyr.	149	ulicinella, Staud.	63	versicolorella, Kearf.	47
trimolybda, Meyr.	120	umbrata, Wals.	254	versicolorella, Walk.	187
trinella, Fuchs	96	umbriferella, Herr.-Sch.	42	versutella, Zell.	83
trinervis, Meyr	248	umbripennis, Wals.	85	verticosa, Meyr.	156, 213
trinotata, Meyr.	185	umbrosella, Zell.	74	veruta, Meyr.	154
trinotella, Busck	196	unctulella, Zell.	84	vestita, Wals.	254
trinotella, Coq.	176			veteranella, Zell.	45

LEPIDOPTERA HETEROCHERA

Pages	Pages	Pages			
veterascens, Meyr.	68	wacoella, Chamb.	81	xuthostola, Wals.	196
vetustella, Herr.-Sch.	44	Wagae, Now.	70	xyloides, Meyr.	214
vetustella, Walk.	195	Walkeri, Wals.	134	xyloglypta, Meyr.	80
viatrix, Meyr.	99	walkeriella, Dougl.	36	xylogramma, Meyr.	235
vicana, Meyr.	45	Walsinghami, Dietz.	79	xylophaea, Meyr.	81
vicaria, Meyr.	92	washingtoniella, Busck	90, 196	xylotripta, Meyr.	225
vicinella, Dougl.	96	wilkella, Linn.	47	Xystophora, Hein.	41
vicinella, Herr.-Sch.	96	Wollastoni, Wals.	113		
victimella, Wals.	203			youngella, Kearn.	47
victrix, Meyr.	138	xanthaspis, Meyr.	101	yuccacella, Busck	183
vicularis, Meyr.	256	xanthastis, Low.	141		
viduella, Fabr.	76	xanthaula, Meyr.	210	zachroa, Meyr.	193
viduellus, Reb.	203	xanthocarpa, Meyr.	115	zalias, Meyr.	24
vigilans, Meyr.	191	xanthocephala, Meyr.	149	Zalithia (genus), Meyr.	132
vigilax, Meyr.	230	xanthochalca, Meyr.	239	zapryrodes, Turn.	138
vilella, Zell.	85	xanthoclista, Meyr.	258	zebrella, Treitsch.	79
villosula, Zell.	61	xanthocosma, Meyr.	239	zebrina, Wals.	121
vinella, Banks	112	xanthodeta, Meyr.	179	zeitunella, Reb.	201
vinitincta, Wals.	142	xanthodora, Meyr.	103	zelosaris, Meyr.	52
vinolenta, Meyr.	71	xanthographa, Wals.	221	Zelasyne (genus), Wals.	135
violacea, Tengst.	79	xanthograpta, Meyr.	257	zemiodes, Durr.	85
violacella, Chrét.	34	xanthomorpha, Meyr.	232	sephyrella, Treitsch.	75
violaceofusca, Zell.	81	xanthophanes, Meyr.	126	zetterstedtiella, Zell.	79
violaria, Meyr.	195	xanthophilella, Barnes	77	zinckenella, Hübn.	47
virescens, Wals.	250	xanthoria, Meyr.	251	zingarella, Wals.	193
viretella, Zell.	125	xanthoselena, Wals.	183	Zizyphia (genus); Chrét.	36
virgella, Thunb.	79	xanthostoma, Wals.	131	Zomeutis (genus), Meyr.	194
virgineilla, Reb.	202	xanthoteles, Meyr.	235	zomias, Meyr.	175
viridans, Meyr.	124	xanthotricha, Meyr.	58	zonaea, Meyr.	223
viridella, Snell.	136	Xenolechia, Meyr.	69	zonariella, Wals.	112
viridescens, Meyr.	132	xerastis, Meyr.	248	zonella, Chrét.	78
viscariella, Staint.	95	xerochroa, Meyr.	38	zonias, Meyr.	231
vitiosella, Zell.	201	xerodes, Wals.	196	zonodesma, Low.	231
vittella, Busck	117	Xerometra (genus), Meyr.	170	zonostoma, Meyr.	178
voltinella, Chrét.	95	xerophaga, Meyr.	247	zopherota, Meyr.	152
volubilis, Meyr.	121	xerophylla, Meyr.	93	zophochalca, Meyr.	63
voluta, Meyr.	148	xeropis, Meyr.	31	zulu, Wals.	138
vorticella, Scop.	112	xestobyrsa, Meyr.	191	zygophora, Meyr.	174
vulgella, Hübn.	71	xestolitha, Meyr.	148	zygophyllea, Reb.	92
Wachtli, Rog.	47	xuthias, Meyr.	149	zygotoma, Meyr.	87
		xuthochyta, Turn.	176	zymotis, Meyr.	68

EXPLANATION OF PLATES

PLATE I

- Fig. 1. *Anomoxena spinigera*, Meyrick.
 — 2. *Apatetris isonira*, Meyrick.
 — 3. *Nealyda bifidella*, Dietz.
 — 4. *Oecocecis guyonella*, Guenée.
 — 5. *Metzneria inflamatella*, Christoph.
 — 6. *Trichembola segnis*, Meyrick.
 — 7. *Isophrictis magnella*, Busck.
 — 8. *Pycnostola operosa*, Meyrick.
 — 9. *Megacraspedus pityritis*, Meyrick.
 — 10. *Photodotis prochalina*, Meyrick.
 — 11. *Empedaula phanerozona*, Meyrick.
 — 12. *Aristotelia roseosuffusella*, Clemens.
 — 13. — *centrosema*, Lower.

- Fig. 14. *Aristotelia citrocosma*, Meyrick.
 — 15. *Stenolechia orsicoma*, Meyrick.
 — 16. *Helice constrictella*, Zeller.
 — 17. *Sympnactis hetaera*, Meyrick.
 — 18. *Smenodoca erebenna*, Meyrick.
 — 19. *Recurvaria thujella*, Kearfott.
 — 20. *Oxylechia confirmata*, Meyrick.
 — 21. *Epithestis attributella*, Walker.
 — 22. *Zelosyne olga*, Meyrick.
 — 23. *Parapsectris neograpta*, Meyrick.
 — 24. *Dissostila crocodora*, Meyrick.
 — 25. *Telphusa callitechna*, Meyrick.

PLATE 2

- Fig. 26. *Gelechia cercerella*, Chambers.
 — 27. — *ochrias*, Meyrick.
 — 28. — *trisignis*, Meyrick.
 — 29. — *blandulella*, Walker.
 — 30. *Platyedra gossypiella*, Saunders.
 — 31. *Stegasta variana*, Meyrick.
 — 32. *Parastega chionostigma*, Walsingham.
 — 33. *Mometa zemiodes*, Durrant.
 — 34. *Phthorimaea synecta*, Meyrick.
 — 35. *Gnorimoschema aquilina*, Meyrick.
 — 36. — *gallaesolidaginis*, Riley.
 — 37. *Sarotorna eridora*, Meyrick.
 — 38. *Thysostoma glaucitis*, Meyrick.

- Fig. 39. *Semnoscota barathrota*, Meyrick.
 — 40. *Thiotricha glenias*, Meyrick.
 — 41. — *polyaula*, Meyrick.
 — 42. *Hierangela erythrogramma*, Meyrick.
 — 43. *Crambodoxa platyaula*, Meyrick.
 — 44. *Polyhymno palinorsa*, Meyrick.
 — 45. *Idiophantis chiridota*, Meyrick.
 — 46. *Stomopteryx polychromella*, Rebel.
 — 47. *Inotica gaesata*, Meyrick.
 — 48. *Calliprora pentagramma*, Meyrick.
 — 49. *Commatica cyanorrhoa*, Meyrick.
 — 50. *Battaristis ardiophora*, Meyrick.

PLATE 3

- Fig. 51. *Anacampsis prasina*, Meyrick.
 — 52. *Compsolechia perlatella*, Walker.
 — 53. — *solidella*, Walker.
 — 54. — *mniocosma*, Meyrick.
 — 55. *Desmaucha chrysostoma*, Meyrick.
 — 56. *Charistica callichroma*, Meyrick.
 — 57. *Strobisia spintheropis*, Meyrick.
 — 58. *Onebala symbolica*, Meyrick.
 — 59. *Teuchophanes leucopleura*, Meyrick.
 — 60. *Acompsia formosella*, Hübner.
 — 61. *Sphaleractis parasticta*, Meyrick.
 — 62. *Protolechia ceramicata*, Meyrick.
 — 63. — *xanthocephala*, Meyrick.

- Fig. 64. *Orthoptila abruptella*, Walker.
 — 65. *Hyodectis crenoides*, Meyrick.
 — 66. *Anarsia molybdota*, Meyrick.
 — 67. *Chelaria corynetis*, Meyrick.
 — 68. *Dactylethora candida*, Stainton.
 — 69. — *incondita*, Meyrick.
 — 70. *Sophronia chilonella*, Treitschke.
 — 71. *Brachyacma epiochra*, Meyrick.
 — 72. *Dichomeris oceanis*, Meyrick.
 — 73. *Eunebristis zachroa*, Meyrick.
 — 74. *Atasthalistis gnophrina*, Felder.
 — 75. — *euchroa*, Lower.

PLATE 4


- Fig. 76. *Musurga sandycitis*, Meyrick.
 — 77. *Cymotricha miltophragma*, Meyrick.
 — 78. *Trichotaphe flavigostella*, Clemens.
 — 79. *Taphrosaris malthacopa*, Meyrick.
 — 80. *Tisis chalybaeella*, Walker.
 — 81. *Mnesteria pharetrata*, Meyrick.
 — 82. *Timyra machlas*, Meyrick.
 — 83. *Crocanthes sidonia*, Meyrick.
 — 84. — *pyrochorda*, Meyrick.
 — 85. *Frisilia rostrata*, Meyrick.
 — 86. *Narthecoceros platyconta*, Meyrick.
 — 87. *Heliangara lampetis*, Meyrick.
 — 88. *Habrogenes eupatris*, Meyrick.

- Fig. 89. *Nosphistica erratica*, Meyrick.
 — 90. *Lecithocera hemichrysa*, Meyrick.
 — 91. — *claustra*, Meyrick.
 — 92. *Deltoplastis cremnaspis*, Meyrick.
 — 93. *Pharangitis spathias*, Meyrick.
 — 94. *Idiopteryx proaula*, Meyrick.
 — 95. *Deocloca xanthoselena*, Walsingham.
 — 96. *Hylograptis thryptica*, Meyrick.
 — 97. *Anaptilora isocosma*, Meyrick.
 — 98. *Sarisophora leptoglypta*, Meyrick.
 — 99. *Crocanthes carcharias*, Meyrick.
 — 100. *Torodora typhlopis*, Meyrick.


PLATE 5

- Fig. 101. *Brachmia phryganitis*, Meyrick.
 — 102. *Apethistis episticta*, Meyrick.
 — 103. *Autosticha aspasta*, Meyrick.
 — 104. — *naulychna*, Meyrick.
 — 105. *Demiophila psaphara*, Meyrick.
 — 106. *Encrasima elaeopis*, Meyrick.
 — 107. *Protobathra leucostola*, Meyrick.
 — 108. *Strenophilà hyptiota*, Meyrick.
 — 109. *Oecia oecophila*, Staudinger.
 — 110. *Holcopogon scaeocentra*, Meyrick.
 — 111. *Symmoca amphicalyx*, Meyrick.
 — 112. — *rhodota*, Meyrick.
 — 113. *Streniastis thermaea*, Lower.
 — 114. *Encolpotis xanthoria*, Meyrick.
 — 115. *Anomoxena spinigera*, Meyrick, forewing.
 — 116. *Aristotelia paphia*, Meyrick, hindwing.
 — 117a. *Gelechia fluvialella*, Busck, forewing.
 — 117b. — — — — head.
 — 117c. — — — — hindwing.

- Fig. 118a. *Lecithocera geraea*, Meyrick, forewing.
 — 118b. — — — — hindwing.
 — 119a. *Crocanthes prasinopsis*, Meyr., forewing.
 — 119b. — — — — hindwing.
 — 120. *Protolechia xanthocephala*, Meyr., hindw.
 — 121a. *Strobisia sapphiritis*, Meyrick, forewing.
 — 121b. — — — — hindwing.
 — 121c. — — — — head.
 — 122a. *Helice constrictella*, Zeller, forewing ♂
 (after Braun).
 — 122b. *Helice constrictella*, Zeller, forewing ♀
 (after Braun).
 — 123a. *Helice pallidochrella*, Chamb., hindw. ♂
 (after Braun).
 — 123b. *Helice pallidochrella*, Chamb., hindw. ♀
 (after Braun).
 — 124. *Trichotaphe flavigostella*, Cl., labial palp.
 — 125. *Dichomeris ligulella*, Hübner, —
 — 126. *Chelaria caryodora*, Meyrick, —
 — 127. *Trichembola segnis*, —




FAM. GELECHIDÆ





GENERAL INSECTORUM

LEPIDOPTERA HETEROCERA




FAM. GELECHIDÆ





FAM. GELECHIDÆ



DIPTERA
FAM. EMPIDIDÆ



DIPTERA

FAM. EMPIDIDÆ

by Axel Leonard MELANDER, Sc. D.

WITH 4 COLORED

AND 4 MONOCHROME PLATES

INTRODUCTION

HE present study was inaugurated in 1900 at the suggestion of Dr. William M. Wheeler, who had been gathering together a collection of Empididæ during the ten previous years. A paper on the Central American species was issued in 1901 with Dr. Wheeler as joint author. The next year was published a monograph of the North American Empididæ exclusive of the genus *Rhamphomyia*, and this has been followed by several short papers in the intervening time. A sojourn of twenty years near the Pacific Coast, where species of Empididæ are particularly numerous, has enabled the writer to get together an extremely rich collection of this family, numbering many hundreds of species and many thousands of specimens. This has been supplemented by specimens received from the following collectors, to whom, naturally, the warmest gratitude is due. Their liberality in furnishing specimens has made possible much of the following contribution.

C. F. Adams, Atherton, Missouri; J. M. Aldrich, Washington, D. C.; C. F. Baker, Los Banos, Philippine Islands; Nathan Banks, Cambridge, Massachusetts; Germain Beaulieu, Montreal, Canada; William Beutenmueller, New York, N. Y.; J. Chester Bradley, Ithaca, New York; C. S. Brimley, Raleigh, North Carolina; Charles T. Brues, Boston, Massachusetts; Gustav Chagnon, Montreal, Canada; T. D. A. Cockerell, Boulder, Colorado; F. R. Cole, Redlands, California; A. B. Cordley, Corvallis, Oregon; R. A. Cooley, Bozeman, Montana; E. T. Cresson, Jr., Philadelphia, Pennsylvania; C. R. Crosby, Ithaca, New York; W. Dietz, Hazleton, Pennsylvania; R. W. Doane, Stanford, California; Charles T. Greene, Washington, D. C.; H. S. Harbeck, Philadelphia Pennsylvania; S. Henshaw, Cambridge, Massachusetts; J. S. Hine, Columbus, Ohio; S. J. Hunter, Lawrence, Kansas; Eldred L. Jenne, deceased; O. A. Johannsen, Ithaca, New York; Charles W. Johnson, Boston, Massachusetts; O. B. Johnson, deceased; Trevor Kincaid, Seattle, Washington; Frederick Knab, deceased;

Adolphus Lutz, Sao Paulo, Brazil; William M. Mann, Washington, D. C.; C. L. Metcalf, Urbana, Illinois; W. L. McAtee, Washington, D. C.; Harold Morrison, Ithaca, New York; William Nason, deceased; H. S. Parish, Toronto, Canada; C. V. Piper, deceased; F. Rogers, Kansas City, Missouri; Pablo Schild, La Suiza de Turrialba, Costa Rica; R. C. Shannon, Washington, D. C.; F. Sherman, New York, N. Y.; Mrs. Annie T. Slosson, New York, N. Y.; A. H. Sturtevant, New York, N. Y.; Roland Thaxter, Cambridge, Massachusetts; E. S. Tucker, deceased; M. C. Van Duzee, San Francisco, California; O. S. Westcott, deceased; William M. Wheeler, Boston, Massachusetts; P. W. Whiting, Boston, Massachusetts; A. J. Wiedt, Newark, New Jersey; M. A. Yothers, Yakima, Washington.

A most important set of specimens was secured in the unworked material belonging to the United States National Museum and that of the Bureau of Biological Survey. This represents the Empid collecting of the Government entomologists and contains much material from the following: J. M. Aldrich, H. S. Barber, G. W. Belfrage, F. C. Bishopp, Robert Brown, Owen Bryant, August Busck, A. N. Caudell, T. D. A. Cockerell, D. W. Coquillett, Crevecoeur, R. P. Currie, R. A. Cushman, E. Daecke, George Dimmock, H. G. Dyar, A. K. Fisher, O. Heidemann, L. O. Jackson, C. R. Jones, J. R. Malloch, W. L. McAtee, J. D. Mitchell, H. K. Morrison, W. D. Pierce, F. C. Pratt, C. V. Riley, S. A. Rohwer, E. A. Schwarz, R. C. Shannon, Mrs. Annie T. Slosson, E. S. G. Titus, C. H. T. Townsend, L. M. Turner, W. V. Warner and E. B. Williamson.

Through the courtesy of C. Howard Curran the unworked Empid material belonging to the Canadian National Museum has been procured and this collection has added several species to the following pages.

A splendid reference collection of about five hundred determined species from Europe and other foreign places has been gathered together through the interest of the following well known dipterists, and has furnished a basis for much of the work.

Theo. Becker, Liegnitz, Germany; Mario Bezzi, Torino, Italy; E. Brunetti, Calcutta, India; E. O. Engel, Dachau, Germany; Richard Frey, Helsingfors, Finland; F. Hermann, Berlin, Germany; F. W. Hutton, New Zealand (deceased); K. Kertész, Budapest (deceased); William Lundbeck, Copenhagen, Denmark; David Miller, Wellington, New Zealand; Lorenz Oldenberg, Berlin, Germany; Gabriel Strobl, Admont, Austria.

In the preparation of this work there have been studied the types of eighty-five per cent. of the North American species, amounting to nearly four hundred species. The remainder, the types of the species described by Say, Macquart, Bigot, Walker and Osten-Sacken, are either no longer in existence or are not available to the American student.

The following pages include a discussion of the genera of the Empididae, a list of the known species together with their bibliography, descriptions of new forms, and here and there tabular keys to species, particularly to those of North America. Unless otherwise designated the types of the new species are in the author's collection. Where no individual is mentioned by name as collector for the new species, the material has been collected by the author.

This study was presented at Harvard University in 1914 in fulfillment of the thesis requirement for the degree of Doctor of Science. At that time the breaking out of the World War and its centralization in Belgium precluded publication in Wytsman's *Genera Insectorum*, and the report has been held in abeyance during the meantime. It is now presented with the inclusions and modifications necessitated by the lapse of ten years.

Contribution no 240 from the Entomological Laboratory of the Bussey Institution for Applied Biology, Harvard University.

Phylogeny. — « These of course are not all the lines of evolution in Diptera, but I believe that they are all irreversible, that evolution has never recovered anything once functionally lost. Moreover, all, or nearly all these lines of evolution are polyphyletic, resulting in numerous cases of parallel resemblances which must be taken into account in any attempt at *true* classification. » (S. W. Williston, Manual of Diptera, 3d. ed. p. 61 [1908].)

The Empididæ include more diverse forms than any other family of Diptera. There is no conservatism to an Empid type, indeed there is no Empid type. Between the full neuration of *Brachystoma* and the reduced venation of the highly specialized *Tetraneurella* there is so great a range of wing structure that one wonders why such diverse forms should be associated.

By usual assent the Empididæ have been grouped with the Dolichopodidæ, forming a division of predatory flies called by Brauer the Orthogena and by Verrall the Microphona. More or less nearly related to these insects are such families as the Asilidæ, Bombyliidæ and Therevidæ, and more remotely in their ancestry such as the Leptidæ and Rhypgidæ. Characters occurring in the Empididæ and also possessed in common by this series of families may usually be regarded as primitive. Postulating thus, the ancestral Empid probably had wings with broadly rounded anal angle, the costa encompassing the entire wing and margining the large alula and calypter as well, the auxiliary vein distinct and complete, ending independently in the costa near or beyond the middle of the wing, the first and second veins simple and long, terminating toward the end of the wing, the third vein with a long fork, the pedicel of the second and third veins arising well toward the root of the wing, the discal cell complete and emitting three simple posterior veins, a strong thickening or fold in the wing membrane beneath the humeral crossvein, the basal cells long, the anal cell acute at its end since the so-called anal crossvein was not at all recurved. Moreover the following characters were probably indicated : head globular, eyes dichoptic and bare, proboscis short, antennæ short, distinctly three-jointed, the last joint ovate-conical and terminated by a three- or four-jointed short style. Thorax small, the metathorax larger than the scutellum, no discal bristles, mesopleuræ hairy, metapleuræ bare, but velvety in front of the spiracle ; abdomen slender, long and cylindrical, pygidium symmetrical and small, no telescopic chitinized ovipositor. Legs simple, slender and elongate, without bristles or apical spurs.

There is of course no living Empid that combines these characters, nor is any such extinct form known, but it is not difficult to picture a delicate bodied insect four or five millimeters in length, with feeble powers of flight and with the structure given above. It would not be overstepping the bounds of analogy to predicate further that this insect was of cinereous brownish coloration, had yellowish legs and hyaline wings, unmarked by a stigma, that it was predatory on smaller weaker insects and lived in shady woods to the North, the adults appearing shortly after the winter snows had disappeared.

Evolution of the genera of the Empididæ has proceeded along certain lines recognized as a usual trend in the development of Diptera. Holopticism or the further development of contiguity of the eyes of both sexes appear again and again. There is a frequent tendency to robustness, with gibbosus thorax, to the introduction of leg armature, specialization of the genitalia, development of metapleural setulæ and of thoracic bristles, fusion of the basal joints of the antennæ, a lengthening of the style into an arista, and a decided lengthening and specialization of the proboscis. In the wings there is a marked tendency toward reduction of the neuration, a shortening of the anterior and of the anal veins, thinning of the hind margin, suppression of the alula and diminution of the anal angle in one direction and lobation of it in another, and a shortening of the pedicels of the second and fourth veins and of the fork of the third. It is such trends as these that Dr. Williston had in mind in his words quoted at the beginning of this section. These tendencies toward specialization are not reversible : a fly with a forked third vein has not descended from one in which this vein is simple, for example, nor

DIPTERA

has a genus with slender femora been produced by such forms as *Edalea* or *Lactistomyia*. But an orthogenetic development of the highly specialized hind legs of these two genera does not necessarily bespeak a phyletic relationship between the two, even though the resemblances are continued to such details of structure as the setosity of the under side of the femur, the patellar bend at the knee, the shortening of the tibia and lengthening of the trochanter and even the compression of the tibia to a flexor edge. The similarities must be interpreted as parallel evolution, for *Edalea* judged by its three posterior veins and dichoptic condition, is more generalized than the thick-legged Hybotinæ, and yet possesses the specialized anal cell and antennæ of the Ocydromiinæ. Both of these genera are better regarded as highly specialized convergent developments of two subfamilies, the result of an orthogenetic evolution. In a group of predatory insects, such as the Empididæ, it would be natural to look for raptorial legs, and such an adaptation does occur in each of the main groups. In *Hoplocyrtoma*, *Scelolabes* and *Palaeoleptopesa* of the Ocydromiinæ the hind legs are fitted for grasping, and in the last genus the adaptation dates back to the Lower Oligocene. In *Platypalpus* the middle legs are modified, even to the patellar bend, and a similar but incipient tendency is noticeable in others of the Tachydromiinæ as in certain species of *Elaphropesa* and *Tachypesa*. In the Hemerodromiinæ it is the front pair that is modified, and so successful has this adaptation proved when on the first pair of legs that the coxae have become lengthened to extend the reach. Of the Empidinæ, *Haplomera*, *Pachymeria*, *Enoplempis* and certain species of *Rhamphomyia* and *Hilara* present thickened femora.

It is my belief that the following tabulation of characters shows the principal lines of evolution in the Empididæ; that development has proceeded from the first set to the second and not in the reverse direction except in some cases of retrograde specialization; and that when two related genera are compared the one possessing a preponderance of the first listed characters is the more generalized and is ancestrally older. Quite obviously no present day Empid is the direct ancestor of another, but with due reservation, particularly with respect to homoplastic characters the tabulation will prove helpful in plotting the phylogeny of the several groups.

Generalized condition.

Head round.
Dichoptic.
Facets uniform.
Eyes bare.
Face narrowly quadrate.
Antennæ three-jointed.
Third joint conical.
Basal joints simple and similar.
Style (terminal).
Proboscis short, vertical.
Palpi two-jointed, bristly.
Body delicate.
POLLINOSA.
Slightly convex thorax.
With few small bristles.
Metanotum long.
Metapleuræ bare.
Pygidium small and symmetrical, short penis,
small valves.

Specialized condition.

Head hemispherical or lengthened.
Holoptic.
Area of large facets.
Eyes pubescent.
Face broad or excised.
Antennæ two-jointed.
Third joint elongate or short.
First joint setulose, second joint with bristle.
Arista (dorsal).
Proboscis long, no labella, inflexed or porrect.
Palpi one-jointed, bare.
Body heavily chitinized.
PRUINOSE, or polished.
Hunchbacked.
With many strong bristles.
Metanotum short and declivous.
Metapleuræ with hairs.
Pygidium large or asymmetrical, long penis,
large valves.

Generalized condition.	Specialized condition.
Ovipositor non-telescopic.	Ovipositor lengthened.
Legs simple.	Legs armed.
Femora and tibiæ with simple setæ.	Femora and tibiæ with scales, or ciliate.
Pulvilli.	No pulvilli.
Costa around entire wing.	Hind margin not thickened, anterior veins stronger, i. e. better fliers.
Anal angle full but not lobose.	Anal angle weak or else lobose.
Alula and large rimmed calypteres.	No alula, calypteres small.
Humeral crossvein present.	Absence of humeral crossvein.
Subhumeral crossvein.	No thickening under humeral crossvein.
Full neuration.	Neuration incomplete.
Veins fixed for the genus.	Veins unstable within the genus, adventitious veins.
Veins straight.	Veins undulating.
Auxiliary vein distinct, ending in costa.	Auxiliary vein near first, evanescent at end.
First and second veins long.	First vein shortened.
Third vein forked.	Third vein simple or with shortening fork.
Pedicel of second and third vein long.	Pedicel of second and third vein short.
Basal cells long.	Basal cells short.
Anal cell long, with acute end.	Second basal shortened at base.
Anal vein strong.	Anal crossvein perpendicular or reflexed.
Anal vein continuous with under side of anal cell.	Anal vein abbreviate.
	Anal vein distinct from under side of anal cell.

While it is a general principle that an organ once functionally lost is never to be regained there are however several instances in the preceding list of characters that appear confusing. For instance, metapleural hairs occur in so many groups of flies, even when the rest of the pleura is bare, that they would seem to be an atavistic index. In the Empididæ the metapleuræ bear setæ or hairs in the genera centering about *Empis*, *Clinocera* and *Chelipoda*, all of which are specialized in their respective subfamilies. Moreover in the forms near *Hybos* and *Ocydromia* the pollen of the pleura becomes villous on the metapleura. These hairs probably bear some deep-seated relation to the little-understood structures at the base of the wing and are also influenced by the metathoracic spiracle, which, in their best developed form, they seem to protect. However in a single genus, e. g. *Empis*, the metapleural hairs range in number from a bushy coarse aggregation to but two or three fine hairs. Those species of *Empis* with but few hairs are in other respects not more primitive than *Oreogeron* which has a great cluster of metapleural hairs, and more likely are losing these structures as the result of a recent retrograde specialization. While such species will not be expected to give rise in the future to hairy forms, it is not so easy to be assured that some of the bare species, like *Toreus* or *Hesperempis*, possess a primitive lack of metapleural hairs or have lost them through secondary specialization.

An ancient stereotyped character that has much value in classification is to be found in the course of the so-called anal crossvein. Originally this posterior branch of the cubitus ended independently in the wing margin. Then for mechanical reasons it became flexed back, narrowing, closing and finally shortening the anal cell. Among living diptera every transition can be found from the open anal cell of *Rhyphus* to the minute vestige of a cell, as in the Acalypterates. In many forms the anal cell

completely disappears. Within the family Empididae most of these transitions occur. *Brachystoma* has a long pointed anal cell, closed toward the margin of the wing, the posterior cubitus continuing its basal direction beyond the furcation. This is the primitive condition, possessed by the more generalized families. In *Empis* the posterior branch of the cubitus is abruptly reflexed and forms the hind margin of the anal cell. In *Clinocera* it is recurved rather than reflexed, but still is continuous with the under side of the anal cell. In *Anthalia* this vein forms a true crossvein, perpendicular to the two sides of the anal cell, apparently an intermediate condition. While this shortening of the anal cell is a mark of specialization it does not mean that the reflexed vein of *Empis* has necessarily passed through a perpendicular stage like that of *Anthalia*. It probably had a sudden origin, or possibly passed through the rounded condition observable in *Clinocera*. The closure of the anal cell affords a most valuable character for the determination of the subfamilies of the Empididae. Those species with acute anal cell invariably belong to either the Hybotinae or the Brachystomatinae. The Ocydromiinae, except for *Bicellaria*, always have the perpendicular crossvein and this is likewise true of all the Tachydromiinae that have an anal cell. The reflexed anal vein is preeminently an Empidinae characteristic, occurring outside of this subfamily only in one line of the Clinoceratinæ and in *Bicellaria*.

There are numerous tendencies evident in the Empididae to develop parallel structures. A strongly gibbous thorax, so characteristic of the Hybotinae, appears independently in *Microphorus*, *Bicellaria* and *Anthalia*. The first vein is setulose in *Oreogeron* and *Phlebotena*, which are not close relatives. The pygidium is bent over the abdomen, forming an epipygium, in the groups about *Clinocera*, *Hilara* and *Microphorus*. Holopticism has been produced in numerous unrelated genera. The Empididae are further interesting in having extended the holoptic condition to the female, as in *Symbalophthalmus* and in most of the Hybotinae. Moreover the eyes may be enlarged below the antennæ instead of above. This tendency is noticeable in the Tachydromiinae, Hemerodromiinae and Clinoceratinæ, and in the extreme cases, such as *Hemerodromia* and *Stilpon*, the eyes quite touch on the face. The formation of an arista is another example of homoplastic structures. The antennæ throughout the family show much variation and afford a series of good taxonomic characters, which however should not be stressed as much as was done by Coquillett. In their generalized condition the antennæ probably terminated in a short three-jointed thickened style. The basal joint of the style disappeared by coalition with the intermediate section, or apparently sometimes by fusion with the apex of the antenna, and in most forms the terminal part remained as a bristle-like segment, shorter than the preceding. This style has been replaced by an arista in many of the genera, but the mechanism of the change is obscure. The arista usually manifests a small basal joint and an attenuated outer portion, which corresponds to the original middle piece of the style. Among the most specialized genera of each of the subfamilies a lengthened arista is to be found.

Metamorphosis. — The early stages of Empididae are relatively little known. The larvae live as predators or scavengers in the ground or in rotting wood. Some have been found in the forests where the adults occur, some are semiaquatic, developing in mud. The species hibernate in the larval stage, pupating in early spring.

The larvae are cylindrical, more or less spindle-shaped, and comprise twelve segments. The head is small, retractile, eyeless, with the two-jointed antennæ small but well developed. The labial plates and the longitudinal rods of the head meet angularly so that in profile they appear bent; mandibles lunate, maxillary palpi small, labium comprising two arcuate bands contiguous and angulate anteriorly. Prothoracic spiracles are small. Abdomen devoid of pseudopods or other appendages, most segments with transverse ventral swellings, the locomotor spinules forming bands on the anterior margins of the segments, the last segment more or less rounded, usually with a tooth or wart below on the hind edge and with a pair of large well separated and sometimes more or less elevated spiracles

above the lower protuberances. In the aquatic *Hemerodromias* the terminal spiracles are replaced by a bunch of delicate external tracheæ.

The pupæ are free, not encased in the last larval skin. The head bears two small carinated setigerous tubercles on the upper anterior part; antennal sheaths raised above the level of the face, directed downward and slightly outward and tapering apically; proboscis elongated. The thorax is generally furnished with long bristles but not with thorns. The respiratory organ may consist of short stalks (*Rhamphomyia*) or be greatly elongated (*Drapetis*). Prothoracic spiracles are present. The hind tarsi extend much beyond the wings. The abdomen may be provided above with girdles of small teeth alternating with bristles and ventrally with hairs alone, or may have a dense dorsal covering of small spinules. Seven pairs of abdominal spiracles are present. In *Hemerodromia* eight pairs of very long filiform tracheal tubes extend from the position of the spiracles.

Ethology and Occurrence. — Empididæ are especially abundant in the spring and early summer in the undergrowth of shady woods. They are prone to occur in mountainous districts and in rather humid regions. While such is the general distribution several of the genera have adopted specialized habitats. For example, most of the Clinoceratinæ frequent running water, some resting beside waterfalls or on water-splashed rocks in swift streams or even amphibiously entering the water; species of the Hemerodromiinæ are to be found in the shady foliage overhanging the banks of streams; *Tachypæza* stalks over the trunks of smooth-barked trees; *Tachydromia* nimbly runs over rocks in search of its minute prey; *Platypalpus* occurs in the summer and is pratal; while *Chersodromia* and its allies are dwellers of the sea shore and as a consequence of strong sea breezes are loath to take to the wing, in some instances having the wings abortive. In dry regions Empididæ are rare, having given place to their more powerful predatory relatives, the Asilidæ. As a group the Empididæ are boreal, though this impression may be the result of the large number of species recorded from the mountainous regions of Europe and North America. Species occur in the tropics, notably the yellow-bodied *Drapetis* and *Syneches*, and a not inconsiderable number are known from the southern hemisphere. The small size, soft consistency and usual lack of distinction make the Empididæ less desirable objects of collection than the more striking exotic species, which may explain why some regions have produced a meager Empid fauna to date.

Feeding Habits. — Most adult Empididæ are predatory on small insects, notably the Diptera, the proboscis being developed for piercing the bodies of their victims. To assist in the capture of prey one or the other pairs of legs are sometimes modified into raptorial organs fitted with holding spines. Various insects are caught for their body-juices, but principally Diptera are selected. The larger species of *Empis* prefer the Bibios, the smaller species of *Empis* procure Cecidomyias. The prey of *Hybos* is usually hymenopterous. Many Empididæ are cannibalistic, feeding either on smaller related forms or even on members of their own species. An interesting way of procuring a miscellaneous lot of insects is to imprison a swarm of dance-flies in the net and sort over the victims dropped by the Empids. Borboridæ, Chironomidæ, Blepharoceridæ, Bibionidæ, Psychodidæ, other Empididæ, etc., sometimes rare species, are in the haul, and nearly always not having been crushed by their captors are in good taxonomic condition.

Many Empididæ frequent flowers. Of the Anthophilous species, some, like *Anthalia*, have a soft proboscis and perhaps are dependent on nectar, while others, like *Empis* or *Drapetis*, use their mouthparts also for the capture of insect prey.

Anæmotropism. — Because of the strange habit of some species of *Rhamphomyia*, *Empis* and *Hilara* of hovering and zig-zagging in the air in swarms, the family designation of Dance-flies has

been bestowed on the Empididae. *Rhamphomyia sociabilis* is one of the most striking of the early summer species in Washington. By thousands it dances up and down, in and out, in an immense swarm, sometimes in the open and sometimes in the shade of a large tree. Some species of *Empis* skim back and forth in numbers among the tree tops. Reacting to air currents the group rises or lowers in a synchronized aerial dance. Other species of *Empis* and *Rhamphomyia* perform their dance closer to the ground, among underbrush or in the open spaces beneath forest trees. The species of *Hilara* and *Rhamphomyia* that hover over water weave in and out, in leisurely or in rapid erratic flight, barely missing the surface. The large metatarsi of the male *Hilaras* may touch the water but the insects are not wetted. In this surface dance the *Hilaras* may congregate on some small floating object and drift down stream, shortly to separate and return. Usually males predominate in the air-dances of the Empids.

Courtship. — No group of the Diptera offers more interesting and varied details of pygidial structure than the Empididae. With the elaboration of primary and secondary sexual structures in pygidium, legs, wings, mouth-parts and vestiture, there has evolved a complex series of mating habits. The significance of many of the cases of sex dimorphism is not understood, since the mating of most of the species has not been observed. The procuring of insect prey and the anæmotropic dances just described are manifestations of the courtship instinct. Hamm and Poulton have suggested an evolutionary sequence in the complexity of behavior, the stages in the use of prey being somewhat as follows.

1. Prey devoured by both sexes independently of mating. Examples, *Tachydromia*, *Hybos*, some species of *Empis*.
2. The prey provided by the male as a gift anticipatory to copulation is devoured, or sucked by the female during amplexus. Examples, *Pachymeria*, *Rhamphomyia*, *Empis* spp.
3. The prey or object provided by the male is not devoured by the female, but acts as a love-charm or stimulus to insure amplexus. Examples, *Hilara*, *Enoplempis*.

Where prey is passed from the male to the female, as in the second and third stages, the act is regarded as a physiological necessity for copulation. Since Empididae are cannibalistic the offering of prey to the female may have first functioned to divert her attention from her consort. Many anthophilous females have never been seen to imbibe insect juices except at the moment of copulation, dropping the prey furnished by their mates as soon as the sexes separate. With these species the use of insect prey is to be regarded as a philter, a stimulus rather than nourishment for the female.

A. T. Hamm has made extensive observations on the mating habits of English Empididae. While there are of course species differences in behavior, the descriptions of two types observed by him are here given as characteristic. *Empis trigramma* is one of the species that does not depend on the transfer of a love-token during the wooing process. A male alights on a leaf near a female, flutters the wings and raises and waves the front legs. The female responds with the same actions. The male then rubs the front tarsi together while vibrating the wings, and the female repeats these motions. As the two approach each other they caress each other's front tarsi, whereupon the female elevates the tip of the abdomen and the male deftly flies to her back and copulation ensues.

Among the species where the male lures his mate with prey Mr. Hamm observed the male of several species of *Empis* pounce on a fly, holding it with the posterior legs while it inserted the proboscis into the neck to pierce the ganglia behind. With this quieted offering the male either zig-zags over the resting females, or, in the case where the females are engaged in their aerial dance, flies into the circle, bearing the victim encircled by his middle legs. The selected female is chased, overtaken, and during a struggle of a few seconds duration the offering is transferred to the female, and the pair settles in copula to the herbage below. The male is apt to support his burden by hanging to a leaf by his front legs, tightly clasping his mate with the posterior legs. While copulating the female squeezes the

prey continuously, methodically inserting the proboscis in various places, but after a few minutes, dependent upon the size of the prey, she terminates the copulation, drops the prey, cleans her proboscis, and the two fly away.

The male of *Empis aerobatica* uses a small midge as a nucleus about which it fashions a large frothy balloon. Flying with this structure between its hind legs it attracts a female who alights on the back of her selected mate. The two then settle slowly to the ground, and after copulation has been completed discard the balloon. Some of the European species of *Hilara* have developed almost exactly the same method of balloon construction during courtship. In these cases it is the balloon rather than the original midge about which it was constructed that furnishes the stimulus for sexual selection because sometimes the minute insect is omitted. Balloons that have been dropped, probably after pairing, are utilized again by other males.

The appearance of so specialized a habit as balloon construction in *Empis* in America and in *Hilara* in Europe might suggest phyletic significance. It is more plausible, however, to regard the habit as a simultaneous outgrowth from the general habit of offering prey for sexual selection, the result of the ability to blow viscous bubbles.

TABLE OF THE SUBFAMILIES OF THE EMPIDIDÆ

- | | |
|--|------------------------|
| 1. <i>Discal cell always united with the second basal, three posterior cells, anal cell and the anal vein wanting or incomplete, auxiliary vein always vestigial and imperfect or wanting, third vein always simple, basal cells often large, alula absent, anal angle reduced, costa stopping at the fourth vein, usually no stigma; antennæ two- or three-jointed, with a long terminal or subdorsal two-jointed arista; proboscis short, vertical or inflexed, palpi one-jointed, more or less incumbent on the proboscis and often broad; pleuræ bare of hairs; pygidium asymmetrical, the uppermost valve unpaired; coxæ not elongate, femora often thickened and mucronate beneath; calyptères closely united to the base of the wing and with a small fringe; cursorial, terrestrial species.</i> | Subfam. TACHYDROMIINÆ. |
| <i>Anal cell and discal cell complete, or if either is incomplete the front coxæ are greatly lengthened and the front legs are raptorial or else the anal angle of the wing is rectangular, intercalary vein usually present thus making four posterior cells, auxiliary vein more or less distinct, third vein often furcate; proboscis often lengthened and palpi often porrect; pygidium with paired upper valves, except in Hybos, etc.</i> | 2. |
| 2. <i>Anal angle of the wing not projecting, the outline of the wing more or less cuneiform, costa continuing around the hind margin of the wing, anal crossvein acute, perpendicular or rounded, very rarely forming an obtuse angle, sometimes wanting, usually no fold in the wing under the humeral crossvein, no alula, calyptères closely united to the base of the wing, with a straight edge and with a small fringe; proboscis short, never longer than the head, either thick and fleshy or sharp and incurved; arista or style terminal; eyes broadly separated on the front, often pubescent, the face narrower than the front; thorax elongate, not highly arched, not pubescent, mesopleuræ obliquely longer than vertically high; front coxæ always longer than the posterior ones, no apical tibial spurs, empodium usually quite distinct although small; gressorial species.</i> | 3. |

Anal angle of the wing almost always more or less projecting, often rectangular, rarely the wings uniformly tapering to the base, in which case the high mesopleura or thin hind margin of the wing are distinctive, a transverse fold in the wing membrane between the origin of the fourth vein and the humeral crossvein; males often holoptic; proboscis usually rigid; mesopleura distinctly higher than long; front coxae not elongate nor the front legs raptorial, empodium at most microscopically setuliform, usually invisible; flying species 4.

3. *Front legs raptorial, located near the head and distant from the posterior pairs, the front coxae greatly lengthened, nearly as long as their femora and two or more times as long as the posterior coxae, the front femora more or less thickened and setulose and setose beneath; palpi minute; antennae inserted below the middle of the head; body and legs often yellow; neuration often reduced, pedicel of the second and third veins arising nearer the anterior crossvein than to the humeral crossvein*

Subfam. HEMERODROMIINÆ.

Legs slender, not raptorial, the front pair not distant from the others, the mesosternum never greatly longer than the prosternum, neither the front coxae greatly lengthened nor the front femora greatly thickened; antennæ usually above the middle of the head; body often olivaceous black, thorax flattened before scutellum; neuration complete, pedicel of the second and third veins arising nearer the humeral crossvein than to the anterior crossvein, auxiliary vein always distinct

Subfam. CLINOCERATINÆ.

4. *Anal crossvein recurved or abruptly reflexed and confluent with the under side of the anal cell, the anal vein usually represented as an independent fold disjoined from the anal cell, the crossvein perpendicular only in Parathalassius, auxiliary vein strong and always distinct from the first vein, third vein forked or not, if not the metapleura setose or the basal cells very small or the anal cell very narrow, always three posterior veins, basal cells long and coextensive, except in the Microphorus group; antennæ obviously three-jointed, except in Hormopeza, the basal joints often more or less setose, the first joint rarely shorter than the second; proboscis often elongate, usually bent back, rarely projecting, palpi often long and hairy, especially toward the base; eyes of the female always broadly separated, fronto-orbitals often present; face quadrate or arched, almost never very narrow; metapleura often with bristles; posthumerals and intra-alars usually present, unless all the bristles are secondarily reduced; no ovipositor, the female with two styles; middle tibiae usually with an apical bristle inside, hind tibiae often with extensor bristles.*

Subfam. EMPIDINÆ.

Anal crossvein forming a distinct angle, either acute, right or slightly obtuse, with the under side of the anal cell which is continuous with the anal vein, recurved only in Anomalempis, third vein forked only in some Brachystomatinae, often only two posterior veins; base of the antennæ without strong setæ; proboscis rarely longer than the length of the head, often porrect, palpi short, at most with a few setæ toward the tip; face often narrow; no fronto-orbital or intra-alar bristles, posthumerals present only in Trichina and Bicellaria; thorax often large and highly arched, meta-

- pleura always devoid of bristles, sometimes with short pile in front of the spiracle.* 5.
5. *Anal cell shorter than or about as long as the second basal, its posterior angle obtuse, or the anal crossvein perpendicular to the anal vein, third vein not forked, basal cells moderately long, costa stopping at the fourth vein, auxiliary vein weak and lying close to the first vein; proboscis usually short, rarely porrect; thorax seldom excessively convex.* Subfam. OCYDROMIINÆ.
- Anal cell as long as the second basal or longer, its outer angle acute, the anal crossvein never reflexed but closing the anal cell toward the margin of the wing, basal cells typically long; auxiliary vein distinct and separate from the first vein.* 6.
6. *Discal cell emitting two veins, if three the basal cells are very short, auxiliary vein continuing beyond the middle of the wing, costa stopping at the third or the fourth vein, anal angle of the wing rectangular, alula sometimes present, third vein simple, basal joints of the antennæ connate, the antennæ therefore apparently two-jointed, a long arista usually present; proboscis rigid, projecting forward; eyes of both sexes typically meeting, upper facets largest, a horizontal line on the eye at the level of the antennæ; ocellar triangle prominent and located on the vertex; thorax greatly arched; hind legs usually stout and setose.* Subfam. HYBOTINÆ.
- Discal cell emitting three veins, auxiliary vein ending in the costa before the middle of the wing, costa continuing around the entire wing, anal angle of the wing weak or wanting, third vein usually forked; no ocellar prominence; antennæ three-jointed; proboscis short, sharp and incurved; upper facets smallest, but no horizontal differentiating line; thorax comparatively small, the abdomen elongate and in the female apically fimbriate.* Subfam. BRACHYSTOMATINÆ.

SUBFAMILY BRACHYSTOMATINÆ

Characters. — Elongate slender, nearly glabrous species of moderately large size. Head globose, eyes bare, with uniform facets, nearly or quite contiguous on the face; antennæ located high up on the head, projecting horizontally, three-jointed; proboscis very short. Thorax much shorter than the abdomen. Legs long and slender, sometimes differing in structure in the two sexes, the femora always slender, the tibiæ of the males sometimes apically swollen or deformed, no tibial spurs. Wings cuneiform and slender, costa encompassing the entire margin, the anal angle not or but little developed, auxiliary vein distinct, first vein ending beyond the middle of the wing, stigma normally present, third vein furcate or simple, the fork extending obliquely to the costa or perpendicularly to the second vein, basal cells very long, discal cell complete, emitting three simple posterior veins, anal cell longer than the basals, its crossvein continuing toward the margin, rounding or acutely closing the anal cell, anal vein strong, usually abruptly ending before the margin, no alula; calypteres minute.

TABLE OF THE GENERA OF THE BRACHYSTOMATINÆ

1. *Third vein forked; anal crossvein acutely closing the anal cell; stigma present; palpi broadened.* 2.
- Third vein simple; anal cell with rounded extremity; eyes widely separated on face and front (female); no stigma; palpi slender and setose, antennæ elongate and with long style (PI. 5, FIG. 38). Genus ANOMALEMPIS, nov. gen.*

I. GENUS BRACHYSTOMA, MEIGEN

Brachystoma, Meigen, Syst. Beschr. Vol. 3, p. 12 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 344 (1834); Zetterstedt, Ins. Lapp. p. 557 (1838); Blanchard, Hist. Nat. Ins. Vol. 3, p. 582 (1840); Zetterstedt, Dipt. Sc. Vol. 1, p. 360 (1842); Boitard, Nouv. Man. Vol. 3, p. 321 (1843); Rondani, Dipt. Ital. Prodr. Vol. 1, p. 150 [1856] (*Brachistoma*); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Schiner, Fauna Austr. Dipt. Vol. 1, p. 117 (1862); Lioy, Atti Inst. Venezia 1864, p. 603 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 121 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 259 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 247, 262, 263 (1903); Melander, Williston's 3d. Man. p. 224 (1908); Kertész, Cat. Dipt. Vol. 6, p. 1 (1909); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 516 (1910).

Blepharoprocta, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 194 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 28, p. 389 (1895); Williston, Manual, p. 74 (1896) (*Blepharoproctus*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 261 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 246, 263 (1903); Melander, Williston's 3d. Man. p. 224 (1908); Kertész, Cat. Dipt. Vol. 6, p. 2 (1909); Bezzı, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 514 (1910).

Heterophlebus, Philippi, Verh. Zoo.-bot. Ges. Wien, Vol. 15, p. 764 (1865).

Characters.—No ocellar prominence, the vertex rounding into the occiput; eyes subcontiguous on the face, distinctly separated above the antennæ, the sides of the front parallel to the ocelli; basal joints of the antennæ together longer than the third which is conical and shorter than the slender terminal arista; proboscis thick, inflexed, palpi small and broad. Thorax short, rather flattened above, the scutellum small, legs close together, no pubescence nor bristles beyond a few microscopic prescutellar dorsocentral hairs and single supraalar and postalar bristles, scutellum with two or more hairs; pleuræ bare but pollinose. Seventh tergite of the female abdomen typically inflated to a large translucent bladder, open apically as a vertical slit, the seventh sternite closing the vesicle below, but in *Blepharoprocta* this segment is compressed and not larger than the preceding and has a more strongly ciliated opening; pygidium small, retracted or exposed, with central filament and two pairs of small narrow lateral valves, sometimes enclosed in a projecting ventral fringe. Legs elongate, slender, femora usually with some setæ beneath, tibiae and tarsi of males of *Blepharoprocta* sometimes greatly deformed. Wings with reduced anal angle, third vein forked, basal cells large, four posterior cells, anal cell long and acute, no basal bristle on the costa.

The sexual dimorphism of *B. serratula* is most striking. Besides the genital differences, the legs of the male are curiously modified and even its neuration is changed. In this species some of the hairs of the hind margin of the wing are bent forward on the inner surface to form a long curved fringe.

NOMENCLATURE

Type species : *B. vesiculosum*, Fabricius (Pl. I, Fig. 2). Westwood in 1840 designated *longicornis* Meigen as the type, a species which Rondani later made the type of his genus *Trichopeza*, but Blanchard, also in 1840, selected *vesiculosum* as the type of *Brachystoma*, a procedure which has been since followed.

There seems to be no satisfactory distinction between *Brachystoma* and *Blepharoprocta*. The latter group, as exemplified by *binummus*, has the marginal cell closed, the end of the female abdomen not inflated, the male legs distorted and there is a prominent fringe on the pygidium. None of these characters occur in *Brachystoma*. However, in *Robertsonii*, which was described as a *Brachystoma*, the marginal cell is open, but the pygidial fringe is lacking and the abdomen of the female is narrow. In *nigrimana*, which Coquillett considered the type of *Blepharoprocta*, the legs are simple and the pygidial fringe is quite inconspicuous. I have a specimen of *nigrimana* lacking the fork of the third vein in one wing, showing that this character also is unstable. *Serratula* and *binummus* are the female and male respectively of the same species. *Heterophlebus* was proposed by Philippi shortly after the publication of *Blepharoprocta*, and was based on the same character, the closure of the marginal cell.

Geographical distribution.

SUBGENUS BLEPHAROPROCTA, LOEW

1. *B. ambigua*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 764 [1865] Chile.
(*Heterophlebus*).
2. *B. melanogastra*, Philippi, ibidem, Vol. 15, p. 764 [1865] (*Heterophlebus*). Chile.
3. *B. nigrimana*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 194: Cent. 2, no 17 E. United States.
[1862] (*Brachystoma*); Melander, Trans. Amer. Ent. Soc. Vol. 28,
p. 261 (1902).
4. *B. serratula*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 319, Vol. 6, p. 194: E. United States.
Cent. 1, no 23 [1862] (*Brachystoma*); Melander, Trans. Amer. Ent.
Soc. Vol. 28, p. 262 (1902). — Pl. 8, Fig. 77.
binummus, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 193 : Cent. 2, no 16 [1862]
(*Brachystoma*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 261, f. 91, 92
(1902).
5. *B. thoracica*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 764 [1865] Chile.
(*Heterophlebus*).

SUBGENUS BRACHYSTOMA, MEIGEN

6. *B. bicolor*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 130 (1889). Chile.
- . *B. distinguendum*, Schiner, Fauna Austr. Dipt. Vol. 1, p. 117 (1862) no description.
7. *B. fuscum*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 763 (1865). Chile.
8. *B. leptidium*, Philippi, ibidem, Vol. 15, p. 762 (1865). Chile.
9. *B. nemorale*, Philippi, ibidem, Vol. 15, p. 764 [1865] (*Heterophlebus*). Chile.
10. *B. nigricorne*, Philippi, ibidem, Vol. 15, p. 763 (1865). Chile.
11. *B. obscuripes*, Loew, Neue Beitr. Dipt. Pt. 4, p. 37 (1856). Sardinia.
12. *B. occidentale*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 260, pl. 8, f. 89 (1902). W. United States.

13. *B. ? quadricinctum*, Fabricius, Syst. Antl. p. 148 [1805] (*Damalis*); Wiedemann, Auss. Zweifl. Ins. Vol. 1, p. 537 [1828] (*Hybos*); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 314 [1909] (*Syneches*). South America.
14. *B. Robertsonii*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 393 (1896); Illinois, Tennessee, Ohio. Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 260 (1902).
15. *B. * spinulosum* Loew, Berendt, Organ. Reste Bernstein, Vol. 1, p. 57 (1845) unnamed; Loew, Bernsteinfauna, p. 41 (1850) undescribed; Giebel, Ins. Vorwelt, p. 209 (1856); Meunier, Miscell. Ent. Vol. 7, p. 178 (1899). Baltic Amber. Lower Oligocene.
16. *B. stigmaticum*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 763 (1865). Chile.
17. *B. testaceum*, Philippi, ibidem, Vol. 15, p. 763 (1865). Chile.
18. *B. vesiculosum*, Fabricius, Ent. Syst. Vol. 4, p. 299 [1794] (*Syrphus*); Syst. Antl. p. 200 [1805] (*Baccha*); Meigen, Syst. Beschr. Vol. 3, p. 13, pl. 22, f. 8, 9 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 344, pl. 8, f. 5 (1834); Boitard, Man. Ent. Vol. 3, p. 321 (1843); Schiner, Fauna Austr. Dipt. Vol. 1, p. 117 (1862); Girschner, Ent. Nachr. Vol. 20, p. 63, 244 (1894); Mik, Wien, Ent. Zeit. Vol. 16, p. 36, fig. (1897). — **Pl. I, Fig. 2.**
var. *flavicollis*, Mik, Wien, Ent. Zeit. Vol. 6, p. 103 (1887); Strobl, Mitth. Naturw. Ver. Steiermark, Vol. 29, p. 42 (1893). S. Europe.
19. *B. vittigerum*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 765 [1865] Chile.
(*Heterophlebus*).

2. GENUS HOMALOCNEMIS, PHILIPPI

Homalocnemis, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 752 (1865); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 121 (1889); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301 (1909); Kertesz, Cat. Dipt. Vol. 6, p. 8 (1909).

Characters. — Species of five or six mm. length; head globose, thorax very strongly gibbous, abdomen slender. Eyes contiguous above the antennæ, separated below; antennæ three-jointed, the third joint long and tapering and merging into the long apical aristæ; proboscis and palpi small and retracted. Legs slender, the femora not thickened, in the genotype the apex of the tibiæ and the basal joints of the tarsi of the male dilated. Wings dark-colored, stigma strong and long, second vein curving forward at tip to meet the costa, third vein forked, discal cell large, complete, emitting three veins, basal and anal cells long.

Type species: *Homalocnemis nigripennis* Philippi, from South America (**Pl. 5, Fig. 41**). The recently described *Brachystoma adelensis* Miller from New Zealand, apparently belongs here. The full venation is somewhat suggestive of Bombyliidæ but the habits of the New Zealand species, in darting hither and thither over the surface of pools of water, show the Empidine affinities.

Geographical distribution.

1. *H. adelensis*, Miller, Trans. New Zeal. Inst. Vol. 45, p. 203-206, f. 11-14, New Zealand.
pl. 1, f. 2 [1913] (*Brachystoma*).
2. *H. nigripennis*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 752, pl. 29, Chile.
f. 56 (1865). — **Pl. 5, Fig. 41.**

3. GENUS ANOMALEMPIS, NOV. GEN.

Characters. — Head globose, no ocellar prominence; front (female) occupying about one-sixth the width of the head, slightly widening toward antennæ, face somewhat broader; occiput bristly

with stiff hairs; facets nearly uniform; antennæ porrect, the basal joints minute, together about one-third as long as the tapering third joint which ends in a thickened style three-fourths as long as the third joint; palpi slender and setose, proboscis inflexed, not tapering, as long as antennæ. Thorax arched, about ten hairs in the dorsocentral series, one notopleural, two supra-alar and one intra-alar bristles developed, scutellum quadrisetose. Seventh abdominal segment (female) long and tubular, apically fringed as in *Blepharoprocia*. Legs simple, middle femora with a few flexor setæ, hind tibiæ with extensor setæ. Wings hyaline, costa encompassing hind margin, auxiliary vein extending straight into costa, third vein not forked, discal cell emitting three posterior veins, basal cells long, the anal cell with rounded apex, the anal crossvein extending towards margin but with recurved tip, anal vein beyond crossvein, represented by a fold, anal angle of wing shallowly rounded but evident.

Genotype the following species.

Geographical distribution.

1. *A. tacoma*, nov. sp. — Pl. 5, Fig. 38 (1).

Washington.

SUBFAMILY HYBOTINÆ

Characters. — Thorax greatly convex, head hemispherical, eyes large, bare, contiguous above the antennæ at least in the male, so as to crowd out the front, the ocelli located on a prominent tubercle situated on the very summit of the head, upper facets larger than the lower, a distinct horizontal line extending across the eyes at the level of the antennæ; proboscis porrect, usually styliform, short in the aberrant Tasmanian genera *Ironomyia* and *Sciadocera*, palpi never broad, sometimes linear and projecting, sometimes short; antennæ small, three-jointed, but the basal joints immovably fused, no strong setæ on the antennæ, third joint more or less elongate oval, arista long, very thin and bare, usually terminal, rarely subdorsal, in *Meghyperus* sometimes replaced by a thick style; no cephalic bristles, a pair of small ocellar hairs. Humeri pronounced, disk of the mesonotum devoid of true bristles, sometimes pilose, sometimes a prescutellar pair of reduced dorsocentrals, two or three notopleural bristles present, scutellum margined with hairs of which the apical two are sometimes bristle-like; pleuræ entirely devoid of hairs. Abdomen comprising eight segments and the pygidium or true ovipositor, the pygidium small and bilaterally constructed in the *Syneches* and *Meghyperus* groups, or usually larger and consisting of opposing upper and lower valves in the *Hybos* group. Hind legs usually large, with their coxæ the strongest pair, their femora incrassate and spinose beneath and longer than their tibiæ; anterior tibiæ and tarsi often furnished with long bristles, hind tibiæ clavate in *Meghyperus* and *Syndyas*; pulvilli large, empodium minute and bristly. Calypteres rather large, separated from the root of the wing and densely fringed. Wings rather broad, the anal angle strongly developed, rectangular, costal margin hairy, costal bristle small, costa continuing to the fourth vein, humeral crossvein well formed, auxiliary vein straight, usually parallel with and close to the first vein and vanishing before it attains the costa, first vein ending beyond the middle of the wing, stigma usually more or less evident, sometimes prominent with the second vein curving down around it, discal cell usually large and elongate, situated

(1) *Anomalempis tacoma*, nov. sp. — Female. Length 2.5 mm. Black species, occiput subshining, front and face brown pollinose, mouthparts and antennæ black. Mesonotum polished, its hairs black, pleuræ entirely dull pollinose. Abdomen polished, its fine hairs pale. Halteres pale yellow, calypteres and fringe sordid. Wings hyaline, only the faintest indication of a stigma, costal sections 1 : 1 : 0.6 : 0.8 (end of third vein a little beyond wing-tip) : 0.3 : 0.4 : 0.4, underside of discal cell 3 : 2, sections of fifth vein equal.

A single specimen of this curious fly was discovered during August 1917, in Paradise Park on Mt. Rainier, Washington.

in the middle of the wing, emitting two posterior veins of which the anterior is furcate in two of the genera, in the aberrant Tasmanian genera *Ironomyia* and *Sciadocera* there are three posterior veins and the basal cells are small, otherwise the basal cells long, nearly equal in extent as the anterior crossvein is placed well toward the base of the discal cell, anal crossvein never recurved but continuing perpendicularly to the wing axis toward the hind margin, the anal cell, always formed, is thus longer than the second basal and its outer lower angle is acute. A distinct alula is present in *Meghyperus*.

This subfamily is the best circumscribed of all. By the earlier writers it was considered as a distinct family, the *Hybotidæ*, but during the last half century it has been amalgamated with the *Empididæ*. In the *Hybotinæ* the anal crossvein, the chief character of the subfamily, continues more or less parallel with the hind margin so as to elongate the anal cell, but in *Meghyperus* it is so curved as to meet the anal vein almost at a right angle.

Meghyperus forms an aberrant genus in the *Hybotinæ*. The presence of a strong alula, the separation of the auxiliary vein, the forked fourth vein, the broad front of the female and the lack of a long arista, are all weighty characters at variance with the other genera and suggestive of the genus *Prorates* Melander, which is discussed at the conclusion of this fascicle as being removed to the *Bombyliidæ*.

TABLE OF THE GENERA OF THE HYBOTINÆ

1. Basal cells long, discal cell present and emitting 2 posterior veins; proboscis projecting forward	2.
Basal cells short, 3 posterior veins, discal cell present or open; arista long and filiform; femora simple; proboscis short; wings broad, costa convex	12.
2. Antenna with a long filiform arista; eyes contiguous above the antenna in both sexes; hind femora often swollen; discal cell emitting two simple posterior veins, auxiliary vein very close to the first vein, no alula	3.
Antenna with a style, eyes of ♀ broadly separated on the front; legs without bristles, hind femora slender; fourth vein forked, auxiliary vein distinctly separated from the first vein which passes through the stigma, alula present; pygidium with lateral valves (Pl. 5, Fig. 40)	Genus MEGHYPERUS, Loew.
3. Pedicel of the second and third veins long, arising before the middle of the basal cells; pygidium with lateral valves; scutellum with several marginal bristles or hairs	4.
Pedicel of the second and third veins short, arising beyond the middle of the basal cells; pygidium with dorsal and ventral valves; scutellum with two or but few marginal bristles	7.
4. Hind femora slender, not thicker nor longer than their tibiae, elongate and not armed beneath with setæ or tubercles; arista usually subapical; middle tarsi ♂ sometimes deformed; dorsocentral, scutellar and tibial bristles undeveloped (Pl. 1, Fig. 7)	Genus PARAHYBOS, Kertész.
Hind femora more or less swollen, thicker and longer than their tibiae, and variously armed beneath with setæ, spines or tubercles (Genus SYNECHES, Walker)	5.
5. Third and fourth veins parallel or diverging; color of the body usually dark or black and the wings sometimes spotted (Pl. 1, Fig. 4)	Subgenus SYNECHES, Walker.
Third and fourth veins converging, the first posterior cell narrower at its apex than opposite the posterior crossvein; color often yellow and wings not spotted	6.

6. *Hind femora setose beneath but not armed with spinose tubercles* Subgenus EPICEIA, Walker.
Hind femora ♂ armed beneath with tubercles which bear spines or strong setæ, hind tibiae ♂ curved and tuberculate (Pl. I, Fig. 6) Subgenus HARPAMERUS, Bigot.
7. *Vein between the first and second basal cells distinct, discal cell never much shorter than the second posterior cell, usually much longer; tibiae not clavate* 8.
Vein between the first and second basal cells very weak, the first basal broader than the second, discal cell much shorter than the second posterior; hind tibiae usually clavate (Pl. 5, Fig. 39) Genus SYNDYAS, Loew.
8. *Hind femora not thickened and entirely unarmed; arista stiff and shorter than the broadly conical third joint; first and second veins ending close to the tip of the wing; proboscis horizontally porrect and slender (Pl. 5, Fig. 42)* Genus ACARTERUS, Loew.
Hind femora more or less thickened and armed beneath with spinous bristles; arista hair-like and much longer than the third joint of the antennæ 9.
9. *Third and fourth veins parallel or diverging; eyes separated on the face; proboscis slender, projecting, about as long as the head, palpi prominent; mesonotum and pleuræ usually pollinose, the pubescence of the thorax sparse, usually a pair each of dorsocentral and scutellar bristles; usually no ovipositor; hind tarsi with minute black thorns beneath, hind femora not extraordinarily thickened (Pl. I, Fig. 5)* Genus HYBOS, Meigen.
Third and fourth veins somewhat convergent, the first posterior cell narrowed; eyes contiguous or nearly so beneath the antennæ in both sexes; no dorsocentral bristles; last segments of ♀ abdomen lengthened forming an ovipositor 10.
10. *Third joint of the antennæ elongate ovate; tibiae without tooth; proboscis projecting forward; head not unusually small; hind femora often tuberculate and always greatly swollen* 11.
Third joint of the antennæ semicircular, the convex side up; front tibiae with a strong subbasal flexor tooth; proboscis perpendicular (abnormal?) palpi small; shining metallic species with very small head Genus CERATHYBOS, Bezzi.
11. *Body covered with tomentum, disc of the mesonotum bare of pile; proboscis about as long as the head, slender for piercing, palpi elongate (Pl. 5, Fig. 45)* Genus LACTISTOMYIA, Melander.
Body shining, often metallic, devoid of tomentum, the mesonotum more or less densely pilose; proboscis short, not constructed for piercing, the labellæ when appressed forming a compressed lamella, palpi short (Pl. I, Fig. 3) Genus EUHYBOS, Coquillett.
12. *Second and third veins approximate and parallel, a heavy stigma present beyond end of first vein, discal cell complete, anal cell longer than basals; thorax slightly arched, its bristles weak (Pl. 8, Fig. 86)* Genus IRONOMYIA, White.
Second and third veins diverging, no stigma, discal cell absent, anal cell shorter than basals; eyes of female widely separated; thorax greatly arched, its bristles pronounced Genus SCIADOCERA, White.

I. GENUS HYBOS, MEIGEN

Hybos, Meigen, Illiger's Mag. Ins. Vol. 2, p. 269 (1803); Syst. Beschr. Vol. 2, p. 261 (1820); Macquart, Mem. Soc. Sc. Lille, 1823, p. 143 (1823); Dipt. N. France, Vol. 3, p. 147 (1827); Hist. Nat. Dipt. Vol. 1, p. 318 (1834); Curtis, Brit. Ent. Vol. 8, p. 661 (1837); Zetterstedt, Fauna Ins. Lappon, p. 535 (1838); Westwood, Gen. Syn. p. 133 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 233 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 120 (1851); Rondani, Dipt. Ital. Vol. 1, p. 153 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 559 (1857); Schiner, Fauna Austr. Dipt. Vol. 1, p. 77 (1862); Lioy, Atti Inst. Venet. 1864, p. 719 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 115 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Wheeler & Melander, Biol. Centr.-Amer. Dipt. Suppl. p. 372 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 245 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 251, 259 (1903); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 324 (1904); Vol. 3, p. 425 (1905); Melander, Williston's Man. N. Amer. Dipt. 3 ed. p. 224 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301, 305 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 553 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 8 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 42, 46 (1910); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 336 (1920).

Acromyia, Bonelli, in Latreille (not Leach, Lioy or Coquillett). Gen. Crust. Ins. Vol. 4, p. 305 (1809); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 329 (1904).

Noeza, Meigen, Nouv. Classif. Mouches, p. 27 (1800); Hendel, Verh. Zool.-bot. Gez. Wien, Vol. 58, p. 56 (1908); Kertesz, Cat. Dipt. Vol. 6, p. 3 (1909); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 453 (1912); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 576 (1910).

Characters. — Black-colored species measuring four to five millimeters, with downward curved abdomen, light pollinose coating and long, spinose hind femora. Head rather globular, the eyes large, meeting above the antennæ up to the prominent ocellar triangle which is placed on the summit of the vertex, below the antennæ the eyes are approximated but leave a distinct shortened face; no cheeks; proboscis slender, projecting forward as far as the length of the head, the palpi long and slender; antennæ short, located below the middle of the head, the basal joints fused, the third joint oval and with a long slender terminal arista; no cephalic bristles. Thorax highly arched, less convex before the scutellum, prothorax quadrangular, appearing as a neck; rows of hair-like acrostichal and dorsocentral bristles, humeri with hairs, two or three notopleural, two scutellar and one small postalar bristles present; pleuræ pollinose but bare of hairs. Abdomen cylindrical, the pygidium more or less globose and large, asymmetrical, with dorsal and ventral convex distorted plates; eighth segment of the female normally retracted giving a blunt termination to the abdomen. Anterior legs hairy and bristly, especially in the male, hind legs elongate, the femora rather clavate and spinose beneath, hind tibiæ shorter than their femora, devoid of spurs, anterior tibiæ usually with apical bristles. Wings with parallel margins, anal angle prominent and rectangular, veins strong, costal margin closely short-hairy, very rarely spinose, second vein ending near the tip of the wing, stigma weak, pedicel of the second and third veins arising beyond the middle of the second basal cell, third and fourth veins both curving backward, the first posterior cell not narrowed in the margin, discal cell narrow and long, complete, sending two simple posterior veins to the margin, basal cells long and equal, the anal crossvein forming an acute angle with the anal vein, no alula. Halteres most often light colored.

Type species : *H. grossipes* Linnaeus. Meigen's first species was *funebris*, which is synonymous with *grossipes*. Curtis, in his British Entomology (1837), designated this as the type.

Hendel would overthrow the classic name *Hybos* in favor of the doubtful name *Noeza*, and in this course unfortunately he has been followed by Kertész, Bezzi and Coquillett. This opens up the question of the general acceptance of the abandoned names of Meigen's 1800 paper, which have been contested by many zoologists. As in this early paper Meigen mentioned no species by name, thus disregarding binomial nomenclature, as identification through his brief and broadly applicable descriptions is uncertain, and as it is impossible to determine the genotypes from the data in this paper alone, it appears that nothing is to be gained by resurrecting the names of 1800 but much is to be lost by the wholesale replacement of scores of generic names in common usage by names of questionable application. An extended discussion to this effect has already been penned by the writer (see, Melander: Psyche, Vol. 17, p. 41-47 (1910). It is unfortunate that when the International Commission on Zoological Nomenclature had this specific case under consideration it decided in Opinion 28 (Smithsonian Publication 1989, p. 66, Oct. 1910) only that Meigen's 1800 paper had actually been published. However, in Opinion 46, regarding the status of genera for which no species was distinctly named in the original publication, the Commission has rendered its opinion that a genus becomes a *genus dubium* when it is impossible to recognize any one of the original species for the genotype from the original description. It is better to let these questionable genera remain doubtful than to attempt at this late date to fix their genotypes, with the consequent overthrow of well known names of such dominant genera as *Rhamphomyia*, *Platypalpus*, *Hybos* and *Clinocera*. Therefore throughout this fascicle Meigen's « Erstlingsarbeit » will not be seriously considered.

The species of *Hybos* are æstival, but occur among the shrubbery of woods. There are relatively but few species, one only in the United States and three in Europe.

TABLE OF THE NEW WORLD SPECIES OF HYBOS

1. Costa nearly straight, if bowed the costal cell is broadened, no costal hooks	2.
Costa rounded, parallel with first vein and bearing strong curved spines.	H. SPINICOSTA, Wheeler & Melander.
2. Front metatarsi longer than or nearly as long as their tibiæ, hind femora not spinose, hind metatarsi not spinulose, middle tibiæ with very long setæ.	3.
Front metatarsi evidently shorter than their tibiæ, hind femora with spines, hind metatarsi with small black thorns below (as far as known).	5.
3. Coxæ and legs wholly yellowish; third antennal joint elongate oval; no dorsocentrals.	H. XANTHOPODUS, nov. sp.
Coxæ and legs almost wholly black; two dorsocentrals	4.
4. Two intermediate flexor setæ on middle tibiæ; third antennal joint subulate; halteres whitish.	H. TYPICUS, Wheeler & Melander.
One flexor seta at middle of middle tibiæ; third antennal joint ovate; halteres piceous.	H. ECTORUS, nov. sp.
5. Thorax more or less dusted; eyes separated on the face.	6.
Thorax shining; eyes contiguous or subcontiguous below antennæ; halteres dark	11.
6. Halteres whitish; dorsocentrals present.	7.
Halteres blackish; base of hind tibiæ yellowish; no dorsocentrals.	H. LURIDUS, Bezzi.

7. Dorsocentrals strong and black; setæ of middle tibiæ very long 8.
 Dorsocentrals weak and pale; setæ of middle tibiæ short, none
 on flexor side (*H. reversus* Walker) 10.
8. Anterior femora, hind knees, middle tibiæ and posterior tarsi
 luteous. 9.
 Femora black, tibiæ and tarsi yellowish; wings subhyaline *H. HALTERALIS*, Bezzii.
9. Wings brown; ten spines in anterior row under hind femora.
 Wings hyaline; seven or eight spines in anterior row under
 hind femora *H. SCIAPTERUS*, nov. sp.
10. Wings of male brownish gray on apical two-thirds
 Wings of both sexes grayish hyaline except for the stigma. *H. CONIATUS*, nov. sp.
11. Legs black, only the tarsi yellow. *H. REVERSUS*, Walker, s. str.
 Knees and base of hind tibiæ yellowish. var. *SLOSSONÆ*, Coquillett.
H. MELLIPES, Wheeler & Melander.
H. NEOTROPICUS, Bezzii.

Geographical distribution.

1. *H. apicis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 12 (1913); Fauna Brit. Burma.
 Ind. Dipt. Vol. 1, p. 340, f. 26, 27 (1920).
2. *H. auripes*, Brunetti, Rec. Indian Mus. Vol. 9, p. 14 (1913); Fauna Brit. India.
 Ind. Dipt. Vol. 1, p. 341 (1920).
3. *H. Bezzii*, Kertész, Termes. Fuezet. Vol. 22, p. 175 (1899); Bezzii, Ann. New Guinea.
 Mus. Hungar, Vol. 2, p. 328 (1904).
4. *H. bisetosus*, Bezzii, ibidem, Vol. 2, p. 324 (1904), Vol. 10, p. 454 [1912] East Indies.
 (*Noesa*); Suppl. Ent. Berlin, Vol. 3, p. 67 [1914] (*Noesa*); Brunetti,
 Fauna Brit. Ind. Dipt. Vol. 1, p. 338 (1920).
5. *H. brachialis*, Rondani, Ann. Mus. Stor. Nat. Genova, Vol. 7, p. 446 Borneo.
 (1875); Bezzii, Ann. Mus. Hungar, Vol. 2, p. 326, note (1904).
6. *H. brachystigma*, Bezzii, ibidem, Vol. 2, p. 327 (1904). New South Wales.
7. *H. brunnipes*, Brunetti, Rec. Indian Mus. Vol. 9, p. 12 (1913); Fauna
 Brit. Ind. Dipt. Vol. 1, p. 343 (1920). India.
- . *H. brunnipes*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 260 (1829) no
 description.
8. *H. coniatus*, nov. sp. (1). Costa Rica.
9. *H. culiciformis*, Fabricius, Syst. Ent. p. 796 [1775] (*Asilus*); Fabricius,
 Spec. Ins. Vol. 2, p. 466 [1781] (*Asilus*); Fabricius, Mant. Ins.
 Vol. 2, p. 361 [1787] (*Asilus caliciformis*); Olivier, Encycl. Méth.
 Ins. Vol. 4, p. 270 [1799] (*Asilus*); Fabricius, Ent. Syst. Vol. 4,
 p. 389 [1794] (*Asilus*); Schrank, Fauna Boica, Vol. 3, p. 161 [1803]
 (*Asilus*); Fabricius, Syst. Antl. p. 172 [1805] (*Dasypteron*); Schiner,
 Fauna Austr. Dipt. Vol. 1, p. 78 (1862); Strobl, Mitteil. Naturw.
 Ver. Steiermark, Graz, Vol. 29, p. 44 (1892); Lundbeck, Dipt. Dan.
 Vol. 3, p. 15, 17 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 46
 (1910); Carter, Ent. Mag. Vol. 47, p. 161 (1911), Vol. 48, p. 59
 (1912); Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37(3), p. 8 (1913). Europe.
- funebris*, Fallen (not Meigen), Empid. Suec. var. p. 5 (1815).
- infuscatus*, Zetterstedt, Dipt. Scand. Vol. 1, p. 234 (1842), Vol. 8, p. 2993
 (1849); Wahlgren, Ent. Tidskr. Vol. 31, p. 46 (1910).

(1) *Hybos coniatus*, nov. sp. — Very close to *sciapterus*, differing as follows: Length 4 mm.; third antennal joint orbicular, arista three and one-half times antennal length; fourth abdominal segment with dorsal patch of dust in place of the band; hind femora much more slender, usually with eight spines in anterior row beneath; pygidial fringes usually less pronounced; wings clear hyaline, stigma weak; often the flexor seta at basal third of middle tibia is quite long.

Six males, nine females, La Suiza de Turrialba, Costa Rica, Pablo Schild, collector.

- rufitarsis*, Zetterstedt, Dipt. Scand. Vol. 8, p. 2994 (1849); Lundbeck, Dipt. Dan. Vol. 3, p. 18 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 46 (1910).
vitripennis, Meigen, Syst. Beschr. Vol. 2, p. 348 (1820); Macquart, Dipt. N. France, Vol. 3, p. 149 (1827); Hist. Nat. Dipt. Vol. 1, p. 319 (1834); Curtis, Brit. Ent. Vol. 8, p. 661 (1837); Zetterstedt, Fauna Ins. Lappon, p. 535 (1838); Dipt. Scand. Vol. 1, p. 234 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Zetterstedt, Dipt. Scand. Vol. 8, p. 2994 (1849); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 485 (1849); Dahlbom, Svenska Vet. Akad. Handl. p. 160 (1851); Scholz, Zeitsch. Ent. Breslau, Vol. 5 (17), p. 48 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 121 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 4267 (1852); Vol. 13, p. 4978 (1859).
10. *H. discoidalis*, Meijere, Tijdsch. v. Ent. Vol. 56, Suppl. p. 66 (1914). Java.
11. *H. ectorus*, nov. sp. (1). Costa Rica.
12. *H. *exilis*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 91, 109, pl. 7, f. 8, 9 (1908). Baltic Amber.
13. *H. femoratus*, Mueller, Zool. Dan. Prodr. p. 181 [1776] (*Asilus*); Walker, Ins. Brit. Dipt. Vol. 1, p. 121 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 78 (1862); Giard, Traité Ent. Vol. 3, p. 992, pl. 108, f. 6 (1885); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 44 (1892); Becker, Mitteil. Mus. Naturk. Berlin, Vol. 2, p. 47 (1902); Czizek, Zeitschr. Mähr. Landesmus. Brünn, Vol. 7, p. 165 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 13 (1910); Walhgren, Ent. Tidschrift, Vol. 31, p. 46 (1910); Carter, Ent. Mag. Vol. 48, p. 59 (1912); Frey, Acta Soc. Sc. Fenn. Vol. 37 (3), p. 7 (1913). Europe, Egypt.
- flavipes*, Meigen, Klassif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 241 (1804); Fabricius, Syst. Antl. p. 145 (1805); Fallen, Empid. Suec. p. 5 (1815); Meigen, Syst. Beschr. Vol. 2, p. 348, pl. 21, f. 20 (1820); Macquart, Mém. Soc. Sc. Lille, p. 145 (1823); Dipt. N. France, Vol. 3, p. 149 (1827); Hist. Nat. Dipt. Vol. 1, p. 319, pl. 7, f. 13 (1834); Zetterstedt, Fauna Ins. Lappon, p. 535 (1834); Curtis, Brit. Ent. Vol. 8, p. 661 (1837); Zetterstedt, Dipt. Scand. Vol. 1, p. 235 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Gimmerthal, Bull. Soc. Natur. Moscou, Vol. 20 (2), p. 163 (1847); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 485 (1849); Zetterstedt, Dipt. Scand. Vol. 13, p. 4978 (1859); Pipping, Not. Sällsk. Fenn. Förh. Vol. 4, p. 114, (1858); Bonsdorff, Finl. Tvärv. Ins. Dipt. p. 142 (1861); Siebke, Nytt Mag. Naturvid. Vol. 14, p. 380, 384, 399 (1866).
- var. *fumipennis*, Meigen, Syst. Beschr. Vol. 2, p. 349 (1820); Ahrens & Germar, Fauna, Ins. Eur. Pt. 12, p. 25 (1829); Curtis, Brit. Ent. Vol. 8, p. 661 (1837); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 486 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (17) p. 48 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 121 (1851); Pipping, Not. Sällsk. Fenn. Förh. Vol. 4, p. 114 (1858); Bonsdorff, Finl. Tvärv. Ins. Dipt. p. 142 (1861); Schiner, Fauna Austr. Dipt. Vol. 1, p. 78 (1862); Neuhaus, Dipt. March. p. 68 (1886); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 44 (1892); Czizek, Zeitschr. Mähr. Landesmus. Brünn, Vol. 7, p. 165 (1907); Wahlgren, Ent. Tidskr. Vol. 31, p. 46 (1910); Kuntze, Deutsche Ent. Zeitschr. p. 548 (1913). Europe.

(1) **Hybos ectorus**, nov. sp. — Male. Length 4 mm. Third antennal joint ovate, one and three-fourths times long as wide, arista three times antennal length; upper facets enlarged. Thorax moderately dusted with brown, leaving a pair of short vittæ and a rounded side spot shining, one pair of dorsocentrals large, apical scutellars long, pleure cinereous. Abdomen dusted with brown, leaving the hind margin of the last three segments shining; hairs sparse and yellow; pygidium not fimbriate, the upper valve deeply cleft, apically ending in two triangular processes of which the right one is smaller and narrower, lower valve with spoon-like termination, intermediate valve, on right side, large and bearing a flattened claw which fits into the cleft of the upper valve. Legs black, anterior coxae and base of hind coxae cinereous, front metatarsi slender, four-fifths as long as tibiæ, middle tibiæ with a long seta at middle of flexor face, at basal fifth and two-fifths of extensor face, hind femora rather slender, hairs black becoming longer and yellow below, a dorsal bristle before knee, six spinous bristles in lower anterior row, ventral denticles arising from minute tubercles, geniculation of hind tibiæ brown, no thorns on metatarsal sole. Knob of halteres brown, wings gray-hyaline, stigma weak, sections of fifth vein 2 : 1.

Female. Dust forming broad triangles on abdominal segments.

Eight males, eleven females, La Suiza de Turrialba, Costa Rica, Pablo Schild, collector.

- ? *rufitarsis*, Roser, Correspondenzbl. Landw. Var. Württemb. Stuttgart, Vol. 1, p. 53 (1840).
14. *H. flavipes*, Brunetti, Rec. Indian Mus. Vol. 9, p. 14 (1913); Fauna India. Brit. Ind. Dipt. Vol. 1, p. 342 (1920).
15. *H. gagatinus*, Bigot, Ann. Soc. Ent. France (6) Vol. 9, p. 127 (1889); India. Bezzi, Ann. Mus. Hungar. Vol. 2, p. 326, note (1904); Brunetti, Rec. Indian Mus. Vol. 9, p. 15 (1913); Fauna, Brit. Ind. Dipt. Vol. 1, p. 339, pl. 4, f. 6 (1920).
16. *H. geniculatus*, van der Wulp, Termész. Fuezet. Vol. 20, p. 137 (1897); Tijdschr. v. Ent. Vol. 42, p. 49 (1899); Bezzi, Ann. Mus. Hungar, Vol. 2, p. 326 (1904); Meijere, Tijdschr. v. Ent. Vol. 54, p. 322 (1911); Bezzi, Ann. Mus. Hungar, Vol. 10, p. 454 [1912] (*Noesa*); Suppl. Ent. Berlin, Vol. 3, p. 67 [1914] (*Noesa*); Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 68 (1914); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 340 (1920). East Indies, Ceylon, Java.
17. *H. grossipes*, Linnaeus, Syst. Nat. Vol. 12, p. 988 [1767] (*Musca*); Gmelin, Syst. Nat. Vol. 5, p. 2878 [1790] (*Musca*); Haliday, Stettin. Ent. Zeit. Vol. 12, p. 136 (1851); Walker, Ins. Brit. Vol. 1, p. 120, pl. 5, f. 1 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 78 (1862); Neuhaus, Dipt. March. p. 68 (1886); Lundbeck, Dipt. Dan. Vol. 3 p. 12 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 46, f. 2 (1910); Carter, Ent. Mag. Vol. 48, p. 59 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 7 (1913). Europe.
- asiliiformis*, Bonelli, in Latreille, Gen. Crust. Ins. Vol. 4, p. 305 [1809] (*Acromyia*) no description; Nouv. Dict. Hist. Nat. 2^e éd. (1818) no description; Bonelli, in Macquart, Dipt. N. Fr. Vol. 3, p. 148 [1827] (*Acromyia*) no description.
- claripennis*, Strobl, Mitth. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 43 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 18 (1910).
- clavipes*, Fabricius, Ent. Syst. Vol. 4, p. 403 [1794] (*Empis*); Syst. Antl. p. 138 [1805] (*Empis*).
- culiciformis*, Gmelin, Syst. Nat. Vol. 5, p. 2900 [1790] (*Asilus*).
- funebris*, Meigen, Klassif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 240 (1804); Fabricius, Syst. Antl. p. 145 (1805); Latreille, Gen. Crust. Ins. Vol. 4, p. 305 (1809); Fallen, Empid. Suec. Pt. 5, p. 5 (1815); Meigen, Syst. Beschr. Vol. 2, p. 347 (1820); Macquart, Mem. Soc. Sc. Lille, p. 145 (1823); Dipt. N. France, Vol. 3, p. 148, pl. 4, f. 6 (1827); Hist. Nat. Dipt. Vol. 1, p. 318 (1834); Curtis, Brit. Ent. Vol. 8, p. 661 (1837); Zetterstedt, Fauna Ins. Lappon. p. 535 (1838); Dipt. Scand. Vol. 1, p. 234 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Zetterstedt, Dipt. Scand. Vol. 8, p. 2993 (1849); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 485 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (17) p. 48 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 4267 (1852); Pipping, Not. Sällsk. Fenn. Förh. Vol. 4, p. 114 (1858); Zetterstedt, Dipt. Scand. Vol. 13, p. 4977 (1859); Bonsdorff, Finl. tvåv. Ins. Vol. 1, p. 142 (1861); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 379 (1866); Leunis, Syn. Zool. Vol. 2, p. 401 (1886).
- pilipes*, Meigen, Syst. Beschr. Vol. 2, p. 349 (1820); Curtis, Brit. Ent. Vol. 8, p. 661 (1837); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 486 (1849); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 43 (1893); Lundbeck, Dipt. Dan. Vol. 3, p. 18 (1910).
- vitripennis*, of authors
Pipping, Not. Sällsk. Fenn. Förh. Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Vol. 1, p. 142 (1861); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 379, 398 (1866).
18. *H. halteralis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 310 (1909). Bolivia.
- . *H. leachianus*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 260 (1829) no description.
19. *H. luridus*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 319 (1909). Bolivia.

20. *H. major*, Bezzi, Ann. Mus. Hungar, Vol. 10, p. 454 [1912] (*Noeza*); Formosa. Suppl. Entom. Berlin, Vol. 3, p. 66 [1914] (*Noeza*); Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 67 (1914).
- . *H. marshamanus*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 260 (1829) no description.
21. *H. mellipes*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, p. 373 Mexico. (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 247, pl. 7, f. 78 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 (1909).
22. *H. neotropicus*, Bezzzi, Ann. Mus. Hungar, Vol. 3, p. 426 (1905); Nova Bolivia, Peru. Acta Akad. Naturf. Halle, Vol. 91, p. 310 (1909).
- . *H. nervosus*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 260 (1829) no description.
23. *H. niger*, Brunetti, Rec. Indian Mus. Vol. 9, p. 12 (1913); Fauna Brit. India. Ind. Dipt. Vol. 1, p. 340 (1920).
24. *H. nigronitidus*, Brunetti, Rec. Indian Mus. Vol. 9, p. 11 (1913); Fauna India. Brit. Ind. Dipt. Vol. 1, p. 342 (1920).
25. *H. nitens*, Brunetti, Rec. Indian Mus. Vol. 9, p. 13 (1913); Fauna Brit. India. Ind. Dipt. Vol. 1, p. 341 (1920).
26. *H. pallipes*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 323 (1911); Bezzzi, Java. Suppl. Ent. Berlin, Vol. 3, p. 67 [1914] (*Noeza*); Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 68 (1914).
27. *H. papuanus*, Kertesz, Termesz. Fuezet. Vol. 22, p. 175, f. 1 (1899); van der Wulp, Tijdschr. v. Ent. Vol. 47, p. 49 (1899); Bezzzi, Ann. Mus. Hungar, Vol. 2, p. 328 (1904); ibidem, Vol. 10, p. 454 (1912) (?). New Guinea.
28. *H. plumicornis*, Bezzzi, Suppl. Ent. Berlin, Vol. 3, p. 67 [1914] (*Noeza*). Formosa.
29. *H. pollinosus*, Bezzzi, Ann. Mus. Hungar, Vol. 2, p. 326 (1904). Australia.
30. *H. reversus*, Walker, List. Dipt. Brit. Mus. Vol. 3, p. 487 (1849); Osten-Sacken, Cat. Dipt. N. Amer. p. 240 (1878); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 437 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 248 (1902). — **Pl. I, Fig. 5.** North America.
var. *Slossonæ*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 437 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 247 (1902). E. United States.
31. *H. sciapterus*, nov. sp. (1). Costa Rica.
32. *H. setosus*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 324 (1911); Bezzzi, Suppl. Ent. Berlin, Vol. 3, p. 66 [1914] (*Noeza*); Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 68 (1914). East Indies.
33. *H. spinicosta*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, Mexico. p. 374 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 246, f. 77 (1902); Bezzzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 304 (1909).
34. *H. striatellus*, Villeneuve, Feuille Jeun. Natur. Paris, Vol. 43, p. 111 (1913). France.

(1) **Hybos sciapterus**, nov. sp. — Male. Length 5 mm. Third antennal joint short oval, arista three times antennal length; upper facets slightly enlarged; occiput cinereous. Mesonotum coated with yellow dust leaving three rounded shining spaces on each side, dorsocentral and distant apical scutellars strong; pleuræ whitish pruinose. Abdomen with light marks of yellow pollen, forming dorsal spots on segments two and three, and basal bands on four following segments; pygidium with fringes of conspicuous black hair. Coxæ black, anterior femora luteous, with a dark streak above near base, front femora with six long fine setæ below, front tibiæ and tarsi jet black, the metatarsus three-quarters length of the tibia, hind femora robust, ten black spinous bristles forming anterior flexor row, hairs yellow, longer below, giving the lower posterior face a ciliate appearance, flexor denticles arising from small tubercles, hind knees broadly luteous, hind tibiæ brownish except the lighter ends, sole of hind tarsi with black denticles, middle tibiæ with very long black setæ, one below knee, one at basal third, one at middle within and one at apex. Wings large of strong brown color, veins black, sections of fourth vein 2 : 1.

Holotype, La Suiza de Turrialba, Costa Rica, Pablo Schild, collector.

35. *H. sydneyensis*, Schiner, Novara Reise, Dipt. p. 203 (1868); Bezzia, Ann. N. S. Wales. Mus. Hungar, Vol. 2, p. 327 (1904).
36. *H. tenuipes*, Brunetti, Rec. Indian Mus. Vol. 9, p. p. 13 (1913); Fauna India. Brit. Ind. Dipt. Vol. 1, p. 338, pl. 4, f. 7, 8 (1920).
37. *H.* tenuis*, Meunier, Ann. Sc. Nat. (Zool.), Vol. 7, p. 91, 109, pl. 7, Baltic Amber. f. 6, 7 (1908).
38. *H. tibialis*, Bezzia, Ann. Mus. Hungar, Vol. 10, p. 455 [1913] (*Noesa*); Formosa. Suppl. Ent. Berlin, Vol. 3, p. 67 [1914] (*Noesa*).
39. *H. typicus*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, p. 373 Mexico. (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 246 (1902); Bezzia, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 (1909).
40. *H. vagans*, Loew, Jenaische Zeitschr. Ges. Naturw. Jena, Vol. 43, p. 417 Persia. (1874).
41. *H. xanthopodus*, nov. sp. (1). Costa Rica.

4. GENUS EUHYBOS, COQUILLETT

Euhybos, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390, 437 (1896); Melander, Trans. Amer. Soc. Vol. 28, p. 248 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 250, 259, 264 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 541 (1910); Bezzia, Ann. Mus. Hungar, Vol. 10, p. 456 (1912).

Characters. — Hunchbacked, shining black, pilose species with strong, spinose hind femora, contiguous eyes, long ovipositor and reduced mouthparts. Head globular, largely composed of the eyes, the occiput broadly conical and hairy but not bristly, ocellar prominence on the summit of the head, no front, face or cheeks, due to the large eyes, rarely, however, a small narrowly triangular subantennal space representing the face, sometimes continued as a line to the oral margin, antennal incision of the eyes small, uppermost facets enlarged, a horizontal line extending through the middle of the eyes; antennæ three-jointed but the basal joints connate, the second joint with a circle of small hairs, the third joint elongate oval or subulate, two to three times as long as broad, the terminal arista very slender, thread-like, bare and averaging twice the antennal length; proboscis not half as long as the head, extending forward, when closed the two valves of the labella come together as a thin vertical lamella, palpi minute, scarcely projecting, each with a preapical hair. Thorax greatly convex, humeri prominent, prothorax quadrate when viewed from above, more or less densely and long pilose, typically no bristles, except two notopleural and a pair of apical scutellars; pleuræ bare, the mesopleuræ and sternopleuræ shining, otherwise the pleuræ lightly coated with pollen. Abdomen elongate, cylindrical, devoid of bristles but with long loose pale hairs toward base; pygidium terminal, rather large, ovoid in outline, comprising an upper and a lower opposing convex valve, each of which terminates in a complicated manner, between these a small elliptical plate on the left side and a delicate process on the right, an upcurved terminal filament, which is flattened, ribbon-like, hairy, and usually is long and attenuated

(1) *Hybos xanthopodus*, nov. sp. — Female. Length 4 mm. Third antennal joint three times as long as wide, arista one and three-quarters times antennal length; upper facets large. Mesonotum very thinly dusted, a patch of white dust near notopleural suture, dorsocentral bristle minute, scutellar pair small; pleuræ white-dusted. Abdominal hairs sparse and yellow. Coxæ and legs yellow, hind tibiae and tarsi and tips of anterior tarsi with brownish tinge, a flexor yellow seta at basal one-third of anterior tibiae, middle tibiae with two long extensor setæ on proximal part, hind femora relatively slender, three fine fulvous setæ toward knee on lower anterior side, flexor denticles black, hind edge ciliate with sixteen delicate yellow setæ, no denticles on metatarsal sole. Halteres, calypteres and root of wing yellow; wings hyaline, stigma faint, not filling marginal cell, veins black, first vein ending in middle of stigma, sections of fifth vein 2 : 1.

Two specimens, La Suiza de Turrialba, Costa Rica, April, November, 1922, Pablo Schild, collector.

or ending in a spearhead point; eighth segment of the female often widened or distorted, the ninth segment elongate, drying in a characteristic way in museum specimens usually by becoming compressed and showing chitin ridges above. Hind legs robust, the hind coxae larger than the others, the trochanters as large as the middle coxae and in the male often armed with a spine beneath, the hind femora clavate, bearing strong spinous bristles beneath, which may arise from more or less developed tubercles, more pronounced in the male, above often with two subapical bristles, middle tibiae with a series of extensor bristles, long apical bristles present on the anterior tibiae, anterior tarsi bearing long bristles both above and below, hind tibiae curved, geniculate at the knee, the inner surface of the hind tibiae edged and in the males of the *purpureus* group excavated also, pulvilli large, empodium hair-like. Halteres black in all the species; calypteres large and heavily fringed. Wings with large anal angle, often darkened in color toward the base, costa stopping at the fourth vein, no basal bristle, costal hairs long and abundant, costal cell sometimes widened; auxiliary vein straight, lying close to the first vein into which it vanishes at the middle of the wing, first vein ending considerably beyond the middle of the wing, second vein usually including a darkened stigma, third vein simple, the pedicel of the second and third veins short, arising beyond the middle of the elongate basal cells, discal cell large, elongate, located beyond the middle of the wing, complete, emitting two simple posterior veins, anal crossvein curved but continuing toward the hind margin, anal vein weak but complete.

Type species: *E. purpureus* Walker (**Pl. I, Fig. 3**), Coquillett's designation, 1903. Coquillett separated this genus from *Hybos* because of the contiguous eyes, short proboscis, lack of setulae beneath the hind tarsi and black halteres. Additional characters are to be found in the more or less densely pilose mesonotum which shows no trace of dorsocentral or serial bristles and is devoid of pollinose coating. The ovipositor is drawn out, unlike in *Hybos*, and the pygidium is complex. In all the species the first posterior cell is narrower at the apex than opposite the posterior crossvein.

Geographical distribution.

The species are confined to North and South America. They are æstival, occurring most abundantly in meadowlands, and are not common in shady mountainous regions. The males of some species exhibit peculiar characters in the hind legs, which are more robust and spinose than in the female, and in some species the femora bear stout spinigerous tubercles, which however, are unlike those of *Lactistomyia* in being confined to a short extent and not distributed along the whole under surface. The females of many of the species resemble each other so closely that they can be identified only with difficulty. Identification characters in the genus are largely drawn from the secondary sexual structures of the male. Variation is great, however, including even the bristles and tubercles of the hind legs and the conformation of the pygidium, which structures afford the most dependable indices to the species. The known species are related to each other in the manner shown by the following table.

TABLE OF THE KNOWN SPECIES OF EUHYBOS

(BASED MAINLY ON MALE CHARACTERS)

1. Pulvilli normal; second joint of anterior tarsi shorter than first; no dorsocentral bristles	2.
Anterior pulvilli greatly lengthened; first and second joints of anterior tarsi equally long; posterior dorsocentral developed	38.

2. Inner flexor edge of hind femora with a series of strong spinigerous tubercles, the corresponding surface of hind tibiæ concave and margined.	3.
Femora not armed with conspicuous spinigerous tubercles, the hind tibiæ not excavated nor prominently edged.	8.
3. The stout tubercles of inner flexor edge of hind femora arising near middle of the hind tibiæ, as seen when inflexed, and extending quite to the knee; hairs of thorax dark; basal half of wing infumated	E. PURPUREUS, Walker.
The stout tubercles not extending nearer the knee than the basal third of the hind tibiæ; hairs of thorax yellow	4.
4. Five to seven strong tubercles present; wings hyaline.	5.
But a single stout tubercle present	6.
5. Principal tubercles of hind femora elevated on a crested swelling opposite middle of tibia; thorax jet black	E. STRUMATICUS, nov. sp. (1).
Tubercles separated, opposite basal third of tibia which bears a pair of distally directed processes; thorax violaceous	E. OCREATUS, nov. sp. (2).
6. Hind tibiæ with prominent tubercle on extensor side; wings clear	E. VERRUCICRUS, nov. sp. (3).

(1) *Euhybos strumaticus*, nov. sp. — Male. Length 3 mm. Black, without metallic tinge, anterior tibiæ and hind knees brown, basal two joints of the tarsi yellow; hind femora with a row of strong spine-tipped tubercles; wings hyaline. Third joint of the antennæ bluntly oval, two and one-half times as long as broad and one-third as long as the arista. Hairs of the notum moderately abundant, long, yellow, scutellum with two bristles and a few small hairs. Upper valve of the pygidium longer than wide, continued as a long broad process which furcates and straddles the terminal valve, the terminal valve tipped with a fringe of curled yellow hairs, right side of the lower valve bearing a long process which is broader at the middle and is fringed posteriorly with long curled yellow hairs. Middle tibiæ with about five moderate extensor bristles and a long apical yellow bristle within, anterior tarsi with long yellow flexor bristle toward the base, and an apical brown extensor bristle, hind trochanters with thorn-like spine, hind femora very robust and spinose beneath, the spines growing from small tubercles which are stronger along the apical third, seven of those of the inner flexor edge grouped on a swelling and bearing very robust but short spines, this cluster corresponding to about the middle third of the tibia which is excavated and sharply margined. Halteres black. Wings hyaline or with very faint uniform brown tinge, stigma very weak but filling out the marginal cell, costal cell narrow and straight in front.

Female. The females supposedly of this species have the abdomen ending in a curious ovipositor and lack the heavy armature of the legs. The eighth segment is elongate, bearing near the middle on each side a strong backward-projecting spur, before which the segment is depressed and trapezoidal and beyond which the segment becomes more or less tubular.

Type : Plummer's Island, Maryland, R. C. Shannon, collector, in the U. S. National Museum. Paratypes from South Carolina, Georgia, Alabama, Louisiana and Texas.

(2) *Euhybos ocreatus*, nov. sp. — Male. Length 4 mm. Black, notum and tergum violaceous, tarsi yellow. Outer antennal joint oval, two times long as wide, arista two times antennal length; occiput brownish gray. Notal hairs yellowish in front, dark behind, apical scutellars strong, three lateral hairs on scutellar margin. Hairs at base of abdomen long and yellow as usual; pygidium small, upper valve ending on left side in a flat hook and on the right side in two ribbon-like processes, lower valve ending in a long ribbon-like process. Front tibiæ with a small pimple on basal third of flexor side, middle tibiæ with three outstanding setæ, hind femora strongly spinose beneath, four or five spines arising from strong close-set tubercles toward the knee opposite a flat expansion at basal two-fifths of the tibia. This expansion projects distally on each side as a prong and beyond it, near the middle of the tibia, is a distinct bend. The flexor surface of the tibia from the bend to the apex is concave, with an edge along each side. Hind metatarsi with two long fine dorsal setæ; last three tarsal joints and apex of claws black. Wings narrow, clear hyaline, stigma distinct and brown, veins thin, discal cell very long, the sections of fifth vein 4 : 1.

Female. Seventh abdominal segment transversely creased at base, then semicircularly bulged and compressed at apex, ovipositor long. Legs normal, two anterior and one superior bristle near hind knees. Stigma faint, discal cell shorter.

One specimen of each sex, La Suiza de Turrialba, April, 1922, Pablo Schild, collector.

(3) *Euhybos verrucicrus*, nov. sp. — Male. Length 5 mm. Black, the anterior legs and hind tarsi yellowish brown with last three joints of all tarsi blackish. Occiput lightly cinereous; last antennal joint conical, two times as long

Hind tibiæ without external tubercle; wings dark at base.	7.
7. Femoral tubercle located at three-fifths the length and beyond middle of hind tibiæ which are uniformly piceous	E. COQUILLETTI, nov. sp. (1).
Femoral tubercle located toward the knee, hind tibiæ paler in middle.	E. DENTIPES, Wiedemann.
8. Hind metatarsi with several pronounced spinous bristles along outer flexor edge	9.
Tarsi not provided with spinous bristles	13.
9 Anterior legs entirely black, with black bristles, hind legs black or at most their metatarsi yellow, hind metatarsi with two or three spines; wings dusky toward base	10.
Tarsi yellow, their hairs and bristles in part pale, pulvilli yellowish, hind metatarsi with four or five spines; apical part of terminal filament slender	12.
10. Terminal filament spatulate, abruptly tapering at apex, middle prong of left or lower valve projecting; spines of hind metatarsi present in female also.	11.

as wide, arista three times antennal length. Notum polished, with faint violaceous tinge, very lightly dusted above wings and in back, hairs scattered, pale in front, black behind, apical scutellars developed, no dorsocentrals; meso- and sternopleura polished, remainder gray-coated. Abdomen polished, faintly violaceous, basal hairs long and whitish, genitalia short, lightly dusted, the right valve deeply excavated below and bearing a long flat process in the excavation. Coxæ black, base of anterior femora fuscous, no excessively long setæ, a small flexor spur near base of front tibiæ, four extensor bristles but no flexor bristle along middle tibiæ, hind trochanters with spinous bristles, hind femora robust, with stout spinous bristles as follows: one preapical dorsal, four anterior, nine anterior-flexor, several ventral toward knee, five posterior-flexor on basal half, of which the third and fourth arise together from a strong tubercle, two posterior at two-thirds the length of the femur, the lower bristles mostly arising from small tubercles, hind tibiæ gently arcuate, a strong tubercle at two-thirds length of outer side, two smaller preapical tubercles on inner side, pulvilli normal. Halteres wholly black. Wings nearly hyaline, veins blackish, stigma long, blackish, filling end of marginal cell, sections of costa 1 : 0.3 : 0.4 : 0.1, of fourth vein subequal, of fifth vein 1 : 0.25, third and fourth veins convergent.

Female. Ovipositor slender; hind legs normal, bristles of hind femora reduced.

Two specimens, La Suiza de Turrialba, Costa Rica, November, 1922, Pablo Schild, collector.

(1) **Euhydos Coquillettii**, nov. sp. — Male and female. Length 4 mm. Body brownish black. Third antennal joint three times as long as broad and nearly half as long as the arista; occipital hairs blackish. Hairs of the mesonotum long, moderately abundant and yellow, scutellum with two strong and two weak bristles and about four additional marginal hairs; notum and abdomen with slight purplish reflection; seventh segment of the female abdomen rather globose, the base of the eighth trapezoidal and separated from the compressed apical part by a groove, the upper part of the apical portion bears an inverted Y-shaped thickening, the lower part becomes pale apically, the ninth segment long, slender, opaque and hairy; upper valve of the pygidium in profile longer than deep, polished on the outer half, not margined with long hairs, bifurcate apically the right fork long and slender, the left broadened obconically to cover the base of the lengthened, curved, loosely hairy, terminal filament. Tarsi and anterior tibiæ testaceous, anterior femora brownish, posterior legs otherwise brownish black, middle tibiæ with about six extensor bristles and a long yellow preapical flexor bristle, middle metatarsi with two long brown bristles on the extensor side and a pair of shorter yellow bristles on the flexor side, hind trochanters ♂ with a strong thorn beneath, hind femora robust, strongly spinose beneath, the spines growing from stubby tubercles, in the male a single strong tubercle tipped with two thorns located at three-fifths the length of the under-inner side, a strong subdorsal bristle before the hind knee, hind tibiæ ♂ conspicuously curved, the femoral tubercle coming beyond the middle. Halteres black. Wings blackish to beyond the middle of the discal cell, costal cell convex in front, a little broader than the first basal.

North Carolina. Type in the U. S. National Museum. This species was associated with specimens from North Carolina determined by Coquillett as *purpureus* (Proc. Ent. Soc. Wash. Vol. 5, p. 264, 1903). The *purpureus* form differs in having about six strong tubercles in the apical part of the hind femora ♂, the pygidium relatively shorter with the upper valve almost globular and margined with hairs, and the pubescence of the mesonotum almost black.

- Terminal filament slender, pointed, middle process of left valve broad and rounded; metatarsal spines absent in female. E. SORDIPES, nov. sp. (1).
11. Tarsi entirely black, their spongy pubescence dark gray, pulvilli dusky E. NIGRIPES, nov. sp. (2).
- Hind metatarsi yellow, their spongy pubescence concolorous, all pulvilli whitish E. METATARSALIS, nov. sp. (3).
12. Pile of mesonotum black; wings usually quite dark toward base; ovipositor cylindrical; metatarsal spines present in female. E. SUBJECTUS, Walker.
- Pile of mesonotum yellow; wings uniformly subhyaline; base of ovipositor greatly swollen and deformed; hind metatarsi of female not spinose. E. GENITIVUS, nov. sp. (4).

(1) **Euhybos sordipes**, nov. sp. — Male. Length 3.5 mm. Black, including nearly all of legs, base of wings infuscated, hind metatarsi of male trispinose. Third antennal joint elliptical, twice as long as wide, arista two and one-half times antennal length; occiput fulvo-pollinose. Notal hairs long, sparse and blackish, four lateral hairs on scutellum, dust at back of mesonotum fulvous. Abdomen bronzed, hairs whitish, pygidium subglobose, black hairy, the flat elliptical piece of the left side between the upper and lower valves large, extending fully half-way to the end of the pygidium, left valve with lobe above at apex, terminal filament very thin, sharply pointed, hairy, its base abruptly deformed and expanded, right valve sharply excised at apex. Legs mostly black, base of anterior tarsi piceous, base of hind tarsi brown, hairs mostly black, those of anterior tibiae and tarsi brown, of hind tarsi reddish, of hind coxae yellow, hind trochanters with single spine, hind femora robust and spinose, with nine spines in front below, two anterior and three dorsal, none on middle posterior face, flexor spines irregular, arising from small tubercles, geniculation of the simple, curved hind tibiae dark castaneous, pulvilli gray. Wings normal, the basal darkening merging with the hyaline apex, stigma dark but almost continuous with the basal darkening, sections of fourth vein 2 : 1.

Female. Hind femora relatively slender, less spinose, no dorsal, one anterior spine; apical segments of abdomen simple.

Four specimens, Mer Bleue, Ottawa, 11 August, 1923, F. Ide. Type and allotype in Canadian National Museum, two male paratypes retained.

(2) **Euhybos nigripes**, nov. sp. — Length 3.5 mm. Shining jet black with entirely black legs, the hind metatarsi with two or three spinous black bristles on the outer side, wings fuliginous toward the base. Eyes narrowly separated below the antennae, the face long and linear; third joint of the antennae twice as long as wide, the arista less than twice as long as the antenna. Mesonotal hairs long, rather abundant and black, scutellum with two bristles and about five pairs of hairs, pile of the abdomen long and whitish. Upper valve of the pygidium but slightly convex, loosely hairy, shining only at the truncated tip and without a subapical tooth on the right side, terminal filament obliquely truncated at the end; ovipositor more or less comical, usually showing a Y-shaped carina and polished trapezoidal basal piece, and pollinose below. Knees black like the remainder of the legs, middle tibiae with about six extensor bristles, the apical bristles long and black, bristles of the tarsi black, but the shorter hairs of the anterior tibiae and tarsi pale, hind femora moderately clavate and spinose beneath, hind metatarsi bearing a couple of long slender extensor bristles and two or three stout bristles on the outer side, one or two of the latter located near the base and one at the tip; pulvilli dusky. In the female the spines are less pronounced, the spur of the hind trochanters absent, the spinous bristles of the hind femora reduced and the spines of the hind metatarsi present only as finer bristles. Wings smoky to beyond the middle, passing gradually into the gray tip, stigma weak, filling the marginal cell, costal cell not bowed.

Numerous specimens: Vermont, Massachusetts (G. de N. Hough and C. W. Johnson), New York (C. T. Brues and O. A. Johannsen), New Jersey (C. T. Greene), Pennsylvania (E. T. Cresson, Jr.), Virginia (N. Banks), Illinois, Missouri (C. F. Adams).

(3) **Euhybos metatarsalis**, nov. sp. — A series of specimens from the Okefenokee Swamp, Georgia, collected by J. C. Bradley, and also from Tifton, Georgia, received from Dr Hough, Sumter, South Carolina (Sturtevant) and Kushla and Saraland, Alabama (Sturtevant), differ from *nigripes*, in their smaller size, measuring three millimeters or less, and in having the hind metatarsi and more or less of the soles of the following joints yellow. The pubescence of the sole is yellow and not grayish as in the black-footed form. The pulvilli are yellowish. In addition, the notal hairs are shorter and less dense, the slender bristles of the anterior tibiae and tarsi are shorter, the hind metatarsi are a little stouter and their spines stronger. The metatarsal spines are evident in the female also, as in *nigripes*.

(4) **Euhybos genitivus**, nov. sp. — Length 3.5 mm. This species is essentially distinct in its curious ovipositor; in other respects it closely resembles *subjectus*, as identified by Coquillett. Eyes contiguous except for a small subantennal triangular space; third joint of the antennae two and one-half times as long as wide, the arista slightly more than twice the

13. Costal cell much widened	14.
Costal cell scarcely or not at all widened	17.
14. Mesonotum with blue lustre	15.
Mesonotum jet black.	16.
15. Mesonotum vittate; middle tibiæ yellow; wings uniformly darkened	E. INTEGER, Walker.
Mesonotum not vittate; middle tibiæ dark; wings with hyaline apex.	E. EURYPTERUS, Bezzi.
16. Abdomen with bronze lustre; wings apically hyaline	E. DIMIDIATUS, Walker.
Abdomen black; wings dark at apex.	E. LATIPENNIS, Bezzi.
17. Base of wings distinctly darker than apical portion	18.
Wings entirely hyaline or subhyaline to the base, the stigma usually present	32.
18. Stigma distinctly limited.	19.
Stigma obsolete, sometimes the entire marginal cell carries forward the basal darkening of the wings	23.
19. Mesonotum purple, blue or green.	20.
Body jet black.	22.
20. Front tibiæ black	E. PILOSUS, Schiner.
Front tibiæ yellowish.	21.
21. The fine villous coating at sides of mesonotum gray.	E. LOEWI, Wheeler & Melander.
The fine villous coating brown	E. ELECTUS, Melander.
22. Legs mostly black.	E. TABASCENSIS, Wheeler & Melander.
Legs brownish yellow	E. CRASSIPES, Fabricius.
23. Brown of base of wings invading discal cell.	24.
Discal cell wholly hyaline	27.
24. Geniculation of hind tibiæ reddish; pygidium globose	25.
Hind knees black; pygidium longer than wide, with long apical process from ventral valve; base of ovipositor distorted, bullate (<i>E. triplex</i> , Walker).	26.
25. End processes of pygidium short; ovipositor simple.	E. DUPLEX, Walker.
End processes as long as body of pygidium, that of upper	

antennal length. Hairs of the notum sparse and yellow, scutellum with two bristles and four minute hairs, hairs of the abdomen sparse. Abdomen with a slight purple tinge, the eighth segment of the female large, jet-black, its base twice as wide and thick as the preceding segment, underneath with a strong broad tubercular basal swelling, beyond the basal third the segment is suddenly constricted, the small tip being pollinose beneath; pygidium small, upper valve shining, its marginal hairs rather long, its apical process notched, terminal filament slender and attenuated to a point. Legs piceous, the anterior tibiae and tarsi and the hind tarsi brownish yellow, the long bristles of the anterior legs yellow, hind femora moderately thickened, spinose below, hind metatarsi of the male with four brown spinous bristles along the outer side, entirely lacking in the female, pulvilli pale yellow. Wings uniformly with a faint brownish tinge, the stigma slightly darker, costal cell straight.

White Mountains, New Hampshire, H. K. Morrison, collector. Type in the U. S. National Museum. In *subjectus*, as determined by Coquillett, the hairs of the thorax are black, the base of the wings more or less brownish, especially noticeable in the male, the ovipositor simple and slender and much less produced, and the hind metatarsi of the female bear small spinous bristles on the outer side. Some specimens of *subjectus* have these spines black and others yellow in either sex. Where they are yellow the base of the wings is less dark. I have specimens of *subjectus* from New Hampshire, Massachusetts, New York, New Jersey, District of Columbia and Georgia.

- valve like a large hooked beak; base of ovipositor strongly constricted in the middle. E. *GRYPHUS*, nov. sp. (1).
26. Apical process of ventral valve single-pointed, the prong at middle of left side of ventral valve directed obliquely backward and broadly rounded at apex. E. *TRIPLEX*, nov. var. *SIMPLEX* (2).
- Apical process more or less deeply furcate, the middle prong on left side with acute tip. E. *TRIPLEX*, Walker, s. str.
27. Body with cyaneous tinge E. *PILOSIFORMIS*, Bezzii. 28.
- Body black, without metallic lustre
28. Mesonotal pile grayish in front; in specimens identified as this species the right valve of pygidium is tipped with a pencil of black setæ E. *SMITHI*, Wheeler & Melander.
- Mesonotal pile brownish to black. 29.
29. Hind femora and tibiæ brown; in specimens identified as this species the hairs of the hind femora are dense and coarse. E. *BAKERI*, Wheeler & Melander.
- Hind tibiæ and femora black (*yucatanus* included here but in absence of type its location can not be more definitely given). 30.
30. Veins thin, first basal cell two-thirds as long as second; pygidial hairs short and sparse, middle prong of left or lower valve shifted apical to terminal one and ribbon-like, end-process of right valve very large and leaf-like. E. *LEPTONEURA*, nov. sp. (3).

(1) ***Euhybos gryphus*, nov. sp.** — Male. Length 4.5 mm. Black, tarsi yellowish, wings infumated, especially at base, no stigma. Third antennal joint conical, a little more than twice as long as wide, arista two and one-half times antennal length; occiput brownish cinereous. Notal hairs black, two pairs of scutellars, pollen above wings brown. Hairs of first segment of abdomen whitish, of remainder black or piceous; right valve of pygidium dusted with brownish, its apical half formed as a strong wide beak, its basal half quadrate when viewed from above, the inner distal angle bearing a strong spur directed to the left; left valve stout, its middle process located at the tip and rather small and acute, its articulate terminal process swollen near the base and thence curved and talon-like. Knees brownish, hind femora robust, slightly shorter than tibia and metatarsus together; femoral hairs black, eight stout, spinous bristles arising from low tubercles along lower posterior face, two along middle of this face, one extensor toward knee, two along middle of anterior face and two full irregular rows at lowermost anterior face, spine on hind trochanters strong. The dark color of base of wings extends as far as apex of anal cell, not sharply distinguished from the scarcely lighter apical portion, fourth vein deeply bent at posterior crossvein, sections of fifth vein 5 : 1.

Holotype, Tabernilla, Canal Zone, Panama, 20 June, 1907, August Busck, collector, in U. S. National Museum.

(2) ***Euhybos triplex*, var. *simplex*, nov. var.** — Male. Differs from the familiar and widely distributed *triplex* Walker in the formation of the pygidium, whose left valve ends in a long, curved, pointed, simple, articulate process at the base of which the valve projects obliquely backward as a strong pointed prong. In *triplex* the terminal process is forked, the two parts either equal or the one to the left smaller. The variety *simplex* represents that stage where the left furcation has disappeared. Normally in *triplex* the robust prong at the end of the left valve is more transversely directed.

Type: Greenfield, Massachusetts, June 1, 1914. Additional specimens from Lyndon, Vermont, Long Island, New York, Chester Co., Pennsylvania, Central Illinois, and Austin, Texas.

(3) ***Euhybos leptoneura*, nov. sp.** — Male. Length 4 mm. Black, tibiæ castaneous, base of tarsi yellowish, wings hyaline, brownish at base. Third antennal joint subulate, twice as long as wide, arista two and one-half times antennal length; occipital pollen fulvous. Notal hairs yellow, one pair scutellars, pollen at edge of notum brown. Abdominal hairs golden, base of pygidium globular and lightly dusted with brown, its hairs short and sparse, the right valve deeply emarginate at end above and continued as a large forked leaf-like process, the left valve ending above in an articulate conical process with deflexed tip. Sides of front tibiæ ciliate, middle tibiæ with six extensor bristles, hind femora rather slender, hairs brown, underside with five very small thorn-bearing tubercles close to knee and with a small transverse ridge opposite apical fifth of tibiæ, anterior face with two bristles near knee, one extensor, two on posterior face near apical third,

Veins not delicate, first and second basal cells subequal; pygidial hairs long, middle prong of left valve not ribbon-like.	31.
31. Excavation at upper inner middle of right valve narrow and tipped on its proximal corner with a spine-like process, inside conical prong distal to excavation.	E. SCHILDI, nov. sp. (1).
Excavation of right valve broad and not spined, inside conical prong arising from center of excavation.	E. DINOPUS, nov. sp. (2).
32. Legs wholly black; wings pure hyaline.	E. HYALOPTERUS, Bezzii.
Legs more or less yellow.	33.
33. Crossveins clouded with brown.	E. STIGMATICUS, Schiner.
Crossveins not clouded.	34.
34. Hind femora strongly and abundantly spinose.	E. THRIXOTHRIX, nov. sp. (3).
Hind femora less spinose, no spines on upper posterior face.	35.

hind tibiæ flattened opposite femoral ridge, hind trochanters without spine. Wings narrow, no stigma, veins very thin, first basal cell about two-thirds as long as second, section of fifth vein 4 : 1. Female. Ovipositor simple.

Two specimens, Frontera, Tabasco, Mexico, March 29, C. H. T. Townsend, collector, holotype in U. S. National Museum. The structure of the hind legs indicates relationship with the *purpureus* group, but as the ridge of the tibiæ and the depression of the femora are not prominent the species has been placed in the latter part of the key.

(1) ***Euhybos Schildi*, nov. sp.** — Male. Length 3.5 mm. Black, the geniculation of the hind tibiæ and the basal two joints of all tarsi brownish yellow, basal third of wings infumated, stigma obsolete. Third antennal joint ovate, a little less than twice as long as wide, arista two and one-half times antennal length; occipital coating brownish gray. Notal hairs black and abundant, the sparse dust at sides and rear brownish, apical scutellars approximate, six setiform hairs on each side of scutellar margin. Abdominal hairs black, basal tergite pollinose, pygidium short, hairy apex contorted, left valve with strongly concave inner margin and terminated by a long slender finger-like process, right valve narrowly and deeply excised at the middle, the proximal corner of the excision tipped with a spine, the distal corner covering a stout tooth, terminal process of right valve undeveloped but a finger-like process extending from apex below. Six extensor setæ on middle tibiæ, hind femora stout and strongly spinose, hairs black, seven irregularly placed spinous bristles on posterior face, no extensors or anterior bristles, eight stout spinous bristles in posterior flexor row, other flexors irregular, hind tibiæ more strongly curved than usual, hind trochanters with a spine. Fourth vein deeply bent at posterior crossvein, sections of fifth vein 3.5 : 1.

Female. Hind femora slender, without the strong bristles on posterior face; ovipositor with deformed base, the seventh abdominal segment large, proximally fringed with long thin black setæ and transversely depressed across the middle and again before apex, the intervening portion bullate.

Two males and five females, La Suiza de Turrialba, Costa Rica, Pablo Schild, collector. The paratype male has the hind tibiæ less curved, as in the females.

(2) ***Euhybos dinopus*, nov. sp.** — Male. Length 3.5 mm. Black, abdomen with slight bronzed lustre, tarsi yellow, basal third of wings deep brown, remainder hyaline. Third antennal joint subulate, slightly over twice as long as wide, arista twice the antennal length; occiput brownish-gray pollinose. Notal hairs blackish, four scutellars. Abdomen short, hairs brown, pygidium longer than wide, greatly contorted, left valve terminated by the broad middle process, which is tipped with a spine, the apical projection articulate and long, right valve excavated along the inside and bearing a stout conical projection at the middle of the excavation, apically broadly expanding and notched for the reception of the termination of the opposing valve, below this notch continued as a spoon-shaped projection. Seriate bristles on outside of middle tibiæ strong, hind femora robust, as long as tibia and metatarsus together, hairs black, seven strong bristles along lower part and four along middle of anterior face, three extensor bristles; pulvilli yellowish, claws reddish except black tip. Brown of wings ceasing at base of discal cell, no stigma, basal cells nearly coextensive, sections of fourth vein 4 : 1, fourth vein much bent at posterior crossvein.

Holotype, San Rafael, Vera Cruz, Mexico, C. H. T. Townsend, collector, in U. S. National Museum.

(3) ***Euhybos thrixothrix*, nov. sp.** — Male. Length 4 mm. Occipital dust brown, hairs black, outer antennal joint ovate, twice as long as wide, arista two and one-half times antennal length. Thorax black, notum with slight blue lustre, pile blackish, the finely villous patches above wings brown. Abdomen black with slight bronze tinge, basal hairs long and whitish, pygidial hairs dark, terminal prong strong, obliquely truncate and crenulate at tip. Coxæ, femora, hind tibiæ and apex of tarsi black, anterior tibiæ and all tarsi brown, middle tibiæ with an extensor row of six setæ, terminal and metatarsal setæ long and brown; hind femora robust, a little longer than the simple curved tibia and metatarsus combined, unusually

35. Last posterior cell four times as long as wide; hind femora twice the width of the anterior ones E. ARDOPEODES, nov. sp. (1).
- Last posterior cell three times as long as wide. 36.
36. Upper valve of pygidium ending in a long strong curved spine, lower valve ending in a flat broadly U-shaped process E. SPINIGER, nov. sp. (2).
- Upper valve ending in a flattened piece, lower valve in two large disconnected leaf-like processes. 37.
37. End-piece of upper valve very short, left basal plate undeveloped; hind femora three times width of anterior ones E. CUSPIDATUS, nov. sp. (3)
- End-piece of upper valve very long, left basal plate large and elliptical; hind femora only moderately robust . . . E. BAROPEODES, nov. sp. (4).

strongly spinose, ten spines in outer flexor row, one on outer face at four-fifths the length, two tandem above the last-mentioned, five along inner face, seven in inner flexor row and along the flexor face several irregularly located, the flexor spines arising from small tubercles; hind trochanters spinigerous. Halteres black; wings uniformly but slightly infumated. Stigma brown, sections of fifth vein 3 : 1

Female. Bristles of hind legs reduced, none on trochanters, about five to seven in outer flexor row, one on outer face and one above and three in inner flexor row, all toward knee.

Three males and three females, La Suiza de Turrialba, Costa Rica, Pablo Schild, collector.

(1) *Euhybos ardepeedes*, nov. sp. — Closely related to *E. thrixothrix*, differing in the following characters; last antennal joint slightly more than twice as long as wide; processes of genitalia smaller, the terminal prong slender, curved, pointed and talon-like; hind femora a little shorter than the tibia and trochanter together and with reduced chaetotaxy, two bristles on upper anterior face toward knee, none on posterior face, about ten irregularly placed in each of inferior rows with irregular spines between. In the female the bristles of the hind femora are still further reduced, with seven in the lower anterior row and only apical three in the lower posterior row.

Three males and three females, La Suiza de Turrialba, Costa Rica, Pablo Schild, collector.

(2) *Euhybos spiniger*, nov. sp. — Male. Length 3 mm. Black, anterior tibiae brownish, base of tarsi yellowish, wings hyaline, no stigma, third joint of antennae elongate oval, slightly over twice as long as wide, arista twice antennal length; occiput lightly cinereous. Notal hairs pale, sparse pollen gray, two scutellars. Abdominal hairs all pale, upper valve of pygidium ending in a long curved black spine, which is in reality double but connate, lower valve ending in a large flat process which is deeply excised to admit the opposing spine, the left portion narrow and ribbon-like, the right portion broader and with an acute expansion along left side. Four moderate extensor bristles on middle tibiae, no spine on trochanters, hairs of hind femora pale, one anterior and two extensor bristles near knee, no posterior bristles, flexor spines short and thick arising from roughenings of the surface, seven in anterior row, the others irregular, hind knees paler. Section of fifth vein 2.5 : 1.

Female. Ovipositor simple, hind femora with smaller bristles.

Four specimens, Liguanea Plain, Jamaica, November-December, 1911, C. T. Brues, and one from Utuado, Porto Rico, January 1899, August Busck, the last in the U. S. National Museum.

(3) *Euhybos cuspidatus*, nov. sp. — Male. Length 3.5 mm. Body, femora and tibiae piceous black, knees and tarsi yellowish; wings uniformly smoky hyaline, the elliptical stigma brown. Arista elongate oval, two and one-half times antennal length; occiput dusky with yellowish gray. Notal hairs brownish, paler in front and darker in back, four scutellars, pollen above wings yellowish. Abdominal hairs pale, pygidial hairs long, the middle and apical processes of left or lower valve very large, leaf-like, pointed and together terminating the pygidium one from each side. Hind femora robust, in length equal to tibia and metatarsus together, its hairs yellow, no spinous bristles on front or back faces, one above toward knee, about ten in flexor rows. Submarginal cell two-thirds width of stigma, sections of fifth vein 1 : 0.6.

Holotype, Horse Neck Beach, near New Bedford, Massachusetts, August 4, 1896, G. de N. Hough, collector. A female which I collected at Cold Spring Harbor, New York, is not associated with any other species and may possibly belong here. It has hyaline wings, no stigma, black body and naturally more slender hind femora.

(4) *Euhybos baropeedes*, nov. sp. — Male. Length 4 mm. Piceous black, anterior tibiae brown, tarsi testaceous, wings uniformly lightly smoky. Third antennal joint broadly lanceolate, scarcely twice as long as wide, arista two and one-half times length of antenna; occiput lightly cinereous. Notal hairs brown, pollen gray, two scutellars. Abdominal hairs whitish at base, brown on the large globose greatly distorted pygidium, left or lower valve bowl-like, with a pointed process at right end and terminated by two widely separated strong curved flattened convergent finger-like processes, the right valve much smaller and ending in a large L-shaped process the extension of which is directed to the left between the two left processes of the left valve. Three small extensor bristles on middle tibiae, no spine on trochanters;

38. Abdomen not surpassing the wings in length E. DERODACTYLUS, nov. sp. (1).
 Abdomen greatly surpassing the wings E. LEPTOGASTER, nov. sp. (2).

Geographical distribution.

1. <i>E. ardopeodes</i> , nov. sp.	Costa Rica.
2. <i>E. Bakeri</i> , Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, p. 372 [1901] (<i>Hybos triplex</i> , var. <i>Bakeri</i>).	Mexico.
3. <i>E. baropeodes</i> , nov. sp.	New Hampshire.
4. <i>E. Coquilletti</i> , nov. sp.	North Carolina.
5. <i>E. crassipes</i> , Fabricius, Syst. Antl. p. 146 [1805] (<i>Hybos</i>); Wiedemann, Aussereurop. Zweifl. Ins. Pt. 1, p. 539 [1828] (<i>Hybos</i>); Bezzi, Nova Acta Kais. Naturf. Halle, Vol. 91, p. 305 [1909] (<i>Hybos</i>).	South America.

hind femora moderate, no bristles on upper posterior face, one extensor near knee, two on anterior face, ventral bristles normal; hind tibiæ simple. Stigma very faintly indicated, sections of fifth vein 3 : 1, fourth vein not much bent at posterior crossvein.

Holotype, White Mountains, New Hampshire, H. K. Morison, collector, in U. S. National Museum.

(1) ***Euhybos derodactylus*, nov. sp.** — Female. Length 5.5 mm. Occiput black, finely dusted with gray; outer antennal joint two times long as broad, arista three times antennal length. Mesonotum polished, with purple lustre, posterior declivity very finely dusted, setulæ seriate, posterior dorsocentral developed, apical scutellars approximate, about six minute lateral hairs on each side; pleuræ black. Abdomen long and slender, equalling the long wings, polished, the tergites with purple lustre, the eighth segment chitinized, short, tubular, the ninth or ovipositor lengthened, bulbous in the middle, hairs long and white at base, elsewhere very sparse and short. Anterior coxæ brown, hind ones black, anterior legs yellow, last three joints of all tarsi black, middle tibiae bullate at basal third and bearing a small spur within, posteriorly ciliate, two basal joints of anterior tarsi extremely long and slender, together equal to their tibiæ, hind femora strongly clavate, with yellow base and abruptly black club, eight strong bristles in outer flexor row, some short stout bristles underneath near knee, inner flexor bristles more slender, a single extensor preapical bristle, hind tibiæ geniculate, arcuate, simple, yellow on basal three-fifths and black apically; no flexor tibial bristles, three long slender extensor bristles on middle tibiæ, extremely long setæ on anterior metatarsi, pulvilli of anterior legs more than twice as long as hind ones. Halteres with long thin yellow stalk and black knob. Wings hyaline, veins thin and blackish, a strong blackish stigma filling end of marginal cell, costal sections 1 : 0.15 : 0.18 : 0.05, first posterior cell decidedly tapering, sections of fourth vein subequal, of fifth vein 1 : 0.5.

Holotype, La Suiza de Turrialba, Costa Rica, November, 1922, Pablo Schild, collector.

(2) ***Euhybos leptogaster*, nov. sp.** — Male. Length 7 mm. An extremely slender species proportioned as in the asilid genus *Leptogaster*. Black; eyes narrowly separated along the lengthened face; last antennal joint top-shaped, the filiform arista nearly three times antennal length; proboscis very short, broad, palpi narrow; occiput cinereous-dusted, its sparse hairs long, fine and black. Thorax dusted, with brown on mesonotum and with gray on pleuræ, a small denuded spot in dorsocentral rows before base of wings, posthumeral area more densely coated, hairs sparse, posterior dorsocentral, two lateral and the apical scutellar pair distinct. Abdomen polished, remarkably thin, second to fourth segments tubular, each about five times as long as wide, segments six to eight shorter and dusted, genitalia consisting of two large, long, convex, distorted, black-hairy valves. Coxæ black, anterior legs yellowish, last two joints of all tarsi black, hind femora black except narrow knees, hind tibiæ and tarsi luteous; first two joints of anterior tarsi very thin and long, together equalling their tibiæ, anterior pulvilli excessively long, twice the length of their claws, hind pulvilli one-half the length of their claws; hind femora with about ten strong spines in each of the flexor rows each arising from a low tubercle, a small spine above towards the knee, hind tibiæ simple; excessively long setæ present as follows: a pair near end of front tibiæ, a pair at apex of front metatarsi, a flexor at basal third of middle tibiae, two short extensor below middle knees, one at tip of middle tibiæ, one at base of middle metatarsi, one extensor near middle and a pair at apex of same joint. Halteres with thin pale stalk and black knob. Wings hyaline, veins thin and brownish, no stigma, first vein short, ending before posterior crossvein, costal sections 1 : 0.6 : 0.3 : 0.1, third and fourth veins almost parallel, slightly converging, sections of fourth vein 1 : 1.2, of fifth vein 1 : 0.6, pedicel of second and third veins very short.

Female. Alternate bristles of anterior flexor row of hind femora reduced, there being five strong ones left, posterior reduced to three small bristles toward knee. Eighth abdominal segment corneous, bulbous on each side of base and continuing as a long flattened ovipositor.

Two specimens, La Suiza de Turrialba, Costa Rica, Pablo Schild. This curious species departs from the habitus of its congeners, but not sufficiently to be removed to a separate genus. The lengthened abdomen, elongate anterior feet with their peculiar pulvilli, the pollinose thorax with developed dorsocentral bristles constitute a combination of characters disagreeing from other species of *Euhybos*. The nearly obliterated face, short proboscis and chitinized ovipositor would place the species in *Euhybos* rather than in *Hybos*.

6. *E. cuspidatus*, nov. sp. Massachusetts.
7. *E. dentipes*, Wiedemann, Aussereurop. Zweifl. Ins. Pt. 1, p. 538 [1828] (*Hybos*); Macquart, Dipt. Exot. Vol. 1, pt. 2, p. 156 [1838] (*Hybos*). America (locality unknown).
8. *E. derodactylus*, nov. sp. Costa Rica.
9. *E. dimidiatus*, Walker (not Loew), Ins. Saunders. Dipt. p. 205 [1852] (*Hybos*); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 [1909] (*Hybos*). Brazil.
10. *E. dinopus*, nov. sp. Mexico.
11. *E. duplex*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 486 [1849] (*Hybos*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903). E. United States.
- electus, Melander, in part, Trans. Amer. Ent. Soc. Vol. 28, p. 247, var. in part (1902).
12. *E. electus*, Melander, ibidem, Vol. 28, p. 247 [1902] (*Hybos*); Bezzi, Ann. Mus. Hungar., Vol. 3, p. 459 [1905] (*Hybos*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 [1909] (*Hybos*). West Indies.
- dimidiatus, Loew (not Walker), Wien. Ent. Monatschr. Vol. 5, p. 36 [1861] (*Hybos*); Williston, Trans. Ent. Soc. Lond. 1896, p. 307, pl. II, f. 83 [1896] (*Hybos*).
13. *E. eurypterus*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 306 Peru.
- [1909] (*Hybos*).
14. *E. genitivus*, nov. sp. New Hampshire.
15. *E. gryphus*, nov. sp. Panama.
16. *E. hyalopterus*, Bezzi, Ann. Mus. Hungar., Vol. 3, p. 425 [1905] (*Hybos*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 [1909] (*Hybos*). Peru.
17. *E. integer*, Walker, Ins. Saunders. Dipt. Vol. 1, p. 205 [1852] (*Hybos*); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 [1909] (*Hybos*). Brazil.
18. *E. latipennis*, Bezzi, ibidem, Vol. 91, p. 307 [1909] (*Hybos*). Bolivia, Peru.
19. *E. leptogaster*, nov. sp. Costa Rica.
20. *E. leptonera*, nov. sp. Mexico
21. *E. Loewi*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, p. 372 Mexico, Costa Rica.
- [1901] (*Hybos triplex*, var. *Loewi*).
22. *E. metatarsalis*, nov. sp. S. United States.
23. *E. nigripes*, nov. sp. E. United States.
24. *E. ocreatus*, nov. sp. Costa Rica.
25. *E. pilosiformis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 308 Peru.
- [1909] (*Hybos triplex*, var. *Loewi*).
26. *E. pilosus*, Schiner, Novara Reise Dipt. p. 202 [1868] (*Hybos*); Bezzi, Ann. Mus. Hungar., Vol. 3, p. 425 [1905] (*Hybos*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 309 [1909] (*Hybos*). Colombia, Peru.
27. *E. purpureus*, Walker, List. Dipt. Brit. Mus. Vol. 3, p. 486 [1849] (*Hybos*); Coquillett, Proc. U. S. Mus. Vol. 18, p. 437 (1896); Proc. Ent. Nat. Soc. Wash. Vol. 5, p. 264 (1903). — Pl. I. Fig. 3. E. United States.
28. *E. Schildi*, nov. sp. Costa Rica.
29. *E. Smithi*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, p. 372 Mexico.
- [1901] (*Hybos triplex*, var. *Smithi*).
30. *E. sordipes*, nov. sp. Canada.
31. *E. spiniger*, nov. sp. West Indies.
32. *E. stigmaticus*, Schiner, Novara Reise Dipt. p. 202 [1868] (*Hybos*); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 [1909] (*Hybos*). Colombia.
33. *E. strumaticus*, nov. sp. E. United States.
34. *E. subjectus*, Walker, List. Dipt. Brit. Mus. Vol. 3, p. 486 [1849] (*Hybos*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 437 (1896); Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903). North America.

35. *E. tabascensis*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, Mexico.
p. 372 [1901] (*Hybos triplex*, var. *tabascensis*).
36. *E. thrixothrix*, nov. sp. Costa Rica.
37. *E. triplex*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 486 [1849] (*Hybos*);
Osten-Sacken, Cat. Dipt. N.-Amer. p. 99 and 240 [1878] (*Hybos*);
Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 437 (1896); ibidem,
Vol. 22, p. 251 (1900); Wheeler & Melander, Biol. Centr.-Amer.
Dipt. Vol. 1, p. 373 [1901] (*Hybos*); Melander, Trans. Amer. Ent.
Soc. Vol. 28, p. 248, f. 79, 81 [1902] (*Hybos*); Coquillett, Proc. Ent.
Soc. Wash. Vol. 5, p. 264 (1903); Bezzi, Nova Acta Akad. Naturf.
Halle, Vol. 91, p. 309 [1909] (*Hybos*).
38. *E. verrucicrus*, nov. sp. Costa Rica.
39. *E. yucatanus*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, Mexico.
p. 372 [1901] (*Hybos triplex*, var. *yucatanus*).

5. GENUS LACTISTOMYIA, MELANDER

Lactistomyia, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 250 (1902); Coquillett, Proc. Ent. Soc.
Wash. Vol. 5, p. 251 (1903); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301, 311 (1909);
Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 557 (1910).

Characters. — Robust, thorax and abdomen typically densely covered with fine reddish brown tomentum. Head more or less hemispherical, the eyes (♂) narrowly separated below the antennæ, the upper facets larger than the lower; antennæ short, the outer joint elongate oval, with a long slender terminal arista; proboscis slender, horizontally porrect, a little shorter than the head; palpi slender, porrect also; occipital bristles minute. Thorax globose, very prominent, no bristles but with short scattered fine reddish pubescence, longer at the sides and posteriorly and almost bristle-like at the margin of the scutellum. Abdomen stout, cylindrical, deflexed posteriorly; pygidium large, flexed to the right, consisting of large convex apically deformed dorsal and ventral valves and a short thin compressed hastate terminal filament; ovipositor elongate and narrow. Legs short, the hind femora and tibiæ remarkably enlarged, the hind femora strongly tuberculate and spiny beneath, their tibiæ bare, geniculate at the patella, curved, sharp on the inner edge and with flat sides, hind metatarsi longer than the remainder of the tarsus. Wings rather narrowed apically, veins strong, no evident stigma, third and fourth veins converging, the first posterior cell much narrower in the margin than at the posterior crossvein, discal cell long and narrow.

Genotype: *L. insolita* Melander (Pl. 5, Fig. 44), the original species. The genus is restricted to tropical America, with one new species from the Philippines. Coquillett, who had not seen specimens, concluded that *Lactistomyia* was synonymous with *Hybos*, but the genus is more related to *Euhybos*, as shown by the femoral armature and narrowed apical cell. The relation of *Hybos*, *Euhybos* and *Lactistomyia* to each other is closely paralleled by the groups of *Syneches* — i. e. *Syneches*, *Epiceia* and *Harpamerus*, and shows the trend of specialization in the Hybotinæ. There is a tendency toward the closure of the first posterior cell, similar to that which is carried to a greater extent in the Syrphidæ or Tachinidæ, and a corresponding tendency to an elaborate development of the hind legs.

Geographical distribution.

1. *L. dimidiata*, Bellardi (not Walker or Loew), Ditt. Messic. Pt. 2, p. 97 Mexico.
[1861] (*Hybos*); Mem. Accad. Sc. Torino (2), Vol. 21, p. 197 [1865]
(*Hybos*).
sequens, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 249 [1902] (*Hybos*);
Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 [1909] (*Hybos*).

2. *L. hyalina*, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 312 (1909). Peru.
 3. *L. insolita*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 251, pl. 7, f. 82 Brazil.
 (1902); Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 311 (1909).
 4. *L. picea*, Wiedemann, Aussereurop. Zweifl. Ins. Pt. 2, p. 647 [1830] (*Hybos*); Brazil.
 Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 305 [1909] (*Hybos*).
 5. *L. polita*, nov. sp. (1). Luzon.
 6. *L. serrata*, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 312 (1909). Peru.

6. GENUS CERATHYBOS, BEZZI

Cerathybos, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301, 303 (1909).

Characters. — Polished metallic black species of 5.5 millimeters length closely related to *Hybos* but the third joint of the antennæ is very much dilated, semicircular, the convex side above. Head very small and with an evident neck, ocellar tubercle small; eyes contiguous on the face (♀) as well as on the front, the facets large and larger above; arista slender and long; proboscis perpendicular in the only known specimen, as long as the head, palpi small. Thorax rather convex, with short pubescence and with lateral bristles only, scutellum with four marginal bristles; ovipositor present. Anterior legs with long bristles, hind femora incrassate, spinose but not tuberculate beneath; front tibiæ with a strong tooth inside near the knee, hind tibiæ shorter than their femora, somewhat bent. Wings short and rather broad, no evident stigma, costa with abundant short hairs, pedicel of the second and third veins arising near the middle of the wing, fourth vein convergent with the third. Halteres black.

Type species: *C. Schnusei*, Bezz, from South America.

Geographical distribution.

1. *C. Schnusei*, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 303, f. 1 Peru.
 (1909).

7. GENUS SYNDYAS, LOEW

Syndyas, Loew, Oefv. Vet. Akad. Förh. Vol. 14, p. 360 (1857); Dipterenf. Südafr. p. 260, 332 (1860); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 115 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Williston, Man. N. Amer. Dipt. p. 74 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 254 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257, 259 (1903); Bezz, Ann.

(1) **Lactistomyia polita**, nov sp — Female. Length 4 mm Head globose, eyes contiguous above and below the antennæ, divided exactly into four quadrants by the horizontal line; antennæ black, the third joint elongate oval, nearly two times long as wide, arista filiform, terminal, three times antennal length; mouthparts black, projecting straight forward beyond base of antennæ, palpi linear, appressed; occiput lightly coated with fulvous pollen. Body black, polished, the rear of the thorax and the pleuræ brown-pollinose, notal hairs scattered and black, no dorsocentral bristles. Abdominal hairs straggling, long and pale, the integument with bronze lustre, ovipositor long and flattened, styles elongate, the last two segments of the abdomen with the styles as long as segments two and three together. Legs black, the pulvilli and base of claws brownish, all joints more or less furnished with black hair, middle tibiæ and anterior tarsi with a few outstanding thin setæ, hind femora at middle as thick as their coxae, their under face bearing 15 small tubercles each provided with a stout spine, middle tibiæ stout, two-thirds as long and nearly as wide as their femora, hind metatarsi nearly as long as four following joints, bearing three stout spines on exterior face, of which two are near the base and one at the apex. Halteres black. Wings lightly infumated, stigma weakly indicated, filling outer half of marginal cell, veins dark, anal vein faint, first posterior cell coarctate.

Holotype, Mt. Makling, Luzon, P. I., received from Prof. C. F. Baker. The species departs from the typical forms in the reduction of pollinosity and in the less deformed hind legs.

Mus. Hungar, Vol. 2, p. 321 (1904); Melander, Williston's 3d. Man. N. Amer. Dipt. p. 224 (1908); Kertész, Cat. Dipt. Vol. 6, p. 9 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 610 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 42 (1910); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 333 (1920).

Characters. — Jet black species with black legs, hemispherical head, greatly arched thorax, clavate hind tibiæ and weak vein between the basal cells. Head relatively large, consisting almost wholly of the eyes, which in both sexes are subcontiguous below the antennæ and have the facets of the upper half enlarged; outer antennal joint oval, with a long thread-like bare terminal arista; proboscis projecting horizontally but shorter than the head, palpi linear; no cephalic bristles. Thorax sparsely pilose, only a pair of notopleurals and a pair of scutellars developed bristle-like. Abdomen cylindrical, curved downward, pygidium terminal, small and closed; last two segments of the female abdomen tapering and drawn out to form a sort of ovipositor. Legs slender but the hind tibiæ compressed clavate and the hind metatarsi swollen, middle tibiæ with slender bristles, hind femora spinulose beneath, pulvilli rather small. Wings transparent, no stigma, costa with very short hairs and extending to the fourth vein, pedicel of the second and third veins very short, turned up at its origin, third vein simple, first posterior cell narrowed at the apex of the wing, discal cell smaller than usual, vein between the first and second basal cells very weak or wanting, anal cell acutely pointed, anal vein weak, no alula.

Type species: *S. opaca*, Loew's first species, which was Coquillett's designation in 1903. Of the commonest species, *polita*, the males differ from the females in having the tergites largely opaque, but of *dorsalis* the tergites of the female are opaque also. The Oriental species *eumera* possesses greatly enlarged and spinose hind femora, and bears a similar relationship to the other species that *Harpamerus* does to *Syneches*.

Geographical distribution.

- | | |
|--|----------------|
| 1. <i>S. aterrima</i> , Meijere, Bijdr. Dierk. Vol. 19, p. 53 (1913). | Ceram. |
| 2. <i>S. brevior</i> , Meijere, Tijdschr. v. Ent. Vol. 53, p. 68, pl. 4, f. 5 (1910). | East Indies. |
| 3. <i>S. dorsalis</i> , Loew, Berl. Ent. Zeitschr. Vol. 5, p. 320: Cent. 1, n° 26 (1861);
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 254 (1902). | United States. |
| 4. <i>S. elongata</i> , Meijere, Tijdschr. v. Ent. Vol. 53, p. 67, pl. 4, f. 4 (1910). | East Indies. |
| 5. <i>S. eumera</i> , Bezzi, Ann. Mus. Hungar, Vol. 2, p. 323, f. 2 (1904). | New Guinea. |
| 6. <i>S. nigripes</i> , Zetterstedt, Dipt. Scand. Vol. 1, p. 240 [1842] (<i>Ocydromia</i>);
Vol. 8, p. 2996 [1849] (<i>Ocydromia</i>); Scholz, Zeitschr. Ent. Breslau,
Vol. 5 (17), p. 48 [1851] (<i>Ocydromia</i>); Loew, Dipterent. Südafr.
p. 260 (1860); Roeder Wien. Ent. Zeit. Vol. 7, p. 96 [1888]
(<i>Hybos</i>); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29,
p. 45 [1893] (<i>Hybos</i>); Wahlgren, Ent. Tidskr. Vol. 31, p. 47 (1910);
Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 8 (1913). | Europe. |
| 7. <i>S. nitida</i> , Loew, Oefv. Vet. Akad. Förh. Vol. 14, p. 369 (1857); Dipterent. Südafr., Seychelles.
Südafr. Vol. 1, p. 260 (1860); Collin, Ent. Mo. Mag. 699, p. 185
(1922). | |
| 8. <i>S. opaca</i> , Loew, Oefv. Vet. Akad. Förh. Vol. 14, p. 369 (1857); Dipterent. Südafr. Vol. 1, p. 260, pl. 2, f. 44 (1860). | Caffraria. |
| 9. <i>S. parvicellulata</i> , Bezzi, Ann. Mus. Hungar, Vol. 2, p. 321, f. 1 (1904);
Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 333 (1920). | East Indies. |
| 10. <i>S. polita</i> , Loew, Berl. Ent. Zeitschr. Vol. 5, p. 321: Cent. 1, n° 27
(1861); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 254, f. 83
(1902). — PI. 5, FIG. 39. | United States. |

8. GENUS ACARTERUS, LOEW

Acarterus, Loew, Oefv. Vet. Akad. Förh. Vol. 15, p. 340 (1858); Dipterenf. Südafr. p. 262 (1860); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 118 (1889); Bezzi, Ann. Mus. Hungar., Vol. 2, p. 321 (1904).

Characters. — Third antennal joint conical, longer than the apical arista, which is stiff, straight and bare, its microscopic hairs visible only by very strong magnification; proboscis longer than the head, horizontally porrect, slender and sharply pointed, the palpi linear; eyes completely contiguous on the vertex. Body wholly black, subshining, the thorax apparently not greatly convex, abdomen conical. Legs simple, the hind femora neither thickened nor spinose below, underside of the hind tibiae and tarsi with short whitish pile. First vein very long ending near the second, which scarcely bows into the costa, stigma long and brown, basal cells of nearly equal extent, discal cell but little shorter than the second posterior, emitting two simple posterior veins, third and fourth veins slightly convergent.

Type species : *A. unicolor*, Loew (Pl. 6, Fig. 42), the original species. The genus *Acarterus*, founded on a South African fly that possessed only negative characters, has been associated with *Parahybos* by Bezzi. At first, the position of the arista, terminal in *Acarterus* and subapical in *Parahybos*, was thought a sufficient distinguishing trait to maintain the two as valid genera. But Bezzi has recently found a series of East Indian species, evidently closely related, in which the location of the arista varies. However, Loew's figures of *Acarterus* show the pedicel of the second and third veins much abbreviated, the first vein unusually long, the arista decidedly shorter than the third antennal joint and the thorax not greatly convex. The South African species can therefore not be congeneric with the Oriental species placed in *Acarterus* and *Parahybos*, and the latter accordingly have here all been removed to *Parahybos*, leaving *Acarterus* a monotypic genus.

Geographical distribution.

1. *A. unicolor*, Loew, Oefv. Vet. Akad. Förh. Vol. 15, p. 340 (1858); Dipterenf. Cape of Good Hope. Südafr. p. 262 (1860). — Pl. 5, Fig. 42.

9. GENUS SYNECHES, WALKER

Syneches, Walker, Ins. Saunders. Dipt. Vol. 1, p. 165 (1852); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 118 (1889); Schwarz, Proc. Ent. Soc. Wash. Vol. 2, p. 146 [1891] (*habits*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390, 436 (1895); Williston, Man. N. Amer. Dipt. p. 74 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 251 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258 (1903); Bezzi, Ann. Mus. Hungar., Vol. 2, p. 329 (1904), Vol. 3, p. 425 (1905); Melander, Williston's Man. N. Amer. Dipt. 3 ed. p. 224 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301, 313 (1909); Kertész, Cat. Dipt. Vol. 6, p. 9 (1909); Mc Atee, Ent. News, Philad. Vol. 20, p. 359 [1909] (*habits*); Coquillett, Proc. U. S. Nat. Mus. Vol. 38, p. 610 (1910); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 456 (1912); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 327 (1920).

Pterospilus, Rondani, Dipt. Ital. Vol. 1, p. 152 (1856); Schiner, Fauna Dipt. Austr. Vol. 1, p. 77, (1862); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 115 [1889] (*Pterospilus*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 256 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 598 (1910).

Epiceia, Walker, Proc. Linn. Soc. Lond. Vol. 4, p. 149 (1860); Bezzı, Ann. Mus. Hungar., Vol. 2, p. 329 (1904), Vol. 10, p. 457 (1912).

Harpamerus, Bigot, Rev. Mag. Zool. Vol. 11, p. 306 (1859); Bezzı, Ann. Mus. Hungar., Vol. 2, p. 329, 334 note (1904), Vol. 10, p. 457 (1912); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 334 (1920).

Characters. — Greatly hunchbacked, usually opaque dark species, yellow, measuring three to eight millimeters, sometimes wings clear or maculate, the stigma usually prominent, hind femora more or less spinose, the notum not pilose. Head hemispherical, occiput flattened, eyes meeting from the antennæ to the highly perched ocellar tubercle, the upper facets very large, the front part of the eyes somewhat flattened, antennal excision shallow; face narrow, retreating, no cheeks; antennæ two-jointed, the basal joints completely fused, the outer joint oval, with apical or subapical thread-like arista; proboscis styli-form, porrect, about as long as the head, palpi usually linear but short and hairy. Thorax very highly arched, notopleural suture short, pubescence sparse and short, notum more or less covered with close short tomentum so as to take away any shine, the color accordingly variable depending on the angle of view, bristles reduced to two notopleural, one dorsocentral and one postalar, but the scutellum margined with many hair-like bristles; mesopleuræ sometimes shining, as may be the mesonotum also in some species. Abdomen tapering in both sexes, the pygidium very small and bilateral, last two segments of the female forming a sort of short ovipositor. Legs variable, hind coxæ not greatly enlarged, anterior tibiæ sometimes bearing long bristles but usually bristleless, hind femora somewhat enlarged, longer than their tibiæ, usually spinose beneath, always with at least several stiff flexor bristles, hind tibiæ hairy, not curved nor with flattened sides, sometimes more or less swollen at the apex, tarsi not incrassate, usually lengthened but in some species the hind tarsi may be short and stout. Wings rather large, anal angle strongly rectangular, costa continuing to the fourth vein, no basal bristle, costal hairs inconspicuous, pedicel of the second and third veins long, arising toward the base of the second basal cell, third vein unforked, third and fourth veins parallel or converging, discal cell emitting two simple posterior veins.

Type species : *S. simplex*, Walker's original species (**Pl. I, Fig. 4**). This insect was previously described by Walker as *Gloma phthia*, according to Dr. Williston, who examined Walker's type. The enigmatical crossveins in the description of *phthia* refer to the two stigmal markings.

Coquillett, in the paper on the Type Species of American Diptera, page 503, would make *Syneches* a synonym of *Acromyia* Latreille. His argument that Latreille mentions the receipt from Bonelli of a specimen named *Acromyia asiliformis* illustrates the fallacy of overthrowing accepted names on scanty information. *Stomoxyx asiliformis* Fabricius is synonymous with *Asilus muscarius* Fabricius within the present genus *Syneches*, but Bonelli's reference to *asiliformis* is regarded as an erroneous identification for *Hybos grossipes*. Thus *Acromyia* is synonymous with *Hybos* and is not to be linked with *Syneches*.

The species of *Syneches* do not occur in the western part of North America but are not rare east of the Rocky Mountains. They have been found in all the other continents except Australia. These insects frequent meadowlands during the summer months and may be discovered in grass sweepings. They are often quite variable in color within the species. Three artificial subgenera are recognized, which are based on the following characters.

Syneches, sens. str. Hind femora not tuberculate nor greatly enlarged, but bearing several to many bristles or spines beneath; third and fourth veins parallel, the first posterior cell not narrowed; body color usually dark and opaque, wings sometimes maculate.

Type species : *S. phthia*, Walker.

Epiceia, Walker. Hind femora not greatly enlarged nor tuberculate, but bearing numerous spine-like bristles beneath, anterior tibiæ sometimes bristly; third and fourth veins converging, the first poste-

rior cell somewhat narrowed in the margin; body color often yellow, wings immaculate except for the stigma.

Type species: *E. ferruginea*, Walker.

Harpamerus, Bigot. Hind femora of the male bearing strong teeth or tubercles beneath from each of which extends a stout spine, hind tibiae of the male curved and tuberculate; anterior tibiae sometimes bearing long medial bristles; third and fourth veins converging, the first posterior cell somewhat narrowed in the margin; wings immaculate except for the stigma. The species that lack the dorsocentral, scutellar and tibial bristles have a broad abdomen. They may ultimately be segregated as a distinct genus.

Type species: *H. signatus*, Bigot.

Geographical distribution.

SUBGENUS SYNECHES, S. STR.

- | | |
|---|---------------------|
| 1. <i>S. albonotatus</i> , Loew, Berl. Ent. Zeitsch. Vol. 6, p. 195 : Cent. 2, E. United States.
No. 18 (1862); Coquillett, Proc. U. S. Mus. Vol. 18, p. 436 (1895);
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 252 (1902). | |
| 2. <i>S. annulipes</i> , Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 316 (1909). | Bolivia. |
| 3. <i>S. armatus</i> , nov. sp. (1). | Philippine Islands. |
| 4. <i>S. ater</i> , nov. sp. (2). | Pennsylvania. |
| 5. <i>S. barypterus</i> , nov. sp. (3). | Costa Rica. |

(1) **Syneches armatus**, nov. sp. — Male. Length 4.5 mm. Head flattened, upper facets very large; base of antennae yellow, last joint ovate, blackish, the arista three times antennal length; mouthparts testaceous; occiput and hairs black, lower portion lightly dusted. Thorax black, shining, more or less brownish on humeri and in front of posterior callousness, bristles and sparse hairs black, one long thin dorsocentral, scutellum with a pair of fine apical bristles and a few marginal hairs; pleurae mostly shining black, somewhat piceous under wings. Abdomen slender, flattened, basal half of second segment yellow, third segment yellowish at base, remainder of abdomen black with purple tinge, venter yellowish, hairs of basal half of abdomen pale, of apical half black, genitalia small, globular, the apex appearing forcipate. Coxæ and femora except anterior knees black or blackish, remainder of anterior legs and hind tarsi except last joint yellow, hind tibiae black, tibia black, pulvilli pale, hairs and bristles concolorous with integument, front tibiae with a long flexor seta at basal two-fifths, middle tibiae with a very long seta at basal two-fifths of both inner and outer sides and another at apex, hind femora stout, as long as their tibia and metatarsus together, the outer inferior face with seven stout spines arising from small tubercles, the ventral face with four smaller spines, flexor face of hind tibiae denticulate. Halteres with yellow stem and black knob. Wings nearly hyaline, apex smoky, veins coarse and blackish, root of fourth vein and anal vein faint, stigma large, black, filling the wide end of the marginal cell abruptly limited opposite posterior crossvein, third and fourth veins parallel, discal cell rather large, sections of fifth vein 1 : 0.6.

Female. Similar to male.

Mt. Makling, Luzon, C. F. Baker, fourteen specimens. Additional specimens were included in the sending but were destroyed by a dermestid larva during transit.

(2) **Syneches ater**, nov. sp. — Male. Length 3.5 mm. Black, velutinous, only the proboscis, knees narrowly, extreme tips of the tibiae and base of the anterior tarsi brownish-yellow, and the legs a little shining. Wings heavily infumated, the costal and marginal cells even darker, marginal cell as broad as the submarginal opposite the end of the first vein, third and fourth veins parallel beyond the discal cell. Thorax strongly convex, scutellum with about ten marginal hairs. Hind femora slender, pubescent, not spinose but with two long hair-like bristles below, hind tibiae closely pubescent. The insect has the form and proportions of *phthisia*.

A single specimen: Hazleton, Pennsylvania, received from Dr. Dietz, who collected it June 28, 1912. The species is readily recognizable by its dark body, wings and legs. The only other American species with dark legs and wings is *albonotatus* Loew, which has the thorax marked with white pollinose spots and the abdomen overlaid with brown pollen.

(3) **Syneches barypterus**, nov. sp. — Female. Length 4 mm. Black, including antennæ, coxæ, most of hind legs, knob of halteres, veins and stigma; the tip of proboscis, root of wings and of halteres, anterior legs beyond the coxæ, basal one-half of hind femora and basal fourth of hind tibiae yellowish. Third antennal joint short oval, subequal to second joint, the arista three times the antennal length. Thorax shining, one dorsocentral and one pair apical scutellar bristles, ten fine long marginal hairs on scutellum. Abdomen long and narrow, much surpassing the hind femora, its hairs pale on basal part and becoming dark on apical segments, middle tibiae with two extremely long setæ at basal third and a moderate one at tip, hind femora relatively slender, beneath with seven setæ of which the proximal four are widely spaced, long and

6. *S. Bigoti*, Bezzi, Ann. Mus. Hungar, Vol. 2, p. 360 (1904), Vol. 10, India.
 p. 456 (1912); Brunetti, Rec. Indian Mus. Vol. 9, p. 16 (1913);
 Fauna Brit. Ind. Dipt. Vol. 1, p. 329, pl. 4, f. 3 (1920).
bicolor, Bigot (not Walker), Ann. Soc. Ent. France (6), Vol. 9, p. 127
 [1889] (*Pterospylus*); Bezzi, Ann. Mus. Hungar, Vol. 2, p. 330 (1904).
7. *S. curvineura*, nov. sp. (1). Costa Rica.
8. *S. curvipes*, Fabricius, Syst. Ant. p. 147 [1805] (*Damalis*); Wiedemann, South America.
 Zool. Mag. Vol. 1, p. 1, 60, pl. 2, f. 3 [1817] (*Damalis*); Aussereurop.
 Zweifl. Ins. Vol. 1, p. 537, pl. 4, f. 9 [1828] (*Hybos*); Bezzi, Nova
 Acta Akad. Naturf. Halle, Vol. 91 p. 314, note 2 (1909).
9. *S. debilis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 436 (1895); S. and E. United States.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 253 (1902).
10. *S. dichætophorus*, Bezzi, Ann. Mus. Hungar, Vol. 2, p. 330 (1904), Vol. 10, East Indies.
 p. 456 (1912).
11. *S. dichogenus*, nov. sp. (2). Costa Rica.
12. *S. dichrous*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 318 (1909). Bolivia.
13. *S. duplex*, nov. sp. (3). Costa Rica.

slender, and the distal three are close-set on distinct tubercles, hind tibiæ somewhat clavate but otherwise simple, pubescence of legs conspicuous, hind metatarsi swollen and one-third as long as their tibiæ. Wings subhyaline somewhat dusky at apex, sections of costa nearly $5:2:2:1$, stigma beyond posterior crossvein, large and completely filling end of marginal cell, second vein curving so that the submarginal cell is two-thirds the width of the marginal at base of stigma, last two sections of fourth vein subequal, of fifth vein $1:0.6$.

Three specimens, La Suiza de Turrialba, Costa Rica, Nov 1922, Pablo Schild.

Related to *repletus* Bezzi, but differs in that all coxae and the hind tarsi are black and the stigma is transversely limited within.

(1) ***Syneches curvineura*, nov. sp.** — Length 3.5 mm. Closely related to *S. longiventris*, differing as follows: stigma larger, almost square, the marginal cell at the stigma two times as wide as the submarginal due to a pronounced curving of the second vein, veins thinner, wings subhyaline with cloudings about the anterior, posterior and anal crossveins; abdominal hairs relatively short and sparse, particularly on posterior portion of venter; thorny bristles before hind knee better developed, three or four in number.

Three males, three females, Turrialba, Costa Rica, April-August, Pablo Schild, in Melander collection.

(2) ***Syneches dichogenus*, nov. sp.** — Male. Length 3.3 - 4 mm. Occiput, knob of halteres and abdomen piceous, notum opaque blackish brown when viewed from in front, reddish brown but not cinereous when viewed from behind, antennæ, mouthparts, legs, stem of halteres, and most of pleuræ yellowish. Upper facets much smaller than ocelli, palpi linear. Mesonotum hemispherical, posteriorly not excavated, its bristles small and black, pleuræ with more or less evident piceous spots above posterior coxae. Abdominal hairs small and sparse, valves terminal shining thin and forcipate. Leg bristles small and brownish, no inside bristle on middle tibia, the extensor bristle located at two-fifths length of middle tibia one-third as long as tibia, hind femora without dorsal bristles, with four small equidistant bristles on anteroflexor face, tarsi slender. Wings uniformly smoky, the long elliptical stigma somewhat darker and extending below ends of first and second veins, veins thin and blackish, third and fourth veins nearly parallel.

Female. Mesonotum fuscous; wings hyaline, veins brown, stigma faintly yellow.

Eighteen males, fifteen females, received from P. Schild who collected them near Turrialba, Costa Rica, during July and August 1921. Types in Melander collection.

(3) ***Syneches duplex*, nov. sp.** — Male. Length 4.5 mm. Related to *simplex* and *stigmatical* in wing structure but differing in the larger size, larger ommatidia, shortened tarsi, the thickened and not slender hind tarsi and the more bristly hind femora. Piceous brown, including antennæ, halteres and coxae, palpi slender and yellow, proboscis and legs brownish-yellow; last joint of hind tarsi black. Upper facets nearly as large as ocelli. Thorax strongly convex, posteriorly declivous, subshining, the declivity gray when viewed from behind, but two distant vittæ, a triangular presacular spot and supra-alar marks faintly gray when viewed from above and back, bristles and sparse hairs black. Abdomen darker apically, its basal long thin hairs yellowish to black, genitalia small and terminal, claspers bluntly triangular, not forcipate. Bristles and hairs of legs pronounced, four equidistant extensor bristles on front tibiae, extensor bristle at two-fifths of middle tibiae and about one-half length of tibia, a shorter preapical bristle, postero-extensor row of four bristles on middle tibiae, middle femora with about eight bristles in each flexor row, hind femora with three stiff bristles on upper posterior face toward knee, with a row of six bristles beneath and with shorter flexor bristles, hind tibiae with long extensor bristles at middle and before apex, hind tarsi robust, the metatarsus with closely pubescent sole and with five black spines on posterior face, third and fourth joints broader than long. Wings lightly brownish, veins brownish yellow, a diffused quadrate stigmal

DIPTERA

14. *S. elevatus*, Beazzi, Ann. Soc. Ent. Belg. Vol. 52, p. 380 (1908). Congo.
 15. *S. fratellus*, Brunetti, Rec. Indian Mus. Vol. 9, p. 18 (1913); Fauna Brit. W. Himalayas.
 Ind. Dipt. Vol. 1, p. 332 (1920).
 16. *S. fuliginosus*, Meijere, Tijdschr. v. Ent. Vol. 58, p. 24 (1915). Simalur.
 17. *S. fuscescens*, Beazzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 319 (1909). Bolivia.
 18. *S. fuscipennis*, Brunetti, Fauna Brit. India, Vol. 1, p. 333 (1920). India.
 19. *S. hyalinus*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 437 (1895); E. United States.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 252 (1902).
 20. *S. immaculatus*, Brunetti, Rec. Indian Mus. Vol. 9, p. 18 (1913); Fauna Ceylon.
 Brit. Ind. Dipt. Vol. 1, p. 328, f. 24 (1920).
 21. *S. inaequalis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 19 (1913); Fauna India.
 Brit. Ind. Dipt. Vol. 1, p. 330 (1920).
 22. *S. insignis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 19 (1913); Fauna Brit. W. Himalayas.
 Ind. Dipt. Vol. 1, p. 332, f. 25 (1920).
 23. *S. lividus*, nov. sp. (1). Luzon.
 24. *S. longipeunis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 346 (1902). North Carolina.
 25. *S. longiventris*, nov. sp. (2). Costa Rica.
 26. *S. luteus*, Wiedemann, Aussereurop. Zweifl. Ins. Pt. 2, p. 647 [1830] Brazil.
 (Hybos); Beazzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 314 (1909).
 27. *S. minutus*, Brunetti, Rec. Indian Mus. Vol. 9, p. 20 (1913); Fauna Burma.
 Brit. Ind. Dipt. Vol. 1, p. 330 (1920).
 28. *S. muscarius*, Fabricius, Ent. Syst. Vol. 4, p. 390 [1794] (*Asilus*); Meigen, S. & C. Europe.
 Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 271 [1804] (*Asilus*);
 Coquebert, Illustr. Icon. Ins. p. 114, pl. 25, f. 12 [1804] (*Asilus*);
 Fabricius, Syst. Antl. p. 151, [1805] (*Dioctria*); Latreille, Gen. Crust. Ins. Vol. 4, p. 305 (1809); Consid. Gén. p. 443 [1810]
 (Hybos); Meigen, Syst. Besch. Vol. 2, p. 264, pl. 21, f. 21 [1820]

spot below end of first vein, apex of marginal cell with more or less evident secondary stigmal mark, submarginal cell slightly narrower than marginal at stigma, third and fourth veins apically parallel.

Female. Facets slightly smaller, styles small and black, wings subhyaline, the stigmal spots as in male.

Two males and two females, La Suiza de Turrialba, Costa Rica, April-July, collector Pablo Schild; in Melander collection.

(1) ***Syneches lividus*, nov. sp. — Male.** Length 4.5 mm. Almost wholly black, the anterior femora and tarsi, root of wings and of halteres, pleural sutures, postalar callosities and tip of proboscis brownish. Third antennal joint oval, nearly two times long as wide, the subterminal arista three times antennal length; sides and lower part of occiput brown-dusted. Two fine dorsocentrals and two scutellars, about ten minute marginal hairs; mesonotum very finely dusted, subshining. Abdomen long and slender, its hairs pale, becoming blackish at apex, genitalia small. Hind femora slender, only a little thicker than their straight tibiæ, underneath with seven moderate bristles, and towards the apex with four thorn-bearing flexor denticles, the usual single setæ of front tibiæ long and the three setæ of the middle tibiæ moderate; pulvilli yellowish, tips of claws black. Wings lightly infumated, darker at tip, stigma distinct, elongate oval, filling the end of the marginal cell beyond the first vein, marginal and submarginal cells of equal width, third and fourth veins gently converging, discal cell moderate, posterior crossvein opposite end of first vein, sections of fifth vein 1 : 0.7.

Three specimens, Mt. Makling, Luzon, Philippine Islands, C. F. Baker, collector.

(2) ***Syneches longiventris*, nov. sp. — Male.** Length 4.2 mm. Shining black with almost entirely black legs, dusky wings, black stigma and elongate abdomen. Antennæ black, outer joint orbicular, the arista sub-apical, palpi and proboscis testaceous. Thorax not excessively convex, declivous but not linearly impressed posteriorly, the dorsum with inconspicuous fulvous pile, scutellum with two strong and eight weaker bristles. Legs slender, the hind femora not greatly thickened, black pilose, the anterior knees, basal fifth of hind tibiæ, apical half of anterior tibiæ and the anterior tarsi brownish yellow, remainder of legs shining black, hind femora loosely seriatly bristly within, on flexor surface with four long bristles and with two preapical thorns, middle tibiæ with two very long bristles attached at proximal third, as in *dichetophorus* and *repletus*, each as long as the tibiæ, and with additional long bristles at middle and at apex, front tibiæ swollen at basal two-fifths. Abdomen slender, three times as long as the thorax, with loose long black hairs, continuing on the venter to the small pygidium. Halteres black, calypteres blackish and with black hairs. Wings smoky, veins firm and black, stigma large, filling tip of marginal cell and sharply margined within, third and fourth veins parallel.

Female. Thorns not differentiated before hind knees; styles slender and black.

One male and five females, La Suiza de Turrialba, Costa Rica, Pablo Schild collector; in Melander collection.

- (*Hybos*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 319 [1834] (*Hybos*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (17), p. 48 [1851] (*Hybos*); Schiner, Fauna Dipt. Aust. Vol. 1, p. 77 [1862] (*Pterospilus*); Neuhaus, Dipt. March. p. 68 [1886] (*Pterospilus*).
asiliformis, Fabricius (not Bonelli), Ent. Syst. Vol. 4, p. 395 [1794] (*Stomoxys*); Syst. Antl. p. 282 [1805] (*Stomoxys*).
hybos, Lamarck, Hist. Nat. Anim. sans Vert. Vol. 3, p. 404 [1816] (*Asilus*).
29. *S. natalensis*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 126 [1889] S. Africa.
(*Pterospilus*).
30. *S. nebulosus*, Loew, Oefv. Vet. Akad. Förh. Vol. 14, p. 360 (1857); Dip- terenf. Südafr. Vol. 1, p. 259 (1860).
31. *S. obeliscus*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 315 (1909). Bolivia, Peru.
32. *S. palliditarsis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 18 (1913); Fauna Brit. Ind. Dipt. Vol. 1, p. 331, pl. 4, f. 4 (1920). India.
33. *S. periscelis*, nov. sp. (1). Luzon.
34. *S. phaopterus*, Bezzi, Ann. Mus. Hungar, Vol. 3, p. 427 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 315 (1909). Peru.
35. *S. phthisia*, Walker, List. Dipt. Brit. Mus. Vol. 3, p. 492 [1849] (*Gloma*). E. North America.
— PI. I, Fig. 4.
- punctipennis*, Van der Wulp, Tijdschr. Ent. Vol. 10, p. 139, pl. 3, f. 18-21 (1867).
simplex, Walker, Ins. Saunders. Dipt. Vol. 1, p. 165, pl. 5, f. 7 (1852); Loew, Jenaische Zeitschr. Naturw. Jena, Vol. 36, p. 115 (1870); Wulp, Tijdschr. v. Ent. Vol. 25, p. 119 (1882); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 254, f. 86 (1902).
36. *S. planiceps*, nov. sp. (2). Costa Rica.
37. *S. præstans*, Bezzi, Ann. Mus. Hungar, Vol. 10, p. 457, 458 (1912). Formosa.
38. *S. pusillus*, Loew, Berl. Ent. Zeitsch. Vol. 5, p. 320: Cent. 1, No. 25 (1861); Williston, Trans. Ent. Soc. Lond. p. 308, pl. 11, f. 84 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 253 (1902). North America.
39. *S. pyramidatus*, Bezzi, Ann. Mus. Hungar, Vol. 3, p. 428 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 315 (1909). Peru.

(1) **Syneches periscelis**, nov. sp. — Male. Length 3.8 mm. Occiput black, cinereous below, the orbital fringe consisting of fine black hairs; antennæ yellow, the apical two-thirds of the small oval last joint brownish, the subterminal black arista four times the antennal length; mouthparts testaceous. Mesonotum subshining, testaceous, an indication of a broad median vitta on forward portion, hairs and bristles black, scutellum with two bristles and ten marginal hairs; pleurae yellowish, shining, more or less brown on meso- and sternopleuræ. Abdomen dull black, venter yellowish, hairs black, genitalia small and closed. Front coxae brown, posterior coxae pale yellow, the remainder of legs testaceous with a black band encircling apical third of hind tibiæ; bristles and hairs concolorous with integument, middle tibiæ with a relatively small seta at basal third of outer side and another at apex; hind femora rather slender, not reaching to end of abdomen, with seven spines on outer inferior face of which only the distal three are pronounced, the flexor face with six small thorn-bearing tubercles along the distal third, hind tibiæ simple, a little swollen on black portion; tarsal claws with black tip, pulvilli yellowish. Halteres black, the root yellow. Wings lightly infumated, stigma obsolete, veins mainly blackish, marginal cell narrow, third and fourth veins parallel, discal cell small, anterior crossvein at its base, the first basal cell therefore shorter than the second, sections of fifth vein subequal.

Holotype, Mt. Makling, Luzon, Philippine Islands, Prof. C. F. Baker, collector.

(2) **Syneches planiceps**, nov. sp. — Female. Length 4 - 4.5 mm. Black, the mouthparts, front tibiæ, base of hind tibiæ, all tarsi and root of halteres yellowish; thorax almost shining, the coating of golden pubescence sparse and not forming definite marks when viewed from different angles, abdomen polished, hairs and bristles of body fine, sparse and black. Bristles of front legs delicate and yellow, the strongest being at base of metatarsus, middle tibiæ with two very long black bristles at basal two-fifths in addition to other moderately long irregular delicate bristles, hind femora elongate, with eight flexor bristles in anterior row, the last six arising from pronounced tubercles, the largest measuring half the width of the femur, hind metatarsi somewhat swollen. Wings hyaline, veins narrow and black, stigma oval almost filling end of marginal cell, black, marginal and submarginal cells of equal width, first posterior cell widest opposite stigma, its bounding veins slightly converging and then becoming parallel at tip of wing; calypteres and fringe yellowish.

Six specimens secured from Pablo Schild who collected them near Turrialba, Costa Rica, during April and July. The tuberculate hind femora are suggestive of *Harpamerus*, but the distinct posterior dorsocentral bristles, the apical scutellars, the very long setæ of middle tibiæ and the relatively slender hind femora and simple tibiæ are as other species of *Syneches*, s. str.

40. *S. quadrangularis*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, Mexico.
p. 374 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 253,
f. 85 (1902); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91,
p. 314 (1909).
41. *S. repletus*, Bezzii, ibidem, Vol. 91, p. 317 (1909). Peru.
42. *S. ruficollis*, Walker, Ins. Saunders. Dipt. Vol. 1, p. 206 [1852] (*Hybos*); Brazil.
Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 315 (1909).
43. *S. rufus*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 320 : Cent. 1, No. 24 E. United States.
(1861); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 253 (1902).
44. *S. rusticus*, Brunetti, Rec. Indian Mus. Vol. 9, p. 20 (1913); Fauna Brit. India.
Ind. Dipt. Vol. 1, p. 331 (1920).
45. *S. semibrunneus*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 325 (1911). Vol. 56, Java.
supp. 68 (1913).
46. *S. stigmatical*, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 316 (1909). Bolivia.
47. *S. testaceus*, nov. sp. (1). Missouri.
48. *S. thoracicus*, Say, Jour. Acad. Nat. Sc. Philad. Vol. 3, p. 76 [1823]
(*Hybos*); Wiedemann, Aussereurop. Zweifl. Ins. Vol. 1, p. 538
[1828] (*Hybos*); Macquart, Dipt. Exot. Vol. 1, p. 272, pl. 13, f. 1
[1838] (*Hybos*); Say, Compl. Writ. Vol. 2, p. 68 [1859] (*Hybos*);
Glover, Manuscr. Notes 26, pl. 3, f. 21 [1874] (*Hybos*); Coquillett,
Proc. U. S. Nat. Mus. Vol. 18, p. 436 (1896); Melander, Trans.
Amer. Ent. Soc. Vol. 28, p. 254, f. 84 (1902). United States.
49. *S. varus*, nov. sp. (2). Luzon.
50. *S. vittatus*, Walker, Trans. Ent. Soc. Lond. N. S. Vol. 5, p. 286 [1860] Natal.
(*Hybos*); Bezzii, Bull. Soc. Ent. Ital. Vol. 37, p. 294 (1906).

SUBGENUS EPICEIA, WALKER

1. *E. bicolor*, Walker (not Bigot), Proc. Linn. Soc. Lond. Vol. 3, p. 91 (1859), East Indies.
Vol. 5, p. 237 [1861] (*Hybos*); Osten-Sacken, Ann. Mus. Stor. Nat.
Genova, Vol. 16, p. 434 [1881] (*Hybos*); Kertész, Termes. Fuezet.
Vol. 22, p. 174 [1899] (*Hybos*); Bezzii, Ann. Mus. Hungar. Vol. 2,
p. 331 [1904] (*Syneches*), Vol. 10, p. 457 (1912).

(1) **Syneches testaceus**, nov. sp. — Female. Length 5.5 mm. Resembling *rufus* Loew, but the marginal cell not widened at the stigma. Entirely testaceous, including the antennæ, mouthparts, halteres and legs, but the tarsi flavous and the last two segments of the abdomen fuscous, stigma strongly brown, elliptical, nearly filling out the end of the marginal cell. Mesonotum covered with yellow tomentum but when viewed from above appearing subshining, mesopleuræ shining. Hind femora spinose beneath. Wings hyaline, veins strong and brown, marginal cell rather acutely pointed, at the stigma not much wider than the submarginal cell.

One specimen : Kansas City, Missouri, June, received from F. Rogers.

(2) **Syneches varus**, nov. sp. — Male. Length 4.5 mm. Occiput black, lightly brown pollinose except above neck ; basal joints of antennæ yellow, third joint blackish, the subdorsal brown arista three times antennal length ; mouth-parts yellowish. Mesonotum testaceous, lightly coated with yellow dust, nearly bare of hairs, bristles black, two scutellars and ten fine marginal hairs ; upper pleurae light brown, lower yellow. Second abdominal segment testaceous, third piceous, remainder black, shining, hairs pale, genitalia rather small and forcipate, its hairs blackish, venter yellowish becoming brown posteriorly. Coxæ brown, femora black, the anterior knees lighter, anterior tibiae and tarsi yellow, hind tibiae brown at base remainder black, tarsi yellowish, the long seta at basal two-fifths of front tibiae very thin, the long setæ near middle and apex of middle tibiae not excessive ; hind femora stout, the outer lower face with ten stout spines, of which the distal six arise from distinct tubercles, four smaller spines on inner lower face ; hind tibiae strongly curved, with two denticles at two-fifths and three-fourths their extent. Knob of halteres piceous. Wings narrow, hyaline, apex lightly infumated, veins brown, stigma fuliginous, detached from the gently curving second vein and abruptly cut off within, third and fourth veins parallel, discal cell moderate, the sections of fifth vein 1 : 0.8.

Holotype, Mt. Makling, Luzon, Philippine Islands, C. F. Baker, collector. The parallel-sided apical cell excludes this species from *Harpamerus*.

2. *E. deficiens*, Walker, Proc. Linn. Soc. Lond. Vol. 3, p. 129 [1859] (*Hybos*); East Indies.
Osten-Sacken, Ann. Mus. Stor. Nat. Genova, Vol. 16, p. 434 [1881]
(*Hybos*); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 332 [1904] (*Syneches*),
Vol. 10, p. 457 (1912).
3. *E. eustylata*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 126 [1889] East Indies.
(*Pterospilus*); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 332 [1904]
(*Syneches*), Vol. 10, p. 457 (1912).
4. *E. ferruginea*, Walker, Proc. Linn. Soc. Lond. Vol. 4, p. 149 (1860), East Indies.
Vol. 5, p. 283 (1861), Vol. 7, p. 209, 233 (1864); Bezzi, Ann. Mus.
Hungar. Vol. 2, p. 331, note (1904), Vol. 10, p. 457 (1912).
5. *E. helvola*, Frey, Oefv. Finska Vet. Soc. Helsingfors. Vol. 59 A, Ceylon.
No. 20, 5 (1917).
6. *E. hyaloptera*, Bezzii, Ann. Mus. Hungar. Vol. 2, p. 331 (1904), Vol. 10, New Guinea.
p. 457 (1912).
7. *E. luctifera*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 457, 458 (1912). Formosa.
8. *E. minor*, Bezzii, Ann. Mus. Hungar. Vol. 2, p. 332 (1904), Vol. 10, New Guinea.
p. 457 (1912).
9. *E. pulla*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 457, 460 (1912), Suppl. Formosa.
Ent. Berlin, Vol. 3, p. 68 (1914).
10. *E. stigma*, Walker, Proc. Linn. Soc. Lond. Vol. 8, p. 111 [1865] (*Hybos*), New Guinea.
Vol. 9, p. 15 (1865); Bezzii, Ann. Mus. Hungar. Vol. 2, p. 333,
note (1904), Vol. 10, p. 457 (1912).

SUBGENUS HARPAMERUS, BIGOT

1. *H. Bakeri*, nov. sp. — PI. I, Fig. 6 (1). Philippines.
2. *H. dinoscelis*, Bezzii, Ann. Mus. Hungar. Vol. 2, p. 333 (1904), Vol. 10, New Guinea, Java.
p. 457 (1912).
3. *H. signatus*, Bigot, Rev. Mag. Zool. (2), Vol. 12, p. 306 (1859); Bezzii, Ceylon.
Ann. Mus. Hungar. Vol. 10, p. 457 (1912); Brunetti, Fauna Brit.
Ind. Dipt. Vol. 1, p. 334 (1920).
4. *H. velutinus*, Meijere, Expéd. Néerl. Guinée, Vol. 5, Dipt. 79 (1906); New Guinea.
Bezzii, Ann. Mus. Hungar. Vol. 10, p. 457 (1912).

10. GENUS PARAHYBOS, KERTÉSZ

Parahybos, Kertész, Termes. Fuezet. Vol. 22, p. 176 (1899); Bezzii, Ann. Mus. Hungar. Vol. 2,
p. 321 (1904), Vol. 5, p. 565, note (1907); Kertesz, Cat. Dipt. Vol. 6, p. 8 (1909); Bezzii, Ann.
Mus. Hungar. Vol. 10, p. 456 (1912); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 343 (1920).

Acarterus auctorum, Meijere, Tijdschr. v. Ent. Vol. 54, p. 328 (1911); Brunetti, Fauna Brit. Ind.
Dipt. Vol. 1, p. 334 (1920).

(1) **Syneches (Harpamerus) Bakeri**, nov. sp. (PI. I, Fig. 6). — Male. Length 6.5 mm. Black, dull. Upper facets but slightly larger than lower ones; face pollinose black; occiput silky black; basal two joints of antennæ yellowish, third joint elliptical, two and one-half times long as deep, black except at extreme base, arista black, three times length of antennæ; palpi black, proboscis piceous. Humeri and scutellum reddish, twelve marginal scutellar black bristles; middle of pleurae shining; hairs of posterior notum long and black. Abdomen much depressed, yellowish at base, genitalia small. Anterior legs including coxae yellowish, tarsi apically dusky, hairs mainly yellowish, hind legs entirely black, hind coxae swollen, hind femora hairy, incrassate, beneath in an anterior row with five very large setigerous prongs, the outermost of which is longer than the diameter of the femora, posteriorly with two smaller prongs, hind tibiæ slender, strongly curved, four-fifths length of femora, tarsi closely hairy. Halteres, calypteres and fringe yellowish. Wings blackish, becoming less dark posteriorly, fourth vein rounding toward tip.

Holotype, received from Prof. C. F. Baker who sent it as the most curious fly he had ever seen, Kolambungan, Mindanao, Philippine Islands.

Characters. — Thorax very strongly convex, subopaque, eyes separated on the face, arista often subdorsal, hind femora not setose, slender, no longer than their tibiæ, middle tarsi of the male sometimes deformed, pygidium bilateral. Head hemispherical; proboscis horizontally porrect, as long as the head, slender and pointed, palpi small; eyes with a horizontal line through the middle, but the upper facets not abruptly enlarged. Thorax without bristles except laterally, scutellum with several fine marginal hairs but no bristles; abdomen of the male somewhat upturned at the end, of the female tapering. Legs slender, often bearing curious bunches of hairs in the male sex, no extremely long tibial setæ such as in *Syneches*, underside of the hind femora completely unarmed, middle tarsi of male often shortened and distorted. Wings not large, stigma weak or absent, first posterior cell narrower in the margin than opposite the posterior crossvein, discal cell elongate, no alula.

Type of the genus: *P. iridipennis* Kertész, the original species. The genus *Parahybos* includes Oriental species only. For a time it was considered as a very close relative of *Acarterus*, the latter name being used for those species with subdorsal arista. As the recent discovery of additional species showed that this character was too elusive to use, the name *Parahybos* was dropped. However, as explained under the genus *Acarterus*, it seems that there has been a misconception of the African genus, and in the present paper all the Oriental species are grouped together under the name *Parahybos*.

Geographical distribution.

1. *P. chatoproctus*, Bezzı, Ann. Mus. Hungar, Vol. 5, p. 565 (1907), Vol. 10, Formosa.
p. 458, 463, f. 1 (1912), Suppl. Ent. Berlin, Vol. 3, p. 68 (1914).
2. *P. chiragra*, Bezzı, Ann. Mus. Hungar, Vol. 10, p. 458, 463, f. 2 (1912), Formosa
Suppl. Ent. Berlin, Vol. 3, p. 68 (1914). — Pl. I, Fig. 7.
3. *P. flavipes*, Brunetti, Rec. Indian Mus. Vol. 9, p. 21 (1913); Fauna Brit. India.
Ind. Dipt. Vol. 1, p. 343 (1920).
4. *P. fuscipennis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 21 [1913] (*Acarterus*); Ceylon.
Fauna Brit. Ind. Dipt. Vol. 1, p. 335, pl. 4, f. 5 [1920] (*Acarterus*).
5. *P. incertus*, Bezzı, Ann. Mus. Hungar, Vol. 10, p. 458 (1912). Formosa.
6. *P. infuscatus*, Meijere, Tijdschr. v. Entom. Vol. 54, p. 326 (1911), Vol. 56, Java.
Suppl. 71 (1914).
7. *P. iridipennis*, Kertesz, Termes. Fuzet, Vol. 22, p. 176 (1899); Wulp, New Guinea.
Tijdschr. v. Ent. Vol. 42, p. 49 (1899); Bezzı, Ann. Mus. Hungar,
Vol. 2, p. 336 (1904), Vol. 10, p. 457 (1912); Collin, Ent. Mo. Mag.
699, p. 186 (1922).
8. *P. luteicornis*, Frey, Oefv. Finska Vet. Soc. Helsingfors, Vol. 59, A, Ceylon.
nº 20, p. 6 (1917).
9. *P. melas*, Bezzı, Ann. Mus. Hungar, Vol. 10, p. 458, 461 (1912). Formosa.
10. *P. orientalis*, Meijere, Tijdschr. v. Ent. Vol. 50, p. 250, pl. 6, f. 17 [1907] Java.
(*Acarterus*); Bezzı, Ann. Mus. Hungar, Vol. 10, p. 456, 458 (1912);
Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. 71 (1914).
11. *P. ornatipes*, Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. 69, pl. 2, f. 8 (1914). Java.
12. *P. pallipes*, Bezzı, Ann. Mus. Hungar, Vol. 2, p. 335 [1904] (*Acarterus*), India.
Vol. 5, p. 566, note (1907); Meijere, Tijdschr. v. Ent. Vol. 50, p. 251
[1907] (*Acarterus*). Vol. 54, p. 328 (1911); Bezzı, Ann. Mus. Hungar, Vol. 10, p. 458 (1912); Brunetti, Fauna Brit. Ind. Vol. 1,
p. 335 [1920] (*Acarterus*).
13. *P. pusillus*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 327 (1911), Vol. 56, Java.
Suppl. 71 (1914).
14. *P. Sauteri*, Bezzı, Ann. Mus. Hungar, Vol. 10, p. 458, 464, f. 3 (1912); Formosa.
Suppl. Ent. Berlin, Vol. 3, p. 68, 69 (1914).
15. *P. simplicipes*, Bezzı, Ann. Mus. Hungar, Vol. 10, p. 458, 462 (1912). Formosa.

III. GENUS MEGHYPERUS, LOEW

Meghyperus, Loew, Stettin Ent. Zeit. Vol. 11, p. 303 (1850); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 560 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 78 (1862); Bigot, Ant. Soc. Ent. France (6), Vol. 9, p. 118 [1889] (*Meghyperus*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 255 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253, 259 (1903); Melander, Williston's Man. Dipt. N. Amer. 3 ed. p. 224 (1908); Kertesz, Cat. Dipt. Vol. 6, p. 2 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 566, (1910).

Characters. — Small shining black species with forked fourth vein, conical style-bearing outer antennal joint and well developed alula. Head globular, eyes of the male contiguous from the antennæ to the vertex, facets of the lowest third minute, eyes of the female broadly separated, the front shining, face short, cheeks very narrow; proboscis of *sudeticus* short, but in the American species the proboscis of the male slender, projecting horizontally forward, slightly longer than the head, the palpi linear elongate and porrect, and of the female the proboscis three times as long as the head, porrect, tipped with a distinct labella, the palpi short and clavate, attached to the proboscis near its base; antennæ two-jointed, the thickened style of the outer joint in the American species slightly less than one-half the length of that joint and tipped with a minute bristle, but in *sudeticus* the style is aristiform, tapering and longer than the antennæ. Mesonotum moderately arched, its hairs roughly arranged in acrostichal and dorsocentral rows, the hindermost dorsocentrals bristle-like, no hairs on the posterior declivity, about six marginal scutellar bristles, three notopleurals; pleuræ entirely pollinose and bare. Abdomen with loose long white hairs; male abdomen cylindrical, pygidium small, bilateral, the lateral valves fleshy and reniform; female abdomen tapering and less hairy. Anterior legs slender, no bristles, hind femora scarcely thickened and not at all spinose, hind tibiae somewhat clavate, particularly in the male, hind metatarsi somewhat swollen, pulvilli small. Wings hyaline, a more or less distinct oval stigma surrounding the end of the first vein, veins strong, costa straight, continuing to the anterior branch of the fourth vein, basal bristle small, costal hairs poorly developed, auxiliary vein distinctly separated from the first vein and vanishing at its end, third vein simple, petiole of the fourth vein as long as or longer than the posterior crossvein, basal and anal cells coextensive, anal crossvein rounded so as sometimes to meet the anal vein at a right angle, anal angle of the wing prominent, rectangular, alula of male much larger than that of the female.

Type species: *M. sudeticus*, the original species. The genus is interesting in the kind of sexual dimorphism it displays; the broad front of the female, her lengthened proboscis and shortened palpi and the difference in the size of the alula are unique in the Hybotinæ. The European *sudeticus* differs rather strangely from the American species in having a very short proboscis, a lengthened arista and, in the female, a bulbous face.

Geographical distribution.

1. *M. nitidus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 255, pl. 7, f. 72, W. United States.
73 (1902).
2. *M. occidens*, Coquillett, Proc. U. S. Nat. Mus. 18, p. 435 (1896); Melander, W. United States.
Trans. Amer. Ent. Soc. Vol. 28, p. 255 (1902). — Pl. 5, Fig. 40.
3. *M. sudeticus*, Loew, Stettin. Ent. Zeit. Vol. 11, p. 303, pl. 1, fig. 40, 42 C. Europe.
(1850); Schiner, Fauna Dipt. Austr. Vol. 1, p. 78 (1862).

12. GENUS IRONOMYIA, WHITE

Ironomyia, White, Pap. Proc. Roy. Soc. Tasmania, 1916, p. 216 (1917).

Characters. — Head broader than the thorax; eyes of male contiguous; proboscis scarcely projecting, palpi rounded and shorter than the proboscis; antennæ about one-fourth the length of the head, the first joint hardly distinguishable, the second a little longer than broad, the third rounded, arista thread-like, about twice the antennal length. Thorax only slightly arched, with a few weak posterior bristles, scutellum with weak marginal bristles. Abdomen short conical, male genitalia inconspicuous. Legs short, practically bare, tibiæ and tarsi subequal, tibiæ thin at the base, gradually thickening to the apex, unguis unusually large, three equal bristles between the well developed pulvilli. Wings broad, costa convex and serrated, auxiliary fused with first vein, which is long, reaching to the sharply defined stigma, second vein unusually long, at first well separated from the third after which the two veins converge, almost meeting below the stigma, and then slightly diverge, reaching the margin close together a little above the wing-tip, third vein unforked, discal cell long, emitting three veins, four posterior cells, the three basal cells short, the first a fraction longer than the second, the anal cell a little longer than the first, hind angle of wing strongly developed.

Genotype : *I. nigromaculata*, White, the only known species of this curious genus. This species is black, with abdomen centrally gray but marked with a dorsal row of black spots and was taken on a tree trunk in Tasmania.

Geographical distribution.

1. *I. nigromaculata*, White, Pap. Proc. Roy. Soc. Tasmania, 1916, p. 217, Tasmania.
f. 39 (1917). — Pl. 8, Fig. 86.

13. GENUS SCIADOCERA, WHITE

Sciadocera, White, Pap. Proc. Roy. Soc. Tasmania, 1916, p. 218 (1917).

Characters. — Head placed low down below the greatly arched thorax; eyes of female very widely separated; proboscis concealed within the oval aperture; antennæ so short that the very large, almost quadrangular terminal joint seems to lie directly against the face, the third joints of the two antennæ touching on the inner margin and reaching to the eyes on the outer margin, covering the face and completely concealing the preceding joints, arista thread-like; front with six stiff bristles — a divergent pair at the center of the vertex, a single one on each side close to the eyes, and one lower down above each antenna. Thorax with three complete rows of small dorsal bristles and well marked humeral, posthumeral, notopleural, supra-alar and postalar bristles, scutellum with four long marginal bristles. Abdomen about equal in breadth to, but a little longer than the thorax, altogether without bristles. Legs of medium length, bearing a few bristles, coxae considerably lengthened. Wings large, unusually broad, the costa strongly convex, auxiliary vein fused with the first vein, which ends at middle of wing, third vein not forked, discal cell open above and outwardly, four posterior cells indicated at the margins, the first two posterior veins incomplete basally, basal cells short, equal, anal cell a little shorter, no stigma.

Genotype : *S. rufomaculata*, White, the only known species. This is a yellowish-brown insect, four millimeters in length, with the black abdomen marked with four pairs of orange-red spots.

Geographical distribution.

1. *S. rufomaculata*, White, Proc. Roy. Soc. Tasmania, 1916, p. 219, f. 40 (1917). Tasmania.

SUBFAMILY OCYDROMIINÆ

Characters. — Small, delicate species, usually with shining body and yellow legs, bristles never strongly developed; thorax rarely very gibbous, shorter than the narrow cylindrical abdomen. Head more or less globular, rarely somewhat hemispherical; eyes never hairy, of the males broadly contiguous on the front (except in *Leptometopiella*), the facets of the upper two-thirds usually large, sometimes the eyes are contiguous in the females also (e. g. *Leptopeza*, etc.) in which case they meet on the face as well; antennæ often elongate due to a tendency toward enlargement of the third joint, the first joint usually abortive or fused with the second, sometimes (*Œdalea*) the third joint is remarkably lengthened, sometimes (*Anthalia*) this joint is broadly orbicular; the antennæ terminate in a long bare filiform arista, subdorsal in *Ocydromia*, or in a short, thickened two-jointed style, the latter rarely may be entirely absent; proboscis usually very short, scarcely protruding, sometimes as long as the head and more or less porrect; palpi small, single-jointed, usually narrowly cylindrical and with a few setæ, rarely the palpi are dilated; ocellar tubercle prominent in the male and then located on the very summit of the head, in dichoptic females the ocelli are not raised and are placed in front of the vertex; ocellar, vertical and occipital bristles never strongly developed although present. Thorax without discal bristles, the lateral bristles comprising a few notopleural, humeral, postalar and prescutellar bristles, several marginal but no discal bristles to the scutellum; pleuræ always devoid of bristles, but in *Leptopeza*, etc., somewhat villose in front of the metathoracic spiracle. Abdomen of the male with loose long fine hairs, pygidium small to minute, symmetrical, closed, the lateral valves largest, the dorsal valves reduced; represented by filament-like appendages, penis short and thick; an ovipositor usually developed from the last three or four segments. Legs typically slender and simple, sometimes (*Œdalea*, *Scelolabes*) there is a tendency for the hind legs to become raptorial similar to the parallel development in the Hybotinæ; coxæ never lengthened, tibiæ without bristles, except in *Leptopeza* and *Hoplopeza*, tarsi normal, pulvilli small, empodium microscopic. Calypteres very small but with a well developed fringe. Wings large, anal angle prominent and rectangular, except in *Leptometopiella*, no alula, costa evanescent beyond the fourth vein, the hind margin of the wing never thickened, basal bristle of costa usually present, marginal hairs weak, no setulæ; auxiliary vein rather weak, straight, parallel with and close to the first vein, evanescent before its end, first vein ending between three-fifths and three-fourth the wing-length, pedicel of the second and third vein arising over the basal half of the second basal cell, except in *Stenoproctus*, third vein always simple, discal cell large, located near the middle of the wing, complete or rarely opening into the third posterior cell, intercalary vein present except in *Stenoproctus* and *Scelolabes*, sometimes the fourth vein is lacking beyond the discal cell (*Leptopeza*, etc.) in which case the intercalary vein functions as the fourth vein; basal cells subequal in length, of moderate size, never very short or very long, the second basal sometimes broader than the first; anal cell shorter than the basals, truncate apically, the anal crossvein reflexed, nearly straight and abruptly meeting the anal vein, the latter always faint but often attaining the wing-margin.

This group is an offshoot of the Hybotinæ. It is quite probable that some ancestral form in the Ocydromiinæ was responsible for the Tachydromiinæ, and that of the modern genera *Trichina*, or better still, *Allanthalia pallida*, is structurally nearest to that form.

SYNOPSIS OF THE GENERA OF THE OCYDROMIINÆ

- A. Discal cell emitting three simple posterior veins, rarely the discal cell open in which case the intercalary vein is joined to the fourth, first vein ending near three-fifths the wing-length, stigma usually dis-

- tinct; females dichoptic, male eyes meeting on the front only; antennæ with a style; middle tibiae without bristles, hind femora ciliate with erect hairs 2.
- B. Discal cell emitting only two posterior veins, the fourth vein obsolete, first vein ending at three-fourths the wing-length, costa evanescent beyond third vein; eyes of both sexes contiguous on the face and usually on the front as well; antennæ with a lengthened arista; proboscis and palpi very small 7.
- C. Discal cell emitting only two posterior veins, the intercalary vein wanting; antennæ with short arista; proboscis and palpi usually projecting horizontally forward; hind femora incrassate and spinose below, hind tibiae geniculate 11.
- D. Discal cell absent, confluent with posterior cells, the anterior crossvein nearly or quite joined to the second basal cell, anal crossvein abruptly reflexed as in most Empidinae; face narrow 12.
2. Hind legs slender and not spinose; basal joint of antennæ imperfect, the third joint short or long 3.
- Hind femora thickened and spinose below, hind tibiae shortened, bowed near the knee; antennæ three-jointed, the third joint elongate, cylindrical and with a short thick style; upper facets of male eyes small, scarcely larger than the lowermost; ovipositor present.
(Pl. I, Fig. 9).
- Genus *ŒDALEA*, Meigen.
3. Proboscis rigid, lengthened, directed obliquely forward, palpi linear, face not deeply excised; front of female broad; antennæ three-jointed, the third joint elongate oval; thorax shining, no posthumeral bristles; ovipositor present; second basal cell very wide, its crossvein perpendicular (Pl. 8, Fig. 82).
- Genus *EUTHYNEURA*, Macquart.
- Proboscis very short or moderately lengthened; basal joint of the antennæ quite vestigial; thorax often more or less pollinose, posthumeral bristles present; second basal cell not greatly widened, its crossvein oblique 4.
4. Species measuring nearly five mm.; anal cell squarely truncate, discal cell narrow and much longer than second posterior, fourth vein interrupted, first vein ending close to second, third vein ending beyond tip of wing; third antennal joint oboviform; proboscis retracted; thorax moderately elevated; venter of male with sparse long hairs (Eocene)
- Genus **PROTOŒDALEA*, Cockerell.
- Species about two mm. in length, anal crossvein more retracted so as to be in line with the crossvein of the second basal cell, discal cell about as long as the second posterior, intercalary vein sometimes interrupted but not the fourth, first vein shorter, third vein ending at wing-tip 5.
5. Eyes of both sexes widely separated, upper facets not enlarged; antennæ lacking the style; proboscis retracted within the wide oral cavity, palpi bulbous; pygidium large and erect, ovipositor lengthened; yellow species with yellow bristles
- Genus *ALLANTHALIA*, nov. gen.

- | | |
|--|-----------------------------------|
| Eyes of male broadly contiguous above eyes, the upper facets more or less enlarged; antennæ ending in a style; proboscis more or less projecting; pygidium not surpassing diameter of abdomen, ovipositor not lengthened. | 6. |
| 6. Antennæ located near the middle of the head, the third joint elongate; proboscis very short, not porrect; front of female narrow, face not excised to the antennæ; dorsocentral bristles usually rather well developed; pygidium relatively large. (Pl. I, Fig. 8) | Genus TRICHINA, Meigen. |
| Antennæ located much below the middle of the head, the third joint short and very broad; proboscis short or somewhat lengthened, directed obliquely forward; front of female broad, face usually excised up to the antennæ; only the posterior two dorsocentrals distinctly differentiated; pygidium very small. | Genus ANTHALIA, Zetterstedt. |
| 7. Hind femora not thickened and not spinose, hind tibiæ not geniculate. Hind femora thickened, spinulose and spinose below, hind tibiæ geniculate and shortened; third antennal joint slender and elongate; pedicel of second and third veins short. (Pl. 5, Fig. 43) | Genus SCLOLABES, Philippi. |
| 8. Wings broad toward the base, the anal angle rectangular; discal cell shortened; eyes contiguous above and below the antennæ in both sexes; antennæ inserted at the middle of the head | 9. |
| Wings cuneiform, no anal angle developed, discal cell elongate, anal crossvein abortive; antennæ three-jointed | 10. |
| 9. Third antennal joint conical, the arista terminal; ovipositor ensiform; middle tibiæ with several sets of bristles. (Pl. 8, Fig. 79) | Genus LEPTOPEZA, Macquart. |
| Third antennal joint oval, the arista subdorsal; no ovipositor; middle tibiæ with only short apical bristles. (Pl. I, Fig. 10) | Genus OCYDROMIA, Meigen. |
| 10. Pedicel of the second and third veins long, anal vein incomplete; eyes separated on the front; legs simple; arista thin and bare | Genus LEPTOMETOPIELLA, nov. gen. |
| Pedicel of the second and third veins short, anal vein complete; eyes contiguous above as well as below the antennæ; hind tibiæ with apical spines, hind metatarsi spinulose above; arista thick and pilose. | Genus HOPI OPEZA, Bezzi. |
| 11. Pedicel of the second and third veins short, first basal cell shorter than second; females dichoptic; third antennal joint conical | Genus LAMACHELLA, nov. gen. |
| Pedicel of the second and third veins long, arising before the middle of the basal cells, the first basal longer than the second; eyes of female contiguous above the antennæ; third antennal joint more nearly cylindrico-conical. (Pl. 5, Fig. 44) | Genus STENOPROCTUS, Loew. |
| 12. Basal cells attaining middle of wing, second vein ending at two-thirds wing-length, sections of third vein proportioned 1 : 4, crossveins distinctly separated on fourth vein, intercalary vein entirely wanting; arista lengthened; thorax gibbous, without bristles but pubescent; legs long and slender, the front femora thickened basally. (Tertiary) | Genus *ELECTROCYRTOMA, Cockerell. |
| Anterior crossvein located very close to origin of third vein and nearly or quite touching the basal crossvein, second vein ending toward tip of wing, fourth and intercalary veins interrupted proximally, | |

- auxiliary vein separate from first; hind legs strongest, the anterior femora not thickened; thorax gibbous* 13.
 13. *Hind legs not raptorial. (Pl. 5, Fig. 46; Pl. 8, Fig. 72).* . . . Genus **BICELLARIA**, Macquart.
Hind femora much thickened, spinose, hind tibiae geniculate. (Pl. 2, Fig. 11). Genus **HOPLOCYRTOMA**, nov. gen.

I. GENUS EUTHYNEURA, MACQUART

Euthyneura, Macquart, Ann. Soc. Ent. France (1), Vol. 5, p. 517 [1836] (*Euthyneura*); Walker, Ins. Brit. Dipt. Vol. 1, p. 111 (1851); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 559 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 82 (1862); Loew, Beschr. Eur. Dipt. Vol. 2, p. 250, note (1871); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 117 (1889); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 256, 347 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 250, 260 (1903); Melander, Williston's Man. N. Amer. Dipt. 3 ed. p. 225 (1908); Kertesz, Cat. Dipt. Vol. 6, p. 109 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 543 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 209 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 43, 102 (1910).

Anthalia, Coquillet, Proc. Ent. Soc. Wash. Vol. 5, p. 246 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 507 (1910).

Characters. — Small, rather slender, shining black species with yellow legs, the abdomen sometimes pruinose. Head globular, as broad as the thorax, eyes bare, of the male broadly contiguous on the front, the upper facets enlarged or not, eyes of the female widely separated; face short, excised by the oral opening but the excision not reaching to the antennæ, cheeks obliterated or narrow; ocellar triangle of the male prominent, vertical and ocellar bristles not pronounced; antennæ inserted low down on the head, lengthened, plainly three-jointed, the last joint large, compressed, conical, with a short terminal style; proboscis as long as or longer than the head, rigid, projecting obliquely downward, palpi elongate and setose. Thorax not gibbous, bristles not pronounced, no posthumeral, one strong notopleural, one dorsocentral, several scutellars; pleurae shining. Abdomen of male cylindrical, furnished with long hairs, pygidium moderate, comprising a tumid basal part and thick lateral valves clasping the stout penis; of the female the abdomen is tapering, the last segments elongated to form a slender ovipositor. Legs simple, slender, bristleless, hind femora ciliate above. Wings large, sometimes clouded, costa vanishing beyond the fourth vein, its basal bristle present, auxiliary vein weak and evanescent, extending close to the first, the first vein ending at three-fifths the wing-length, discal cell large, located at the middle of the wing, emitting three posterior veins, sometimes open, in which case the fourth vein is furcate, basal cells elongate, the second basal broad and shorter than the first since its crossvein is nearly perpendicular, anal crossvein reflexed, abruptly meeting the nearly complete anal vein.

Type species: *E. myrtilli* (Pl. 8, Fig. 82), Macquart's only species. As noted under *Anthalia*, Coquillett would make *Gyllenhali* the type of that genus, thus drawing *Anthalia* into the synonymy.

SYNOPSIS OF THE AMERICAN SPECIES OF EUTHYNEURA

3. Abdomen of male silvery pruinose; hairs of legs and pygidium pale. (Or., Ida., Col.) E. ARGYRIA, nov. sp.
 Abdomen black; hairs of legs and pygidium black. (N. Mex.) E. APERTA, Melander.
4. Hind femora apically blackened and with flexor thorns near the knee; halteres of male with dark knob; third antennal joint short and broad. (Ida., Wash.) E. SPINIPES, nov. sp.
 Hind femora not blackened about the knee nor with thorns 5.
5. Stigma, veins, thoracic bristles, halteres and tarsi blackish; third antennal joint more than twice as long as wide. (Wash.) E. MATERA, nov. sp.
 Stigma, veins, halteres and tarsi pale; third antennal joint less than twice as long as wide. (Ont., Mass., N. H., Pa., Md., Mich., Wisc., Mo., Ida.) E. BUCINATOR, Melander.

Geographical distribution.

1. *E. aperia*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 348 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 [1903] (*Microphorus*). New Mexico.
 2. *E. argyria*, nov. sp. (1). Idaho, Oregon.
 3. *E. brevipes*, Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. 1840, p. 22, f. 28-30 [1840] (*Hemerodromia*), Oken, Isis, Encycl. Zeitschr. Leipzig, Vol. 7, p. 550, f. 28-30 [1840] (*Hemerodromia*); Wien. Ent. Monatschr. Vol. 8, p. 255 [1864] (*Hemerodromia*). C. Europe.
gracilipes, Schiner, Fauna Dipt. Austr. Vol. 1, p. 83, no desc. [1862] (*Hemerodromia*).
 4. *E. bucinator*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 348 (1902). North America.
 5. *E. crocota*. Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 413 [1900] (*Microphorus*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 257 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 [1903] (*Microphorus*). Alaska.
 6. *E. Gyllenhali*, Zetterstedt, Fauna Ins. Lappon. p. 538 [1838] (*Anthalia*); Dipt. Scand. Vol. 1, p. 249 [1842] (*Anthalia*), Vol. 8, p. 2998 [1849] (*Anthalia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 19, 50 [1851] (*Anthalia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 79 [1862] (*Anthalia*); Lundbeck, Dipt. Dan. Vol. 3, p. 211, f. 86, 87 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 80 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 60 (1913). N. and C. Europe.
 7. *E. immatura*, Zetterstedt, Dipt. Scand. Vol. 13, p. 4981 (1859). Scania.
 8. *E. matura*, nov. sp. (2). Washington.

(1) ***Euthyneura argyria*, nov. sp.** — Male. Length 2 mm. Black, abdomen silvery pruinose, legs brownish. Upper facets larger than lowermost; proboscis porrect, nearly as long as height of head, palpi small and black; antennæ with third joint compressed, conical, twice as long as deep, with fine white pubescence, style one-fourth as long as the third joint. Thorax polished, slightly dusky in front of wings, its sparse hairs and fine bristles white, six scutellars. Abdomen with long loose white hairs, the narrow sternites not pruinose, pygidium minute. Legs simple, hairs pale. Halteres white. Wings opalescent, veins white, stigma almost obsolete, discal cell open apically, the fourth vein forked midway its length.

Female. Front broad, at ocelli wider than length of third antennal joint, facets uniform; bristles and veins yellow, wings not opalescent; abdomen not pruinose.

Seven specimens. Mt. Hood, Oregon, elevation 6000 feet, July 30, 1922; Moscow Mountain, Idaho, June 29, 1918 (Melander); Colorado (F. C. Baker.).

(2) ***Euthyneura matura*, nov. sp.** — Male. Length 3 mm. Black, legs brownish, veins and stigma strong. Upper facets large; third antennal joint long, bluntly lanceolate, two and one-half times long as deep, velvety with microscopic pale hairs, style minute one-third as long as width of third joint; proboscis projecting obliquely forward nearly as much as head-height, palpi slender and black. Thorax polished, sparingly pollinose above the notopleural suture and posteriorly, the fine scattering hairs and the bristles blackish, six scutellars. Abdomen shining black, its hairs pale at base and black at apex, pygidium small. Coxæ and legs yellowish brown, the posterior tibiæ and all

9. *E. myrtilli*, Macquart, Ann. Soc. Ent. France (1), Vol. 5, p. 519, pl. 15, A. N. and C. Europe.

f 1-4 (1836); Walker, Ins. Brit. Vol. 1, p. 111, pl. 4, f. 3 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 82 (1862); Loew, Syst. Beschr. Vol. 2, p. 251, note 2 (1871); Beling, Arch. Naturg. Berlin, Vol. 48, p. 234 (1882); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 96 (1893); Wahlgren, Ent. Tidskr. Vol. 31, p. 80 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 61 (1913). —

Pl. 8, Fig. 82.

albibennis, Zetterstedt, Dipt. Scand. Vol. 1, p. 250 (1842).

var. *incompleta*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, p. 208 C. Europe. (1898).

10. *E. simillima*, Strobl, ibidem, Vol. 29, p. 97 (1893), Vol. 34, p. 208 (1898). C. Europe.

11. *E. spinipes*, nov. sp. (1). Western North America.

2. GENUS TRICHINA, MEIGEN.

Trichina, Meigen, Syst. Beschr. Vol. 6, p. 335 (1830), not Owen [1835] (*Nemathelminthes*), not Kirby, [1837] (*Coleoptera*); Zetterstedt, Fauna Ins. Lappon. p. 539 (1838); Rondani, Dipt. Ital. Vol. 1, p. 152 (1856); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 35 (1864); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258 (1903); Melander, Williston, Man. N. Amer. Dipt. 3 ed. p. 225 (1908); Kertesz, Cat. Dipt. Vol. 6, p. 100 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 184 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 43, 77 (1910).

Characters. — Small, slender, shining or subshining black species. Eyes of the male broadly contiguous above the antennæ, subcontiguous on the face, the upper facets large, of the female narrowly separated on both the front and the face; oral opening small and not encroaching on the face, proboscis very small, vertical, not protruding, palpi short and with a subapical hair; antennæ located near the middle of the head, elongate, imperfectly three-jointed, the last joint elongate, rather slender and tipped with a short two-jointed style; vertical and ocellar bristles small. Thorax moderately arched, bristles fine but rather long, humeral, posthumeral, notopleural, dorsocentral and scutellar bristles present,

tarsi blackish, hairs of femora and tibiæ pale, no bristles. Halteres blackish. Wings hyaline, stigma pronounced but not at all filling end of marginal cell, veins strong and brown, discal cell broad and short, the sections of the fourth vein proportioned about 5 : 1 : 5 : 7, intercalary vein weak at tip.

Seven specimens, all males, from Valleyford, Wash., 19 June 1919.

(1) **Euthyneura spinipes**, nov. sp. — Male. Length 2.2 mm. Body polished black, its sparse hairs and bristles yellow, legs including the coxæ pale yellow, the distal third of the hind femora, tibiæ and tarsi blackened, the hind femora apically black-setose. Upper facets large; antennæ black and pubescent, the third joint broadly oval, one-third longer than wide, the underside apically concave, the style one-third as long as the third joint; proboscis projecting obliquely forward about as far as the length of the head, palpi shining and furnished with a few long black hairs; occiput very lightly dusted, its sparse hairs black. Scutellum with four bristles. Hairs of the abdomen sparse and pale, pygidium minute. Hairs of the legs pale, extensor cilia of the hind femora equal to the diameter of the femur, about five black thorn-like setæ before the knee on the anterior side and three on the posterior side beneath. Halteres blackish. Wings slender, veins narrow and dark, a diffused stigma around the end of the first vein, discal cell as long as the wide second basal, emitting three complete posterior veins, its basal crossvein nearly perpendicular, sections of the fourth vein proportioned 1 : 0.2 : 0.9 : 1.2, of the fifth vein, 1 : 0.5 : 0.9 : 0.8, anal vein faint.

Female. Eyes separated, facets uniform; proboscis more nearly vertical; abdomen with conical apex, the last segment forming a lengthened ovipositor; halteres pale yellow.

Numerous specimens, Moscow Mountain and Lookout Mountain, Idaho, June, July; Mount Spokane, Washington; one specimen, Kaslo, British Columbia (U. S. N. M.). The species is distinct from *bucinator*, with which it cohabits, by the color and spines of the hind legs. The latter species is closely related to *myrtilli*, but has a shorter third antennal joint. Western specimens of *bucinator* have the hairs and bristles of the thorax invariably yellow, in Eastern specimens the bristles are almost always black.

discal setulæ almost wanting; pleuræ usually largely shining. Abdomen cylindrical, loosely hairy, pygidium of moderate size, often spinulose at the apex, no drawn-out ovipositor. Legs rather slender, simple, the hind tibiæ sometimes clavate, no true bristles, the hind femora ciliate with erect hairs. Wings large, anal angle prominently projecting, rectangular, costa evanescent beyond the fourth vein, with a basal bristle and rather weak marginal hairs, auxiliary vein extending close to the first which ends at three-fifths the wing-length, discal cell large, placed in the middle of the wing, emitting three posterior veins, basal cells large, subequal, anal crossvein reflexed, abruptly meeting the nearly complete but faint anal vein.

Type species: *T. clavipes*, Meigen (Pl. I, Fig. 8).

The name *Trichina* was used by Owen for the well-known genus of roundworms and as the Dipterous genus was neglected and of no economic importance the homonym has unfortunately come into general usage. This has resulted in a score of names connected with the roundworm disease, such as Trichinosis, Trichinotic, Trichiniferous, Trichinoscope, etc. Loew, in 1864, suggested that Dipterists concede Meigen's right of priority and change the Empid name, but Railliet in 1895 adopted an easier solution of the difficulty by proposing the term *Trichinella* for the conflicting Nematode.

KEY TO NORTH AMERICAN SPECIES OF TRICHINA

1. Third antennal joint subcylindrical, much longer than the style 2.
 Third antennal joint more or less conical, subequal to the style 3.

2. Pleuræ pollinose; pygidium small, side parts rounded; legs black; 8 scutellars (Cal.) T. ATRIPES, Melander.
 Pleuræ polished; side pieces of pygidium terminating above in a strong jet-black claw; legs fuscous; 6 scutellars (Mo.) T. NITIDA, nov. sp.

3. Legs wholly black; scutellum with 8 or 10 bristles. 4.
 Legs largely yellowish; scutellum with 4 or 6 bristles. 5.

4. Wings pale at base, brownish on apical half; acrostichals bristle-like; side pieces of pygidium terminating posteriorly in a pecten. (Cal.). T. BASALIS, nov. sp.
 Wings uniformly dusky; acrostichals not prominent; side pieces of pygidium ending in a finger-like process at the base of which is a reduced pecten (Wash., Mont.) T. PULLATA, nov. sp.

5. Stigma filling end of marginal cell; scutellum with four bristles; front of female about as broad as last antennal joint. (Ont., Me., Mass., N. Y.) T. NURA, Melander.
 Stigma not filling end of marginal cell; 6 scutellars; front of female very narrow. (Eur., Me.) T. FLAVIPES, Meigen.

Geographical distribution.

1. *T. atripes*, Melander, Trans. Am. Ent. Soc. Vol. 28, p. 349 [1902] (*Euthyneura*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 [1903] (*Microphorus*). Western United States.
 2. *T. basalis*, nov. sp. (1). California.

(1) ***Trichina basalis*, nov. sp.** — Male. Length 2 mm. Black, subopaque, completely coated with pollen which is brown on the thorax and gray on the abdomen. Upper facets large, antennæ inserted below the middle of the head, face relatively broad and excised midway to the antennæ; proboscis very short; antennæ with the third joint compressed conical, twice as long as wide and twice as long as the style; ocellar hairs short. Two posthumeral, three notopleural, eight scutellar, nine dorsocentral and four small supraalar bristles, a median double row of five

3. *T. clavipes*, Meigen, Syst. Beschr. Vol. 6, p. 336 (1830); Haliday, Ent. Mag. London, Vol. 1, p. 158 (1833); Macquart, Hist. Nat. Dipt. Vol. 1, p. 346 [1834] (*Microphorus*); Zetterstedt, Dipt. Scand. Vol. 1, p. 254 [1842] (*Microphora*); Walker, List. Dipt. Vol. 3, p. 488 [1842] (*Microphorus*); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 50 [1851] (*Microphora*); Walker, Ins. Brit. Dipt. Vol. 1, p. 114, pl. 4, f. 5 [1851] (*Microphorus*); Bonsdorff, Finl. tvåv. Ins. Vol. 1, p. 144 [1861] (*Microphora*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 79 [1862] (*Microphorus*); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 35, 40 (1864); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 133 (1887); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 93 (1892); Wien. Ent. Zeit. Vol. 18, p. 26 (1899); Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 311 (1906); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 72 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 187 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), 58 (1913).
- minuta*, Fallen (not Fabricius nor Walker) Empid. Suec. 32, p. p. [1816] (*Empis*) [see also *Edalea flavipes* Zett.]; Zetterstedt, Fauna Ins. Lappon. p. 538 [1838] (*Edalea*).
- ? *pallipes*, Zetterstedt, Fauna Ins. Lappon. p. 538, male only [1838] (*Edalea*).
- var. *lissonota*, Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 144 (1899).
- var. *sexsetosa*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 58 (1913).
4. *T. elongata*, Haliday, Ent. Mag. London, Vol. 1, p. 158 (1833); Walker, List Dipt. Brit. Mus. Vol. 3, p. 488 [1849] (*Microphorus*); Loew, Zeitschr. Ent. Breslau, Vol. 14, p. 40 (1863); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 59 (1913).
- fascipes*, Zetterstedt, Fauna Ins. Lappon. p. 540 [1838] (*Trichina*); Dipt. Scand. Vol. 1, p. 256 [1842] (*Microphora*), Vol. 8, p. 2999 [1849] (*Microphora*); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 50 [1851] (*Microphora*); Walker, Ins. Brit. Dipt. Vol. 1, p. 115 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 4982 [1859] (*Microphora*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 79 (1862); Loew, Zeitsch. Ent. Breslau, Vol. 17, p. 46 (1864); Strobl, Mitt. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 92 (1893).
- minuta*, Walker (not Fallen). Ins. Brit. Dipt. Vol. 1, p. 115 [1851] (*Microphorus*).
5. *T. nigripes*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, p. 206 (1898); Jahrb. Naturh. Mus. Kärnten, Klagenfurt, Vol. 47, p. 201 (1901). C. Europe.
6. *T. nitida*, nov. sp. (1). Missouri.

acrostichals; pleurae entirely pollinose. Abdomen with long pale hairs; pygidium rather large, the lateral valves convex, tipped with a row of black denticles, a pair of dorsal filaments present. Legs not shining, simple, the hairs entirely black, front tibiae not noticeably swollen, cilia of the hind femora shorter than the diameter of the femur. Calypteres and halteres piceous brown. Wings broadest in the middle, strongly infumated beyond the basal cells, veins strong, stigma distinct, discal cell slightly shorter than the second basal, petiole of the second and third veins arising over the basal fourth of the second basal cell, sections of the fourth vein proportioned 1 : 0.2 : 1 : 1, of the fifth vein, 1.6 : 0.7 : 1 : 1, last two posterior veins interrupted just before the margin, anal crossvein strongly reflexed, meeting the anal vein at an angle of one hundred twenty-five degrees, the anal vein vanishing much before the margin.

One specimen: Stanford, California, February 4, 1906, received from Dr. Aldrich.

(1) *Trichina nitida*, nov. sp. — Male. Length 2 mm. Black over all, the legs fuscous, paler apically. Occiput fulvous pollinose, its sparse hairs yellowish, third antennal joint four times as long as wide, its basal half barrel-shaped, its apical half cylindrical, the style minute. Mesonotum and most of pleurae polished, the posterior portion of pleurae lightly pollinose, bristles black, six scutellars, acrostichals and notal hairs very sparse and short. Abdominal hairs fine, sparse and pale, genitalia erect, globose, pollinose, the side parts ending in back above in a strong jet-black claw, the point directed forward. Legs simple, the posterior femora and hind tibiae bearing rows of fine hairs. Halteres black. Wings normal, subhyaline, stigma weak not filling the end of the marginal cell.

Female. Front shining, wider than the third antennal joint

Two specimens, Atherton, Missouri, May, received from Dr. C. F. Adams.

7. *T. nura*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 349 [1902] (*Euthyneura*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 [1903] (*Anthalia*). Eastern North America.
8. *T. opaca*, Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 40 (1864); Verrall, C. and S. Europe. Ent. Mag. London, Vol. 48, p. 26 (1912).
9. *T. pullata*, nov. sp. (1). Washington, Montana.

3. GENUS ANTHALIA, ZETTERSTEDT

Anthalia, Zetterstedt, Fauna Ins. Lappon, p. 538 (1838); Dipt. Scand. Vol. 1, p. 249 (1842); Schiner, Fauna Dipt. Austr. Vol. 1, p. 78 (1862); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 119 (1889); Melander, Williston Man. N. Amer. Dipt. 3 ed. p. 225 (1908).

Characters. — Small, robust species, with large wings and large discal cell. Eyes bare, contiguous in the male the whole length of the front except for a small triangular space just above the antennæ, the facets of the upper two-thirds large; face very short and broadly excised usually up to the antennæ, cheeks narrow; basal joint of the antennæ rudimentary, usually not visible, the middle joint very short, the third joint orbicular, compressed, very wide and short, with a terminal short, two-jointed arista which varies in length from as long as the third joint to about one-eighth as long; proboscis lengthened and often horizontally porrect, rarely surpassing the length of the head, rigid, palpi short and slightly swollen; ocellar triangle prominent, located in the male on the vertex, one pair of ocellar and one or two of vertical bristles. Thorax rather or quite prominent, moderately pollinose, more thickly so on the pleuræ, several humeral, posthumeral and notopleural bristles present, only the posterior two dorsocentrals distinct, acrostichals numerous but short, scutellum margined with from four to ten bristles. Abdomen furnished with loose hairs, longer in the male, pygidium small or minute, a short ovipositor formed from the shining last four segments, lateral pits visible. Legs short, simple, front tibiæ usually slightly swollen, hind femora more or less ciliate above and below. Costa stopping at the fourth vein, with a strong basal bristle but with short marginal hairs, auxiliary vein weak and lying close to the first, the latter ending just beyond the middle of the wing, discal cell longer than the basals, emitting three posterior veins, the intercalary sometimes abbreviated and sometimes joined to the fourth vein when the discal cell is open outwardly, anal crossvein abruptly meeting the anal vein which usually attains the margin although quite faint.

Nomenclature and type species : *A. Schoenherri*, Zetterstedt. Zetterstedt had three species when he proposed the genus *Anthalia*, — *Gyllenhalli*, *Schoenherri* and *pallida*. Coquillett (1903) considered the first of these as the type of *Anthalia* and thus threw the genus into the synonymy with *Euthyneura*, since *Gyllenhalli* is a very close relative of *myrtilli*, the monotypical species of *Euthyneura*. This would necessitate bestowing a new generic name on *Schoenherri* and the series of subsequently discovered species related to it. Since Coquillett did not adopt the wisest course we have an instance

(1) **Trichina pullata**, nov. sp. — Male. Length 2.25 mm. Black over all. Occiput lightly pollinose, its hairs fine, abundant, black; third antennal joint broadly lanceolate, subequal to the style. Thorax sparsely pollinose, hairs and bristles blackish, the lateral ones pale, twelve in dorsocentral row, ten pale scutellars, pleuræ pollinose. Abdomen subshining, hairs long and pale, genitalia large, open, extending backward, the lateral valves posteriorly with a pronounced pecten beyond which they continue as a finger-like process. Legs simple, the hind tibiæ slightly clavate, knees narrowly brown, hairs abundant, pale. Halteres black. Wings normal, somewhat smoky, stigma faint, not filling end of marginal cell.

Type, Paradise Park, Mt. Rainier, August, 1917. A male from Gold Creek, Montana, July 29, 1918 and a female from Gardiner, Montana, August 17, 1918, differ in having the mesonotum shining.

where it is just as well not to read the codes too literally, but to utilize Zetterstedt's second species as the genotype and thus preserve a well-known and time-honored name for this series of species. An iron-clad rule that permits any reviewer to fix for perpetuity the types of genera without first-hand knowledge of the species concerned should meet with protest. It were better to modify the rule so as to exclude from consideration species belonging to recognized genera. Zetterstedt's third species, *pallida*, is dichoptic and therefore is not congeneric with either of the others, a fact noted by Zetterstedt. The next following genus, *Allanthalia*, is erected for its reception.

The genera *Anthalia*, *Euthyneura*, *Trichina*, *Microphorus* and *Anthevisorius* have been much confused due to a general resemblance between them. The first three have the anal cell of similar construction, with the anal crossvein bent sharply back and meeting the anal vein in a distinct but obtuse angle, the characteristic neuration of the Ocydromiinae. In this subfamily they form a natural group distinct in having three posterior veins, a short, two-jointed antennal style with the first joint thick and the second joint microscopic, and are evidently related to *Edalea*. Those species with short antennae belong to *Anthalia*, those with lengthened antennae and short proboscis to *Trichina* and those with lengthened antennae and projecting proboscis to *Euthyneura*. As long as it was considered that the genera were defined by single characters there was justification in regarding them synonymous, and as these characters are all quite variable the genera have been merged by various writers. However, as indicated in the key to the genera, the species of each of these groups have a distinctive facies, and undoubtedly the groups as outlined present phyletic lines of descent which may be considered to have generic rank. The commonest American species differ from the typical forms in having a highly arched thorax, a more or less protruding proboscis and a lengthened antennal style, and bear a superficial resemblance to *Anthevisorius*, with species of which genus they associate at the same flowers.

TABLE OF THE NORTH AMERICAN SPECIES OF ANTHALIA

- | | |
|--|-------------------------------------|
| 1. Yellow species with yellow legs and halteres; proboscis very short | 2. |
| Black species, the halteres and legs usually blackish; bristles black,
pale only in <i>femorata</i> | 4. |
| 2. Bristles of the body yellow; head and base of the antennae yellow,
third antennal joint elongate oval; sections of the fifth vein 1 : 1.9. A. <i>GILVIHIRTA</i> , Coquillett. | |
| Bristles of the body black; head and antennae black; sections of the
fifth vein subequal; third antennal joint rotund (<i>bulbosa</i> ♀). | 3. |
| 3. Head entirely black, thorax medially darker, veins brownish | var. <i>BULBOSA</i> , Melander. |
| Front and cheeks, entire body and veins yellow. | var. <i>FLAVA</i> , Coquillett. |
| 4. Upper facets of ♂ eyes scarcely larger than the lower; style nearly as
long as the third antennal joint; proboscis projecting as far as the
length of the head, palpi very narrow; stigma distinct; scutellum
with four bristles | A. <i>MANDALOTA</i> , nov. sp. (1). |

(1) *Anthalia mandalota*, nov. sp. — Male. Length 2 mm. Entirely black including the antennae, mouth-parts, legs and halteres. Head rather hemispherical, eyes contiguous almost to the antennae, narrowly separated on the face and almost touching below, facets uniformly small; antennae short, the first joint minute, the third joint broadly pyriform, one-half longer than deep, the style slender and two-thirds the length of the third joint; proboscis geniculate at the base, slender and projecting forward a distance somewhat greater than the length of the head, the palpi narrow, almost linear and furnished with four small setæ; ocellar and vertical bristles short. Thorax greatly arched, nearly shining, hairs inconspicuous, four notopleurals, one posthumeral, four scutellars; pleurae lightly pollinose, a sternopleural spot shining: pygidium minute, abdominal hairs sparse. Front tibiae slightly swollen, short-ciliate along the extensor edge, extensor cilia of the hind femora as long as the diameter of the femur, of the under side bristle-like. Calypteres dusky.

Upper facets of ♂ eyes conspicuously larger than the lowermost; palpi not linear.	5.
5. Halteres, veins and wings whitish; abdomen sericeous, its hairs white	A. LACTEIPENNIS, nov. sp. (1).
Halteres fuscous to black, wings not opalescent, veins yellow to black.	6.
6. Stigma distinct; legs and halteres black; style distinct; proboscis often projecting; sternopleura more or less shining	7.
Stigma obsolete; legs and halteres brownish, hind femora rarely ciliate below; proboscis and style minute; pleuræ almost wholly opaque	II.
7. Hind femora and tibiæ thickened, the tibiæ rounded at the knee and shortened, the femora not ciliate beneath; hairs of the abdomen and legs and some of the thoracic bristles whitish; abdomen largely shining	A. FEMORATA, nov. sp. (2).

Wings with a very light infumation, veins blackish, stigma distinct, discal cell one-third longer than the rather narrow second basal, its posterior veins reaching the margin, sections of the fourth vein proportioned, $0.7 : 0.3 : 1 : 1$, of the fifth vein, $1 : 0.2 : 1 : 0.8$, anal angle prominent, anal crossvein parallel with the hind margin, anal vein faint.

Female. Eyes broadly separated on the front; abdomen tapering, the last four segments short and shining.

Numerous specimens from Mount Constitution, Vashon, Tacoma, Ilwaco, Sultan and Quilcene, Washington, May-July; and Portola, California, April, the last received from Dr. J. M. Aldrich.

(1) **Anthalia lacteipennis**, nov. sp. — Male. Length 1.8 mm. Body black, very lightly dusted, the abdomen more cinereous, antennæ and mouth-parts black, halteres whitish, wings and veins opalescent white, legs piceous. Eyes broadly contiguous on the front, the upper facets large, mouth-opening narrow, extending quite to the antennæ, epistome shining, cheeks narrow; antennæ two-jointed, the outer joint large and broadly pyriform, scarcely longer than broad and subequal in length to the style; proboscis projecting forward more than the length of the head, palpi slender and furnished with five setæ; vertical bristles small. Thorax highly arched, two pairs of strong prescutellars, four scutellars, one humeral, one posthumeral, three notopleurals and one postalar, acrostichals weak; pleuræ entirely pollinose. Abdomen with sparse long white hairs, pygidium minute, lateral pits distinct. Front tibiæ not noticeably swollen, extensor cilia of the hind femora short, flexor cilia setiform, metatarsi yellowish. Wings with parallel sides, costa yellow, other veins white, costal hairs black, no stigma, discal cell one-half longer than the narrow second basal, sections of the fourth vein proportioned $0.4 : 0.2 : 1 : 1.4$, of the fifth vein, $1.2 : 0.3 : 1 : 1.3$, posterior veins weak but reaching the margin, anal crossvein almost recurved, vanishing, anal vein obsolete.

Female. Eyes broadly separated by the shining front; discal cell longer, twice as long as the second basal, sections of the fourth vein, $0.4 : 0.15 : 1 : 1$, of the fifth vein, $1 : 0.15 : 1 : 1$; terminal segments of the abdomen short and shining.

Numerous specimens; type from Moscow Mountain, Idaho, frequenting flowers of *Pentstemon* on the summit (4900 feet altitude), July 3, 1911. Paratypes from the same place and from Bovill and Waha, Idaho; Pullman, Almota, Glenwood, Husum, Blewett, Seattle, Tacoma, Vashon, South Bend, Nahcotta and Ilwaco, Washington; Kaslo, British Columbia (U. S. N. M.); Yosemite Valley (Cresson), Humbolt Co. (H. S. Barber) and Santa Cruz Mountains (Doane), California.

(2) **Anthalia femorata**, nov. sp. — Male. Length 2.3 mm. Black, subshining, bristles of lower occiput, sides of thorax, abdomen, coxæ and legs whitish, hind legs robust, the tibiæ shortened, abdomen mainly shining and devoid of pollen. Eyes broadly contiguous on the front, the upper facets large, face widely excavated up to the antennæ, proboscis directed forward, but scarcely protruding from the oral opening, palpi with a few long setæ, ocellar bristles stout. Mesonotum lightly dusted, discal setulae pale, four pale humerals, five or six pale posthumerals, three black and two pale notopleurals, two black dorsocentrals and six black scutellar bristles; pleuræ pollinose except most of the sterno- and mesopleuræ. Venter and sides of the tergites shining, the upper side of the abdomen lightly pollinose, leaving however a circular shining spot at the base of each hair; pygidium scarcely longer than the preceding segment, broad. Anterior knees narrowly yellow, front tibiæ not swollen, hind femora much thicker than the others, highly shining, not ciliate below, but above with a row of short white cilia which are two-thirds as long as the thickness of the femur, hind tibiæ three-fourths as long as their femora, likewise incrassate, shining except as the tip, strongly curved at the knee. Halteres black, calypteres blackish and fringed with dusky hairs. Wings nearly hyaline, broader at the base, veins blackish, stigma distinct, discal cell blunt, one-half longer than the rather narrow second basal which is subequal in length to the first, posterior veins complete, sections of the fourth vein proportioned $0.6 : 0.3 : 1 : 1$, of the fifth vein, $1.3 : 0.3 : 1 : 0.9$.

Eight specimens: Kettle Falls, Washington, May 3, 1912, and Moscow Mt., Idaho, June and July. In the structure of the hind legs this species suggests *Œdalea*, but there are no spines and the knee is not angulate. The general habitus, with short thick body, is like the species of *Anthalia*.

- Hind legs not thickened, the femora ciliate beneath, the tibiæ not shortened; thoracic bristles and hairs of abdomen, coxæ and femora black, except in *interrupta*; abdomen pollinose. 8.
8. Scutellum with eight or six bristles; mesonotum only subshining. its bristles strong and abundant 9.
- Scutellum with four or six bristles; mesonotal bristles weak and not conspicuous (*stigmatis*) 10.
9. Pygidium minute; hairs of the abdomen black; intercalary vein complete; style of the antennæ about two-thirds as long as the third joint
- Pygidium relatively larger, obliquely ascending; hairs of the abdomen whitish; second posterior cell very narrow at the base, the intercalary vein abruptly shortened; style nearly as long as the third joint
10. Discal cell complete, emitting three veins of which the intercalary may be evanescent.
- Discal cell open outwardly, the fourth vein forked
- A. *SCUTELLARIS*, nov. sp. (1).
- A. *INTERRUPTA*, nov. sp. (2).
- var. *STIGMALIS*, Coquillett.
- var. *PETIOLATA*, nov. var. (3).

(1) *Anthalia scutellaris*, nov. sp. — Male. Length 2 mm. Entirely black, rather opaque dusted, bristles well developed, wings sybhyaline, stigma distinct. Head rather hemispherical, eyes contiguous along the front, face very short, excised by the mouth-opening up to the antennæ; proboscis retracted, or if porrect, not extending beyond the head, palpi rather long and broad and with three minute setæ; antennæ two-jointed, the outer joint broadly orbicular, but slightly longer than the style; ocellar bristles strong. Thorax bristly, the discal setulae prominent and relatively long, typically the following bristles present, but there may be some variation: four humeral, four posthumeral, seven notopleural, two dorsocentral, one postalar, eight scutellar; pleurae almost entirely dark pollinose. Abdomen subopaque, its hairs black, pygidium minute and not exposed. Cilia of the hind femora longer than the diameter of the femur, tarsi entirely black and with black hairs. Halteres black. Wings with parallel sides, veins blackish, discal cell one-third longer than the narrow second basal, sections of the fourth vein proportioned $0.5 : 0.25 : 1 : 1$, of the fifth vein $1.2 : 0.2 : 1 : 1$, anal crossvein parallel with the hind margin, anal vein thin.

Female. Eyes broadly separated by the pollinose front, face less excised, facets uniform, thoracic bristles less pronounced, last four segments of the abdomen forming a short ovipositor, legs piceous, the front coxæ and femora only fuscous.

Numerous specimens from the flowers of pink currant (*Ribes*), Tacoma and Seattle, Washington, May 12-14, 1913; also from Ilwaco and Mt. Rainier, Washington.

(2) *Anthalia Interrupta*, nov. sp. — Male. Length 1.8 mm. Entirely black, subshining, scutellum with six bristles, abdomen with white hairs, pygidium relatively large, upper facets enlarged, style subequal to the orbicular third joint of the antennæ, stigma distinct, discal cell pointed, intercalary vein shortened. Face broadly excised up to the antennæ, proboscis scarcely projecting from the oral opening, palpi thick; basal joint of the antennæ minute, third joint very broadly oval, scarcely longer than deep and subequal in length to the style; ocellar bristles distinct. Thorax highly arched, its bristles rather prominent and black, about four humerals, five small subhumerals in a transverse row, four notopleurals, two dorsocentrals, six scutellars; pleurae dusted but with a large shining sternopleural spot encroaching on the mesopleura. Pygidium obliquely ascending, as long at the two preceding segments together. Hairs of the coxæ and cilia of the hind femora pale, the cilia equal in length to the diameter of the femur, hind tibiae straight and nearly as long as their femora. Calypteres dusky but fringed with pale hairs. Wings nearly hyaline, veins blackish stigma distinct, discal cell one-half longer than the narrow second basal, broad but tapering almost to a point at the apex, the second posterior cell much narrower than the third, intercalary vein extending two-thirds the distance to the margin, abruptly interrupted, second basal cell longer than the first, sections of the fourth vein proportioned $0.55 : 0.2 : 1 : 1.3$, of the fifth vein $1.4 : 0.4 : 1 : 1.4$, anal angle prominent.

Female. Ovipositor segments very short.

Numerous specimens, Kettle Falls, Washington, May 3, 1912, from flowers of wild cherry.

(3) *Anthalia stigmatis*, var. *petiolata*, var. nov. — *Anthalia stigmatis* is widely distributed along the Pacific Coast. I have over four hundred specimens from Alaska, Idaho, Washington, Oregon and California, the species extending inland as far as Kendrick, Idaho. A series of specimens, ten males and eight females, from the summit of Moscow Mountain and from Craig's Mountain, both in Idaho, uniformly differ from the typical form in having the discal cell apically open, the intercalary vein joining with the fourth vein midway between the anterior crossvein and the tip of the wing. As the variation is constant and is not found in any of the other specimens this form may be

11. Pygidium large, obliquely ascending; intercalary vein interrupted;
thorax shining; legs blackish A. SCHOENHERRI, Zetterstedt.
- Pygidium minute; intercalary vein complete; thorax coated with
pollen, at most subshining; legs fuscous. 12.
12. Style one-sixth as long as the third antennal joint; veins delicate . . A. BULBOSA, Melander ♂.
- Style one-half as long as the third joint; veins coarse and yellowish. A. INORNATA, nov. sp. (1).

Geographical distribution.

1. *A. bulbosa*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 349 [1902] Eastern North America.
(*Euthyneura*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263
[1903] (*Euthyneura*); Slosson, Ent. News, Philad. Vol. 14, p. 267
[1903] (*Euthyneura*).
2. *A. femorata*, nov. sp. Washington, Idaho.
3. *A. flava*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 268 (1903); Slosson,
Ent. News Philad. Vol. 14, p. 266 (1903). Eastern North America.
4. *A. gilvihirta*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 268 [1903] Eastern North America.
(*Microphorus*).
5. *A. inornata*, nov. sp. Washington.
6. *A. interrupta*, nov. sp. Washington.
7. *A. lacteipennis*, nov. sp. Western North America.
8. *A. mandalota*, nov. sp. Western North America.
9. *A. Schœnherri*, Zetterstedt, Fauna Ins. Lappon, p. 539 (1838); Dipt. Sc.
Vol. 1, p. 252 (1842); Frey, Acta Soc. Sc. Fenn. Helsingfors,
Vol. 31 (9), p. 5 [1908] (*Euthyneura*); Strobl, Mitteil. Naturw. Ver.
Steiermark, Graz, Vol. 46, p. 74 [1909] (*Euthyneura*); Wahlgren,
Ent. Tidskr. Vol. 31, p. 80 [1910] (*Euthyneura*); Frey, Acta Soc.
Fenn. Vol. 37 (3), p. 61 [1913] (*Euthyneura*).
10. *A. scutellaris*, nov. sp. Washington.
11. *A. stigmalis*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 268 (1903).
var. *petiolata*, nov. var. Western North America.
Idaho.

4. GENUS ALLANTHALIA, NOV. GEN.

Characters. — Small yellow species with the habitus of *Anthalia*. Head spherical, eyes bare, widely separated on both front and face in both sexes, facets small; antennæ inserted below middle of head, basal joint vestigial, third joint large, elongate oval, twice as long as wide, without style; face

given the varietal name *petiolata*. *Stigmalis* and this variety differ from *scutellaris* in having four scutellar bristles, the meso-, ptero- and sternopleuræ mostly shining, the mesonotum less dusted, the hairs and bristles of the thorax reduced, there being but four notopleural, three humeral and one posthumeral bristles, and the wings pure hyaline.

(1) ***Anthalia Inornata*, nov. sp.** — Female. Length 2.25 mm. Piceous black in color, quite opaque, legs and halteræ fuscous. Head brownish pollinose, subshining; antennæ black, the third joint large, orbicular, as deep as long, its style one-third as long as the joint; mouthparts retracted, black; hairs and bristles black. Thorax dusted excepting a small spot on the sternopleura, hairs and bristles black, five notopleurals, three posthumerals, eight scutellars, discal hairs short but bristly, four rows of acrostichals; abdomen subopaque except the last three segments which are not lengthened, hairs black. Front tibiæ swollen, hairs of the legs rather outstanding, mainly black, those of the tibiæ and tarsi in part paler, extensor cilia of the hind femora shorter than the diameter of the femur, flexor cilia long, bristle-like and about thirteen in number. Wings large, lightly cinereous, veins brownish yellow and rather strong, no stigma, discal cell one-half longer than the narrow second basal, sections of the fourth vein proportioned 0.6 : 0.2 : 1 : 1, of the fifth vein, 1 : 0.2 : 1 : 0.8, anal vein faint, posterior veins attaining the margin, second posterior cell broad at the base, basal cells equal, the discal crossvein oblique.

One specimen : Almota, Washington, May 25, 1913.

short, oral cavity wide, proboscis retracted, palpi spatulate, fleshy, bare; cheeks broad behind. Humeral, posthumeral, notopleural, supra-alar, two posterior dorsocentral and four scutellar bristles present and yellow, notal hairs short. Pygidium large and erect, about twice as high as abdomen, lateral valves prominent; ovipositor elongate, formed from the polished 6-8 segments together with the styles. Legs simple, not ciliate. Wings normal, as in *Anthalia*.

Type species: *A. pallida*, Zetterstedt. Zetterstedt questioned the propriety of including his species in *Anthalia*.

Geographical distribution.

The species occurs in North America as well as in Europe, specimens having been studied from Fall's Church, Va. (Banks); Plummer's Island, Md., from blossoms of wild plum (Shannon); Aylmer and Hull, Quebec (Curran), and Atherton, Mo (Adams). The following is the only known species.

1. *A. pallida*, Zetterstedt, Lappon. p. 539 [1838] (*Anthalia*); Dipt. Sc. Vol. 1, North Europe; North p. 253 [1842] (*Anthalia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 80 America. [1910] (*Euthyneura*).

5. GENUS *ŒDALEA*, MEIGEN.

Œdalea, Meigen, Syst. Beschr. Vol. 2, p. 355 (1820); Macquart, Dipt. N. France, Vol. 3, p. 141 (1827); Hist. Nat. Dipt. Vol. 1, p. 321 (1834); Meigen, Syst. Beschr. Vol. 7, p. 77 [1838] (*Œdalia*); Zetterstedt, Fauna Ins. Lappon. p. 537 (1838); Blanchard, Hist. Nat. Ins. Vol. 3, p. 581 [1840] (*Œdalea*); Westwood, Gen. Syn. p. 133 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 245 (1842); Boitard, Man. Ent. Vol. 3, p. 317 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 111 (1851); Rondani, Dipt. Ital. Prodr. p. 152 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 559 (1857); Schiner, Fauna Dipt. Aust. Vol. 1, p. 80 (1862); Lioy, Atti Instit. Ven. 1864, p. 724 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 115 (1889); Melander, Trans. Am. Ent. Soc. Vol. 28, p. 256 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 254, 260 (1903); Melander, Williston Man. N. Amer. Dipt. p. 224 (1908); Kertesz, Cat. Dipt. Vol. 6, p. 104 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 578 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 193 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 43, 78 (1910).

Xiphidiocera, Macquart, Hist. Nat. Dipt. Vol. 1, p. 356 (1834); Blanchard, Hist. Nat. Ins. Vol. 3, p. 583 [1840] (*Xiphidiocera*); Boitard, Man. Ent. Vol. 3, p. 324 (1843); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 560 [1857] (*Xyphidiocera*); Lioy, Atti Ist. Veneto Sc. Venezia, 1864, p. 722 (1864); Meunier, Bull. Soc. Ent. France, 1894, p. IX (1894); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 259 (1903).

Characters. — Moderately stout, rather small species with robust hind femora and greatly elongated antennæ. Head globose, eyes bare, of the female well separated, of the male contiguous on the front, with the antennæ inserted below the middle of the head, upper facets scarcely larger than the lowermost; basal joints of the antennæ small but distinct, the third joint greatly lengthened, more or less cylindrical and from four to eight times as long as broad, tipped with a short two-jointed style, the apical joint of which is minute and bristle-like, the style variable in length, sometimes almost wanting; proboscis short or protruding obliquely forward nearly as far as the head-height, palpi minute; ocellar triangle not elevated, ocellar and vertical bristles poorly developed. Thorax quadrate, finely pubescent, only the prescutellar, dorsocentral, two or three notopleural, one postalar, and about six scutellar bristles present. Abdomen conical, the female with chitinized ensiform ovipositor, the male with a very small

symmetrical pygidium bearing an erect dorsal basal part. Anterior legs normal, bristleless, hind femora elongate, rather clavate, beneath toward the apex with four rows of thornlike spines, those of the lateral rows longer, hind tibiae geniculate, compressed to an edge opposite the femoral spines, tarsi not swollen. Wings broad, anal angle broadly rectangular, auxiliary vein rather weak and close to the long first vein, costa interrupted at the fourth vein, its basal bristle very weak, basal cells as long as the large discal cell, third vein simple, posterior veins shortened by the enlargement of the discal cell, anal crossvein truncating the anal cell, the anal vein weak but usually attaining the margin.

Type species: *Œ. hybotina*, Fallen, by Westwood's designation. Lundbeck records *flavipes* as having been bred from decaying wood. The species of *Œdalea* have yellow legs, the hind pair more or less darkened. The body is usually shining black with yellow pubescence and bristles. They are sluggish in flight and occur in shady woodlands, but are not commonly met with.

KEY TO THE NEARCTIC SPECIES OF *ŒDALEA*

1. Third antennal joint relatively broad and tapering, four times as long as broad and eight times as long as the style; thorax pollinose; intercalary vein usually short *Œ. PRUINOSA*, Coquillett.
- Third antennal joint elongate cylindrical, much more than four times as long as broad and the style much shorter; thorax shining; intercalary vein complete 2.
2. Antennal style almost invisible; upper facets of ♂ obviously larger; halteres fuscous, wings with strong stigma. *Œ. ASTYLATA*, nov. sp. (1).
- Antennal style visible, about one-fifteenth as long as the third joint; halteres yellowish 3.
3. Body black; discal cell nearly three times as long as wide, its sides parallel; stigma distinct; upper facets of ♂ obviously larger . . *Œ. OHIOENSIS*, Melander.
- Pleuræ at least testaceous; discal cell somewhat widened apically; stigma suffused; upper facets ♂ scarcely larger (*lanceolata*, nov. sp.) 4.
4. Mesonotum, abdomen and hind tibiae blackish. *Œ. LANCEOLATA*, nov. sp. (2).

(1) *Œdalea astylata*, nov. sp. — Male. Length 3 mm. Body polished black, legs including the coxae pale yellow, the hind tibiae brownish; antennal style microscopic; halteres fuscous. Proboscis short, black, scarcely protruding beyond the oral opening; third joint of the antennæ cylindrical, eight times as long as wide and about twenty times as long as the almost invisible style; upper facets conspicuously larger than the lower. Thoracic hairs rather sparse, scutellum bearing six brown bristles, prealar bristle brown; pleuræ highly shining. Pygidium rather long. Hind femora with the usual long pale hairs above and the black setæ beneath, hind tibiae blackened beyond the geniculation, hind tarsi piceous, the metatarsi not paler, apical joints of the anterior tarsi brownish. Wings with light infuscation, stigma very distinct, veins narrow but dark brown, sections of the fourth vein proportioned 0.7 : 0.2 : 1 : 1, of the fifth vein, 1 : 0.6, intercalary vein complete.

One specimen; Black Rock Mountain, Rabun County, Georgia, May, collected by J. Chester Bradley.

(2) *Œdalea lanceolata*, nov. sp. — Length 3 mm. Head, mesonotum and abdomen black, pleuræ and often the humeri yellowish. Antennæ black, the third joint slightly tapering, eight times as long as broad and twelve times as long as the thickened style; proboscis black, projecting nearly as far as the head-height. Thorax shining, mesonotal yellow hairs rather conspicuous. Legs yellow, the hind tibiae beyond the knee, the hind tarsi except the base and the anterior tarsi toward the tip brownish, subgenual carina of the hind tibiae black, the spinous setæ and setulae of the hind femora strong. Halteres pale yellow to brownish. Wings very lightly infumated, the elongate stigma darker, veins dark brown, discal cell widening distally so as to be less than two and a half times as long as wide, the last two sections of the fourth vein proportioned 1 : 0.8, of the fifth vein, 2 : 1, intercalary vein complete.

Numerous specimens; type from Friday Harbor, Washington, June 29, 1909; paratypes from Olga, Tacoma, Everett, Index, Tulalip, Lake Crescent, Pluvius, Ilwaco, Mt. Rainier, La Center, Lake Chelan and Glenwood, Washington; Collins, Moscow Mt. and Priest Lake, Idaho; Perma and Glacier Park, Montana, and Mt. Hood, Viento and Eagle Creek, Oregon.

Thorax entirely yellow, abdomen varying from yellow to brown;
hind tibiæ scarcely darkened var. *TESTACEA*, nov. var. (1).

Geographical distribution.

1. *OE. apicalis*, Loew, Neue Beitr. Dipt. Pt. 6, p. 47 (1859); Schiner, Fauna Dipt. Aust. Vol. 1, p. 81 (1862); Verrall, Ent. Mag. London, Vol. 48, p. 25 (1912). C. Europe.
2. *OE. astylata*, nov. sp. Georgia.
- . *OE. bracata*, Rondani, Dipt. Ital. Vol. 1, p. 152 (1857), no description. Spain.
3. *OE. brevicornis*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 26 (1899). N. and C. Europe.
4. *OE. flavipes*, Zetterstedt, Dipt. Scand. Vol. 1, p. 247 (1842), Vol. 8, p. 2998 (1849); Walker, Ins. Brit. Dipt. Vol. 1, p. 113, pl. 4, f. 4 (1851); Loew, Neue Beitr. Dipt. Vol. 6, p. 48 (1859); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 93 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 197, f. 75, 76, 78 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 78 (1910).
 - minuta*, Fallen (not Fabricius, Empid. Suec. p. 32 [1816]) (*Empis*); Meigen, Syst. Beschr. Vol. 2, p. 356 (1820); Zetterstedt, Fauna Ins. Lappon. p. 538 (1838); Dipt. Sc. Vol. 1, p. 247 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 488 (1849); Ins. Brit. Dipt. Vol. 1, p. 113 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 81 (1862).
 - rufipes*, Macquart, Hist. Nat. Dipt. Vol. 1, p. 357, pl. 8, f. 11 [1834] (*Xiphidicera*); Meigen, Syst. Beschr. Vol. 7, p. 101 [1838] (*Xiphidicera*); Blanchard, Hist. Nat. Ins. Vol. 3, p. 583 [1840] (*Xiphidicera*); Boitard, Man. Ent. Vol. 3, p. 325 [1843] (*Xiphidicera*).
 - ? *tibialis*, Macquart, Dipt. N. France, Vol. 3, p. 142, pl. 4, f. 2 (1827); Hist. Nat. Dipt. Vol. 1, p. 321 (1834); Meigen, Syst. Beschr. Vol. 7, p. 77 [1838] (*Edalia*); Blanchard, Hist. Nat. Ins. Vol. 3, p. 581 [1840] (*Edalea*); Boitard, Man. Ent. Vol. 3, p. 317 (1843); Loew, Neue Beitr. Dipt. Pt. 6, p. 47 (1859).
5. *OE. Holmgreni*, Zetterstedt, Dipt. Scand. Vol. 11, p. 4267 (1852); Loew, Neue Beitr. Dipt. Pt. 6, p. 49 (1859); Strobl, Mitteil. Naturw. Verein. Steiermark, Graz, Vol. 29, p. 93 (1892); Verrall, Ent. Mag. (2), Vol. 10, p. 141 (1894); Lundbeck, Dipt. Dan. Vol. 3, p. 199, f. 200 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 79 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37, p. 59 (1913). N. and C. Europe.
6. *OE. hybotina*, Fallen, Empid. Suec. p. 31 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 2, p. 356 (1820); Macquart, Dipt. N. France, Vol. 3, p. 142, pl. 4, f. 3 (1827); Hist. Nat. Dipt. Vol. 1, p. 321, pl. 7, f. 16 (1834); Zetterstedt, Fauna Ins. Lappon. p. 538 (1838); Dipt. Scand. Vol. 1, p. 245 (1842); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 49 (1851); Loew, Neue Beitr. Dipt. Pt. 6, p. 46 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 80 (1862); Neuhaus, Dipt. March. p. 69 (1886); Lundbeck, Dipt. Dan. Vol. 3, p. 198, f. 77 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 78 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 59 (1913). N. and C. Europe.
7. *OE. inermis*, Becker, Deutsche Ent. Zeitschr. 1910, p. 645 (1910). Corsica.
8. *OE. infuscata*, Loew, Neue Beitr. Dipt. Pt. 6, p. 48 (1859); Schiner, Fauna Dipt. Aust. Vol. 1, p. 82 (1862). C. Europe.
9. *OE. lanceolata*, nov. sp. Western North America.
var. *testacea*, nov. var. Washington.

(1) *Edalea lanceolata*, var. *testacea*, nov. var. — Thorax entirely yellowish, the abdomen brown rather than black, the dark color of the legs less evident.

Four males and eleven females, from Woodland, Seattle, Everett, Mt. Rainier, Pluvius, and Lilliwaup, Washington.

10. *OE. ohioensis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 256, 347, E. United States.
f. 74, 76 (1902).
stigmatella, Coquillett, det. in New Jersey List, p. 654 (1899).
11. *OE. pallipes*, Zetterstedt, Fauna Ins. Lappon. p. 538 (1838); Dipt. Scand. N. and C. Europe.
Vol. 1, p. 248 (1842). Vol. 8, p. 2998 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 49 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 4980 (1859); Schiner, Fauna Dipt. Aust. Vol. 1, p. 81 (1862); Wahlgren, Ent. Tidskr. Vol. 31, p. 79 (1910).
12. *OE. ? pennata*, Gimmerthal, Bull. Soc. Nat. Moscou, Vol. 15, p. 665 (1842). Russia. [States.
13. *OE. pruinosa*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 267 (1903). North-Eastern United
14. *OE. * robusta*, Meunier, Ann. Sci. Nat. (Zool.) Vol. 7, p. 91, 110, pl. 7, f. 10, 12 (1908). Baltic Amber, Lower Oligocene.
15. *OE. stigmatella*, Zetterstedt, Dipt. Scand. Vol. 1, p. 246 (1842), Vol. 8, p. 2998 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 49 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 112 (1851); Loew, Neue Beitr. Dipt. Pt. 6, p. 49 (1859); Schiner, Fauna Dipt. Aust. Vol. 1, p. 81 (1862); Siebke, Nyt. Mag. Naturvid. Vol. 14, p. 387 (1866); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 94 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 198, f. 79 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 78, f. 10 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 59 (1913). Europe.
- fulvipes*, Zetterstedt, olim in litt. Dipt. Scand. Vol. 1, p. 246 (1842).
stigmatica, Boheman, Svenska Vet. Akad. Handl. p. 190 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 81 (1862); Verrall, List Brit. Dipt. p. 15 (1888).
16. *OE. tristis*, Scholz, Zeitsch. Ent. Breslau, Vol. 5 (19), p. 49 (1851); Loew, C. and S. Europe.
Neue Beitr. Dipt. Pt. 6, p. 48 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 80 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 93 (1892); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 145 (1899). — Pl. I, Fig. 9.

6. GENUS LEPTOPEZA, MACQUART

Leptopeza, Macquart, Dipt. N. France, Vol. 3, p. 143 [1827] (*Lemtopeza*); Hist. Nat. Dipt. Vol. 1, p. 320 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 240 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 117 (1851); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 560 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 82 (1862); Lioy, Atti Inst. Ven. Sc. Venezia, 1864, p. 724 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 118 [1889] (*Lemtopeza*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Williston, Man. N. Amer. Dipt. p. 74. (1896); Melander, Trans. Amer. Ent. Vol. 28, p. 257 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252, 259 (1903); Bezzi, Ann. Mus. Hungar, Vol. 2, p. 321 (1904); Melander, Williston Man. N. Amer. Dipt 3 ed. p. 225 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301 (1909); Kertesz, Cat. Dipt. Vol. 6, p. 107 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 558 [1910] (*Lemtopeza*); Lundbeck, Dipt. Dan. Vol. 3, p. 204 (1910); Wahlgren, Ent. Tidsk. Vol. 31, p. 43, 79 (1910); Bezzi, Ann. Mus. Hungar, Vol. 10, p. 453 (1912); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 362 (1920).

Characters. — Slender elongate black or yellow species, variable in color, but fixed in structure. Head globose, eyes large, contiguous above and below the antennæ in both male and female, the lower part of the face visible as a linear separation of the eyes, facets uniform or the uppermost large; antennæ inserted at the middle of the head, extending straight forward, two-jointed, that is, the basal joints are

entirely connate, the last joint elongate conical, with a very slender elongate bare single-jointed terminal arista; mouthparts very short, haustellate, the palpi scarcely projecting and hairy; ocellar triangle small, ocellar bristles reaching halfway to the antennæ, two pairs of vertical bristles of similar length. Thorax less than half as long as the abdomen and broader than the head, with very fine seriate pubescence, lateral bristles long but hair-like, an ascending row of three notopleurals, one supraalar, one postalar, one prescutellar and several scutellars present. Abdomen cylindrical, pygidium small, distorted, asymmetrical, the valves with slender ribbon-like processes; terminal segments of the female usually prolonged to an ensiform chitinized ovipositor. Legs slender, hairy, coxæ short, the front coxæ slightly the longest, femora without bristles, middle tibiae with several sets of long bristles. Wings elongate, anal angle prominent, rounded rectangular, basal bristle of the costa present, costa stopping beyond the third vein, pubescence of the wing-membrane coarse, auxiliary vein rudimentary, lying close to the first vein, which terminates at the outer fourth of the wing, third vein simple, its pedicel long, discal cell emitting two unforked veins which are the fifth and sixth longitudinals, only a spur of the fourth vein sometimes distinguishable beyond the discal cell, basal cells large, the first basal narrower than the second, anal cell shorter than the second basal, obliquely truncate by its reflexed straight crossvein.

Type species: *L. flavipes*, Meigen (Pl. 8, Fig. 19), the only species given in the original description, and which is synonymous with *ruficollis* Meigen. The habits of the species of this genus are unknown, except that the adults occur in open and rather shady spots in woods near streams or lakes. Most of the species, like *ruficollis*, *borealis*, *nigripes*, *disparilis* and *compta* are not clearly limited, and often specimens can be found on the borderland between species, making identification difficult at times. This is due to the variable color within the species and the stereotyped habitus of the genus. *Sphenoptera*, however, departs from the others and several times it has been suggested that this species be removed from *Leptopeza*. It is peculiar in several characters, and in many ways is more closely related to the South American *Hoplopeza* than to *Leptopeza*. This species has here been removed under the generic name *Leptometopiella*.

TABLE FOR THE SEPARATION OF THE NEARCTIC SPECIES OF LEPTOPEZA

1. Thorax devoid of pubescence, lightly pollinose, scutellum with only two bristles; antennæ elongate, arista white, outer joint six times as long as broad and three times as long as the inner joint; no ovipositor; wings strongly infumated; mouthparts black. L. ANTENNALIS, nov. sp. (1).

Thorax shining, with short but evident pubescence, scutellum with more than two bristles; antennæ much shorter, the arista black; a long ensiform ovipositor developed; wings clear. 2.

2. Mouthparts black; antennæ black; hind femora and tibiæ largely black, especially in the male 3.

(1) ***Leptopeza antennalis*, nov. sp.** — Length 3.5 mm. Black, thorax almost devoid of pubescence, lightly pollinose, scutellum with only two bristles; antennæ elongate, the outer joint six times as long as wide and three times as long as the basal joint, arista yellow; no ovipositor. Upper facets of male large, occiput with brownish pollen, mouthparts black, arista about one-third longer than the black antennæ, the outer antennal joint narrow and tapering. Hairs of the mesonotum and scutellum very sparse and microscopic. Abdomen shining, hairs of the first ventral long dense and pale, pygidium small but open, the middle valves strongly forcipate. Coxæ yellow, base of legs yellow, becoming progressively darker distally to the piceous tarsi, the hind legs darkest. Halteres yellow (♀) or with black knob (♂). Wings strongly infumated, no stigma.

Six specimens; Washington, D. C., August 17, 1913, taken by the author near Chain Bridge on the Potomac River. It is strange that this very distinct species was not discovered earlier by the many entomologists who have collected in this locality.

- Mouthparts yellowish, particularly the palpi 4.
3. Dorsal valve of pygidium concave behind; arista about twice the length of last antennal joint; 3 mm. L. BOREALIS, Zetterstedt.
- Dorsal valve with straight hind edge; arista more than twice as long as third joint; 4 mm. L. DISPARILIS, Melander.
4. Antennæ entirely black L. RUFICOLLIS, Meigen.
- Antennæ yellow, the basal joint sometimes black; body varying from yellow to black. L. COMPTA, Coquillett.

Geographical distribution.

1. *L. antennalis*, nov. sp. District of Columbia.
2. *L. bimaculata*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 337 (1904); White, Proc. Roy. Soc. Tasmania, 1916, p. 242 (1917). New South Wales, Tasmania.
3. *L. biplagiata*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 473 (1912). Formosa
4. *L. borealis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 243 (1842), Vol. 8, p. 2997 (1849); Wahlgren, Ent. Tidskr. Vol. 31, p. 80 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 60 (1913). N. Europe, E. United States.
- flavipes*, Zetterstedt (not Meigen), Fauna Ins. Lappon. p. 537, part [1838] (*Ocydromia*). Lower Oligocene, Baltic Amber.
5. *L. *clavigipes*, Loew, Bernstein Fauna, Vol. 1, p. 41 (1850); Giebel, Ins. Vorwelt, p. 208 (1856). United States.
6. *L. compta*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 435 (1895); Melander, Trans. Am. Ent. Soc. Vol. 28, p. 258, f. 70 (1902). Baltic Amber.
7. *L. *concinna*, Meunier, Ann. Sci. Nat. (Zool.) Vol. 7, p. 91, 110, pl. 7, f. 13, 14 (1908). United States.
8. *L. disparilis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 258, f. 69 (1902). N. Europe.
9. *L. flavimana*, Zetterstedt, Dipt. Scand. Vol. 1, p. 244 (1842), Vol. 8, p. 2997 (1849). Java.
10. *L. javana*, Meijere, Tijdschr. v. Ent. Vol. 56, suppl. 71 (1913). Tasmania.
11. *L. levicosta*, White, Proc. Roy. Soc. Tasmania, 1916, p. 244, f. 48 (1917). N. Europe.
12. *L. nigripes*, Zetterstedt, Dipt. Scand. Vol. 1, p. 44 (1842). New South Wales, Tasmania.
13. *L. pulcherrima*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 336 (1904); White, Proc. Roy. Soc. Tasmania, 1916, p. 243 (1917). Cape Horn.
14. *L. rivosa*, Bigot, Miss. Scient. Cap. Horn. (Zool.) Vol. 6, p. 21 [1888] (*Lemtopeza*); Bezzi, Ann. Mus. Hungar, Vol. 3, p. 458 (1905). Tasmania.
15. *L. rubrithorax*, White, Proc. Roy. Soc. Tasmania, 1916, p. 243 (1917). Europe, North America.
16. *L. ruficollis*, Meigen, Syst. Beschr. Vol. 2, p. 353, pl. 21, f. 24 [1820] (*Ocydromia*), Vol. 6, p. 334 [1830] (*Ocydromia*); Loew, Bemerk. Posen Gegend. Art. Zweifl. Gatt. p. 19 (1840); Isis, Vol. 7, p. 554 [1840] (*Ocydromia*); Stæger, Naturhist. Tidskr. Vol. 4, p. 100 (1842); Zetterstedt, Dipt. Scand. Vol. 1, p. 241 (1842); Vol. 8, p. 2996 (1851); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (17), p. 48 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 117, pl. 4, f. 7 (1851); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 94 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 79 (1910). — Pl. 8, Fig. 79.
- flavipes*, Meigen, Syst. Beschr. Vol. 2, p. 353 [1820] (*Ocydromia*); Macquart, Mém. Soc. Sc. Lille, p. 147 [1823] (*Ocydromia*); Ins. Dipt. France, Vol. 3, p. 144, pl. 4, f. 4 [1827] (*Lemtopeza*); Hist. Nat. Dipt. Vol. 1, p. 321, pl. 7, f. 15 (1834); Zetterstedt, Fauna Ins. Lappon. p. 537, part [1838] (*Ocydromia*); Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt. p. 19 (1840); Isis, Vol. 7, p. 554 [1840] (*Ocydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 242 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 488 [1849] (*Ocydromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 2997 (1849); Dahlbom, Svenska, Vet. Akad. Handl.

- p. 160 (1850); Scholz, Ent. Breslau, Vol. 5 (17), p. 48 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 117 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 4267 (1852), Vol. 12, p. 4597 (1855), Vol. 13, p. 4979 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 82 (1862); Raddatz, Arch. Freund. Naturg. Mecklenb. Rostock, Vol. 27, p. 42 (1873); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 94 (1892); Coquillet, Proc. U. S. Nat. Mus. Vol. 18, p. 435 (1895); Strobl, Wien. Ent. Zeit. Vol. 18, p. 27 (1899); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 258, f. 68 (1902); Lundbeck, Dipt. Dan. Vol. 3, p. 206, f. 83, 84 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 79 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 60 (1913).
- tibialis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 242 (1842). Vol. 8, p. 2997 (1849); Bonßdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 143 (1861); Siebke, Nyt. Mag. Naturvid. Vol. 14, p. 380 (1866).
- var. *unicolor*, Strobl, Jahrb. Naturhist. Landesmus. Kärnten, Klagenfurt, Vol. 47, p. 202 (1901).
17. *L. rugosiventris*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, C. Europe. p. 73 (1909).
18. *L. serraticosta*, White, Proc. Roy. Soc. Tasmania, 1916, p. 245 (1917). Tasmania.
19. *L. setigera*, Bezzi, Bul. Soc. Ent. Ital. Vol. 32, p. 82 (1900). S. Europe.
- flavipes*, Bezzi (not Meigen), Bul. Soc. Ent. Ital. Vol. 30, p. 145 (1899).
20. *L. *spinigera*, Loew, Bernstein Fauna, p. 41 (1850); Meunier, Miscell. Ent. Vol. 7, p. 178 (1899). Lower Oligocene,
Baltic Amber.
21. *L. tachydromiaeformis*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 338 (1904). New South Wales.
22. *L. vitripennis*, Brunetti, Rec. Indian. Mus. Vol. 9, p. 31 (1913); Fauna Brit. Ind. Dipt. Vol. 1, p. 363, pl. 4, f. 12, 13 (1920). India.

7. GENUS OCYDROMIA, MEIGEN

Ocydromia, Meigen, Syst. Beschr. Vol. 2, p. 351 (1820); Macquart, Mém. Soc. Sc. Lille, 1823, p. 145 (1823); Dipt. N. France, Vol. 3, p. 144 (1827); Hist. Nat. Dipt. Vol. 1, p. 319 (1834); Zetterstedt, Fauna Ins. Lappon, p. 535 (1838); Blanchard, Hist. Nat. Ins. Vol. 3, p. 581 [1840] (*Ocydromya*); Westwood, Gen. Syn. p. 133 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 236 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 118 (1851); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 560 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 81 (1862); Lioy, Atti Instit. Venet. 1864, p. 723 (1864); Beling, Arch. Naturg. Berlin, Vol. 48, p. 239 (1882); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 117 [1889] (*Ocydromya*); Coquillet, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Williston, Man. N. Amer. Dipt. 3 ed., p. 74 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 259 (1902); Coquillet, Proc. Ent. Soc. Wash. Vol. 5, p. 254, 259 (1903); Melander, Williston, Man. N. Amer. Dipt. 3 ed. p. 225 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301 (1909); Kertész, Cat. Dipt. Vol. 6, p. 105 (1909); Coquillet, Proc. U. S. Nat. Mus. Vol. 37, p. 577 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 200 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 43, 79 (1910).

Characters. -- Slender, shining, blackish or yellow species about three or four millimeters in length. Head globular, eyes contiguous above the antennae and subcontiguous on the face, facets uniformly small; antennae inserted at the middle of the head, the basal joints small and fused, the third joint oval and with a sub-lorsal long slender hair-like arista; proboscis very short, fleshy, scarcely protruding from the oral opening; ocellar triangle not prominent, ocellar bristles small. Thorax short but rather highly arched, devoid of pollen and stiff bristles, its hairs very sparse; scutellum villose and margined with several bristles; pleurae shining and bare. Abdomen twice as long as the thorax, slender, cylindrical, pygidium minute, asymmetrical, abdomen of the female with blunt termination, no

ovipositor. Legs simple, slender, hairy but without bristles. Calypteres with a prominent fringe. Wings rather large, the anal angle broadly rectangular, costa greatly attenuated beyond the third vein, auxiliary vein very close to the first vein into which it ends, third vein simple, its pedicel arising before the middle of the basal cells, basal cells long, coextensive, nearly as long as the discal cell and longer than the anal cell, the discal cell emitting two veins which are the intercalary and the fifth longitudinal, at most a stump of the fourth vein projecting from the discal cell, anal crossvein reflexed and straight, forming an angle of about one hundred degrees with the anal vein, the anal vein weak but extending to the margin.

Type species : *C. glabricula*, Fallen (Pl. I, Fig. 10), Westwood's designation, a species widely distributed over the Northern Hemisphere. The larva and pupa of this species have been described by Beling in the reference above given. The adults occur in shady woodlands where they fly feebly about the vegetation.

Geographical distribution.

- . *O. ? cothurnata*, Bigot, in litt.; Brunetti, Rec. Indian. Mus. Vol. 9, p. 31 (1913).
- 1. *O. elegans*, Bigot, Miss. Scient. Cap. Horn, Vol. 6, p. 21, pl. 3, f. 5 [1888] Cape Horn. (*Ocydromyia*); Bezzii, Ann. Mus. Hungar., Vol. 3, p. 458, note 4 [1905] (? *Euthyneura*).
- 2. *O. fuscipennis*, Macquart, Mém. Soc. Sc. Lille, p. 147 (1823). France.
- 3. *O. glabricula*, Fallen, Empid. Suec. p. 33 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 2, p. 352, pl. 21, f. 23 (1820); Macquart, Mém. Soc. Sc. Lille, p. 146 (1823); Dipt. N. France, Vol. 3, p. 145, pl. 4, f. 5 (1827); Hist. Nat. Dipt. Vol. 1, p. 320, pl. 7, f. 14 (1834); Zetterstedt, Fauna Ins. Lappon. p. 536 (1838); Blanchard, Hist. Nat. Ins. Vol. 3, p. 581 (1840); Stæger, Naturh. Tidskr. Vol. 4, p. 101 (1842); Zetterstedt, Dipt. Scand. Vol. 1, p. 236 (1842); Boitard, Ent. Man. Vol. 3, p. 316 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 487 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 2995 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 48 (1851); Walker, Ins. Brit. Vol. 1, p. 118, pl. 4, f. 8 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 143 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 82 (1862); Beling, Archiv. Naturg. Berlin, Vol. 48 (1), p. 213 (1882); Neuhaus, Dipt. March. p. 69 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 133 (1887); Verrall, The Entom. Vol. 23, p. 153 (1890); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 94 (1892); Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 421 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 259, f. 66, 67 (1902); Czizek, Zeitschr. Mähr. Landesmus. Brünn, Vol. 7, p. 166 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 202, f. 81, 82 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 79, f. 11 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 59 (1913). — Pl. I, Fig. 10.
- ? *coxalis*, Roser, Correspondenzbl. Landw. Ver. Würtenb. Stuttgart, Vol. 1, p. 53 (1840).
- dispar*, Curtis, Brit. Ent. Vol. 8, p. 517 [1834] (*Rhamphomyia*).
- dorsalis*, Meigen, Syst. Beschr. Vol. 6, p. 334 (1830).
- melanopleura*, Loew, Bemerk. Posen, Gegend. Art. Zweifl. Gatt. p. 19 (1840); Isis, Vol. 7, p. 545 (1840); Czizek, Zeitsch. Mähr. Landesmus., Brünn, Vol. 7, p. 166 (1907).
- nigripennis*, Meigen (not Fabricius), Syst. Beschr. Vol. 6, p. 334 (1830); Macquart, Hist. Nat. Dipt. Vol. 1, p. 320 (1834); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 488 (1849);

- Scholz, Zeitschr. Ent. Breslau, Vol. 5 (17), p. 48 (1851); Neuhaus, Dipt. March. p. 69 (1886).
- peregrinata*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 488 (1849).
- ruficollis*, Macquart (not Meigen), Mém. Soc. Sc. Lille, p. 147 (1823); Dipt. France, Vol. 3, p. 146 (1827); Hist. Nat. Dipt. Vol. 1, p. 320 (1834); Boitard, Man. Ent. Vol. 3, p. 316 (1843).
- rufipes*, Meigen, Syst. Beschr. Vol. 2, p. 353 (1820); Zetterstedt, Fauna Ins. Lappon. p. 536 (1838); Stæger, Naturh. Tidskr. Vol. 4, p. 100 (1842); Zetterstedt, Dipt. Scand. Vol. 1, p. 239 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 487 (1849); Zetterstedt, Dipt. Sc. Vol. 8, p. 2996 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (17), p. 48 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 119 (1851); Pipping, Not. Sällsk. Fenn. Förh. Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 143 (1861); Siebke, Nyt. Mag. Naturvid. Vol. 14, p. 399 (1866); Neuhaus, Dipt. March. p. 69 (1886).
- scutellata*, Meigen, Syst. Beichr. Vol. 2, p. 354 (1820); Macquart, Hist. Nat. Dipt. Vol. 1, p. 320 (1834); Zetterstedt, Fauna Ins. Lappon. p. 537 (1838); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 19 (1840); Isis, Vol. 7, p. 554 (1840); Zetterstedt, Dipt. Sc. Vol. 1, p. 237 (1842); Boitard, Man. Ent. Vol. 3, p. 316 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 487 (1849); Zetterstedt, Dipt. Sc. Vol. 8, p. 2995 (1849); Walker, Ins. Brit. Dipt. Vol. 1, p. 119 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 4978 (1859); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 143 (1861); Czizek, Zeitschr. Mähr. Landesmus. Brünn, Vol. 7, p. 166 (1907).
4. *O. hirsutipes*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 123 (1914). E. Africa.
5. *O. Philippii*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 128 (1889); Chile.
- Bezzi, Ann. Mus. Hungar., Vol. 3, p. 458 (1905).

8. GENUS LEPTOMETOPIELLA, NOV. GEN.

Characters. — Eyes separated on the front in both sexes, the lowermost facets largest; antennæ three-jointed, inserted above the middle of the head; discal cell long, anal crossvein interrupted beyond its middle, anal vein reduced to a fold, anal angle of the wing scarcely developed, the wings accordingly cuneiform; notopleural bristles reduced; lateral valves of the pygidium greatly convex. Other characters as described for *Leptopeza*.

Type species: *L. sphenoptera*, Loew. The shortened anal crossvein is found also in *Hoplopeza* and is more or less indicated in true Leptopezas, where the anal vein is weakened and the crossvein becomes attenuated.

Geographical distribution.

1. *L. sphenoptera*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 215 (1873) ♀; Strobl, C. and S. Europe, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 94 (1892) ♂, N. Africa. Vol. 34, p. 207 (1898); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 119 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 208, f. 85 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 26 (1912).
- lonchoptera*, Pokorný, Verh. Zool. bot. Ges. Wien, Vol. 37, p. 394 (1887).

9. GENUS HOPOPEZA, BEZZI

Hoplopeza, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301, 385 (1909).

Characters. — Black shining species, closely related to *Leptopeza*. Head round, occiput with few short bristles, ocellar bristles small; eyes contiguous above and below the antennæ in the male (the female unknown), bare; antennæ three-jointed, the last joint elongate, bearing a thickened pilose termi-

nal arista as long as the antenna; proboscis short and thick, slightly projecting, palpi small and rounded. Thorax but little arched, without discal bristles, two scutellars; pleuræ pollinose except the sternopleura. Abdomen long and cylindrical; pygidium small and rounded, beneath with six long bristles. Hind femora with a knee bristle, front tibiæ fimbriate within with hairs, hind tibiæ with some long apical spines, the lowermost curved, hind metatarsi spinulose above. Wings cuneiform, no anal angle, no stigma, pedicel of the second and third veins short, discal cell long, emitting two posterior veins, anal cell very short, the anal vein reflexed and abortive, anal vein complete.

Type species: *H. chloropa*, Bezzi, the original species.

Geographical distribution.

1. *H. chloropa*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 385, f. 10-12 (1909).

10. GENUS SCELOLABES, PHILIPPI

Scelolabes, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 751 (1865); Reed, Ann. Univ. Chile, Vol. 78, p. 27 [1888] (*Scelobates*); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 116 (1889); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 301 (1909); Kertész, Cat. Dipt. Vol. 6, p. 104 (1909).

Characters. — Polished, testaceous species with raptorial hind legs. Head globose, eyes very narrowly separated on the front, face linear, basal joints of the antennæ short and subcylindrical, the third joint pubescent, long-conical, about twice as long as the basal two and terminated by a pubescent long slender arista; proboscis short and projecting obliquely forward, palpi ovate; ocellar triangle not elevated. Thorax rather stout and strongly arched, almost bare of bristles and hairs, scutellum with two long apical bristles, one long prealar black bristle present. Abdomen subcylindrical. Legs pubescent and bristly, anterior legs simple, hind femora much swollen, with numerous fine setulæ and setæ beneath and with irregular macrochætæ, hind tibiæ geniculate and much shorter than their femora, middle tibiæ with four extensor bristles along basal half, pulvilli large. Wings with a rounded anal angle, costa stopping at the fourth vein, auxiliary vein very close to the first vein, first vein ending much beyond the middle of the wing, opposite end of discal cell, third vein simple, pedicel of the second and third veins very short, arising near the end of the basal cells, the latter longer than the discal cell, nearly coextensive, the vein limiting the second basal cell oblique, discal cell emitting two veins, the fourth vein obsolete, anal crossvein inflexed, anal vein evanescent.

Genotype and only species: *S. bivittatus*, Philippi.

Geographical distribution.

1. *S. bivittatus*, Philippi, Verh. Zool. Bot. Ges. Wien, Vol. 15, p. 751, pl. 28, Chile.
f. 45 (1865); Bezzi, Ann. Mus. Hungar., Vol. 3, p. 458 (1905). —

Pl. 5, Fig. 43.

11. GENUS LAMACHELLA, NOV. GEN.

Characters. — Polished, testaceous species with raptorial hind legs. Head globose, eyes of female separated on front and face, basal joints of antennæ short and cylindrical, third joint lanceolate, arista slender, pubescent, shorter than third joint; proboscis projecting obliquely forward, piercing, palpi slender, long, porrect; ocellar triangle elevated. Thorax strongly arched, almost bare, two small black prealar and two small scutellar bristles present. Abdomen subconical, terminal segments of

female decreasing in size but not forming an ovipositor. Legs pubescent, anterior legs simple, devoid of bristles, hind femora much swollen, with numerous irregular thorn-like setulae and setae beneath, hind tibiae shortened and geniculate, pulvilli large. Wings with prominent anal angle, costa evanescent beyond fourth vein, auxiliary vein very close to the first vein with which it combines, first vein ending much beyond the middle of the wing, third vein simple, pedicel of second and third veins very short, arising near the middle of the basal cells, discal cell emitting two veins, the fourth and fifth, second basal cell slightly longer than the first, its apical vein nearly transverse, anal crossvein perpendicular to the weak but complete anal vein.

Genotype : *L. univittata*, the following species.

Geographical distribution.

1. *L. univittata*, nov. sp. (1).

Kamerun.

12. GENUS STENOPROCTUS, LOEW

Stenoproctus, Loew, Oefv. Vet. Akad. Förh. Vol. 15, p. 340 (1858); Dipterenf. Südafr. Vol. 1, p. 261 (1860); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 116 (1889); Kertész, Cat. Dipt. Vol. 6, p. 12, (1909).

Acanthopeza, Becker, Ann. Soc. Ent. France, Vol. 83, p. 122 (1914).

Characters. — Black shining species measuring three millimeters, with hyaline wings and yellowish legs. Head globose, eyes of the female completely contiguous on the front; third joint of the antennæ cylindrico-conical, twice as long as the combined basal joints, ending in a straight slender, nearly bare arista which is as long as the third joint and is tipped with a bristle; proboscis projecting horizontally forward, nearly as far as the head-height, palpi linear. Thorax rather stout and highly arched, shorter than the abdomen, two scutellar bristles present. Hind legs lengthened, their femora incrassate and spinose beneath, hind tibiae simple and curved, anterior legs simple. Wings with a prominent anal angle, first vein ending beyond the middle of the wing, a pronounced stigma present, second vein broadly sinuous but not steeply curving into the costa, pedicel of the second and third veins long, arising toward the base of the basal cells, discal cell small, emitting two posterior veins, basal cells long, the anterior much longer and narrower than the posterior, anal crossvein perpendicular to the anal vein which attains the margin.

Type species : *S. unipunctatus*, Loew (Pl. 5, Fig. 44), the original genotype.

(1) **Lamachella univittata**, nov. sp. — Female. Length 3 mm. Head black, antennæ and mouthparts yellow, thorax except for a median vitta yellow, abdomen black, legs yellow, halteres blackish. Proboscis one-half as long as the head-height, porrect, constructed for piercing, palpi of the same length, spatulate and porrect. Median vitta of the thorax blackish-brown, not sharply defined, scutellum and metanotum blackish, a small humeral bristle present. Pubescence of the legs black, front tibiae somewhat inflated, a little thicker than the middle pair, all the tibiae with some projecting extensor hairs among the incumbent pubescence, middle tibiae with a small yellow apical flexor bristle, hind femora much swollen, thickest at the middle, studded with numerous setulae beneath and with a row of setae along the posterior edge, hind tibiae three-fourths as long as their femora, tarsi not thickened. Wings hyaline, veins brown, an elongate stigma visible, first four sections of the costa proportioned 1 : 0.3 ; 0.3 : 0.15, basal cells of equal width, second basal longer than the first because of the obliquity of its crossvein, sections of the fourth vein, 1 : 1.1, of the fifth vein, 1 : 0.8.

One specimen, in the Museum of Comparative Zoology, Cambridge, Mass., received through Professor Roland Thaxter, Kamerun, Africa.

Geographical distribution.

1. *St. sylvaticus*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 123 [1914] E. Africa.
(*Acanthopeza*).
2. *St. unipunctatus*, Loew, Oefv. Vet. Akad. Förh. Vol. 15, p. 340 (1856); Cape of Good Hope.
Dipterenf. Südafr. p. 261, f. 45 (1860). — PI. 5, Fig. 44.

13. GENUS BICELLARIA, MACQUART

Bicellaria, Macquart, Mém. Soc. Sc. Lille, 1823, p. 155 (1823); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 122 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 246, 261 (1903); Melander, Williston Man. N. Amer. Dipt. 3 ed. p. 225 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 300 (1909); Kertész, Cat. Dipt. Vol. 6, p. 13 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 514 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 20 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 43, 48 (1910).

Cyrtoma, Meigen, Syst. Beschr. Vol. 4, p. 1 (1824); Macquart, Dipt. N. France, Vol 3, p. 106 (1827); Hist. Nat. Dipt. Vol. 1, p. 359 (1834); Zetterstedt, Fauna Ins. Lappon. p. 533 (1838); Westwood, Gen. Syn. p. 133 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 329 (1842); Boitard, Man. Ent. Vol. 3, p. 326 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 115 (1851); Rondani, Dipt. Ital. Vol. 1, p. 152 (1855); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 564 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 76 (1862); Lioy, Atti Inst. Ven. 1864, p. 722 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 123 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Williston, Man. N. Amer. Dipt. p. 74 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 335 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 530 (1910).

Characters. — Rather small, greatly humpbacked, more or less opaque black species with no discal cell. Head globular, attached low to the thorax; eyes broadly contiguous above the antennæ in both sexes, bare, the upper facets large and separated from the lower by a horizontal line of demarcation; face long and narrow, no cheeks; occiput hairy but not bristly; antennæ located at the middle of the head, two-jointed, due to the fusion of the basal joints, the third joint elongate, somewhat compressed, usually somewhat bulbous below near the base, ending in a two-jointed thickened arista which is more or less shorter than the third joint and consists of a short basal and a lengthened apical segment; proboscis much shorter than the head, vertical, the labrum slightly incurved and pointed, palpi short, broad and decumbent. Thorax very large, provided with a greater or less number of fine bristles, posthumeral and lateral hairs usually quite evident, scutellum with about four bristles, acrostichals loosely biseriate; pleuræ bare. Abdomen slender, tapering, pygidium terminal, small and closed, the lateral valves oval and directed backward, ventral piece cleft, ending in a symmetrical two-tined fork. Legs rather long, hairy, often with long fine bristles, often the hind femora ciliate, the hind legs longest, usually with clavate tibiæ, hind metatarsi sometimes thickened. Anal angle of the wings rectangular, a distinct angulation at the axilla, costa stopping at the end of the third vein, basal bristles present or absent, auxiliary vein lying close to the first vein, third vein simple, anterior crossvein placed close to the fork of the second and third veins, no discal cell, the fourth vein interrupted at the anterior crossvein resuming again toward the tip of the wing as a weakened spur, the intercalary vein likewise weak and evanescent basally, second basal cell longer than the first, its crossvein very oblique, almost in line with the last section of the fifth vein, anal cell wider than the first basal, its crossvein reflexed and broadly angulate with the continuous anal vein.

Type species : *B. spuria*, Fallen (Pl. 5, Fig. 46), redescribed by Macquart as *nigra*, nov. sp. The species of *Bicellaria* are common in meadowlands during the summer month, especially to the North.

TABLE OF THE SPECIES OF BICELLARIA

1. Body brownish in color	2.
Ground-color of body and legs black	3.
2. Legs and halteres yellow	<i>B. RUFÀ</i> , Meigen.
Legs and halteres brown ; tines of pygidial ventral fork thick and truncate	<i>B. UVENS</i> , nov. sp. (1).
3. Hind tibiæ uniformly slender ; mesonotum of male velvety black and very gibbous ; wings of female hyaline.	<i>B. MELÆNA</i> , Haliday.
Hind tibiæ more or less thickened toward apex	4.
4. Hairs of forward part of notum almost as long as those behind, two vittæ more or less indicated	5.
Hairs of forward part of mesonotum short, no vittæ	6.
5. Abdomen dark gray, dull or slightly shining.	<i>B. PILORA</i> , Lundbeck.
Abdomen satiny whitish in certain lights	<i>B. ALPINA</i> , Bezzii.
6. Hind metatarsi not at all thickened, legs rather short, twelve hairs in postero-dorsal row of hind femora ; pygidial fork narrowly U-shaped, tines moderately thick and tipped with a couple of microscopic hairs	<i>B. SPURIA</i> , Fallen.
Hind metatarsi more or less thickened	7.
7. Basal joints of hind tarsi but slightly thickened	8.
Basal joints of hind tarsi of male distinctly thickened ; abdominal hairs black or brown ; legs long, eighteen very long hairs on antero-dorsal row of hind femora	16.
8. Halteres of male and female pale yellow; wings subfuscous; hairs of legs pale ; base of third antennal joint broadly ovate, arista nearly as long as the third joint ; ventral fork of pygidium U-shaped	<i>B. HALTERALIS</i> , Loew.
Halteres of male dark, of female either dark or pale ; arista usually much shorter than third antennal joint	9.
9. Wings white-hyaline, wide in female ; base of third antennal joint broadly oval	<i>B. DISPAR</i> , Oldenberg.
Wings more or less infumated	10.

(1) *Bicellaria uvens*, nov. sp. — Male. Length 3 mm. Blackish-brown, third antennal joint with rounded base and gradually tapering to apex, heavily pubescent ; arista equal to third joint ; occipital hairs black. Thorax completely covered with dark brown pollen, bristles black. Abdomen dull, hairs black, pygidium dusky, the ventral fork deeply excised with the prongs robust and wider than the excision. Legs normal, hairs black, about eighteen in the anterior dorsal row of the hind femora, hind tibiæ gradually clavate with many outstanding extensor setæ, hind metatarsi scarcely thicker than other joints. Halteres black, wings heavily infumated, stigma not differentiated.

Female. Last abdominal segments shining, halteres brown.

Numerous specimens from Fort Chimo, Ungava Bay, Hudson Straits, near northern Labrador (L. M. Turner), type in U. S. National Museum. The specimens appear to have been preserved in liquid which may account for the castaneous color of most of the specimens.

10. Tines of ventral fork of pygidium flattened	III.
Tines of pygidial fork more or less cylindrical	12.
11. Pygidial fork widely and shallowly excised, the tines short triangular; abdominal hairs pale; bristles of hind tibiæ fine	B. BREVIFURCA, nov. sp. (1).
Pygidial fork narrowly and deeply excised, the tines long and blunt; abdominal hairs of male black; at least four outstanding bristles on hind tibiæ	B. PILIPES, Loew.
12. Tines of pygidial fork very long and thin	13.
Tines of pygidial fork shorter and stouter	14.
13. Pubescence black; third antennal joint more than twice as long as wide, its elongate base tapering apically	B. ANGUSTIFURCA, nov. sp. (2).
Hairs of abdomen and femora for most part pale yellow; third antennal joint less than twice as long as wide, its ovate base distinct from the tubular apex	B. FURCIFER, nov. sp. (3).
14. Pygidial fork narrowly U-shaped, its tines tipped with microscopic hair (if tines are stubby, see <i>uvens</i>)	15.
Pygidial fork broadly U-shaped, its tines tipped with a long hair; halteres of both sexes dark; base of third antennal joint ovate, apex tubular	B. LONGIPES, Loew.

(1) **Bicellaria brevifurca**, nov. sp. — Length 3 mm. Black with faint brown coating, hairs and bristles black, the hairs of the abdomen pale. Facets of eyes moderately large above, about sixteen along the line of contiguity; under side of third antennal joint gently rounding into the apex, arista about two-thirds as long as this joint. Notum velvety brown, the anterior hairs short, four scutellars. Ventral fork of pygidium flat, M-shaped in outline. Hind femora rather short-ciliate, eighteen or more setæ in anterior extensor row and twenty or more in anterior flexor row; hind tibiæ subclavate, about twice as broad at end as at base, extensor bristles fine and uniform; hind metatarsi slightly thicker than the following joints. Halteres black. Wings infumated in male, grayish in female, stigma brown, fork of fourth vein usually open, the pedicel shorter than the fork.

About one hundred specimens, alpine parks of Mount Rainier, Washington, July to September.

(2) **Bicellaria angustifurca**, nov. sp. — Male. Length 3.2 mm. Very black, opaque; third antennal joint elongate and large, two times as long as deep, tapering from the rounded base to the apex without tubular tip, arista two-thirds the length of this joint. Hairs and bristles entirely black; pygidium small, ventral fork polished, very deeply cleft, the tines unusually long, slender, parallel on basal half and then diverging. Setæ of both upper and lower surface of hind femora very long and abundant, about twenty-one in upper anterior row; hind tibiæ gradually but decidedly clavate the extensor setæ uniform and moderate, hind metatarsi but slightly thickened. Halteres thickened, wings very strongly infumated almost blackish brown, the stigma scarcely differentiated.

Female. Not opaque black but thinly coated with brown dust on notum. Abdomen subshining, pleuræ cinereous, abdominal hairs brownish, last segment shining.

Types, Olympia, Washington, 22 June, 1920; six paratypes; Bellingham, Washington, Mount Hood, Oregon, Priest Lake and Moscow, Idaho (Melander).

(3) **Bicellaria furcifer**, nov. sp. — Length 2.75 mm. Black with faint brown coating, hairs and bristles generally black, those of lower occiput, abdomen and coxæ and base of femora pale. Facets moderately large above, about sixteen in line of contiguity of the eyes; third antennal joint about twice as long as deep, the ovate base distinct from the tubular apex, arista about half the length of the third joint. Notum subshining, the hairs short in front, four brownish scutellar bristles; hairs of abdomen long. Tines of pygidial fork very long and thin, tipped with a microscopic hair. About fifteen setæ in upper and lower rows on hind femora; hind tibiæ subclavate, about three times as wide toward apex as at base, with two or three of the extensor hairs longer than the others; hind metatarsi scarcely thicker than the next joint. Halteres of male black, of female yellowish brown; hairs of calypteres brownish. Wings grayish hyaline, stigma brown, fork of fourth vein usually complete and shorter than its pedicel.

Type and allotype, Upper St. Regis River, Montana, 28 July, 1918. Nine paratypes, Yellowstone Park, several places, Wyoming; Jefferson Island and Perma, Montana; Craig's Mountain, Idaho; and Nelson, British Columbia, all collected by the author.

15. Base of third antennal joint elliptical; halteres of female pale;
middle tibiae with paired bristles at basal fifth and two-fifths;
four scutellars B. INTERMEDIA, Lundbeck.
- Base of third antennal joint quadrate; bristles of middle tibiae
smaller; six or eight scutellars B. LUGUBRIS, nov. sp. (1)
16. Tines of the U-shaped pygidial fork long, thin and tipped with
microscopic hair B. NIGRA, Meigen.
- Tines stronger and tipped with one long and several short hairs B. PECTINATA, nov. sp. (2)

Geographical distribution.

1. *B. alpina*, Bezzii, Ditterfaun. nivale, p. 75 (1918). Alps.
2. *B. angustifurca*, nov. sp. W. United States.
3. *B. ? aterrima*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 55 [1840] (*Cyrtoma*). C. Europe.
4. *B. brevifurca*, nov. sp. — Pl. 8, Fig. 72g. Washington.
5. *B. ? collina*, Philippi, Verh. Ges. Wien, Vol. 15, p. 768 [1865] (? *Cyrtoma*). Chile.
6. *B. dispar*, Oldenberg, Zool. Jahrb. Vol. 43, Syst. p. 221 (1920). Alps.
7. *B. furcifer*, nov. sp. — Pl. 8, Fig. 72a. W. North America.
8. *B. halteralis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 206 : Cent. 2, E. United States.
No. 46 [1862] (*Cyrtoma*); Melander, Trans. Amer. Ent. Soc. Vol. 28,
p. 335 [1902] (*Cyrtoma*).
9. *B. intermedia*, Lundbeck, Dipt. Dan. Vol. 3, p. 25 (1910); Collin, Ent. N. Europe.
Mag. London (2), Vol. 24, p. 104 (1913).
- nigra*, Zetterstedt, part. Dipt. Scand. Vol. 1, p. 330, part. [1842] (*Cyrtoma*). North America.
10. *B. longipes*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 206 : Cent. 2, No. 47 [1862] (*Cyrtoma*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 336, f. 71 [1902] (*Cyrtoma*). — Pl. 8, Fig. 72b. North America.
11. *B. lugubris*, nov. sp. — Pl. 8, Fig. 72d. W. United States.
12. *B. melana*, Haliday, Ent. Mag. Vol. 1, p. 158 [1833] (*Cyrtoma*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 489 (1849); Ins. Brit. Dipt. Vol. 1, p. 116 [1851] (*Cyrtoma*); Lundbeck, Dipt. Dan. Vol. 3, p. 22 (1910).
simplices, Zetterstedt, Dipt. Scand. Vol. 1, p. 331 [1842] (*Cyrtoma*); Bonsdorff, Finl. tvåv. Ins. Vol. 1, p. 157 [1861] (*Cyrtoma*); Schiner, Fauna Dipt. Aust. Vol. 1, p. 76 [1862] (*Cyrtoma*); Wahlgren, Ent. Tidskr. Vol. 31, p. 48 (1910); Collin, Ent. Mag. London (2), Vol. 24, p. 105 [1913] (*Cyrtoma*). N. and C. Europe.

(1) ***Bioellaria lugubris*, nov. sp.** — Male. Length 2.75 mm. Very black, all hairs and bristles black. Upper facets large, about fourteen along the line of contiguity of the eyes. Base of third antennal joint rather quadrate, about equal in length to the slender apical part, arista more than two-thirds the length of the third joint. Anterior hairs of notum short, posterior four dorsocentrals longer than the others, six to eight scutellars. Pygidium narrowly U-shaped, tines moderately thick, about three times as long as width at middle, tips rounded and bearing a small hair. Hind femora appearing ciliate, with about eighteen setae in anterior flexor row and fifteen in extensor row, hind tibiae subclavate, their extensor hairs irregular with two or three longer than the others, hind metatarsi slightly thicker than the following joint. Halteres black. Wings infumated, stigma brown, fork of fourth vein open, the pedicel shorter than the fourth vein.

Nine males taken along Indian Creek in the northern part of Yellowstone Park, Wyoming, July 8, 1923. A female from the Gallatin Station in the northwestern part of Yellowstone Park, July 28, 1923, is probably the same species.

(2) ***Bioellaria pectinata*, nov. sp.** — Length 3.5 mm. Large black species with all hairs and bristles black. Upper facets large, about twelve along the line of contiguity of the eyes. Third antennal joint slender, the base ovate and rounding into the apical part, arista two-thirds the length of the third joint. Notum subshining, anterior hairs short, two dorsocentrals, four or six scutellars. Pygidium with rounded V-shaped incision on ventral fork, the tines moderately short and strong and tipped with one long and several short hairs. Setae of hind femora very long, about eighteen in the anterior extensor row and twenty-two in the flexor row; hind tibiae distinctly clavate, the extensor hairs irregular; hind metatarsi twice as thick as the second joint. Halteres black. Wings infumated, stigma brown, fork of fourth vein open, the pedicel subequal in length to the fourth vein beyond.

Thirty-four specimens, Ilwaco, Washington, May-September.

13. *B. nigra*, Meigen, Syst. Beschr. Vol. 4, p. 3, pl. 33, f. 5 [1824] (*Cyrtoma*); Europe.
 Macquart, Dipt. N. France, Vol. 3, p. 108 [1827] (*Cyrtoma*); Hist. Nat. Dipt. Vol. 1, p. 360, pl. 8, f. 15 [1834] (*Cyrtoma*); Zetterstedt, Fauna Ins. Lappon. p. 533 [1838] (*Cyrtoma*); Blanchard, Hist. Nat. Ins. Vol. 3, p. 583 [1840] (*Cyrtoma*); Zetterstedt, Dipt. Scand. Vol. 1, p. 330, part. [1842] (*Cyrtoma*); Walker, Ins. Brit. Dipt. Vol. 1, p. 116 [1851] (*Cyrtoma*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 60 [1851] (*Cyrtoma*); Zetterstedt, Dipt. Scand. Vol. 13, p. 4997 [1859] (*Cyrtoma*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 76 [1862] (*Cyrtoma*); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 387 [1866] (*Cyrtoma*); Becher, Denkschr. Akad. Wiss. Wien, Vol. 45, p. 147, pl. 3, f. 12 [1882] (*Cyrtoma*); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 133 [1887] (*Cyrtoma*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 43 [1892] (*Cyrtoma*); Lundbeck, Dipt. Dan. Vol. 3, p. 26 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 48 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 8 (1913).
14. *B. pectinata*, nov. sp. — Pl. 8, Fig. 72c. Washington.
15. *B. pilipes*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 207: Cent. 2, No. 48 [1862] (*Cyrtoma*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 336 [1902] (*Cyrtoma*); Coquillet, Proc. Wash. Acad. Sc. Vol. 2, p. 412 [1900] (*Cyrtoma*). — Pl. 8, Fig. 72e. North America.
16. *B. pilosa*, Lundbeck, Dipt. Dan. Vol. 3, p. 27 (1910); Collin, Ent. Mag. Lond. (2), Vol. 24, p. 104 [1913] (*Cyrtoma*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 9 (1913). N. Europe.
17. *B. rufa*, Meigen, Syst. Beschr. Vol. 4, p. 3 [1824] (*Cyrtoma*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 360 [1834] (*Cyrtoma*); Schiner, Fauna Dipt. Aust. Vol. 1, p. 76 [1862] (*Cyrtoma*); Dahl, Fauna Chorin, p. 464 (1912). C. Europe.
18. *B. spuria*, Fallen, Empid. Suec. p. 33 [1816] (*Empis*); Zetterstedt, Fauna Ins. Lappon. p. 533 [1838] (*Cyrtoma*); Dipt. Scand. Vol. 1, p. 329 [1842] (*Cyrtoma*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 489 [1849] (*Cyrtoma*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3013 [1849] (*Cyrtoma*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 60 [1851] (*Cyrtoma*); Walker, Ins. Brit. Dipt. Vol. 1, p. 116, pl. 4, f. 6 [1851] (*Cyrtoma*); Zetterstedt, Dipt. Scand. Vol. 13, p. 4997 [1859] (*Cyrtoma*); Pipping, Not. Sällsk. Fenn. Förh. Vol. 4, p. 114 [1858] (*Cyrtoma*); Nylander, Not. Sällsk. Fenn. Förh. Vol. 4, p. 247 [1858] (*Cyrtoma*); Bonsdoff, Fin. tvåv. Ins. Vol. 1, p. 156 [1861] (*Cyrtoma*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 76 [1862] (*Cyrtoma*); Siebke, Nyt. Mag. Naturvid. Vol. 14, p. 381, 399 [1866] (*Cyrtoma*); Neuhaus, Dipt. March p. 68, 124 [1886] (*Cyrtoma*); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 133 [1887] (*Cyrtoma*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 42 [1892] (*Cyrtoma*); Wien. Ent. Zeit. Vol. 18, p. 12 [1899] (*Cyrtoma*); Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 305 (1906); Lundbeck, Dipt. Dan. Vol. 3, p. 24, f. 4, 5 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 48, f. 4 (1910); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 464 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 8 (1913); Bezzi, Suppl. Ent. Vol. 3, p. 70 (1914). — Pl. 5, Fig. 46. N. and C. Europe.
- atra*, Meigen, Syst. Beschr. Vol. 4, p. 2 [1824] (*Cyrtoma*); Macquart, Dipt. N. France, Vol. 3, p. 108, pl. 3, f. 3 [1827] (*Cyrtoma*); Hist. Nat. Dipt. Vol. 1, p. 360 [1834] (*Cyrtoma*); Boitard, Man. Ent. Vol. 3, p. 326 [1843] (*Cyrtoma*).
- sulcata*, Zetterstedt, Dipt. Scand. Vol. 1, p. 331 [1842] (*Cyrtoma*); Bonsdorff,

Finl. tvåv. Ins. Dipt. Vol. 1, p. 157 [1861] (*Cyrtoma*) ; Siebke, Nyt. Mag. Naturvid. Vol. 12, p. 108 [1864] (*Cyrtoma*).

nigra, Macquart, Mém. Soc. Sc. Lille, 1823, p. 156 (1823) ; Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 246 (1903).

19. *B. ? tibialis*, Roser, Correspondenzbl. Landw. Ver. Würtenb. Stuttgart, C. Europe. Vol. 1, p. 55 [1840] (*Cyrtoma*).

20. *B. uvens*, nov. sp. — PI. 8, Fig. 72 f. Hudson Straits.

14. GENUS HOPLOCYRTOMA, NOV. GEN.

Characters. — Similar to *Bicellaria* in general structure and in neuration but differing in the possession of stout and spinose hind femora and correspondingly strong hind tibiæ. The hind femora are swollen and abundantly armed beneath with a mixture of spines and thorns. The hind tibiæ are much shorter than the femora, not clavate but nearly straight and cylindrical, geniculate at the knee and carinate along the flexor edge. Apically they are somewhat obliquely truncate and closely fimbriate. The tines of the ventral piece of the pygidium are thick and bluntly rounded, the one on the right longer than the one on the left.

Genotype : *Bicellaria procera*, Loew (PI. 2, Fig. II). The two American species, *procera* and *femorata*, are closely related, differing from the species of *Bicellaria* in the same leg specialization that has been repeatedly developed in various groups of the *Empididae*.

Geographical distribution.

1. *H. femorata*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 84 : Cent. 5, No. 69 E. United States.

[1864] (*Cyrtoma*) ; Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 335
[1902] (*Cyrtoma*).

2. *H. procera*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 85 : Cent. 5, No. 70 Alaska; Washington.

[1864] (*Cyrtoma*) ; Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 335
[1902] (*Cyrtoma*). — PI. 2, Fig. II.

SUBFAMILY EMPIDINÆ

Characters. — Head globular, occiput little convex, in *Xanthempis* drawn out to a sort of cone ; eyes bare except in *Parathalassius* and *Microphorella*, males dichoptic or holoptic, in the former case the facets are uniform or the anterior middle ones slightly enlarged, in the latter case the upper facets are more or less enlarged, eyes of the females always separated ; face broad, often short and receding and then with the oral margin arched, in specialized forms of *Empis* and *Rhamphomyia* the face becomes narrow : front quadrate, often with small orbita ; antennæ three-jointed, the first joint usually as long as or longer than the globular second joint, in *Hormopesa* the basal joints fused, in *Ragas* the basal joints small, third joint conical and cylindrical, or (*Gloma*) reniform, or (*Hormopesa*) orbicular, or (*Apalocnemis*, *Opeatocerata*) oval, tipped with a style of two to four joints, usually the style with a quadrate basal piece ending in a short bristle, sometimes the end piece of the style is longer and tapers, sometimes (*Gloma*, etc.) the antenna is furnished with a long thin arista ; proboscis usually fitted for piercing, corneous, variable in length and direction, sometimes shorter than the head, sometimes in the flower-sucking species nearly as long as the body, palpi single-jointed. Thorax never greatly convex, usually the humeral, posthumeral and supra-alar and almost always the notopleural and scutellar bristles present, the acrostichals and dorsocentrals variable, the former almost never pronounced, the latter usually weak;

metapleural setæ absent or present. Pygidium terminal or more or less reflexed, when reflexed usually somewhat compressed; no ovipositor. Legs of moderate length, coxæ never long, sometimes the hind femora incrassate, middle tibiae usually with apical bristles, and posterior tibiae often furnished with sets of setæ. Wings more or less oblong, the anal angle usually quite distinct though often broadly rounded, alula never large, costa continuing around the wing or stopping at the fourth vein, auxiliary vein strong and always distinctly separated from the first, attaining the costa or distally evanescent, third vein forked or not, discal cell complete and emitting three veins, but in a few species of *Rhamphomyia* the posterior crossvein is absent, anal cell shorter than the second basal, the anal crossvein abruptly reflexed (except in *Parathalassius* where it is perpendicular), anal vein almost always separate from the anal cell, reaching the margin or abbreviated but always weak.

This subfamily connects to an astonishing degree with the Clinoceratinæ. In the *Microphorus* group the enlarged epipygium and small basal cells are quite suggestive of the Clinocerans. *Hilarempis* and the allied genera show so many traits of *Proclinopyga* and *Dipsomyia*, in the type of genitalia, chætotaxy, neuration and structure of wing, that there is certainly a close relationship between them. On the other hand *Proclinopyga* connects still more closely with *Clinocera*, as shown by the V-shaped front, narrowed face, mouthparts, structure of the thorax and abdomen and of the legs.

SYNOPSIS OF THE GENERA AND SUBGENERA OF THE EMPIDINÆ

- Basal and anal cells very small, third vein simple, costa continuing around the wing; male genitalia formed as a very large epipygium which is twisted to the right; thorax longer than the abdomen, robust and gibbous; fronto-orbital, posthumeral and intra-alar bristles present; arista one-fifth to three times the length of the third antennal joint, the latter often suddenly attenuated beyond the base and produced styliform; middle tibiae with an apical spur, hind femora usually conspicuously ciliate* 2.
- Basal and anal cells never markedly short; male genitalia never thrown far forward over the abdomen nor twisted to the right; abdomen at least as long as the thorax; third antennal joint not abruptly constricted just beyond the base; eyes bare.* 5.
2. *Eyes of the male broadly contiguous along the front, bare, the upper facets larger than the lowermost; ocelli approximated; antennæ inserted below the middle of the head, distinctly three-jointed; face more or less excised at the oral margin; anterior dorsocentrals reduced in size; abdomen of the male with long hairs, the last segments of the female short, tubular; discal cell relatively broad, anal crossvein rounding, anal vein rudimentary, anal angle of the wing broadly rectangular, a strong alular fringe (genus *Microphorus*).* 3.
- Eyes of both sexes broadly separated along the front, pubescent, the uppermost facets smaller than the lower; ocelli distant; antennæ inserted above the middle of the head; the basal two joints fused; mouthparts short and vertical, not protruding; anterior dorsocentrals strong; abdomen of the male with short hairs, fifth segment of the female abdomen broadly triangular;*

- discal cell relatively narrow, its basal crossvein weakened in front, no stigma* 4
3. *Proboscis short, sometimes as long as the head, directed horizontally forward; third joint of the antennæ abruptly constricted, second joint setose; female with one small fronto-orbital; two or more notopleurals; costa with a basal bristle, stigma usually distinct. (Pl. 7, Fig. 63).*
- Proboscis short, not protruding beyond the oral margin, vertical; second antennal joint not setose; female with two fronto-orbitals; no basal bristle on the costa, no stigma. (Pl. 7, Fig. 62)*
4. *Anal crossvein rounding, anal vein more or less rudimentary, anal angle poorly developed; eyes finely pubescent; face not broadened below, with only two oral hairs; two scutellars.*
- Anal crossvein perpendicular, base of the anal vein strong, anal angle distinct; eyes densely hairy; face very narrow beneath the antennæ, broadened below, more or less hairy; hairs of the abdomen coarse; four or six scutellars*
5. *Proboscis directed horizontally or obliquely forward, either short or long, palpi porrect; eyes of the male contiguous above the antennæ, the upper facets largest; face broad, arched and receding from directly beneath the antennæ, the oral margin therefore widely arched, facial orbits distinct; auxiliary vein complete, ending in the costa, anal angle strongly rectangular, with an axillar excision, costa usually thin on the hind margin; bristles of the body well developed.* 6
- Proboscis long and thick, directed obliquely forward, palpi incumbent; eyes widely separated, facets uniform; antennæ inserted low, the face short but vaulted, facial orbits not differentiated, third antennal joint large and oval, without arista or style; auxiliary vein incomplete at tip, third vein furcate, anal angle weak, no axilla, costa encompassing wing; bristles of body weak; hind tibiae geniculate; Clinocera-like flies. (Pl. 2, Fig. 16)*
- Proboscis vertical or somewhat reflexed, sometimes slightly projecting, in which case the relatively narrow and lengthened face is characteristic; face not immediately receding but flat or convex below the antennæ; style shorter than the third antennal joint; pygidium usually not open.* 12.
6. *Antennæ apparently with only two joints, the outer joint broadly orbicular; anal crossvein not greatly reflexed, the anal cell as broad as the second basal; tibiae with several sets of bristles; pleurae bare; pygidium globose, the dorsal valves large, erect and simple, the lateral valves very convex and posteriorly with two blunt prongs. (Pl. 8, Fig. 78)*
- Antennæ distinctly three-jointed, the last joint more or less elongate, never orbicular; anal crossvein abruptly reflexed, the*
- Subgenus MICROPHORUS, Macquart.
- Subgenus SCHISTOSTOMA, Becker.
- Genus MICROPHORELLA, Becker.
- Genus PARATHALASSIUS, Mik.
- Genus BROCHELLA, nov. gen.
- Genus HORMOPEZA, Zetterstedt.

- anal cell narrow; pygidium with exposed dorsal processes, the dorsal valves usually narrow and fissate* 7.
7. *First antennal joint longer than the second and both strongly setose; style aristiform, indistinctly four-jointed, the basal joint minute, the terminal like a short hair, the intermediate joints lengthened and moderately thick; palpi strongly setose; basal valve of the pygidium ovate, setose and not fissate; hind margin of the wing somewhat thickened* 8.
- Antennæ not setose, at most with a few fine basal hairs, furnished with a short style or a long arista; palpi not strongly setose; metapleura bare; basal valve of the pygidium irregular and fissate; veins bare, wings scarcely margined behind* 9.
8. *Metapleura bare; veins bare. (Pl. 6, Fig. 53).* Genus TIMALPHES, nov. gen.
- Metapleura with a cluster of setæ; first second and sometimes third veins closely setulose on the upper surface and the second underneath. (Pl. 5, Fig. 50; Pl. 6, Fig. 51)* Genus OREOGETON, Schiner.
9. *Proboscis short, scarcely protruding beyond the head; third antennal joint short, furnished with a lengthened arista which is twice as long as the antenna and whose basal segment is narrowly tubular, followed by a very long geniculate short-pubescent portion and ending in a distinct fine hair; third vein furcate, discal cell of moderate size.* 10.
- Proboscis long, extending obliquely forward, palpi linear; third antennal joint lengthened cylindrical, the terminal style short and thick and tipped with a short but distinct bristle; discal cell large and blunt* 11.
10. *Arista terminal, the third joint of the antennæ small and globular. (Pl. 6, Fig. 56)* Genus APALOCNEMIS, Philippi.
- Arista dorsal, inserted near the base of the reniform third antennal joint; calypteres lobose and thickly margined. (Pl. 6, Fig. 52; Pl. 8, Fig. 73, 74)* Genus GLOMA, Meigen.
11. *Third vein furcate. (Pl. 6, Fig. 59)* Genus ITEAPHILA, Zetterstedt.
- Third vein simple. (Pl. 8, Fig. 80)* Genus ANTHEPISCOPUS, Becker.
12. *Metapleura bare of hairs or bristles; costa encompassing the entire wing, axillary incision rarely distinguishable, usually obtuse or entirely wanting; style ending in a distinct short bristle (except in Philetus); usually the bristles of the body, head and legs weak; male genitalia usually more or less of the form of an epipygium with compressed ventral keel.* 13.
- Metapleura bearing a variable number of hairs or bristles in front of the halteres; costa stopping at the tip of the wing, or at least greatly thinned on the hind margin, auxiliary vein straight, not reaching the costa, axillary incision sharp; eyes of male contiguous above the antennæ, except in some species of Empis and Rhamphomyia; style not ending in a distinct thin short bristle; usually the bristles of the head, body and*

- legs strong; pygidium rarely reflexed and not carinate beneath.
(If costa encompasses wing and front is broadly V-shaped,
see Proclinopyga in the Clinoceratinae) 23.
13. Auxiliary vein complete, ending in the costa 14.
- Auxiliary vein evanescent at the end, not attaining the costa 17.
14. Eyes of the male contiguous, upper facets largest; basal joints
of the antennae minute, first joint of the style very thick; proboscis strong, the labrum inflexed; face short, epistome broadly
receding; anal angle of the wing strong; anal vein meeting the
under side of the anal cell, anal crossvein short; pygidium
showing dorsal appendages. (Pl. 8, Fig. 84) Genus RAGAS, Walker.
- Eyes of the male broadly separated, middle facets largest; first
antennal joint distinct; face not short; pygidium not open 15.
15. Auxiliary vein bending forward at the end to meet the costa, anal
angle prominent, anal crossvein reaching nearly halfway to
the base of the anal cell, anal vein separate from the under
side of the anal cell, fork of the third vein long; at least
fronto-orbital and lateral thoracic bristles evident; front
metatarsi of the male almost always enlarged; pygidium
compressed and directed forward over the abdomen; palpi
cylindrical and upturned or slightly clavate. (Pl. 6,
Fig. 55) Genus HILARA, Meigen.
- Auxiliary vein almost straight, not bending forward to meet the
costa, anal angle broadly rounded, anal crossvein short, anal
vein meeting the under side of the anal cell or obsolete, fork of
the third vein short; metatarsi simple; bristles rarely evident;
pygidium globose 16
16. Antennae tipped with a short two-jointed arista whose basal seg-
ment is one-fourth the distal; dorsocentral, scutellar and
lateral bristles present; ocelli located in front of the vertex,
antennae below the middle of the head; pygidial valves directed
backward. (Pl. 7, Fig. 66; Pl. 8, Fig. 78, 81). Genus PHILETUS, nov. gen.
- Antennae tipped with a short two-jointed style, whose basal seg-
ment is longer than the terminal stubby bristle; thoracic
bristles almost wanting; ocelli located on the vertex, antennae
at the middle of the head; pygidial valves erect. (Pl. 7,
Fig. 61) Genus HESPEREMPI, Melander.
17. First vein thickly beset with hairs along its entire length;
mesonotum clothed with short hairs, mesopleura and ptero-
pleura sometimes hairy; proboscis as long as the head; males
dichoptic; axillary angle of the wing strong, anal angle well
developed. (Pl. 2, Fig. 15) Genus PHLEBOCTENA, Bezzii.
- All the veins bare; mesopleura and pteropleura bare 18.
18. Eyes of the male contiguous on the front; abdomen very short;
third vein forked, anal angle well developed; third joint of
the antennae long and conico-cylindrical, style three-jointed,

- the basal joint small, the middle joint long, the terminal joint short and bristle-like. (Pl. 2, Fig. 14)* Genus DEUTERAGONISTA, Philippi.
- Eyes of both sexes widely separated; abdomen evidently longer than the thorax; anal angle of wing usually broadly rounded.* 19.
19. *Proboscis twice as long as the head; almost entirely devoid of hairs and bristles; third vein forked; third joint of the antennæ conical.* Genus TOREUS, Melander.
- Proboscis about as long as the head; at least the lateral bristles of the thorax more or less developed* 20
20. *Submarginal cell closed by the erect fork of the third vein; third antennal joint expanded nearly oval at base and narrowly drawn out at apex, style very long and aristiform; legs very slender; proboscis with long hairs. (Pl. 8, Fig. 83)* Genus TENONTOMYIA, White.
- Submarginal cell open; antennæ of other conformation, style shortened; proboscis bare* 21.
21. *Hind femora thickened and longer than their geniculate tibiæ; third joint of the antennæ very long, gradually tapering, almost cylindrical. (Pl. 6, Fig. 58)* Genus HAPLOMERA, Macquart.
- Hind femora and tibiæ not modified; third joint of the antennæ conical* 22.
22. *Third vein forked; front metatarsi of male often enlarged. (Pl. 6, Fig. 54)* Genus HILAREMPIS, Bezzi.
- Third vein simple; tibiæ of the female without bristles. (Pl. 6, Fig. 60)* Genus ATRICHOPLERA, Bezzi.
23. *Third antennal joint short-ovate, with aristiform style three times its length; anal angle of wing undeveloped, axillary incision weak, basal cells shorter than discal, first vein ending before middle, third vein with long erect fork, fourth and intercalary veins evanescent; thorax strongly convex, bristles reduced; ovipositor compressed; shining yellow species.* Genus OPEATOCERATA, nov. gen.
- Third antennal joint lengthened, longer than the style; base of wing better developed, the anal angle usually full, the axillary incision usually deep, first vein ending beyond the middle; if yellow species not polished* 24.
24. *First antennal joint nearly or quite as long as the third, the antennæ located high up on the head; brilliant metallic tropical species usually with the legs variously modified* 25.
- First antennal joint not lengthened, much shorter than the third joint, the antennæ located at the middle or but little above the middle of the head.* 26.
25. *Arista much shorter than third antennal joint; body pilose; third vein with erect fork. (Pl. 2, Fig. 13)* Genus LAMPREMPIS, Wheeler & Melander.
- Arista thick and nearly as long as remainder of antenna; thorax not pubescent; third vein simple. (Pl. 2, Fig. 12)* Genus PORPHYROCHROA, nov. gen.
26. *Robust hairy species, the face furnished with hairs; fringe of the calypteres rudimentary* 27.
- Face bare; calypteres fringed with hairs, rarely the fringe short.* 28.

27. *Third vein furcate* Genus **EMPIMORPHA**, Coquillett.
Third vein simple (Pl. 6, Fig. 57). Genus **NEOCOTA**, Coquillett.
28. *Third vein furcate* (Genus **EMPIST**, Linnaeus.) 29.
Third vein simple (Genus **RHAMPHOMYIA**, Meigen.) 41.
29. *Dorsocentrals uniserial, even in the middle of the dorsum, acrostichals usually absent, when the dorsocentrals occur in more than one row the males are dichoptic* 30.
Dorsocentrals bi- or pluriserial, at least along the middle of the row, acrostichals usually present, when but one row of dorsocentrals present the males are holoptic 35.
30. *Eyes of both sexes separated or at most subcontiguous, the facets small and uniform.* 31.
Eyes of the male contiguous, the upper facets enlarged; no acrostichals 34.
31. *No propleural or notopleural bristles, metapleural hairs few, short and stiff, anal lobe of the wing rounded; hind femora not thickened* 32.
Propleural and notopleural bristles present; head round; anal angle of the wing rectangular. 33.
32. *No acrostichal setulae; head almost always pear-shaped, the occiput drawn out; more or less yellow species* Subgenus **XANTHEMPIS**, Bezzi.
Acrostichal setulae occurring as a single median row, rarely absent or closely biserial; head globose, the occiput not conical; cinereous black species, the hind knees of the male usually with strong apophyses Subgenus **ENOPLEMPIS**, Bigot.
33. *Hind femora strongly thickened; metapleural hairs piliform; black species.* Subgenus **ANACROSTICHUS**, Bezzi.
Femora not thickened; metapleural hairs setiform; largely yellow species Subgenus **PYRREMPIS**, nov. subgen.
34. *Discal cell blunt apically; male with silvery pruinosity; scutellar bristles cruciate.* Subgenus **ARGYRANDRUS**, Bezzi.
Discal cell sharp apically; male not pruinose; scutellar bristles not cruciate. Subgenus **HAPLOMERINX**, Bezzi.
35. *Fourth vein interrupted before the margin of the wing, fork of third vein short and nearly erect, discal cell blunt, anal angle rectangular; eyes of the male contiguous; femora not thickened; legs of female pinnate; acrostichals sometimes wanting.* Subgenus **COPTOPHLEBIA**, Bezzi.
Fourth vein reaching the margin, in livida the fourth and intercalary veins interrupted at the margin; acrostichals present, bi- or pluriserial 36.
36. *Anal angle of the wing broadly rounded; face narrow; eyes of both sexes narrowly separated, in the males sometimes subcontiguous; one presutural bristle; body shining black, with fine pile and reduced bristles; hind legs long, the metatarsi more or less swollen* Subgenus **LISSEMPIS**, Bezzi.
Anal angle not obtuse but more or less rectangular; face broad,

<i>eyes of the female broadly separated; usually more than one presutural bristle present</i>	37.
37. <i>Eyes of the male broadly contiguous, the upper facets larger than the lower; hind femora not thickened</i>	38.
<i>Eyes separated, rarely approximated in the male on the middle of the front, always with small uniform facets; hind femora thickened</i>	40.
38. <i>Discal cell apically blunt; legs of the female usually more or less pennate, of the male the front or hind metatarsi or both more or less enlarged; generally smaller species. (Pl. 5, Fig. 48, 49)</i>	Subgenus <i>EMPIIS</i> , Linnæus, s. str.
<i>Discal cell apically more or less narrowed; legs of the female not pennate; of the male the metatarsi not thickened</i>	39.
39. <i>Delicate, slender, long-legged species; face narrow, palpi bare, proboscis very long and thin, projecting obliquely forward; thorax not pubescent, acrostichals few; pygidium small and terminal; leg bristles undeveloped.</i>	Subgenus <i>ACALLOMYIA</i> , nov. subgen.
<i>Robust, generally larger species; face broad, palpi usually heavily hairy; thorax usually pubescent; pygidium stout; legs bristly.</i>	Subgenus <i>PLATYPTERA</i> , Meigen.
40. <i>Segments of the abdomen with strong lateral bristles; humeral, propleural and metapleural bristles very strong</i>	Subgenus <i>POLYBLEPHARIS</i> , Bezzi.
<i>Abdominal segments without strong bristles; humeral, propleural and metapleural bristles not unusually strong. (Pl. 5, Fig. 47)</i>	Subgenus <i>PACHYMERIA</i> , Stephens.
41. <i>Third antennal joint greatly lengthened, the antennæ about twice as long as head; ocellar triangle not raised, the ocellar bristles weak; body narrow, legs slender; no evident dorsocentrals or acrostichals; wings small, with very obtuse to rectangular axillar incision, alulae almost wanting; eyes of male separated.</i>	Subgenus <i>LUNDSTROEMIELLA</i> , Frey.
<i>Third antennal joint not lengthened, the antennæ less than twice as long as the head; ocellar triangle developed, with evident ocellar bristles</i>	42.
42. <i>Small delicate species with long slender legs; dorsocentral bristles in a single row, acrostichals present or absent; color of thorax differing in male and female, or uniformly yellow in both sexes, axillar incision of wings blunt to somewhat pointed; eyes of male contiguous; pygidium small and closed</i>	Subgenus <i>HOLOCLERA</i> , Schiner.
<i>Body more robust; legs stronger, acrostichals present; thorax of male and female similarly colored, rarely (in American species) yellow</i>	43.
43. <i>Axillar incision of wings obtuse to rectangular, rarely somewhat pointed; pygidium usually open, penis often filiform and projecting; hind femora not spinose beneath.</i>	44.
<i>Axillar incision rather acute; pygidium rather small or else strongly bulbous, penis usually short and thick, rarely filiform and extended</i>	45.

44. Eyes of male separated; discal cell of female often more or less broadened; hairs of abdomen mostly white or brown; dorsocentrals usually weak, hair-like and in several rows; legs with few bristles, those of female not feathered Subgenus *MEGACYTTARUS*, Bigot.
- Eyes of male contiguous; discal cell of female not widened; dorsocentrals usually in one or two rows, if short and in several rows then the thorax is shining black; legs of female sometimes feathered Subgenus *PARARHAMPHOMYIA*, Frey.
45. Pygidium vesiculate above or broadly deformed, the seventh tergite usually set in the wide sixth, often with two prongs below and with two bunches of yellow bristles on last sternite; thorax usually opaque, dorsocentrals hair-like, multiserrate, scutellars numerous; wings of female usually rather broadened and brown; hind femora rarely setose below Subgenus *DASYRHAMPHOMYIA*, Frey.
- Pygidium of other form, usually small, half-open, with short thick penis; hind femora of both sexes usually more or less bristly below Subgenus *RHAMPHOMYIA*, Meigen, s. str.

I. GENUS *PARATHALASSIUS*, MIK

Parathalassius, Mik. Wien. Ent. Zeit. Vol. 10, p. 217 (1891); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255, 260 (1903); Melander, Williston N. Amer. Dipt. p. 224 (1908); Becker, Wien. Ent. Zeit. Vol. 28, p. 25-27 (1909); Kertész, Cat. Dipt. Vol. 6, p. 101 (1909).

Characters. — Small robust silvery gray pruinose species with white bristles. Head longer than broad, occiput rather flattened, eyes of both sexes widely separated on the front, approaching just beneath the antennæ so as there to narrow the face, more in the male and less in the female, and then again diverging below, anterior and central facets large, the uppermost facets minute, eyes strongly pubescent, especially below, where in the male the hairs give a glistening sheen to the eyes; face concave above, convex below, the oral margin at the lowest level of the eyes and not excised, fringed especially in the male with white hairs; basal joints of the antennæ fused and nearly bare, the third joint pyriform, with a slender two-jointed arista which is as long as the antenna itself; proboscis very short, tubular and thick, not or scarcely projecting; ocelli widely separated, not elevated, ocellar bristles distant and diverging, a pair of strong converging fronto-orbitals; bristles of the upper occiput moderately strong, below a mystax of coarse white hairs present. Thorax stout, not highly arched, five or six strong dorsocentrals, one posthumeral, several supraalar, two or more scutellars, the middle pair distant; pleuræ uniformly and densely pruinose and devoid of hairs. Abdomen thick and short, densely pruinose, its hairs coarse, only moderately long; epipygium very large, tumid, globose, twisted to the right, its under parts coarse and complicated; fifth segment terminating the abdomen of the female, the apical segments retracted. Legs simple, pruinose, the femora more or less ciliate, middle tibiæ with an apical bristle. Wings narrow, with parallel sides, costa continuing around the entire wing, several basal bristles and costal setulæ present, third vein simple, discal cell narrow and long, its basal crossvein abortive, anal crossvein straight and perpendicular, anal vein angulate at the crossvein and strong halfway to the margin, anal angle broadly rounded, no alula, the alular hairs moderately long; calypters with straight margin.

Type species: *P. Blasigii*, Mik, the original species. Judged by the course of the anal cross-vein this genus should be located in the Ocydromiinae, but as all the other characters clearly point to

the *Microphorus* group it follows that the perpendicular crossvein has been independently developed in *Parathalassius*. The species of the genus frequent the sands of the seashore, which habit accounts for their white bristles and silvery gray color.

Geographical distribution.

1. *P. Aldrichi*, Melander, Ent. News, Philad. Vol. 17, p. 374, fig. (1906). California, Washington.
2. *P. Blasigii*, Mik, Wien. Ent. Zeit. Vol. 10, p. 217 (1891); Becker, ibidem, S. Europe; N. Africa. Vol. 28, p. 29 (1909). *alter*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 119 [1907] (*Schistostoma*).
3. *P. candidatus*, Melander, Ent. News, Philad. Vol. 17, p. 375 (1906). Washington.
4. *P. Melanderi*, Cole, Report Laguna Marine Labor. Vol. 1, p. 154, f. 88 (1912). California.

2. GENUS MICROPHORELLA, BECKER

Microphorella, Becker, Wien. Ent. Zeit. Vol. 28, p. 28 (1909).

Characters. — Small opaque black bristly species with the eyes separated in both sexes and with the anal angle of the wing reduced, the legs and halteres always black. Front broad in both male and female, with a pair of converging strong fronto-orbital bristles, ocellar triangle not elevated, the ocelli widely spaced apart, ocellar bristles strong and distant from each other; face long, with parallel sides, narrower than the front, no cheeks; eyes with microscopic pubescence, the lower facets of the male very slightly larger than the upper; antennæ inserted above the middle of the head, two-jointed as the basal joints are fused, the basal joint without bristles, the outer joint conical, pyriform or cylindrical, pubescent, with a rather thick, two-jointed apical arista, which is geniculate beyond the minute basal joint; mouthparts very short, not protruding, fleshy, palpi small, elongate oval and tipped with a single hair; occiput hairy the upper part setose and with stronger vertical bristles. Thorax large, robust, longer than the abdomen, with a complete series of about six strong dorsocentrals, acrostichals biseriate and biseriate setulæ present between the dorsocentral and supraalar bristles, one strong posthumeral, one notopleural, two scutellar bristles; pleuræ bare, but opaque pollinose. Abdomen thick, its hairs sparse and very short; pygidium enormously enlarged, sometimes as large as the remainder of the abdomen, twisted to the right, globular, completely enclosed above, the fourth ventral more or less excised medially along its hind margin; no ovipositor, the last segments of the female abdomen short and broad and margined with bristles. Legs slender, in the male often deformed, especially the tarsi, middle tibiæ with a short apical spur, hind femora more or less ciliate below. Wings rather pointed, broadest at the middle, the hind angle greatly reduced and not rectangular, costa encompassing the entire wing, its basal bristle strong, no stigma, auxiliary vein distinct, usually shortened, first vein ending beyond the middle of the wing, third vein simple, discal cell narrow, its basal crossvein weakened, the posterior crossvein very oblique, basal cells minute, anal crossvein round, anal vein present or absent, no alula and almost no alular hairs; fringe of the calypteres weak.

Type species : *M. præcox* Loew, apparently a very rare form and the only species known to Becker. The American species are closely related to each other but are easily recognized by the peculiar secondary sexual characters of the males. They occur in the grass and bushes along streams. Becker proposed *Microphorella* as a subgenus of *Microphorus*. As shown in the table of genera the genus is much more closely related to *Parathalassius*, but is distinct from both, and is well entitled to generic rank.

TABLE FOR THE IDENTIFICATION OF MALES OF MICROPHORELLA

1. Third antennal joint long and cylindrical; anal vein completely wanting; auxiliary vein ending opposite the end of the discal cell. *M. PRÆCOX*, Loew.
- Third joint of the antennæ conical or pyriform; base of the anal vein indicated as a fold; auxiliary vein very short, ending before the middle of the discal cell 2.
2. Hind trochanters with a spine-tipped tubercle; hind femora closely ciliate with bristles beneath; third joint of the antennæ more or less conical 3.
- Hind trochanters not tuberculate; hind femora with short or sparse cilia; third antennal joint attenuated at the end; ventral processes of the pygidium yellowish-brown 4.
3. Tubercle of the hind trochanters as long as the trochanter and clavate; hind tibiæ with erect hairs; ventral processes of the pygidium yellowish brown M. TUBERIFERA, nov. sp. (1).
- Tubercle of the hind trochanters small and cylindrical; hairs of the hind tibiæ not unusual; ventral processes of the pygidium black. M. ACROPTERA, nov. sp. (2).
4. Front tarsi with white sole, hind tarsi compressed, the joints lobate above; hind femora with five long flexor bristles; third joint of the antennæ densely pubescent beneath. M. ORNATIPES, nov. sp. (3).

(1) *Microphorella tubifera*, nov. sp. — Male. Length 2 mm. Opaque black with a brownish tinge. Outer antennal joint elongate oval, one-half longer than wide, with very short pubescence, the arista nearly twice as long. Six strong dorsocentrals, the lateral setulae reduced, scutellum with a minute lateral bristle. Abdomen opaque, pygidium large and bulbous, underneath with relatively large brownish processes. Legs hairy, hind trochanters with a long trumpet-shaped tubercle bearing a blunt spine in front, front femora not ciliate, hind femora with fifteen moderate bristles along the antero-flexor edge, middle tibiæ tipped with several bristles, hairs of the hind tibiæ outstanding, tarsi simple but rather stout, the joints decreasing in length but the last joint as long as the third. Wings narrow, with a strong brownish tinge, first vein ending opposite the apex of the discal cell, sections of the fourth vein proportioned 3 : 3 : 10 : 13, of the fifth vein 2 : 1, anal vein short but strong.

Female. Similar except for the blunt abdomen, absence of the tubercle of the trochanters and lack of the femoral bristles.

One specimen of each sex: Palo Alto, California, March, 1895, received from Professor R. W. Doane.

(2) *Microphorella acroptera*, nov. sp. — Male. Length 1.5 mm. Opaque lead-black, the abdomen subshining. Third joint of the antennæ broadly oval, with an apical cylindrical process one-third the length of the basal portion, arista less than twice as long as the third joint. Abdomen with almost a brassy tinge, pygidium rather shining, large and inflated, its ventral processes entirely black. Hind trochanters with a short spine-tipped tubercle, anterior femora not ciliate, hind femora with fourteen bristles which are longer than the diameter of the femur, middle tibiæ with several minute apical bristles, tarsi simple, the joints decreasing in length but the last two subequal. Wings with very light infuscation, two and two-thirds times as long as broad, sections of the fourth vein proportioned 3 : 3 : 10 : 13, of the fifth vein 1 : 0.6, anal vein practically wanting.

Female. Abdomen with bristly blunt termination, femora not ciliate, trochanters not tuberculate.

Type from Pullman, Washington, May 30, 1913. The species occurs also in California. I have received specimens from San Jose and from Berkeley Hills, the latter from E. T. Cresson, Jr.

(3) *Microphorella ornatipes*, nov. sp. — Male. Length 1.75 mm. Black, uniformly coated with opaque olive gray pollen. Third joint of the antennæ densely pubescent with yellow hairs, pyriform, two-thirds longer than broad, constricted on the outer two-thirds, the arista longer than the third joint; palpi with a long brown hair. Six strong dorsocentrals, and two intra-alars besides the usual other bristles. Fifth, sixth and seventh segments of the abdomen progressively smaller, the pygidium lying to the right of these segments and provided underneath with many short filamentous yellow processes. Front femora with about ten cilia beneath, front tibiæ enlarged apically, almost trumpet-like, front tarsi yellowish within, the metatarsi as long as the remainder of the tarsus, slender, sinuous, the last four joints almost equal in length, the fifth joint flattened; hind trochanters not spined, hind femora with about seven long flexor bristles, hind tarsi compressed, the metatarsus slender and nearly as long as the remainder of the tarsus, the other joints subequal and lobosely carinate above. Wings with slight infumation, no stigma, first vein terminating opposite the end of the discal cell.

Female. Abdomen with a blunt tip, legs simple.

One specimen of each sex: Kendrick, Idaho, received from Dr. J. M. Aldrich.

- Front tarsi black, hind tarsi not lobate; hind femora without conspicuous bristles; pubescence of the third antennal joint not very dense 5.
5. Last joint of all the tarsi two or three times as long as the preceding joint, the front tarsi uniformly thickened; base of the anal vein distinct M. LONGITARSIS, nov. sp. (1).
- Last tarsal joint not elongate, subequal to the preceding joint; only the basal joint of the front tarsi swollen; anal vein indistinct M. CHIRAGRA, nov. sp. (2).

Geographical distribution.

1. <i>M. acroptera</i> , nov. sp.	W. United States.
2. <i>M. chiragra</i> , nov. sp.	Washington.
3. <i>M. longitarsis</i> , nov. sp.	Idaho.
4. <i>M. ornatipes</i> , nov. sp.	Idaho.
5. <i>M. præcox</i> , Loew, Zeitschr. Ent. Breslau, Vol. 14, p. 47 [1863] (<i>Microphorus</i>); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 93 [1892] (<i>Microphorus</i>); Becker, Wien. Ent. Zeit, Vol. 28, p. 25, fig. (1909).	C. Europe.
6. <i>M. tubifera</i> , nov. sp.	California.

3. GENUS MICROPHORUS, MACQUART

Microphorus, Macquart, Dipt. N. France, Vol. 3, p. 139 [1827] (*Microphor*); Hist. Nat. Dipt. Vol. 1, p. 345 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 253 [1842] (*Microphora*); Boitard, Man. Ent. Vol. 3, p. 322 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 113 (1851); Rondani, Dipt. Ital. Vol. 1, p. 151 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 563 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 79 (1862); Lioy, Atti Inst. Sc. Veneto, Venezia, 1864, p. 602 (1864); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 33-50 (1864); Nowicki, Gel. Ges. Krakau, Vol. 42, p. 72 (1871); Loew, Syst. Beschr. Eur. Dipt. Vol. 2, p. 250 (1871); Beling, Arch. Naturg. Berlin, Vol. 48, p. 240 (1882); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 118 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 332 (1902);

(1) ***Microphorella longitarsis*, nov. sp.** — Male. Length 1.6 mm. Black, opaque, coated with dark slaty brown pollen. Lower part of the face tuberculate, third joint of the antennæ pubescent beneath, suddenly constricted at its middle, one-half longer than broad and about half as long as the arista. Pygidium shining along the interior edge. Trochanters not tuberculate, front and hind femora weakly ciliate, joints of the front tarsi thickened and quadrate, first three joints together as long as their tibia, the fourth joint shorter than broad and one-third as long as the fifth joint, middle tibiæ with an apical spur, posterior tarsi similar to the front ones but less thickened. Wings long and narrow, nearly three times as long as broad, lightly infumated, sections of the fourth vein proportioned 3 : 3 : 11 : 18, of the fifth vein 1.5 : 1, discal cell four times as long as broad, first vein ending beyond the apex of the discal cell, anal vein distinct halfway to the margin.

Female. Similar except for the genitalia and simple tarsi.

Ten specimens : Moscow Mountain, Idaho, June 12, 1910.

(2) ***Microphorella chiragra*, nov. sp.** — Male. Length 1.6 mm. Black, coated with dark brownish gray pollen, the abdomen subshining. Third joint of the antennæ pubescent beneath, suddenly constricted on its outer third, the basal portion circular, arista one-half longer than the third joint. Seven dorsocentrals present. Fifth, sixth and seventh segments of the abdomen tumid; pygidium very large, flexed to the right, its complicated inner parts large and brownish. Trochanters not tuberculate, femora not ciliate, tibiæ simple, the middle pair with an apical spine, front metatarsi swollen, thicker than their tibiæ and as long as the second to the fourth tarsal joints which are of decreasing length, the fourth joint more than half as long as the flattened fifth joint, posterior tarsi simple. Wings narrow, nearly hyaline, nearly three times as long as wide, sections of the fourth vein proportioned 0.3 : 0.3 : 1 : 1.5, of the fifth vein 1 : 0.7, anal vein short and very faint.

Female. Same, except for the blunt spinose-tipped abdomen and the simple tarsi.

Nine specimens : Almota and Pullman, Washington, May and June.

Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253, 260 (1903); Melander, Williston Man. N. Amer. Dipt. p. 225 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302 (1909); Kertész, Cat. Dipt. Vol. 6, p. 101 (1909); Becker, Wien. Ent. Zeit. Vol. 28, p. 25-28 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 570 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 189 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 43, 77 (1910).

Holoclera, Melander, not Schiner, Trans. Amer. Ent. Soc. Vol. 28, p. 333 (1902).

Schistostoma, Becker, Mitteil. Zool. Mus. Berl. Vol. 2, p. 46 (1902); Wien. Ent. Zeit. Vol. 28, p. 27 (1909); Kertész, Cat. Dipt. Vol. 6, p. 101 (1909).

Characters. — Small usually opaque black hunchbacked species, the males with a twisted epipygium and large contiguous eyes. Head rather globular, the occiput flattened, eyes bare, of the female widely separated, of the male contiguous along the front, the facets of the upper two-thirds larger than below; antennæ inserted rather below the middle of the head, three-jointed, the middle joint typically with strong setæ above and below, the third joint compressed conical or more or less abruptly attenuated beyond the base, with a lengthened terminal arista which varies from one-half to three times as long as the third joint; proboscis usually retracted within the oral cavity, sometimes more or less protruding when the fleshy labium can be seen surrounding the chitinized piercing parts, palpi minute, cylindrical; bristles of the occiput not strong. Thorax greatly arched, robust, about as long as the abdomen, more or less bristly, with two or four rows of acrostichals, a row of dorsocentrals, some post-humerals and supraalars, as well as the usual humeral, notopleural, postalar and scutellar bristles, the last mentioned varying from four to eight in number; pleuræ bare. Abdomen stout, cylindrical, clothed with long thin hairs in the male, epipygium twisted to the right, distorted, often very large, sometimes the last ventral segments of the male are spinose; no distinct ovipositor, but the last segments of the female abdomen compressed or tubular and shining. Legs short, often the femora and tibiæ pectinate with bristles, and in the male, variously furnished with bristles which are lacking in the female. Fringe of the calypteres prominent. Wings large, anal angle full, costa encompassing the entire wing, stigma usually distinct, auxiliary vein distinct from the first and terminating in the costa, basal cells minute, discal cell large, complete, third vein simple, anal crossvein received, forming a jog in the under side of the anal cell, anal vein completely wanting, no alula but the alular cilia long.

Type species : *M. velutinus*, Meigen (Pl. 7, Fig. 63), by Rondani's designation, 1856. Westwood indicated *Trichina clavipes* as the type, but Macquart did not have this species when he erected the genus *Microphorus*. The genus is restricted to the Northern Hemisphere. The species are not rare in shady woods where they are supposed to prey on smaller insects. Beling has described the earlier stages of *M. pusillus*. The sort of sexual dimorphism exhibited by this genus is curious. Here and there on the body, such as on the middle trochanters and femora and on the under side of the abdomen, the males of some species have developed characteristic blunt spinous bristles. Again in other species the hind tibiæ and metatarsi of the males are swollen, but are normal in the females.

SUBGENUS SCHISTOSTOMA, BECKER

Characters. — Proboscis very short, not protruding, vertical, fleshy, palpi short, cylindrical, furnished with five or six setæ; second joint of the antennæ not setose, only some short setulæ present, third joint conical, longer than its arista; front of female with two fronto-orbitals. Only the posterior two or three dorso-centrals lengthened, one posthumeral, one notopleural, the setulæ weak, two scutellars. Pygidium with several long filamentous appendages, ovipositor short and tubular. Alula somewhat distinct, calypteres large, no basal bristle on the costa.

Type species : *S. eremita*, Becker (Pl. 7, Fig. 62). This genus was proposed by Becker for a whitish pruinose Egyptian fly occurring on the sands of the seashore. At first sight its whitish color, light colored legs and pure white bristles were far from suggesting the black species of *Microphorus*, but as recently a white haired *Microphorus* and a black *Schistostoma* have been found, Becker (1909) concluded that *Schistostoma* is but a subgenus of the older *Microphorus*.

SYNOPSIS OF THE NORTH AMERICAN SPECIES OF MICROPHORUS

- | | |
|--|----------------------------|
| 1. Middle trochanter δ of the males with one or two strong spine-like bristles, middle femora δ with some long stiff flexor bristles, middle tibiæ δ more or less excised within before the tip; acrostichals biseriate; fifth ventral δ with apical spines. | 2. |
| Middle trochanters and femora not armed beneath with long stiff bristles, middle tibiæ straight; fifth ventral not with projecting spines. | 5. |
| 2. Hind tibiæ δ clavate and strongly pectinate, hind metatarsi δ much swollen; size about 3 mm. | 3. |
| Hind tibiæ not clavate and not strongly pectinate, hind metatarsi not swollen; size about 2 mm. | 4 |
| 3. Fourth and fifth ventrals δ each with apical spines; middle femora δ with five or six flexor bristles; halteres yellow. | M. ROBUSTUS, nov. sp. (1). |
| Fifth ventral δ with apical spines; middle femora δ with two flexor bristles and one sub-basal arising from the anterior side, hind metatarsi δ pectinate above, halteres black | M. ARMIPES, nov. sp. (2). |
| 4. Third antennal joint conical, as long as its arista; middle femora δ with a group of three strong flexor bristles, inner side of middle tibiæ δ with a bristly callous at three-fifths the tibial length; female cinereous | M. RAVIDUS, Coquillett. |

(1) ***Microphorus robustus*, nov. sp.** — Male. Length 3 mm. Black, coated with opaque brown pollen, halteres brownish yellow. Upper facets enlarged; third joint of the antennæ one-half longer than deep, conical with concave sides, no constricted process, arista twice as long as the third joint; mouthparts not projecting. Thoracic bristles rather sparse, acrostichals biseriate, five pairs in number, about nine dorsocentrals, all but the last two short, two posthumerals, two small supra-alars, four scutellars. Hairs of the abdomen moderately long and black, fourth and fifth ventral segments with a double spine-like bristle on each side of the apical margin, pygidium rather small, twisted. Middle trochanters with a stiff double bristle, front femora with fine long cilia below, middle femora with about six spinous flexor bristles, hind femora with short pectination above and below, middle tibiæ distorted on the apical half, the inner side broadly excised, hind tibiæ clavate and exteriorly pectinate, hind metatarsi globose but not pectinate. Wings dark, stigma distinct, sections of the fourth vein proportioned 1 : 2 : 3, of the fifth vein equal.

Type received from E. T. Cresson, Jr. and deposited in the Philadelphia Academy of Science. Swarthmore, Pennsylvania, June 1, 1909.

(2) ***Microphorus armipes*, nov. sp.** — Male. Length 2.9 mm. Black, including all the appendages. Upper facets of male enlarged; third joint of the antennæ suddenly constricted beyond the base, the constricted portion as long as the basal portion, arista twice as long as the third joint; mouthparts not projecting. Mesonotum lightly coated with brown, subshining, bristles long, the acrostichals biseriate, about seven pairs in number, eight to ten dorsocentrals, one posthumeral, two supra-alar, four scutellars. Abdomen subshining, its hairs sparse, long and black, fifth ventral with four apical spinous bristles; pygidium twisted, penis furcate. Middle trochanters with a double long stiff bristle front femora pectinate below, hind femora pectinate above and below, middle femora with two long strong flexor bristles, near the middle, the distal one blunt and shorter than the proximal, a shorter subbasal bristle arising from the anterior side, hind tibiæ clavate, strongly pectinate along the extensor edge, hind metatarsi swollen and likewise pectinate. Wings darkly infumated, stigma distinct, sections of the fourth vein proportioned 1 : 2 : 3, of the fifth vein equal.

Female. Bristles short, legs not pectinate nor spinose.

Several specimens from Quilcene, Olga and Seattle, Washington, the last mentioned received from Professor Kincaid

- Third antennal joint strongly constricted beyond its base, much shorter than its arista; middle femora ♂ with about six bristles distributed along the under side, middle tibiae not callous; female black M. *SYCOPANTOR*, Melander.
5. Fourth ventral segment ♂ posteriorly expanded into a bilobate process which bears spinous bristles; facets ♂ all moderately small; middle tibiae ♂ with a subapical spur; acrostichals biseriate M. *ISOMMATUS*, nov. sp. (1).
- Ventral segments not thus developed, no ventral spines; upper facets ♂ larger; acrostichals quadriseriate 6.
6. Proboscis projecting forward; posthumerals sparse; third antennal joint conical and longer than its arista; wings hyaline 7.
- Proboscis very short; posthumerals ♂ abundant; third antennal joint suddenly attenuated beyond the base and much shorter than the arista; wings dark M. *ATRATUS*, Coquillett.
7. Thorax ♂ velvety, abdomen ♂ with yellow hairs; hind tibiae and metatarsi ♂ dilated. M. *OBSCURUS*, Coquillett.
- Thorax ♂ with two shining vittæ; hairs of the abdomen black; legs not dilated M. *BILINEATUS*, Melander.

Geographical distribution.

SUBGENUS MICROPHORUS, MACQUART

1. *M. albopilosus*, Becker, Deutsche Ent. Zeitschr. 1910, 644 (1910). Corsica.
2. *M. anomalus*, Meigen, Syst. Beschr. Vol. 4, p. 9 [1824] (*Platypesa*); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 43 (1864); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 92 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 192, f. 73, 74 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 78 (1910); Frey, Acta Soc. Sc. Fenn, Helsingfors, Vol. 37 (3), p. 59 (1913).
- crassipes*, Macquart, Dipt. N. France, Vol. 3, p. 140, pl. 4, f. 1 [1827] (*Microphor*); Hist. Nat. Dipt. Vol. 1, p. 346, pl. 8, f. 7 (1834); Meigen, Syst. Beschr. Vol. 7, p. 78 [1838] (*Trichina*); Boitard, Nouv. Man. Vol. 3, p. 322 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 488 (1849); Ins. Brit. Dipt. Vol. 1, p. 114 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 79 (1862).

(1) ***Microphorus isommatus*, nov. sp. — Male.** Length 2 mm. Entirely black, with a brownish tinge, halteres white. Eyes subcontiguous, touching along the middle of the front, the upper facets scarcely larger than the lower, third joint of the antennæ conical, broad at the base, scarcely longer than wide, not produced as an attenuated portion, its sides slightly concave, arista twice as long as the third joint; mouthparts retracted, only the tip normally exposed; ocellar triangle not prominent. Thoracic bristles sparse, acrostichals biseriate, about five pairs in number, about ten small dorso-centrals, two posthumerals, three supra-alaris and four scutellars. Hairs of the abdomen short, sparse and black, fourth ventral segment polished, broadly emarginate in the middle and produced backward on each side as a large convex lobe whose inner margin is set with six stiff spine-like bristles; pygidium large and twisted, penis ending in two claws. Legs simple, not spinose or thickened, pubescence black, front tibiae with loose extensor cilia, hind femora with a row of twelve bristles beneath which are not longer than the diameter of the femur, middle tibiae with a short apical spur and middle metatarsi with a corresponding spur at the base. Wings dark, rather broad, stigma distinct, third vein sinuous, strongly diverging from the second at the tip, sections of the fourth vein proportioned 0.5 : 0.5 : 1 : 1.5, of the fifth vein equal.

Female. Similar except for the separated eyes and the different genitalia, last three segments of the abdomen simple, broadly compressed, shining.

Several specimens; London Hill Mine, Bear Lake, British Columbia, altitude 7000 feet, July 21, 1903, collected by J. W. Cockle. Type in the U. S. National Museum.

pusillus, Macquart, Dipt. N. France, Vol. 3, p. 140 [1827] (*Microphor*); Hist. Nat. Dipt. Vol. 1, p. 346 (1834); Meigen, Syst. Beschr. Vol. 7, p. 78 [1838] (*Trichina*); Zetterstedt, Dipt. Scand. Vol. 1, p. 256 [1842] (*Microphora*); Vol. 8, p. 2999 [1849] (*Microphora*); Walker, Ins. Brit. Dipt. Vol. 1, p. 115 (1851); Zetterstedt, Dipt. Scand. Vol. 12, p. 4600 [1855] (*Microphora*); Beling, Arch. Naturg. Berlin, Vol. 48 (1), p. 212 (1882).
tarsellus, Zetterstedt, Dipt. Scand. Vol. 1, p. 257 [1842] (*Microphora*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 50 [1851] (*Microphora*); Zetterstedt, Dipt. Scand. Vol. 13, p. 4982 [1859] (*Microphora*).

- | | |
|---|-------------------------------|
| 3. <i>M. armipes</i> , nov. sp. | Washington. |
| 4. <i>M. atratus</i> , Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 412 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 333 [1902] (<i>Holoclera</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 (1903). | Alaska. |
| 5. <i>M. Beckeri</i> , Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 72 (1909). | C. Europe. |
| 6. <i>M. bilineatus</i> , Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 334, f. 99 [1902] (<i>Holoclera</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 (1903). | Louisiana. |
| 7. <i>M. dalmatinus</i> , Strobl, Glasn. Mus. Bosn. Herzeg, Sarajewo, Vol. 14, p. 468 (1902); Mitth. Bosn. Herzeg, Sarajewo, Vol. 9, p. 527 (1904). | S. Europe. |
| 8. <i>M. defunctus</i> , Handlirsch, Contrib. Can. Palæoz. Ins. Vol. 2, p. 124, f. 32, 33 (1910). | Oligocene : British Columbia. |
| 9. <i>M. drapetoides</i> , Walker, List Dipt. Brit. Mus. Vol. 3, p. 489 (1849); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 332 (1902). | Hudson's Bay. |
| 10. <i>M. eocenicus</i> , Meunier, Ann. Soc. Sc. Nat. Bruxelles (8), Vol. 16, p. 402, pl. 2, f. 7, 8 [1902] (<i>Holoclera</i>); Handlirsch, Foss. Ins. p. 1016 [1906] (<i>Holoclera</i>); Meunier, Ann. Sc. Nat. [Zool.] (9), Vol. 7, p. 85, note, 126 [1908] (<i>Holoclera</i>). | Baltic Amber. |
| 11. <i>M. hiemalis</i> , White, Proc. Roy. Soc Tasmania, 1916, p. 240, f. 47 (1917) | Tasmania. |
| 12. <i>M. isommatus</i> , nov. sp. | British Columbia. |
| 13. <i>M. obscurus</i> , Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 268 (1903). | United States |
| 14. <i>M. pilimanus</i> , Strobl, Wien. Ent. Zeit. Vol. 18, p. 26 (1899); Verh. Zool. Bot. Ges. Wien, Vol. 59, p. 177 (1909). | Spain. |
| 15. <i>M. putidus</i> , Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 94, 126, pl. 7, f. 10, 11 (1908). | Baltic Amber. |
| 16. <i>M. ravidus</i> , Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 409 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 333 [1902] (<i>Holoclera</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 (1903). | W. United States. |
| 17. <i>M. robustus</i> , nov. sp. | Pennsylvania. |
| 18. <i>M. rostellatus</i> , Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 42 (1864); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 177 (1909). | S. Europe. |
| 19. <i>M. scapuliferus</i> , Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 129 (1889); Bezzia, Ann. Mus. Hungar, Vol. 3, p. 458 (1905). | Chile. |
| 20. <i>M. semifulvus</i> , Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 129 (1889); Bezzia, Ann. Mus. Hungar, Vol. 3, p. 458 (1905). | Chile. |
| 21. <i>M. sycophantor</i> , Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 334 [1902] (<i>Holoclera</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263 [1903] (<i>Anthalia</i>). | W. United States. |
| 22. <i>M. velutinus</i> , Macquart, Dipt. N. France, Vol. 3, p. 140 [1827] (<i>Microphor</i>); Hist. Nat. Dipt. Vol. 1, p. 346 (1834); Meigen, Syst. Beschr. Vol. 7, p. 78 [1838] (<i>Trichina</i>); Walker, Ins. Brit. Dipt. Vol. 1, p. 114 (1851); Zetterstedt, Dipt. Scand. Vol. 12, p. 4599 [1855] (<i>Microphora</i>); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 144 [1861] (<i>Microphora</i>); Schiner, Fauna Dipt. Austr. Vol. 1, p. 79 (1862); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 45 (1864); Neuhaus, Dipt. March. p. 69 | Europe. |

DIPTERA

[1886] (*Trichina*); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 133 (1887); Mik, Wien. Ent. Zeit. Vol. 6, p. 100 (1887); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 92 (1892); Bezzia, Bull. Soc. Ent. Ital. Vol. 30, p. 144 (1899); Strobl, Wien. Ent. Zeit. Vol. 18, p. 26 (1899); Lundbeck, Dipt. Dan. Vol. 3, p. 191 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 78, f. 9 (1910); Frey, Acta Soc. Sc. Fenn., Helsingfors, Vol. 37 (3), p. 59 (1913); Boitard, Man. Ent. Vol. 3, p. 322 (1843). — PI. 7, Fig. 63.

fuscipes, Zetterstedt (not 1838, which is *Trichina elongata*), Dipt. Scand. Vol. 11, p. 4268 [1852] (*Microphora*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 71 (1909).

? *holosericeus*, Meigen (not 1838, which is *Rhamphomyia gibba*), Classif. Beschr. Eur. Dipt. Vol. 1, p. 231 [1804] (*Empis*); Syst. Beschr. Vol. 3, p. 58 [1822] (*Rhamphomyia*); Vol. 7, p. 89 (1838) (*Rhamphomyia*); Loew, Bemerk. Posen. Gegend Art Zweifi Gatt. 1840, p. 21, f. 22; Isis, Vol. 7, p. 549 f. 22 [1840] (*Rhamphomyia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 425 [1842] (*Rhamphomyia*); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 105 [1849] (*Rhamphomyia*); Scholz, Zts. Ent. Breslau, Vol. 5 (19), p. 50 (*Microphora*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 100 [1862] (*Rhamphomyia*).

? *rufipes*, Meigen, Syst. Beschr. Vol. 7, p. 78 [1838] (*Trichina*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 79 (1862).

vicus, Mik, Wien. Ent. Zeit. Vol. 6, p. 99 (1887).

SUBGENUS SCHISTOSTOMA, BECKER

1. *Sch. eremita*, Becker, Mitteil. Zool. Mus. Berl. Vol. 2, p. 47, pl. 4, f. 5 Egypt. (1902). — PI. 7, Fig. 62).
2. *Sch. nigrescens*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 121 (1907); Wien. Algeria. Ent. Zeit. Vol. 28, p. 25 (1909).
3. *Sch. truncatus*, Loew, Zeit. Ent. Breslau, Vol. 14, p. 48 [1863] (*Microphorus*); C. Europe. Strobl, Jahrb. Naturk. Landesmus. Kärnten, Klagenfurt, Vol. 47, p. 201 [1901] (*Microphorus*); Becker, Wien. Ent. Zeit. Vol. 28, p. 25 (1909); Strobl, Verh. Zool. bot. Ges. Wien, Vol. 59, p. 177 [1909] (*Microphorus*).
4. *Sch. yakimensis*, nov. sp. (1) Washington.

4. GENUS HORMOPEZA, ZETTERSTEDT

Hormopeza, Zetterstedt, Fauna Ins. Lappon. p. 540 (1838); Dipt. Scand. Vol. 1, p. 257 (1842); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 111 (1862); Bigot, Ann. Soc. Ent. France (6), Vol. 9 p. 118 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 273 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 251, 263 (1903); Melander, Williston, Man. N. Amer. Dipt. p. 224 (1908); Kertész, Cat. Dipt. Vol. 6, p. 83

(1) **Microphorus (Schistostoma) yakimensis**, nov. sp. — Male. — Length 2.3 mm. Body entirely black but thickly overlaid with gray pollen, the conspicuous long hairs and bristles white. Third antennal joint one and one-half times as long as broad and equal to the arista in length; proboscis black, extending obliquely forward but not reaching beyond the strongly vaulted oral opening; palpi with white hairs; occiput densely gray pollinose, the orbital bristles long and dense. Thorax faintly bilobate when viewed from in front, the acrostichals forming a double row, the dorsocentrals numerous, scutellum bisetose. Abdomen gray pollinose except for the seventh sternite and the large reflexed end of the epipygium; the penis and black ventral processes long, as in Becker's figure of *S. eremita*. Legs entirely black though overlaid with gray dust, the bristles white, hind femora and tibiae ciliate. Calypteres white, densely fringed; halteres white. Wings hyaline, the veins brown, stigma very faint, no costal bristle.

One specimen taken in April, 1915, at Yakima, Washington.

(1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 553 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 65 (1910).

Characters. — Blackish species measuring three or four millimeters in length, not closely related to any of the other genera. Head almost globular, the occiput rather flattened, eyes bare, of the male broadly contiguous along the front, the ocellar triangle elevated, the upper facets large, of the female widely separated, the front quadrate and bearing six minute fronto-orbitals, face broadly receding but wide, especially in the male; antennæ located below the middle of the head, apparently two-jointed since the basal joints are completely fused into a small globular nearly bare segment, the last joint compressed and broadly orbicular, ending in a short two-jointed style, whose basal joint is quadrate and whose outer joint consists of a short stubby bristle; proboscis short and thick, smaller and retracted in the male, protruding obliquely forward in the female, palpi short, horizontal and gradually thickened; occiput not bristly above but seriatel hairy, one pair of ocellar bristles. Thorax markedly bristly, about four humerals, three posthumerals, five notopleurals and several scutellars, the dorsocentrals numerous, acrostichals biseriate, quite a number of lateral setulæ beyond the dorsocentral rows; metapleuræ bare, pectus with some bristles at the base of the coxae. Abdomen of the female subequal to the thorax in length, of the male one-half longer, eighth segment of the female broadly conical, almost wedge-shaped and strong; pygidium robust, globose, the dorsal valves large and erect, the lateral valves rather hemispherical and with two blunt posterior prongs, penis hidden. Legs strongly bristly, anterior femora of the female with two flexor rows of bristles, tibiæ of both sexes with several sets of bristles, tarsal joints more or less setose, pulvilli large. Wings not colored, rather pointed, anal angle strongly rectangular, costa stopping at the tip of the wing, auxiliary vein distinct and extending into the costa, a thickened stigma between the ends of the auxiliary and first veins, third vein with a long fork, discal cell located before the middle of the wing, the basal and discal cells not large, the sections of the fifth vein nearly 1 : 2, anal crossvein reflexed less abruptly than usual, angulate with the anal vein, the latter confluent with the under side of the anal cell, second basal and anal cells of equal width, a small alula present; calypters with a strong fringe.

Type species: *H. oblitterata*, the only species occurring in Europe. On many occasions I have taken specimens of *Hormopeza* on the window-panes in houses. The affinities of this genus are not clear. In 1908, when stressing the importance of the form of the anal cell, I located *Hormopeza* in the Ocydromiinae, in which group it would be distinct in having the third vein furcate. It seems more likely that the genus is a specialized but ancient offshoot of the Oreogoton branch of the Empidinæ.

KEY TO THE SPECIES OF HORMOPEZA

- Eyes of male separated; joints of anterior tarsi rounded at ends;
basal dorsal process of pygidium rather wide and erect *H. SENATOR*, nov. sp. (1).
- Eyes of male broadly contiguous 2.

(1) ***Hormopeza senator*, nov. sp.** — Male. Length 3 mm. Black, head and thorax entirely covered with cinereous pollen; abdomen shining; first and most of second segment lightly cinereous pollinose, extreme base of segments three and four similarly pollinose. Eyes narrowly separated on the front, the facets uniform, antennæ brown, the third joint short, ovate; proboscis short, black, together with the yellow palpi reaching forward; occipital hairs pale yellow. Thoracic hairs yellow; hairs of abdomen pale; pygidium lightly pollinose, its ventral setæ yellow and apical hairs brown, dorsal process wider than usual and erect, side valves terminating in a very large, oblong process whose lower apical angle is rounded, above at the base of this process the side valve is continued inward as a short, robust tooth. Legs brown, the coxae testaceous, joints of anterior tarsi rounded, oblong; pulvilli whitish, one-third as long as last tarsal joint; bristles of hind tibiæ blackish; hind femora with fringe of short setæ above. Wings hyaline, veins thin and pale, apex of costal cell scarcely differentiated.

Holotype, Washington D. C., 17 August, 1913.

2. Third antennal joint pear-shaped, nearly twice as long as wide; joints of anterior tarsi rounded-oblong; dorsal process of pygidium vertically erect, side valves flattened above and with a rectangularly indent tip H. OBLITERATA, Zetterstedt.
- Third antennal joint short ovate; tarsal joints oblong; dorsal process of pygidium bent posteriorly 3.
3. Abdomen uniformly though lightly cinereous pollinose; dorsocentrals blackish; front of female narrowest just above antennae; pulvilli one-half (male) or one-fourth (female) the length of the last tarsal joint 4.
- Abdomen polished, at most with faint basal bands of pollen on the segments; dorsocentrals yellow; front of female narrowest midway to the ocelli; pulvilli two-thirds the length of the last tarsal joint. 5.
4. Pygidial valves terminated below by an elongate narrow arm and above by a strong quadrate process H. COPULIFERA, nov. sp. (1).
- Pygidial valves lacking the lower narrow arms, but with a long slender erect process arising from base of the L-shaped terminal piece H. VIRGATOR, nov. sp. (2).
5. Stigma and veins fuscous; halteres blackish; legs black except apex of coxae and knees. No stigma, veins thin and pale; halteres and legs yellow to brown 6.
6. Robust, 4.5 mm.; abdomen shining black; third antennal joint broadly oval. H. BULLATA, Melander.
- Smaller, 3-4 mm.; abdomen often brownish-black and with bands of pollen across base of segments; third antennal joint subrotund;

(1) *Hormopeza copulifera*, nov. sp. — Male. Length 4 to 5 mm. Black, eyes contiguous, third antennal joint round, occipital hairs white. Thorax completely coated with cinereous pollen, the prominent setæ of middle of notum blackish, lateral and posterior bristles pale. Abdomen lightly covered with gray pollen, its hairs pale; pygidium mostly shining, lightly pollinose below, with strong convex lateral valves terminating below in a narrow, incurved, parallel-sided brown arm which bears a few long and short setæ, and above by a heavy quadrate process which is somewhat expanded at the tip, within the last mentioned process the valve bears a short but strong conical projection, hairs and setæ of pygidium prominent and blackish. Legs simple, hairs pale in the femoral crests, elsewhere mostly dark, all tibiae bearing dark bristles, pulvilli dusky, about half the length of the last tarsal joint. Knob of halteres blackish, wings cinereous, almost hyaline, veins thin but dark, distal portion of costal cell blackish; sections of fifth vein 2 : 3.

Female. Eyes widely separated, front narrowest above antennæ, ovipositor retracted in the tubular eighth segment. Types taken from a window in a sawmill at Coeur d'Alene, Idaho, 24 August, 1916. Paratypes from Douglas, Alaska (E. L. Jenne), Friday Harbor, Washington (J. M. Aldrich), Banff, Alberta (C. B. D. Garrett); and Port Angeles, Olga, Friday Harbor, Spokane, Washington; and Potlatch and Moscow, Idaho (Melander). Most of these specimens were taken on windows.

(2) *Hormopeza virgator*, nov. sp. — Male. Length 3 mm. Black, eyes contiguous, antennæ black, third joint round; occipital hairs pale; thorax entirely covered with cinereous dust, hairs of middle of notum blackish, lateral and posterior bristles pale; abdomen lightly coated with pollen, subshining, its long, fine hairs white; pygidial valves polished, terminating in a broadly L-shaped inflexed flap from the base of which arises a slender, erect, long prong, a few long dark setæ on lateral valves. Legs simple, hairs yellow, hind femora ciliate above, bristles of hind tibiae black, pulvilli gray less than one-half as long as last tarsal joint. Halteres blackish; wings narrow, nearly hyaline, veins thin and brown, distal portion of costal cell brownish-gray, sections of fifth vein proportioned 5 : 6.

Female. Eyes widely separated, the sides of the front diverging from the antennæ. Last abdominal segment retracted.

Types, Pullman, Washington, 11 September, 1908, taken from a window. Twenty paratypes from Spokane, Washington, and from Potlatch, Chatcolet, and Moscow, Idaho, the last collected by J. M. Aldrich. Most of the paratypes were taken from window-panes, August to October.

pygidial valves terminated by large quadrate infolding flaps, bearing long yellow setæ behind which crisscross beneath the pygidium. H. BREVICORNIS, Loew.

Geographical distribution.

1. *H. brevicornis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 83 : Cent. 5, No. 65 W. North America.
 (1864); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 273, f. 87
 (1902). — **Pl. 8, Fig. 76.**

2. *H. bullata*, Melander, ibidem, Vol. 28, p. 274, f. 88 (1902). Wyoming.

3. *H. copulifera*, nov. sp. W. North America.

4. *H. nigricans*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 83 : Cent. 5, No. 66 W. North America.
 (1864); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 275 (1902).

5. *H. oblitterata*, Zetterstedt, Lapp. p. 540 (1838); Dipt. Sc. Vol. 1, p. 257 N. Europe.
 (1842), Vol. 8, p. 3000 (1849); Bonsdorff, Finl. tvåv. Ins. Vol. 1,
 p. 145 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 111 (1862);
 Wahlgren, Ent. Tidskr. Vol. 31, p. 65 (1910); Frey, Acta Fenn.
 Vol. 37 (3), p. 45, pl. 1, f. 5 (1913); Collin, Ent. Mo. Mag. Vol. 54,
 p. 278 (1918).

6. *H. senator*, nov. sp. E. United States.

7. *H. virgator*, nov. sp. W. United States.

5. GENUS TIMALPHES, NOV. GEN.

Characters. — Very close to *Oreogeton*, but the metapleuræ are bare and none of the veins is setulose. Strongly setose, rather large species. Eyes of the male broadly contiguous on the front, facets uniform and bare, face wide but short, receding into the oral cavity almost immediately below the antennæ; antennæ about as long as the head, distinctly three-jointed, the basal joints setose, the first joint slightly longer than the second, the third joint conical, a little more than twice as long as wide, its sides somewhat concave, arista terminal, longer than the third joint, thickened, geniculate at its articulation, the basal segment very minute and the apical seta microscopic; proboscis horizontally porrect, thick, palpi likewise porrect, thick and cylindrical and strongly setose; occiput with abundant hairs which become biseriate above as the vertical bristles, ocellar bristles four in number and small. Thorax robust, no pubescence, one humeral, one posthumeral, three notopleurals, several supraalars, four scutellars, about ten dorsocentrals, the foremost placed outside of the row, acrostichals strong and biseriate; pleuræ entirely bare. Abdomen strongly cylindrical, somewhat bent downward, abundantly hairy, the basal segments marked with basal transverse, lateral longitudinal and subdiscal oblique series of pittings; pygidium small, the lateral valves with a forcipate pair of posterior prongs, dorsal valves small and hairy, penis hidden. Legs simple, hairy and bristly, femora and hind tibiæ more or less distinctly ciliate with bristles, anterior tibiæ with several sets of extensor and apical bristles, pulvilli large. Wings with nearly parallel sides, anal lobe rectangular, a distinct axillar angle, costa continuing around the entire wing, auxiliary vein complete, ending in the costa at the middle of the wing, third vein with a long fork, sections of the fifth vein subequal, basal cells equal and shorter than the blunt discal cell, anal crossvein abruptly reflexed, one-third as long as the anal cell, anal vein not continuous with the anal cell, extending almost to the margin; alula very weak, calypteres lobose, margined and with a prominent fringe.

Type species: *T. fumosa*, Hutton (Pl. 6, Fig. 53), a species described as a *Clinocera*. It seems quite likely that Meunier's fossil *Brachystoma gracilis* from the Lower Oligocene belongs to this genus.

However, the illustration of the head drawn by Mrs. Meunier suggests a dipteric insect. Loew's species of *Gloma* with slender antennæ, likewise from the Baltic amber, possibly belong here.

Geographical distribution.

1. *T. * ? acuticornis*, Loew, Bernsteinfauna, p. 41 [1850] (*Gloma*); Giebel, Ins. Baltic Amber. Vorwelt, p. 208 [1856] (*Gloma*).
2. *T. fumosa*, Hutton, Trans. New. Zeal. Inst. Vol. 33, p. 31 [1901] (*Clinocera*); New Zealand. Bezzi, Ann. Mus. Hungar. Vol. 2, p. 321, note [1904] (*Heleodromia*); Kertész, Cat. Dipt. Vol. 6, p. 123 [1909] (*Heleodromia*); Engel, Deutsche, Ent. Zeitschr. 1918, p. 47 [1918] (*Hydrodromia*). — **Pl. 6, Fig. 53.**
3. *T. * ? gracilis*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 113, pl. 8, Baltic Amber. f. 10, Vol. 9, p. 2-4 [1908] (*Brachystoma*).
4. *T. * ? palpata*, Loew, Bernsteinfauna, p. 41 [1850] (*Gloma*); Giebel, Ins. Baltic Amber. Voorwelt, p. 208 [1856] (*Gloma*).

6. GENUS OREOGETON, SCHINER

Oreogeton, Schiner, Wien. Ent. Monatschr. Vol. 4, p. 53 (1860); Fauna Dipt. Austr. Vol. 1, p. 112 (1862); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 116 (1889); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 275 (1902); Coquillet, Proc. Ent. Soc. Wash. Vol. 5, p. 254, 262 (1903); Bezzı, Ann. Mus. Hungar. Vol. 3, p. 451 (1905); Melander, Williston, Man. N. Amer. Dipt. p. 225 (1908); Kertész, Cat. Dipt. Vol. 6, p. 80 (1909); Coquillet, Proc. U. S. Nat. Mus. Vol. 37, p. 580 (1910).

Characters. — Rather large black or yellow more or less pollinose species somewhat resembling *Leptis* in appearance. Head small, eyes of the male contiguous on the front but widely separated below, facets nearly uniform, bare; antennæ inserted below the middle of the head, the face small and quadrate, receding below as the pollinose broad epistome; basal joints of the antennæ short and setose, third joint conical, terminated by a thick arista longer than the third joint and tipped with a short hair; proboscis very short and fleshy, directed forward, palpi long, clavate, hairy, porrect; occipital hairs abundant, not differentiated into bristles above; ocellar triangle of the male a little elevated. Thorax bristly, dorsocentrals numerous, several humerals, posthumerals, notopleurals, supra-alars and scutellars, the acrostichals biseriate; pleurae pollinose, metapleuræ setose. Abdomen hairy and bristly, twice as long as the thorax, rather slender and cylindrical, pygidium not large, bilateral, lateral valves oblong, pointed at the upper apex and bearing a dorsal tooth, dorsal valves small and cerciform; abdomen of the female somewhat depressed, tapering at the apex. Legs slender, setose, in the male the middle pair sometimes ornate with bunches of bristles, the tibiae bearing sets of apical as well as intermediate bristles, pulvilli large. Wings large, veins strong, the first, second and sometimes the third veins bearing setulae on the upper surface and the second setulose on the lower side, the setulae of the first vein dense, costa continuing around the entire wing but weakened on the hind margin, auxiliary vein distinct, ending in the costa, third vein forked, the upper branch short and oblique, basal cells shorter than the apically blunt discal cell, anal crossvein abruptly reflexed, continuous with the underside of the anal cell, anal vein obsolete, anal angle of the wing strongly rectangular, alula distinct; calypteres large, margined and densely fringed.

Type species: *O. basalis*, Loew (Pl. 5, Fig. 50). There is a tendency in the genus for the base of the abdomen to be pellucid and for the females to be yellowish in color. In neuration and body-color

the species are quite constant. The following table shows the variations in arrangement of the special bristles of the males. Females are more difficult to determine specifically than the males. The male of *rufus* is not known and the species is therefore omitted from the table. Both sexes of *obscurus* are dark, and the species is thus distinct from *rufus*.

KEY TO THE SPECIES OF OREOGETON, BASED ON MALES

1. Front tarsi with last two joints flat, black and densely fringed with flat setæ	2.
Front tarsi without expanded and pennate tip	3.
2. Front tibiæ with dense cluster of black setæ around tip	O. SCOPIFER, Coquillett.
Front tibiæ not setose around tip	O. CYMBALLISTA, nov. sp. (1).
3. Some of the flexor setæ near middle of middle tibiæ projecting far beyond the others	4.
Middle tibiæ nearly uniformly setose	O. BASALIS, Loew.
4. Third vein setulose above; central seta of middle femora stronger than the others; arista arising from above apex of antenna.	O. CAPNOPTERUS, nov. sp. (2).
Third vein bare, rarely with one or two setulae; more than one long seta under middle femora; arista terminal	5.
5. Three or four outstanding setæ under middle femora; setæ of middle coxæ pointed	6.
About eight outstanding setæ under middle femora; apical setæ of middle coxæ strong and blunt	O. MITREPHORUS, nov. sp. (3).

(1) **Oreogeton cymballista**, nov. sp. — Male. Length 7 mm. Head and thorax black, base of abdomen pellucid yellow, remainder of abdomen piceous, legs, halteres and base of wings yellow. Last two joints of front tarsi flat, forming a black disk heavily fringed with black scale-like setæ, last two joints of posterior tarsi likewise contrastingly black but not discoid, middle femora with five very long setæ near middle beneath and with a comb of seven shorter setæ toward knee along postero-flexor face, middle tibiæ bent, swollen at three-fifths their length where they bear a dense brush of flexor setulæ, also seriatel setulose toward knee beneath, along extensor side with five scattered long setæ. Wings with reddish brown tinge.

Holotype, Nelson, British Columbia, 17 July, 1910 (Melander).

(2) **Oreogeton capnopterus**, nov. sp. — Male. Length 5.5 mm. Head black, thorax piceous, base of abdomen pellucid yellow, remainder of abdomen piceous, coxæ and anterior legs yellow, hind legs brown, halteres with yellow stalk and piceous knob, calypteres yellow. Last two joints of tarsi small, blackish but not contrasting with remainder of legs; central flexor seta of middle femora projecting beyond the others, the next one distal to it nearly as long, followed by six shorter not crowded setæ toward the knee; middle tibiæ straight, finely and nearly uniformly setulose and with an extensor seta at three-fifths the length and a small pair near knee. Wings with blackish smoky tinge, third vein setulose above. The arista is attached just above the apex of the antennæ.

Female. Occiput, front, third antennal joint and last two tarsal joints black, otherwise the insect is yellow, becoming brownish on posterior part of abdomen. Femora without specialized setæ. Wings less infumated. Arista arising from tip of antennæ.

Four males, one female, south side of Mt. Rainier, Washington, August, 3500-5500 feet altitude (Melander); 1 male Banff, Alberta, 12 July, 1922 (Garrett), in Canadian National Museum.

(3) **Oreogeton mitraphorus**, nov. sp. (PI. 6, FIG. 51.) — Male. Length 5.5 mm. Head and thorax black, the sutures rufous, second and third segments of abdomen pellucid yellow, remainder piceous, legs, halteres, and base of wing yellowish. Tarsi apically blackened, not expanded or pennate; middle coxæ with five long blunt apical setæ; middle femora with about eight long flexor setæ extending from before the middle almost to knee; middle tibiæ almost straight, very slightly swollen at two-thirds the length where there are a few erect flexor setulæ; one extensor seta near knee, two near middle and two near apex of middle tibiæ; apical setæ of hind coxæ small, hind femora more or less ciliate above and below. Wings with yellowish tinge, third vein not setulose.

Female. Somewhat darker than male, only the second abdominal segment yellowish, coxæ piceous, femora more or less brown, specialized leg bristles absent, third vein setulose.

Five males, thirty-six females. Moscow Mountain and Lookout Mountain, Idaho, June-August (Melander).

6. Hind coxae with long setæ reaching nearly to end of trochanters;
 females yellow O. HETEROGAMUS, nov. sp. (1).
 Setæ of hind coxae much shorter than trochanters; females blackish. O. OBSCURUS, Loew.

Geographical distribution.

1. *O. basalis*, Loew, Neue Beitr. Dipt. Pt. 4, p. 36 [1856] (*Gloma*); Schiner, C. Europe.
 Fauna Dipt. Austr. Vol. 1, p. 112 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 133 (1887); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 87 (1893). — PI. 5, Fig. 50.
2. *O. capnopterus*, nov. sp. W. North America.
 3. *O. cymballista*, nov. sp. British Columbia.
 4. *O. heterogamus*, nov. sp. Oregon.
 5. *O. mitraphorus*, nov. sp. (PI. 6, Fig. 51). Idaho.
 6. *O. obscurus*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 84 : Cent. Vol. 5, p. 63 [1864] (*Gloma*); Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 412 [1900] (*Gloma*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 276, f. 93 [1902] (*Gloma*); Coquillett, Harriman Exped. Vol. 9 (2), p. 26 [1904] (*Gloma*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 451 (1905). E. United States; Alaska.
7. *O. rufus*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 84 : Cent. Vol. 5, p. 67 [1864] (*Gloma*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 275 [1902] (*Gloma*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 451 (1905). E. United States.
8. *O. scopifer*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 412 [1900] (*Gloma*); Alaska. Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 276 [1902] (*Gloma*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 451 (1905); Coquillett, Harriman Exped. Vol. 9 (2), p. 26 [1904] (*Gloma*).

7. GENUS APALOCNEMIS, PHILIPPI

Apalocnemis, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 752 (1865); Bezzi, Ann. Mus. Hungar., Vol. 3, p. 424, 451 (1905); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 117 (1889); Melander, Williston, Man. N. Amer. Dipt. p. 226 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 383 (1909); Kertész, Cat. Dipt. Vol. 6, p. 111 (1909).

Characters. — Small reddish or blackish species measuring three to five millimeters. Head globose, occiput hairy, eyes of the male contiguous on the front; antennæ short, basal joints brief and nearly bare, third joint rotund and short, with a lengthened terminal arista which is obviously longer than the antenna and has a distinct basal segment and an apical seta; proboscis very short, horizontal but not protruding beyond the oral margin, palpi large and briefly hairy. Thorax strong, bristly, a cluster of humerals, several notopleural, scutellar and supra-alar bristles, acrostichals short or long, dorsocentral series complete, the posterior bristles long; pleuræ bare. Abdomen rather cylindrical, pygidium small, bearing four small dorsal appendages. Legs slender, simple, short-hairy, only the male

(1) **Oreogoton heterogamus**, nov. sp. — Male. Length 5.5 mm. Head and thorax black, second and third abdominal segments pellucid yellowish. Last two tarsal joints black, of the anterior legs slightly expanded but not pennate; middle femora with four long flexor setæ near middle, then a short interrupted space followed by four seriate setæ toward knee; middle tibiae bent at middle and with a few erect flexor setulae at two-thirds the length, exteriorly with two setæ; apical setæ of hind coxae long, curved, reaching nearly to end of trochanters; hind femora ciliate above and below with fine setæ. Wings with light yellowish tinge, third vein bare. The thoracic bristles are long.

Female. Occiput, front, third antennal joint and last two tarsal joints black, remainder testaceous yellow, abdomen light brown on apical one-half, the ovipositor yellowish. Specialized leg bristles absent. Third vein setulose. Two males, three females. Near the mouth of Eagle Creek, Cascade Mountains, Oregon. July 1, 1917 (Melander).

sometimes with femoral bristles. Anal angle well developed, costa encompassing the entire wing, basal bristle present, stigma present or absent, third vein acutely forked, the anterior branch not short, discal cell rather small, anal vein obsolete or thin, not attaining the margin.

Type species: *A. obscura*, Philippi (Pl. 6, Fig. 56), the original species. The genus is related to *Glooma* but differs in the structure of the antennæ. It appears to be restricted to the West coast of South America.

Geographical distribution.

1. *A. breviventris*, Philippi, Verh. Zool.-bot. Wien, Vol. 15, p. 762 [1865] Chile.
(? *Hilara*); Bezzia, Ann. Mus. Hungar, Vol. 3, p. 444 (1905).
2. *A. cingulata*, Bezzia, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 383, f. 9 Peru.
(1909).
3. *A. obscura*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 753, pl. 29, Chile.
f. 55 (1865); Bezzia, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 383
(1909). — Pl. 6, Fig. 56.
4. *A. variegata*, Bezzia, Ann. Mus. Hungar, Vol. 3, p. 450, f. 2 (1905); Nova Peru.
Acta Akad. Naturf. Halle, Vol. 91, p. 383 (1909).

8. GENUS GLOOMA, MEIGEN

Glooma, Meigen, Syst. Beschr. Vol. 3, p. 14 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 345 (1834); Westwood, Gen. Syn. p. 132 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 439 (1842); Boitard, Man. Ent. Vol. 3, p. 321 (1843); Rondani, Dipt. Ital. Vol. 1, p. 149 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Lioy, Atti Inst. Sc. Veneto, Venezia, 1864, p. 602 (1864); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 131 (1887); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 120 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 250, 263 (1903); Bezzia, Ann. Mus. Hungar, Vol. 3, p. 451 (1904); Kertész, Cat. Dipt. Vol. 6, p. 111 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 85 (1910).

Characters. — Opaque blackish species of about four millimeters length. Eyes bare, of the male contiguous on the front, the upper facets enlarged, of the female widely separated, ocellar triangle of the male prominent, located on the vertex, ocellar bristles long, no fronto-orbitals, face very short and broad, receding into the epistome, the mouth-opening strongly arched, occiput at least of the male with abundant hairs; antennæ shorter than the head, apparently two-jointed due to the close growth of the last two joints, the first joint small, the second cyathiform, with a crown of long setæ, the third joint reniform, bent down at the end and bearing a basal dorsal slender two-jointed arista, which is nearly three times as long as the antenna and is geniculate at the end of its first segment; proboscis short, fleshy, thick, horizontal, scarcely protruding from the oral opening, palpi short, cylindrical and with short hairs. Thorax not convex, devoid of pubescence, bristles long, one humeral, two posthumerals, seven or more dorsocentrals, three notopleurals, four or six scutellars, the acrostichals long and arranged as an irregularly biserrate median row; pleuræ bare. Abdomen twice as long as the thorax, hairy, especially in the male, pygidium small, directed backward, complex with small processes and provided with two short basal prongs and two trumpet-shaped posterior processes; abdomen of the female with a broad blunt termination. Legs slender, simple, in the male more elongate, more hairy, the hind femora ciliate and the hind tibiæ and metatarsi lengthened and slightly swollen, in the female certain of the extensor hairs of the tibiæ almost bristle-like. Wings very broad, the anal angle prominent,

costa stopping at the third vein, auxiliary vein distinct, straight, vanishing before entering the costa, third vein forked, discal cell of moderate size, located a little before the middle part of the wing, anal crossvein abruptly recurved and continuous with the underside of the anal cell, anal vein obsolete, alula scarcely noticeable and with a close fringe; calypteres large, strongly margined.

Type species: *G. fuscipennis*, Meigen (Pl. 6, Fig. 52; Pl. 8, Fig. 74). The genus is distinct in the Empididae in having a truly dorsal arista. Most of the species described as *Gloma* do not belong to this genus. Individuals of *Gloma* are rarely found. They live in subalpine forests and occur in Europe and North America.

Geographical distribution.

1. *G. fuscipennis*, Meigen, Syst. Beschr. Vol. 3, p. 14, pl. 22, f. 11 (1822); Europe.
Macquart. Hist. Nat. Dipt. Vol. 1, p. 345, pl. 8, f. 6 (1834); Blanchard, Hist. Nat. Ins. Vol. 3, p. 582 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 439 (1842); Boitard, Man. Ent. Vol. 3, p. 321 (1843); Schiner, Fauna Dipt. Austr. Vol. 1, p. 112 (1862); Verrall, Ent. Mag. London, Vol. 19, p. 225 (1883); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 87 (1893); Wahlgren, Ent. Tidskr. Vol. 31, p. 85 (1910). — Pl. 6, Fig. 52; Pl. 8, Fig. 74.
2. *G. halterata*, Becker, Mission Arc Méridien Amér. Sud, Vol. 10, p. 168 Ecuador.
(1919).
3. *G. *? hirta*, Loew, Bernsteina fauna, p. 41 (1850); Giebel, Ins. Vorwelt, Baltic Amber.
p. 208 (1856).
4. *G. luctuosa*, nov. sp. (1). — Pl. 8, Fig. 73. W. North America.
5. *G. ossicula*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 132, fig. (1887). Alps.

9. GENUS ITEAPHILA, ZETTERSTEDT

Iteaphila, Zetterstedt, Fauna Ins. Lappon. p. 540 (1838); Dipt. Scand. Vol. 1, p. 258 (1842); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 559 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 110 (1862); Loew, Beschr. Eur. Dipt. Vol. 2, p. 250 (1871); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 117 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389 (1895); Williston, Man. N. Amer. Dipt. p. 74 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 331 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 251, 263 (1903); Melander, Williston, Man. N. Amer. Dipt. p. 226 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302 (1909); Kertész, Cat. Dipt. Vol. 6, p. 80 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 556 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 64 (1910).

(1) ***Gloma luctuosa*, nov. sp. (Pl. 8, Fig. 73).** — Male. Length 3-4 mm. Closely related to *G. fuscipennis*, Meigen, but entirely black and with different proportions to the parts of the pygidium. The basal dorsal process is relatively robust, more setose and its distal fork shorter than in the European species. The main end-valve is broad, stout and roughly triangular, the posterior valve is heavily setose and with crenulate posterior margin. In *fuscipennis* the main valve is narrow, inverted bootshaped and the posterior valve has only two apical and about two basal setae. A pair of anteriorly directed delicate sinuous processes with deflected tip arise from the middle of the pygidium between the dorsal and main valves, and these are straight in *fuscipennis*.

Four males and four females, taken by the author in the heavy forest on the Fairfax Trail just Northwest of Indian Henry's Hunting Ground on Mount Rainier, Washington, August 9, 1922.

Another mountain species of *Gloma* has been collected on Mount Constitution and along Hoods Canal, Washington, on Mount Hood, Oregon, and on Lookout Mountain near Priest Lake, Idaho. As all the specimens are female the species is not described here. This form has the legs, halteres, abdomen and base of wings largely yellowish, as in *fuscipennis*, but it is only two-thirds as large as that species.

Sphicosa, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 751 (1865); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 116 [1889] (*Spicosa*); Kertész, Cat. Dipt. Vol. 6, p. 81 (1909).

Steleocheta, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 129 (1887); Wien. Ent. Zeit. Vol. 9, p. 32 (1890), Vol. 13, p. 156-159 (1894); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257 (1903); Kertész, Cat. Dipt. Vol. 6, p. 80 [1909] (*Steleocheta*); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 608 (1910).

Characters. — Rather slender, usually black, more or less opaque species measuring from two to six millimeters. Head globular, eyes bare, of the male broadly contiguous on the front, the upper facets enlarged, the lowermost minute, of the female widely separated, the front with a few small fronto-orbitals; face receding between the orbits, cheeks narrow but distinct; antennæ inserted at the middle of the head, very long, plainly three-jointed, the basal joints equal, not strongly setose, the third joint cylindrical, somewhat tapering, blunt apically, about five times as long as wide, the terminal style very short, consisting of a broad quadrate basal segment and a minute end-bristle; proboscis extending obliquely forward, about as long as the head, rather thick, the labellar lobes distinct, palpi long and slender, more hairy and longer in the male; hairs of the occiput dense, not seriate, ocellar bristles fine. Thorax rather strong, its hairs abundant and replacing the discal bristles, lateral and posterior bristles fine, scarcely differentiated, scutellum with a fine marginal fringe; pleuræ entirely bare but pollinose. Abdomen slender, about twice as long as the thorax, shorter in the female; pygidium of moderate to rather large size, the middle valves directed backward, the dorsal valves erect and bearing a posterior appendix, penis arcuate, sometimes quite long; abdomen of the female tapering, the terminal styles long. Legs slender, simple, femora of the male more or less ciliate with fine bristles. Wings a little tapering apically, anal lobe prominently rectangular, axillar angle pronounced, wings of the male darker than those of the female, stigma visible, hind margin thin, no basal bristle, marginal hairs minute, auxiliary vein distinct, ending in the costa at the middle of the wing, third vein with a rather short oblique fork, discal cell large and apically blunt, placed in the middle of the wing, distinctly longer than the equal basal cells, last section of the fifth vein much shorter than the preceding, anal crossvein abruptly reflexed, one-third as long as the anal cell, anal vein rather distinct, almost reaching the margin and separate from the under side of the anal cell, alula small; calypteres with a straight edge and a dense fringe.

Type species : *I. Macquarti* (Pl. 6, Fig. 59), Zetterstedt's first species. These flies appear in early spring and frequent the flowers of willow, currant, etc.

Geographical distribution.

1. *I. ? bicolor*, Bigot, Mission Sc. Cap Horn, Vol. 6, p. 19 [1888] (*Sphicosa*); Cape Horn.
Bezzi, Ann. Mus. Hungar, Vol. 3, p. 458, note [1905] (? *Syneches*);
Brethes, An. Mus. Hist. Nat. Buenos-Aires, Vol. 16, p. 290 [1907]
(? genus).
2. *I. conjuncta*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 411 [1900] (*Empis*); Alaska.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 277 [1902] (*Ragas*);
Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 [1903] (? *Empis*);
Melander, Ent. News, Philad. Vol. 17, p. 378 (1906).
3. *I. coriacea*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 126 (1889); Bezzi, Chile.
Ann. Mus. Hungar. Vol. 3, p. 458 (1905).
4. *I. cormus*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 496 [1849] (*Empis*); Canada.
Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 400 [1895] (*Empis*);
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 296 [1902] (*Empis*).
5. *I. italica*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 216 (1873). S. Europe.

6. *I. luctuosa*, Kirby, Fauna N. Amer. (Zool.) Ins. Vol. 4, p. 311 [1837] (*Empis*); Canada.
 Walker, List Dipt. Brit. Mus. Vol. 3, p. 496 [1849] (*Empis*);
 Bethune, Canad. Ent. Vol. 13, p. 165 [1881] (*Empis*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397, 400 [1895] (*Empis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 296 [1902] (*Empis*).
geniculata, Kirby, Fauna N. Amer. (Zool.) Ins. Vol. 4, p. 311 [1837] (*Empis*); Bethune, Canad. Ent. Vol. 13, p. 165 [1881] (*Empis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 296 [1902] (*Empis*).
7. *I. Maackii*, Loew, Beschr. Eur. Dipt. Vol. 2, p. 252 (1871). Siberia.
8. *I. Macquarti*, Zetterstedt, Fauna Ins. Lappon. p. 541 (1838); Dipt. Scand. Vol. 1, p. 258 (1842). Vol. 7, p. 3001 (1849). Vol. 11, p. 4269 (1852); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 145 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 111 (1862); Siebke, Cat. Dipt. Norv. p. 19 [1877] (*Steaphila*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 331 (1902); Wahlgren, Ent. Tidskr. Vol. 31, p. 64 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 36, pl. 1, f. 2 (1913). Europe, Siberia, North America.
- PI. 6. Fig. 59.**
- ? *geniculata*, Zetterstedt, Dipt. Scand. Vol. 1, p. 375 [1842] (*Empis*); Siebke, Nyt. Mag. Naturvid. Vol. 12, p. 153 (1864); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 61 (1867).
setacea, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 130 [1887] (*Steleocheta*); Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 27 [1900] (*Steleocheta*).
9. *I. nigra*, Philippi, Verh. Zool. bot. Ges. Wien, Vol. 15, p. 751, pl. 28, f. 48 [1865] (*Sphicosa*); Bezz, Ann. Mus. Hungar. Vol. 3, p. 458 [1905] (*Sphicosa*).
10. *I. nitidula*, Zetterstedt, Fauna Ins. Lappon, p. 451 (1838); Dipt. Scand. Vol. 1, p. 260 (1842); Wahlgren, Ent. Tidskr. Vol. 31, p. 65 (1910). C. and N. Europe.
meridionalis, Becker, Wien. Ent. Zeit. Vol. 11, p. 126 [1892] (*Steleocheta*).
styriensis, Becker, Wien. Ent. Zeit. Vol. 10, p. 284 [1891] (*Steleocheta*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, p. 201 (1898).
11. *I. obscura*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3018 [1849] (*Hilara*); Strobl, N. Europe.
 Verh. Zool. bot. Ges. Wien, Vol. 42, p. 180 [1892] (*Hilara*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 37, pl. 1, f. 3 (1913).
 var. *fuscipennis*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 39 (1913).
12. *I. orchestris*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 354 (1902). New Mexico.
13. *I. triangula*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 410 [1900] (*Empis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 296 [1902] (*Empis*); Ent. News, Philad. Vol. 17, p. 378 (1906). W. North America.

10. GENUS ANTHEPISCOPUS, BECKER

Anthepisopus, Becker, Wien. Ent. Zeit. Vol. 10, p. 281 (1891); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 246, 260 (1903); Bezz, Ann. Mus. Hungar, Vol. 2, p. 320 (1904); Melander, Williston, Man. N. Amer. Dipt. p. 225 (1908); Kertész, Cat. Dipt. Vol. 6, p. 14 (1909).

Characters. — Rather slender black species about three millimeters in length, the males frequently opaque, the females often more shining, bristles relatively fine but often long and abundant. Head globular, eyes of the male broadly contiguous on the front, bare, the upper facets moderately enlarged, of the female the eyes round, widely separated, facets uniform; antennæ inserted at the middle of the head, elongate, plainly three-jointed, the basal joints equal, not strongly setose, the third joint somewhat compressed cylindrical, slightly tapering, blunt at the apex, three to five times as long as wide, the terminal style very short, two-jointed, the outer joint bristle-like; proboscis extending obliquely forward, longer than the head, palpi very long, linear, more or less hairy; occiput hairy, ocellar

bristles fine, female with several reduced fronto-orbitals. Thorax usually with many hairs and bristles more or less definitely arranged, several humeral, posthumeral, notopleural and supra-alar bristles, acrostichals and anterior dorsocentrals short but numerous, scutellum with a fringe of marginal bristles; pleuræ bare of hairs, pollinose. Abdomen slender, twice as long as the thorax, pygidium small, with a pair of erect dorsal valves, penis short, more or less arcuate, female abdomen tapering. Legs slender, simple, posterior femora of the male more or less ciliate with fine bristles. Wings large, hind angle rectangular, costa vanishing on the hind margin, no basal bristle, marginal hairs microscopic, auxiliary vein complete, distinct from the first, stigma usually present, third vein not forked, basal cells three-fourths as long as the large but blunt discal cell, second posterior cell sessile, much narrower than the third, anal crossvein abruptly reflexed and continuous with the under side of the anal cell, anal vein more or less indicated, no alula, alular hairs short; calypters margined with fine hairs.

Type species : *A. ribesii*, Becker, selected by Coquillett in 1903. The species of this genus are all vernal and show a partiality to the flowers of *Ribes*. There is a possibility that *Microphorus drapetoides* Walker belongs to this genus.

KEY TO THE NORTH AMERICAN SPECIES OF ANTHEPISCOPUS

1. Bristles of the body yellow; proboscis shorter than the head; knobs of the halteres black; femora and tibiæ yellowish; sections of the fifth vein ♂ proportioned 4 : 1 A. FLAVIPILOSUS, Coquillett.
- Bristles black; proboscis longer than the head; last section of the fifth vein not less than one-third the preceding 2.
2. Entirely black including the legs, halteres and calypters, ♂ ♀ 3.
- At least the knees yellowish, ♀; stem of the halteres and calypters paler, ♀; style distinct 5.
3. Sections of the fifth vein 3 : 2; style distinct; penis trumpet-shaped; ten scutellar bristles; femora ciliate. — PI. 8, FIG. 80 A. STENTOR, Melander.
- Sections of the fifth vein 2 : 1; style minute; penis flexed but not enlarged at the end 4.
4. The two prongs of the dorsal valve of the pygidium subequal; lateral bristles of the thorax sparse; scutellum with about ten bristles; flexor cilia of the hind femora inconspicuous; last joint of antennæ ♀ three times as long as broad; palpi ♀ rounded at the end A. POLYGYNUS, nov. sp. (1).

(1) **Anthepisopus polygnus**, nov. sp. — Male. Length 3 mm. Entirely black including the legs, halteres and calypters. Head subopaque, black, upper facets moderately large; third joint of the antennæ but little tapering, four times as long as broad, the style very short, its basal joint square and one-fifth as wide as the antenna; proboscis projecting as far as the head-height, palpi one-half as long as the proboscis and furnished with a few long hairs; occipital hairs abundant. Mesonotum opaque black, but when viewed from in front brownish gray, its bristles long but sparse, two posthumerals, four notopleurals, acrostichals biseriate, eight to ten scutellar bristles. Abdomen nearly opaque, with long black hairs, pygidium rather small, the furcation of the dorsal valve distinct, the posterior branch nearly as long as the anterior, penis shaped like an interrogation-mark (?). Wings with uniform light infumation, stigma distinct, veins blackish, sections of the fourth vein proportioned 0.7 : 0.5 : 1 : 1.5, of the fifth vein 2 : 1, anal vein faint but almost reaching the margin.

Female. Facets uniform; third joint of the antennæ broader, less than four times as long as wide, the style smaller; palpal hairs short; front shining, with two small fronto orbitals. Thorax and abdomen subshining, bristles shorter, sections of the fourth vein proportioned 0.5 : 0.4 : 1 : 1.3.

One male and seven females; Deer Park, Washington, May 5, 1912 (Melander).

- Posterior prong of the dorsal valve minute; lateral bristles of the thorax ♂ abundant; scutellum ♂ with about twenty bristles; flexor cilia of the hind tibiae ♂ longer than the diameter of the femur; third joint of antennae ♀ four times as long as wide; palpi ♀ pointed A. HIRSUTUS, nov. sp. (1).
5. ♀ palpi very slender, longer than the head; halteres, calypters and legs including the coxae yellow; sections of the fifth vein 5 : 2; thorax shining A. LONGIPALPIS, nov. sp. (2).
- Palpi much shorter than the head; knob of the halteres black 6.
6. Thorax shining; coxae and femora black or brown; sections of the fifth vein 5 : 2; stigma distinct. A. NUPTUS, nov. sp. (3).

(1) **Anthepisopus hirsutus**, nov. sp. — Male. Length 3.2 mm. Entirely black including the legs, halteres and calypters, almost opaque. Upper facets slightly enlarged; third joint of the antennae five times as long as broad, slightly tapering, blunt at the end, style very short, the basal segment square and one-fifth as wide as the base of the third joint; proboscis one-half longer than the head, palpi linear, elongate, two-thirds as long as the proboscis, loosely provided with long black hairs; hairs of the occiput rather abundant. Thorax showing a blackish gray median vitta when viewed from in front, its bristles long and numerous, about eight humerals, three posthumerals, ten notopleurals, five supra-alaris, twenty scutellars, the acrostichals biserrate. Abdomen with moderate long fine blackish hairs; pygidium of moderate size, dorsal valves with a minute prong at the middle of the posterior edge, penis rectangularly bent at the middle, the apical part curled on itself. Hairs of the legs black, femoral hairs long, abundant on the middle femora, hind femora ciliate, the flexor hairs nearly twice as long as the diameter of the femur and fifteen in number. Wings nearly hyaline, stigma distinct, veins black, discal cell blunt, three times as long as broad, sections of the fourth vein proportioned 0.6 : 0.4 : 1 : 1.6, of the fifth vein 5 : 3.

Female. Subshining, lightly dusted; hairs and bristles reduced in size and number; facets uniform; palpi shortened. Five humerals, five notopleurals, supra-alaris indistinct; femora not ciliate.

Type from Chehalis, Washington, March 19, 1911. Paratypes from Chehalis and Seattle, Washington, March 13 to 24; seven specimens, taken on *Salix* blossoms by the author.

(2) **Anthepisopus longipalpis**, nov. sp. — Female. Length 2.5 mm. Piceous black, shining, legs, halteres and root of the wings yellow; bristles black. Head shining, two fronto-orbitals; third joint of the antennae thick but bluntly tapering, about three times as long as broad and five times as long as the style; proboscis nearly twice as long as the head, palpi linear, one-half longer than the third joint of the antennae and two-thirds as long as the proboscis; occipital hairs sparse but long. Dorsocentral and acrostichal setulae sparse long and fine, the latter biserrate, two small posthumerals, three long and several short notopleurals, ten scutellar bristles. Abdomen and its hairs brownish. Legs slender, yellow, including the coxae, but the tips of the tarsi darkened, the hairs yellow. Wings with a cinereous tinge, no stigma, veins narrow and brownish yellow, discal cell two and a half times as long as broad, posterior crossvein transverse, sections of the fourth vein proportioned 0.6 : 0.3 : 1 : 1.3, of the fifth vein 5 : 2, anal vein very weak.

One specimen, Arlington, Washington, May 21, 1908, found on a window (Melander).

(3) **Anthepisopus nuptus**, nov. sp. — Male. Length 3.8 mm. Shining, head and thorax very lightly pollinose, entirely black including the halteres and calypters, the knees and the anterior femora and tibiae brownish. Upper facets moderately enlarged; third joint of the antennae three and a half times as long as the width at the base, narrowed beyond the basal fourth, tapering only at the tip, the basal joint of the style oblong, one-third as long as the width of the base of the third antennal joint; proboscis one-half longer than the head, palpi elongate, linear, the hairs sparse. Thoracic bristles long, but the hairs fine and short, about five small humerals, five long notopleurals, twelve scutellars, the dorsocentrals not differentiated excepting the posterior pair, irregularly biserrate, acrostichals irregularly quadriserrate. Hairs of the abdomen long, fine and abundant; pygidium moderately large, the lateral valves circular except for their broad posterior process, dorsal valves short, chitinized, the posterior prong short and placed rather near the tip, penis slender, only gently sinuous. Hairs of the legs short, anterior legs broadly fuscous about the knees, middle femora with sparse hairs, hind femora ciliate below only toward the knee, the hairs there about as long as the diameter of the femur. Wings with a light infumation, stigma distinct, veins blackish, discal cell very blunt, two and a half times as long as broad, sections of the fourth vein proportioned 2.1 : 1 : 3 : 4.5, of the fifth vein 2 : 1.

Female. Pollen coating of the thorax a little more evident; facets uniform; palpi shortened; bristles of the thorax short and less numerous, there being but three distinct notopleurals, ten scutellars, etc.; hairs of the abdomen short; fuscous color of the legs more extended, no ciliation; base of the halteres and calypters fuscous; wings subhyaline, veins brown.

Numerous specimens of each sex from flowers of *Ribes*; Tacoma, Washington, April 12, 1913. Three females also from Olga, Washington, May 18, 1910.

Thorax cinereous dusted, opaque; coxae. anterior femora and base
of the hind femora pale yellow; sections of the fifth vein 2 : 1,
no stigma. A. FLAVICOXA, nov. sp. (1).

Geographical distribution.

- | | |
|--|-------------------|
| 1. <i>A. antipodus</i> , Bezz, Ann. Mus. Hungar, Vol. 2, p. 339, f. 3, 4 (1904). | New South Wales. |
| 2. <i>A. cælebs</i> , Becker, Wien. Ent. Zeit. Vol. 10, p. 283 (1891); Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 27 (1900). | Alps; Siberia. |
| 3. <i>A. consobrinus</i> , Zetterstedt, Fauna Ins. Lappon, p. 541 [1838] (<i>Rhamphomyza</i>).
<i>myrica</i> , Haliday, Walker, Ins. Brit. Dipt. Vol. 1, p. 111 [1851] (<i>Euthyneura</i>);
Loew, Syst. Beschr. Vol. 2, p. 251, note 2 [1871] (<i>Euthyneura</i>); Lundbeck,
Dipt. Dan. Vol. 3, p. 211 [1910] (<i>Euthyneura</i>); Wahlgren, Ent. Tidskr.
Vol. 31, p. 80, f. 12 [1910] (<i>Euthyneura</i>); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 60 [1913] (<i>Euthyneura</i>).
<i>rostratus</i> , Zetterstedt, Dipt. Scand. Vol. 1, p. 250, 415, note 2 [1842] (<i>Anthalia</i>),
Vol. 8, p. 2998 [1849] (<i>Anthalia</i>); Loew, Beschr. Eur. Dipt. Vol. 2, p. 251
[1871] (<i>Euthyneura</i>). | C. & N. Europe. |
| 4. <i>A. flavicoxa</i> , nov. sp. | Washington. |
| 5. <i>A. flavipilosus</i> , Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 413 [1900]
(<i>Microphorus</i>); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 257
[1902] (<i>Euthyneura</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 263
[1903] (<i>Microphorus</i>). | British Columbia. |
| 6. <i>A. hirsutus</i> , nov. sp. | Washington. |
| 7. <i>A. longipalpis</i> , nov. sp. | Washington. |
| 8. <i>A. nuptius</i> , nov. sp. | Washington. |
| 9. <i>A. ædalinus</i> , Zetterstedt, Fauna Ins. Lappon, p. 570 (1838) ♀ (<i>Rhamphomyza</i>);
Dipt. Scand. Vol. 1, p. 413 [1842] (<i>Rhamphomyia</i>); Bonsdorff,
Finl. tvåv. Ins. Dipt. Vol. 1, p. 172 [1861] (<i>Rhamphomyia</i>); Wahlgren,
Ent. Tidskr. Vol. 31, p. 57 [1910] (<i>Rhamphomyia</i>); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 34, pl. 1, f. 1 (1913).
<i>fraternellus</i> , Zetterstedt, Fauna Ins. Lappon, p. 570 [1838] (<i>Rhamphomyza</i>);
Dipt. Scand. Vol. 1, p. 419 [1842] (<i>Rhamphomyia</i>); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 172 [1861] (<i>Rhamphomyia</i>); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 18, note [1908] (<i>Rhamphomyia</i>); Wahlgren, Ent. Tidskr. Vol. 31, p. 57 [1910] (<i>Rhamphomyia</i>).
<i>?furcatus</i> , Zetterstedt, Dipt. Scand. Vol. 1, p. 252 [1842] (<i>Anthalia</i>); Schiner,
Fauna Dipt. Austr. Vol. 1, p. 110 [1862] (<i>Empis</i>); Loew, Beschr. Eur. Dipt. Vol. 2, p. 253, note [1871] (<i>Iteaphila maackii</i> ♀ ?); Kertész, Cat. Dipt. Vol. 6, p. 81 [1909] (<i>Euthyneura</i> ?); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 35 (1913).
<i>niger</i> , Zetterstedt, Dipt. Scand. Vol. 1, p. 251 [1842] (<i>Anthalia</i>); Wahlgren, Ent. Tidskr. Vol. 31, p. 80 [1910] (<i>Euthyneura</i>). | N. Europe. |
| 10. <i>A. polygynus</i> , nov. sp. | Washington. |

(1) *Anthevisorius flavicoxa*, nov. sp. — Female. Length 2.8 mm. Opaque, cinereous black; base of the legs yellow; bristles black. Head lightly cinereous, five fronto-orbital setulae; third joint of the antennae broad, only slightly tapering, almost four times as long as broad, style nearly invisible; proboscis as long as the head, thick, palpi narrowly clavate, blunt at the tip, three-fourths as long as the third joint of the antennæ. Three posthumerals, three notopleurals, ten scutellars, anterior dorsocentrals setuliform, arranged in a definite row, similar to the biseriate acrostichal setulae. Coxæ, anterior femora and tibiæ and base of the hind femora clear yellow, remainder of the legs blackish brown, the hairs black. Calypteres and stem of the halteres yellow, knob blackish. Wings hyaline, veins narrow but blackish, stigma distinct, discal cell three times as long as broad, one-half longer than the second basal cell, sections of the fourth vein proportioned 0.6 : 0.4 : 1 : 1.5, of the fifth vein, 1 : 0.6, anal vein indicated by only a weak fold.

One specimen; Kettle Falls, Washington, May 3, 1912.

11. *A. ribesii*, Becker, Wien. Ent. Zeit. Vol. 10, p. 282, pl. 3, f. 1-3 (1891); C. Europe.
 Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, p. 201 (1898).
 var. *nigripes*, Strobl, Glasn. Bosn. Herzeg. Sarajevo, Vol. 10, p. 402 (1898); Hercegovina.
 Mitteil. Bosn. Herzeg. Sarajevo, Vol. 7, p. 564 (1900).
12. *A. stentor*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 348 [1902] New Mexico.
 (*Euthyneura*). — PI. 8, Fig. 80.
13. *A. Zontaki*, Nowicki, Rocznik tow. nauk. Krakow, Vol. 42, p. 73 [1871] C. Europe.
 (*Microphorus*); Loew, Beschr. Eur. Dipt. Vol. 2, p. 249 [1871] (*Micro-*
phorus), Vol. 3, p. 320 [1873] (*Microphorus*).

III. GENUS BROCHELLA, NOV. GEN.

Characters. — Heavily pruinose species suggestive of *Clinocera*, with long projecting proboscis, no antennal style and with geniculate hind tibiæ. Head horizontally lengthened, occiput very convex, subconical, eyes of both sexes widely separate, bare, facets uniform, ocellar triangle not elevated, face very short, vaulted over the mouth-opening, facial orbits not differentiated, sides of face continuous with the narrow cheeks; antennæ large; joints distinct, the first joint short, cylindrical, nearly bare, the second joint globose, with a whirl of inconspicuous setulæ, the third joint compressed oval, much longer and deeper than the basal joints together, with scattered brief pubescence, no trace of style or arista; proboscis twice as long as head, thick and straight, projecting obliquely forward, palpi ribbon-like, bisetose, attached at basal one-fourth of proboscis, incumbent; ocellar, postocellar and the single vertical seta short, occiput with scattered short bristles, lower side of head with straggling isolated hairs. Thorax somewhat flattened before the scutellum, devoid of pubescence but with setæ as follows; a complete row of 8-12 uniform dorsocentrals, no acrostichals, three small humeral, three small posthumeral, one small notopleural, two supra-alar, one intra-alar, one postalar, two converging scutellar; pectus with weak straggling hairs, a cluster of a dozen metapleural setulæ, no other pleural hairs. Abdomen depressed, seven segments before the large pygidium of the male, last three segments of female short, telescopic within the seventh segment, second segment twice girdled by closely placed shining pittings, less conspicuous in female, hairs not conspicuous; pygidium globose, erect, dorsal valves oval directed posteriorly and parallel, side valves large petaloid, somewhat truncate apically, inferior valves small, rounded, penis straight, extending backwards, compressed and furnished with a fine velvety brush underneath. Legs pollinose, not bristly, hind femora and tibiæ robust, coxæ with scattered hairs, front femora with fine setæ, posterior femora short-setose beneath, hind tibiæ geniculate at the very base, straight, shorter than the femur, in male tipped with an umbo which fits into a depression of the femur next to the trochanter and fimbriate exteriorly to the umbo; tarsi normal, pulvilli distinct, empodia microsetiform.

Wings fully developed, no basal seta, costa continuing around entire margin, auxiliary vein straight, stopping opposite the anterior crossvein, third vein forked before end of second vein, discal cell large, located at middle of wing, emitting three veins, basal cells long, coextensive, anal crossvein reflexed, nearly in line with the lower-outer edge of the second basal and continuous with the under vein of the anal cell, the anal vein represented only by a disjointed fold, anal vein weak, no axillar incision; calypteres with weak fringe.

Type species: *Brochella monticola*, nov. sp. The genus has been placed with the Empidinæ even though it has no near relatives in that subfamily. The habits and habitus of the species are suggestive of the *Clinocera* group, but there also it would be isolated. The low-placed antennæ, lack of style,

long proboscis, reflexed anal vein, and distinct fold under the humeral crossvein at the origin of the second basal cell do not indicate close kinship to the Clinoceratinæ.

Geographical distribution.

1. *B. monticola*, nov. sp. (1). — **Pl. 2, Fig. 16.**

Washington.

12. GENUS RAGAS, WALKER

Ragas, Walker, Ent. Mag. London, Vol. 4, p. 229 (1837); Westwood, Gen. Syn. p. 133 (1840); Walker, Ins. Brit. Dipt. Vol. 1, p. 102 (1851); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 561 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 111 (1862); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 121 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 256 (1903); Melander, Ent. News, Philad. Vol. 17, p. 378 (1906); Kertész, Cat. Dipt. Vol. 6, p. 84 [1909] (*Rhagás*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 5 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 65 [1910] (*Rhagás*).

Characters. — Small dark-colored rare species of generalized structure. Eyes of the male subcontiguous above the antennæ, separated only by a line, the facets uniform and bare, of the female broadly separated on the short front, female with two small fronto-orbitals, face very short and broad, the large quadrate epistome receding, cheeks linear; antennæ short, basal joints minute and globular, third joint long and conical, style very thick, nearly half as long as the third joint, its basal segment four times as long as broad and tipped with a short bristle; proboscis short and chitinized, the labrum incurved, broad at the base and pointed, palpi very short, retracted, horizontal. Thoracic bristles short, one humeral, three posthumeral, about eleven dorsocentral and three notopleural, but the scutellum with six strong bristles; pleuræ bare. Abdomen twice as long as the thorax, tapering in the female and tipped with two long styles, in the male with a terminal small open pygidium which bears a pair of lateral slender curved pointed valves and a pair of erect dorsal processes. Legs simple, bristleless, hairs sparse and inconspicuous. Wings broad, anal angle strong, no axillar excision, costa enclosing the entire wing, no humeral bristle, auxiliary vein distinct, complete and ending by a gentle curve in the costa, fork of the third vein long, contact of the discal cell with the third posterior cell longer than with the fourth, anal crossvein reflexed, anal vein continuous with the under side of the anal cell.

Type species: *R. unica*, Walker. — **Pl. 8, Fig. 84.**

Geographical distribution.

1. *R. *? generosa*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 94, 129, pl. 12, Baltic Amber.
f. 7-9 (1908).
2. *R. unica*, Walker, Ent. Mag. London, Vol. 4, p. 229 (1837); Ins. Brit. Dipt. Europe.
Vol. I, p. 102, pl. 3, f. 3 (1851); Schiner, Fauna Dipt. Austr. Vol. 1,
p. 111 (1862); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9),
p. 3 (1908), Vol. 37 (3), p. 46 (1913). — **Pl. 8, Fig. 84.**

(1) **Brochella monticola**, nov. sp. (**Pl. 2, Fig. 16.**) — Male. Length 4.5 mm. Black in groundcolor, heavily pruinose or pollinose. Front brown, face gray, proboscis silvery white, occiput brownish gray. Notum brownish gray becoming brown on dorsocentral rows, scutellum, pleuræ and abdomen cinereous; pygidial valves hairy. Wings uniformly infumated, veins brown, no stigma, a crossvein near basal third of discal cell.

Female. Hind femora less robust than in male, the tibæ without umbo or fringe at apex.

Types Seventeen males, seven females, found by the writer along the rills above Paradise Park on Mt. Rainier, Washington, during the middle of August, 1917. A visit to the same locality in the latter part of August of the same year and again in July, 1922, failed to disclose additional specimens, although special effort was made to discover them.

13. GENUS PHILETUS, NOV. GEN.

Characters. — Black opaque species with slender black legs. Head produced obliquely downward, the antennæ placed low and the ocelli far forward; eyes of both sexes widely separated, bare, facets almost uniform; sides of the short front slightly diverging behind, two pairs of small fronto-orbitals, face square then receding as the slightly longer epistome; antennæ shorter than the head, the first joint minute, the second joint globular and with minute setulae, the third joint ovate-conical ending in a short thick arista whose short basal segment is one-fifth as long as the subulate distal portion; proboscis one-half as long as the head, extending downward and slightly forward, the labrum needle-like, the labium fleshy, palpi short and rather spatulate; hairs of the occiput sparse, becoming gradually stronger above, one pair of parallel ocellars and one of diverging post-ocellars. About eight pairs of dorsocentrals, one humeral, two posthumeral, three notopleural, two supra-alar and four scutellar bristles; pleurae bare but entirely pruinose. Abdomen slender, one-half longer than the thorax, second segment with a double transverse row of pittings and the second to the sixth segments with two pits on each side; pygidium large, globose, terminal, dorsal valves long, narrow, and recumbent, upper lateral valves deeply and horizontally cleft from the convex sides, ventral part not compressed, bearing the short curved slender penis; abdomen of the female entirely tapering, ending in two styles. Legs slender, simple, not bristly, pulvilli small. Wings with nearly parallel sides, slightly broadest beyond the middle, two and two-thirds times as long as broad, anal angle broadly rounded, costa continuing around the entire wing, basal bristles small, auxiliary vein distinct, complete, ending in the costa at two-fifths the length of the wing, third vein arising at one-third the distance between the humeral and anterior crossveins, its distal fork rather large, first basal cell a little longer than the second and as long as the discal cell, sections of the fifth vein equal, anal crossvein abruptly recurved, confluent with the underside of the anal cell but reaching only one-fourth the distance back, anal vein obsolete, no alula; calypters with a straight edge and the usual fringe.

Type species: *Ph. memorandus*, nov. sp. Two subalpine North American species have been discovered by the author.

Geographical distribution.

1. *Ph. memorandus*, nov. sp. (1). — PI. 7, Fig. 66; PI. 8, Fig. 78.
2. *Ph. schizophorus*, nov. sp. (2). — PI. 8, Fig. 81.

W. United States.
Washington.

(1) **Philetus memorandus**, nov. sp. (PI. 7, Fig. 66; PI. 8, Fig. 78). — Male. Length 2.75 mm. Black, the front, face, pleurae and abdomen cinereous pollinose. Third joint of the antennæ two and a half times as long as wide, the arista two-thirds as long as the third joint; mouthparts black, face square. Dorsum of the thorax olivaceous brown. Hairs of the first segment of the abdomen rather dense, pygidium black, coated with gray villosity, the two upper pairs of processes narrow, parallel and pruinose, the second jointed with the middle valve, penis curving to meet the second process. Wings hyaline, veins dark brown, second submarginal cell three times as long on the costa as the first, sections of the fourth vein proportioned 1 : 0.3 : 1 : 1.6, vein between the basal cells weak, discal cell two and a half times as long as broad, anal vein indicated by a weak transparent fold, hairs of the hind margin shorter than the anterior crossvein, alular hairs short and not dense.

Female. Pruinosity of the abdomen less evident, face above the epistome less deep than broad.

Type and allotype, Mount Constitution, Washington, the male collected July 22, 1909, the female, May 17, 1910. Five paratypes collected by the author on the South slope of Mount Rainier, Washington, and another at the Canyon Camp, Yellowstone Park, Wyoming.

(2) **Philetus schizophorus**, nov. sp. (PI. 8, Fig. 81.) — Differing from the preceding species in the genitalia but otherwise similar. The uppermost pair of pygidial processes shining, broadly quadrate at base and then deeply and widely excised so as to terminate in two long thin divaricating fingers, the second process arising from the dorsal by a narrow polished stem, penis very short and thin, not attaining the second process.

Two specimens, Van Trump Creek, above Christine Falls, Mount Rainier, 1 September, 1917 (Melandier).

14. GENUS HESPEREMPIST, MELANDER

Hesperempis, Melander, Ent. News, Philad. Vol. 17, p. 377 (1906); Williston, Man. N. Amer. Dipt. p. 226 (1908); Kertész, Cat. Dipt. Vol. 6, p. 84 (1909); Coquillett, Proc. U. S. Mus. Vol. 37, p. 551 (1910).

Ragas, Melander (not Walker), Trans. Amer. Ent. Soc. Vol. 28, p. 276 (1902).

Characters.— Bristleless, entirely opaque dusted, blackish or yellow species with yellow legs. Eyes separated, broadly so on the front, and in the male narrowly just beneath the antennæ, the sides of the face diverging below, facets bare, of uniform size in the female or the anterior slightly larger in the male; face of the female short and quadrate, the lower part retracted as the pollinose epistome; antennæ inserted above the middle of the head, distinctly three-jointed, the first joint longer than the second, the basal joints with only a few microscopic hairs, the third joint compressed conical, terminated by a short two-jointed style whose basal segment is thick and whose outer segment is short and bristle-like; proboscis very short and thick, the labella broad fleshy and hairy, palpi flattened, as long as the proboscis and incumbent on it, spongy pubescent but almost devoid of hairs; no cheeks; ocellar triangle not elevated, ocellar and vertical bristles reduced in size so as to be almost invisible. Thorax not much elevated, entirely without bristles, scutellum with six or eight short marginal hairs, metapleuræ bare, at most with a couple of microscopic hairs. Abdomen slender, longer than the thorax, depressed, hairs short and sparse, a transverse series of small round pits present at the base of the second segment; pygidium terminal globular, with a flat dorsal valve and moderately large convex lateral valves, penis short and thick, no projecting appendages, abdomen of the female tapering to the end styles. Legs slender, simple, rather sparsely provided with fine pubescence, but entirely devoid of bristles; pulvilli small. Calypteres with a straight edge and reduced fringe. Wings long and slender, anal angle obtusely rounded, not prominent, costa extending around the entire wing, no basal bristle, auxiliary vein straight, distinct, ending in the costa just before the middle of the wing, a very faint stigmal spot, third vein forked, basal cells elongate and equal, discal cell rather narrow, often pointed apically, sections of the fourth vein proportioned $0.8 : 0.2 : 1 : 1.5$, of the fifth vein, $1.4 : 0.5 : 1 : 1.2$, anal crossvein abruptly recurved and fused with the under side of the anal cell but not reaching back to the middle of the cell, anal vein evanescent.

Type species : *H. Mabelæ* (Pl. 7, Fig. 61), by original designation. This species occurs sparingly in the deep shade of fir forests, flying close to the ground.

Geographical distribution.

1. *H. Mabelæ*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 277, f. 98 [1902] Idaho. Montana.
(*Ragas*); Ent. News, Philad. Vol. 17, p. 377, fig. (1906). — **Pl. 7,**
Fig. 61.
2. *H. sanduca*, nov. sp. (1). California.

(1) **Hesperempis sanduca**, nov. sp. — Male. Length 2.6 mm. Entirely pale yellow except the eyes and the last two joints of the hind tarsi, not pollinose but only subshining, hairs and bristles yellow. Third joint of the antennæ bluntly conical, scarcely tapering, twice as long as wide and six times as long as the brown style. About fifteen short dorsocentral hairs, six scutellæ; metapleuræ with two microscopic hairs. Hairs of the abdomen rather long. Legs simple, bristleless. Wings clear hyaline, veins weak and yellow, second posterior cell sessile, the vein between the discal and the third posterior cells angulate at two-thirds its length and sending a short spur into the discal cell, anal vein nearly obsolete, its base continuous with the under side of the anal cell.

Type, San Francisco, California, collected by Charles Fuchs. A female from Forest Grove, Oregon, June 2 (F. R. Cole).

15. GENUS HILARA, MEIGEN

Hilara, Meigen, Syst. Beschr. Vol. 3, p. 1 (1822); Curtis, Brit. Ent. Vol. 8, p. 130 (1826); Macquart, Dipt. N. France, Vol. 3, p. 108 (1827); Hist. Nat. Dipt. Vol. 1, p. 340 (1834); Zetterstedt, Fauna Ins. Lappon, p. 554 (1838); Westwood, Gen. Syn. p. 132 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 335 (1842); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 97 (1851); Rondani, Dipt. Ital. Vol. 1, p. 151 [1856] (*Hylara*); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 112 (1862); Lioy, Atti Inst. Sc. Veneto, Venezia, 1864, p. 601 (1864); Beling, Arch. Naturg. Berlin, Vol. 48, p. 240 (1882); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 117 (1889); Strobl, Verh. Zool.-bot. Ges. Wien. Vol. 42, p. 85, 182 (1892); Becker, Wien. Ent. Zeit. Vol. 13, p. 156, 159 (1894); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389, 394 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 262 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 250, 263 (1903); Bezzı, Ann. Mus. Hungar. Vol. 2, p. 320, 341, note (1904); Melander, Williston, N. Amer. Dipt. Man. p. 226 (1908); Bezzı, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 375 (1909); Kertész, Cat. Dipt. Vol. 6, p. 84 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 552 (1910); Grünberg, Süßwasserf. Deutschl. 2a, Vol. 1, p. 154 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 140 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 65 (1910); White, Proc. Roy. Soc. Tasmania, 1916, p. 220 (1917); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 357 (1920).

Characters. — Medium-sized species, usually black or blackish in color although sometimes yellowish, distinguished by the anterior bend at the end of the auxiliary vein and by the almost universally enlarged front metatarsi of the male. A genus rich in species and hence presenting a wide range in the variability of characters, particularly of the chaetotaxy and the secondary sexual devices. Head globular, eyes separated in both sexes, except in a few species where the males are holoptic and have the upper facets enlarged, otherwise the facets are uniform; front and face each quadrate, fronto-orbital bristles usually prominent, cheeks linear; antennae inserted at the middle of the head, distinctly three-jointed, as long as the head, the basal joints short and with few setulae, the first joint cylindrical, the second globular, the third compressed, conical, ending in a lengthened three-jointed style, whose basal segment is minute and quadrate and whose apical part is a short bristle; proboscis as long as the head or somewhat shorter, vertical, the labrum sharp, chitinized, a little incurved, the labium shorter and with prominent labellar lobes, palpi porrect, up-curved and hairy; occiput more or less hairy, ocelli not elevated, distant, the ocellar bristles separated, thorax rather robust, wider than the head, the metathorax short and declivous, bristles usually reduced, the rows of discal bristles usually replaced by hairs, pronotum often with a lateral bristle, notopleural bristles evident but variable; pleuræ pollinose, meta-pleuræ not setose. Abdomen more or less cylindrical, the seventh and eighth segments of the male usually small to accommodate the reflexed epipygium; abdomen of the female tapering, ending in two thin styles; epipygium usually compressed, the ventral keel ending in a curved tapering process, the lateral valves extending forward, variously fissate and spiny at the anterior end, dorsal valves and penis hidden. Legs not lengthened, rather hairy, usually with variable bristles on the tibiæ, femora rarely thick, front metatarsi of the male almost always greatly swollen and often cristate with long hairs, pulvilli of moderate size, empodium distinct. Anal angle rather prominent but rounded, costa continuing around the entire margin, stigma formed, auxiliary vein complete and distinct, curving forward at the end to enter the costa, third vein with a long fork, basal cells coextensive, discal cell as long as

the basals, anal crossvein abruptly reflexed, usually extending back half the length of the anal cell, the anal vein usually arising at the juncture of the crossvein with the under side of the cell.

Type species : *H. maura*, Fabricius (Pl. 6, Fig. 55), selected by Curtis in 1838. The species of *Hilara* are common along small streams, often skimming over the surface in an aerial dance. The males of some species have the habit of enveloping their prey in a web spun from the mouth. The genus is best developed in the Northern Hemisphere, although a few species are known from South of the equator.

Geographical distribution.

1. *H. abdominalis*, Zetterstedt, Fauna Ins. Lappon, p. 555 (1838); Dipt. N. Europe.
Scand. Vol. 1, p. 337 (1842); Strobl, Verh. Ges. Zool.-bot. Wien.
Vol. 42, p. 180 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910).
2. *H. aberrans*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 376 (1909). Bolivia.
3. *H. abnormis*, Bezzi, ibidem, Vol. 91, p. 377 (1909). Peru.
4. *H. aeronetha*, Mik, Wien. Ent. Zeit. Vol. 11, p. 81 (1892); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 158, note (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1893); Mik, Wien. Ent. Zeit. Vol. 13, p. 283 (1894); Lundbeck, Dipt. Dan. Vol. 3, p. 162, f. 59 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912). C. Europe.
var. *angustifrons*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 132, 158, note (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1892).
5. *H. aethiops*, Zetterstedt, Fauna Ins. Lappon, p. 556 (1838); Dipt. Scand. N. Europe.
Vol. 1, p. 347 (1842); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 132, note (1892).
6. ? *H. albipennis*, Roser, Correspondenzbl. Landw. Würtemb. Stuttgart, C. Europe.
Vol. 1, p. 53 (1840).
7. *H. albitarsis*, Roser, ibidem, Vol. 1, p. 53 (1840). C. Europe.
8. *H. albocingulata*, Wood, Ent. Mag. Lond. Vol. 49, p. 13 (1913). England.
9. *H. algaeivascensis*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 24 (1899). Spain.
10. *H. almeriensis*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 310 (1906). Spain.
11. *H. amaranta*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 123 (1907). Tunis.
12. *H. anglokanica*. Lundbeck, Naturh. Medd. Kjöbenhavn, Vol. 64, p. 325 (1913). Denmark.
13. *H. atra*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 205 : Cent. 3, No. 42 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 266 (1902). United States.
14. *H. aurata*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 411 (1900); Alaska.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 269 (1902).
15. *H. baculifer*, Melander, ibidem, Vol. 28, p. 271, f. 96 (1902). Georgia.
16. *H. balnearia*, White, Proc. Roy. Soc. Tasmania, 1916, p. 223, f. 41b (1917). Tasmania.
17. *H. barbipes*, Frey, Medd. Soc. Sc. Fenn. Helsingfors, Vol. 33, p. 67 (1907); Finland.
Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 56, f. 9 (1913).
18. *H. bares*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 491 (1849); Brunetti, E. Indies.
Fauna Brit. Ind. Dipt. Vol. 1, p. 359 (1920).
19. *H. basalis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 206 : Cent. 3, No. 45 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 267 (1902). Illinois.
20. *H. Beckeri*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 143 (1892); C. & N. Europe.
Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 167, f. 61 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Verrall, Ent. Mag. Lond. Vol. 48,

- p. 25 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 49 (1913).
quadrivittata, Zetterstedt (not Meigen), Dipt. Scand. Vol. 1, p. 339 (1842); ibidem, Vol. 8, p. 3015 (1849).
21. *H. bella*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 271, f. 94 (1902). Massachusetts.
 22. *H. bistriata*, Zetterstedt, Dipt. Scand. Vol. 1, p. 340 (1842); ibidem, Europe.
 Vol. 8, p. 3015 (1849); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 145 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 170, f. 63 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 50 (1913).
 brevivittata, Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 161 (1861).
 var. *cantabrica*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 22 (1899). Spain.
 var. *flavohalterata*, Strobl, Glasnik. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 412 (1898); Mitteil. Mus. Bosn. Herceg. Sarajevo, Vol. 7, p. 572 (1900). S. Europe.
23. *H. bivittata*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 113 (1892); Europe.
 Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 151, f. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 56 (1913).
24. *H. borealis*, Oldenberg, Arch. Naturg. Berlin, Vol. 81, A (9), p. 172 Alps, Lappland.
 (1916).
- Czernyi*, Strobl, part. Verh. Zool-bot. Ges. Wien, Vol. 59, p. 176, part. (1909).
25. *H. borina*, Becker, Deut. Ent. Zeitsch. p. 644 (1910). Corsica.
26. *H. Braueri*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 135 (1892); C. & N. Europe.
 Verrall, Ent. Mag. London, Vol. 48, p. 25 (1912).
 argyrosoma, Strobl, Verh. Zool.-bot. Ges. Wien, vol. 42, p. 136 (1892); Meijere, Tijdschr. v. Ent. Vol. 50, p. 177 (1907); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).
 niveipennis, Zetterstedt, var. Dipt. Scand. Vol. 1, p. 352, var. a and b. (1842).
27. *H. brevipila*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 204 : Cent. 3, No. 41 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 266 (1902). C. United States.
28. *H. brevivittata*, Macquart, Dipt. N. France, Vol. 3, p. 114 (1827); Hist. Europe.
 Nat. Dipt. Vol. 1, p. 342 (1834); Meigen, Syst. Beschr. Vol. 7, p. 80 (1838); Zetterstedt, Dipt. Scand. Vol. 1, p. 357 (1842); ibidem, Vol. 8, p. 3018 (1849); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 146 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).
29. *H. cærulescens*, Oldenberg, Arch. Naturg. Berlin, Vol. 81, A (9), p. 168 Tirol.
 (1916).
30. *H. cana*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 395 (1895); California.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 269 (1902).
31. *H. canescens*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3014 (1849); ibidem, C. & N. Europe
 Vol. 11, p. 4269 (1852); Strobl, Verh. Ges. Zool.-bot. Wien. Vol. 42, p. 164 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 91 (1892); Verrall, Ent. Mag. London (2), Vol. 5, p. 141 (1894); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 47 (1913).
32. *H. capensis*, Schiner, Novara Reise, Dipt. p. 206 (1868). Cape Good Hope.
33. *H. carbonaria*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 272 (1902). Massachusetts.
34. *H. carbonella*, Zetterstedt, Dipt. Scand. Vol. 1, p. 359 (1842); ibidem, N. Europe.
 Vol. 13, p. 5003 (1859); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 162 (1861); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 181, (1892).

35. *H. carinthiaca*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 144 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 168, f. 62 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 25 (1912).
quadrivittata, Becker (not Meigen), Berl. Ent. Zeitschr. Vol. 31, p. 127 (1887).
36. *H. castanipes*, Loew, Ofvers. Vet. Akad. Förh. Vol. 14, p. 371 (1857); Dipt. Südafr. Vol. 1, p. 268 (1860); Beazzi, Ann. Mus. Hungar., Vol. 3, p. 443 (1905).
37. *H. certa*, Walker, Ins. Saunders, Dipt. Vol. 1, p. 204 (1852). Tasmania.
38. *H. chiragrîca*, Speiser, Kilimandjaro-Meru Exped. Vol. 10, p. 107 (1910). Africa.
39. *H. chorica*, Fallen, Empid. Suec. p. 24 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 4 (1822); Macquart, Dipt. N. France, Vol. 3, p. 111 (1827); Hist. Nat. Dipt. Vol. 1, p. 340 (1834); Zetterstedt, Fauna Ins. Lappon, p. 556 (1838); Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt. p. 20, f. 14 (1840); Isis, Vol. 7, p. 546, f. 14 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 357 (1842); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 490 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3018 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 99 (1851); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 162 (1861); Siebke, Nyt. Mag. Naturvid. Vol. 14, p. 381 (1866); Leunis, Syn. Zool. Vol. 2, p. 402 (1886); Neuhaus, Dipt. March. p. 70 (1886); Strobl, Verh. Ges. Zool.-bot. Wien, Vol. 42, p. 108 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892); Grünberg, Süsswasserf. Deutschl. 2a, Vol. 1, p. 155 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 154, f. 53 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 56 (1913).
nana, Macquart, Mém. Soc. Sc. Lille, p. 161 [1823] (*Empis*); Dipt. N. France, Vol. 3, p. 111 (1827); Hist. Nat. Dipt. p. 341 (1834); Meigen, Syst. Beschr. Vol. 7, p. 80 (1838); Walker, List Dipt. Brit. Mus. Vol. 3, p. 491 (1849); Ins. Brit. Vol. 1, p. 101 (1851).
nitens, Macquart, Mém. Soc. Sc. Lille, p. 162 [1823] (*Empis*).
40. *H. ciliipes*, Meigen, Syst. Beschr. Vol. 3, p. 3, pl. 22, f. 3 (1822); Curtis, Brit. Ent. Vol. 8, p. 130, pl. (1826); Macquart, Dipt. N. France, Vol. 3, p. 110, pl. 3, f. 4 (1827); Hist. Nat. Dipt. Vol. 1, p. 340 (1834); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 490 (1849); Ins. Dipt. Brit. Vol. 1, p. 98, pl. 3, f. 2 (1851); Glover, Manuscr. notes, pl. 6, f. 6 (1874); Neuhaus, Dipt. March. p. 70 (1886); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 171 (1892).
ciliata, Macquart, Hist. Nat. Dipt. Vol. 1, pl. 8, f. 4 (1834).
? clavipes, Harris, Engl. Ins. p. 150, pl. 44, f. 3 [1782] (*Empis*).
41. *H. cinerea*, Macquart, Hist. Nat. Dipt. Vol. 1, p. 342 (1834); Meigen, Syst. Beschr. Vol. 7, p. 81 (1838); Boitard, Man. Ent. Vol. 3, p. 321 (1843); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 181 (1892). C. Europe.
42. *H. cinereomicans*, Strobl, ibidem, Vol. 42, p. 156 (1892); Verrall, Ent. Mag. London, Vol. 48, p. 25 (1912). C. Europe.
- var. *trigemina*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 175 (1909). Spain.
43. *H. cingulata*, Dahlbom, Svenska Vet. Akad. Handl. p. 160 (1850); Zetterstedt, Dipt. Scand. Vol. 11, p. 4270 (1852); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 159 (1892); Mitteil. Mus. Bosn. Europe.

- Herceg. Sarajevo, Vol. 7, p. 573 (1900); Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 311 (1906); Lundbeck, Dipt. Dan. Vol. 3, p. 180 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 25 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 46 (1913).
? Sturmii, Meigen, Syst. Beschr. Vol. 3, p. 5 (1822); Schiner, Fauna, Dipt. Austr. Vol. 1, p. 113 (1862).
var. morena, Strobl, Wien. Ent. Zeit. Vol. 18, page 25 (1899). Spain.
 44. *H. clypeata*, Meigen, Syst. Beschr. Vol. 3, p. 4 (1822); Macquart, Dipt. N. France, Vol. 3, p. 111, pl. 3, f. 5 (1827); Hist. Nat. Dipt. Vol. 1, p. 341 (1834); Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt. p. 20, f. 16, 17 (1840); Isis, Vol. 7, p. 546, f. 6, 17 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 356 (1842); Macquart, Explor. Algérie (Zool.) Vol. 3, p. 444 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 99 (1851); Loew, Dipterent. Südafr. Vol. 1, p. 262 (1860); Schiner, Fauna Dipt. Austr. Vol. 1, p. 114 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 384, 399 (1866); Neuhaus, Dipt. March. p. 70 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 129 (1887); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 106 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 87 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 150, f. 49 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 56 (1913). Europe, N. Africa.
algira, Macquart, Dipt. Exot. Vol. 1, Pt. 2, p. 280 (1838).
pinstorum, Becker (not Zetterstedt), Berl. Ent. Zeitschr. Vol. 31, p. 126, part. (1887); Silen, Medd. Soc. Sc. Fenn. Helsingfors, Vol. 32, p. 107 (1906). C. Europe.
var. brevifurca, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 107 (1892). C. Europe.
var. longifurca, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 107 (1892). C. Europe.
 45. *H. compacta*, Brunetti, Rec. Indian. Mus. Vol. 9, p. 30 (1913); Fauna Brit. Ind. Dipt. p. 358, f. 32 (1920). W. Himalayas.
 46. *H. confirmata*, Walker, Ins. Saunders, Dipt. Vol. 1, p. 205 (1852). Tasmania.
 47. *H. congregaria*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 272 (1902). W. United States.
 48. *H. coracina*, Oldenberg, Arch. f. Naturg. Berlin, Vol. 81, A (9), p. 170 (1916). C. Europe.
H. coracula, Lundbeck, Dipt. Dan. Vol. 3, p. 149 (1910). Denmark.
 50. *H. cornicula*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 42 (1873); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 105 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 87 (1892); Verrall, Ent. Mag. London (2), Vol. 5, p. 141 (1894); Becker, Deutsche Ent. Zeitschr. Vol. 643 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 148, f. 48 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 55 (1913). C. & S. Europe.
chorica, Schiner (not Falien), Fauna, Dipt. Austr. Vol. 1, p. 115 (1862).
? lugubris, Meigen, Sys. Beschr. Vol. 3, p. 10 (1822).
 51. *H. cuneata*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 43 (1873); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 180 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 71 (1909). Hungary.
 52. *H. curvipes*, Siebke, Nyt Mag. Naturvid. Vol. 12, p. 109 (1864); ibidem, Norway. Vol. 14, p. 399 (1866); Cat. Dipt. Norv. p. 29 (1877).
 53. *H. Czernyi*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 176, part. (1909); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 69 (1909). Spain.
 54. *H. dalmatina*, Strobl, Glasnik. Bosn. Herceg. Sarajevo, Vol. 10, p. 412 (1898); Mitth. Bosn. Herceg. Sarajevo, Vol. 7, p. 572 (1900). S. Europe.
 55. *H. dimidiata*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 128 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1892). C. Europe, Alps.

56. *H. discoidalis*, Lundbeck, Dipt. Dan. Vol. 3, p. 151, f. 50, 51 (1910); Denmark, Finland. Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3) p. 56 (1913).
57. *H. discolor*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 165 (1892). C. & S. Europe.
58. *H. diversipes*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 125 (1892); C. Europe. Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1892); Verrall, Ent. Mag. Vol. 48, p. 24 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3) p. 55 (1913).
- pinetorum*, Becker (not Zetterstedt), Berl. Ent. Zeitschr. Vol. 31, p. 126, part (1887).
59. *H. efficiens*, White, Proc. Roy. Soc. Tasmania, 1916, p. 222, f. 1a (1917). Tasmania.
60. *H. ephippium*, Scholz, Zeitsche Ent. Breslau, Vol. 5 (19), p. 51 (1851); C. Europe. Schiner, Fauna, Dipt. Austr., Vol. 1, p. 116 (1862); Strobl, Verh. Ges. Zool.-bot. Wien, Vol. 42, p. 180 (1892).
61. *H. escorialensis*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 175 (1909). Spain.
62. *H. eumera*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 43 (1873); Strobl, C. Europe. Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 178 (1892).
63. *H.* exilis*, Meunier, Ann. Sc. Nat. (Zool.), Vol. 7, p. 93, 121, pl. 11, f. 4 [1908] (*Empis*). Baltic Amber.
64. *H. fasciata*, Meigen, Syst. Beschr. Vol. 3, p. 11 (1822); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 181 (1892).
65. *H. femorata*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 202 : Cent. 3, No. 35 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 264 (1902). E. United States.
66. *H. flava*, Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 174 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 92 (1892). C. Europe.
67. *H. flavipes*, Meigen, Syst. Beschr. Vol. 3, p. 11 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 342 (1834); Guérin, Icon. Régne Anim. (Ins.) p. 537, pl. 94, f. 6 [1835] (*Hybos*) ; Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 52 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 113 (1862); Beling, Arch. Naturg. Berlin, Vol. 48, p. 221 (1882); Verrall, Ent. Mag. London, Vol. 22, p. 202 [1886] (*Oreogeton*) ; Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 160 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 181 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 25 (1912); Engel, Deutsche Ent. Zeitschr. 1918, p. 6, note [1918] (?*Atalanta*). C. Europe.
- ? *acephala*, Panzer, Fauna Germ. Vol. 54, pl. 24 [1798] (*Empis*) ; Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 234 [1804] (*Empis*).
- ? *gracilipes*, Boheman, Svenska Vet. Akad. Handl. p. 191 (1851); Zetterstedt, Scand. Vol. 12, p. 4607 (1855); Strobl, Verh. Ges. Zool.-bot. Wien, Vol. 42, p. 161, 180 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).
- ? *obscura*, Meigen, Syst. Beschr. Vol. 3, p. 11 (1822); Walker, Ins. Brit. Dipt. Vol. 1, p. 102 (1851); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 180 (1892).
68. *H. fortis*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 124 (1914). E. Africa.
69. *H. fulvibarba*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 23 (1899); Collin, Ent. Mag. London (2), Vol. 24, p. 106 (1913). W. Europe.
70. *H. fuscipes*, Fabricius, Ent. Syst. Vol. 4, p. 406 [1794] (*Empis*) ; Meigen, Europe. Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 233 [1804] (*Empis*) ; Fabricius, Syst. Antl. p. 144 [1805] (*Tachydromia*) ; Meigen, Syst. Beschr. Vol. 3, p. 6 (1822); Macquart, Dipt. N. France, Vol. 3, p. 113 (1827); Hist. Nat. Dipt. Vol. 1, p. 343 (1834); Zetterstedt, Fauna Ins. Lappon, p. 555 (1838); Dipt. Scand. Vol. 1, p. 338 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 491 (1849);

- Zetterstedt, Dipt. Scand. Vol. 8, p. 3015 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19) p. 51 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 100 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 144 (1862); Neuhaus, Dipt. March. p. 70 (1886); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 154 (1892); Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 91 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 6, 183 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).
- albida*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 227 [1804] (*Empis*).
intermedia, Fallen, Empid. Succ. p. 23, part. [1816] (*Empis*).
plumbea, Fabricius, Ent. Syst. Vol. 4, p. 406 [1794] (*Empis*); Meigen, Classif. Beschr. Eur. Ins. vol. 1, p. 233 [1804] (*Empis*); Fabricius, Syst. Antl. p. 144 [1805] (*Tachydromia*).
71. *H. fusitibia*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 20 (1899); Glasn. Mus. S. Europe.
 Bosn. Herceg. Sarajevo, Vol. 14, p. 471 (1902); Mitteil. Mus. Bosn. Herceg. Sarajevo, Vol. 9, p. 530 (1904).
72. *H. gallica*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 222 C. & N. Europe. [1804] (*Empis*); Fallen, Empid. Suec. p. 23 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 9 (1822); Macquart, Dipt. N. France, Vol. 3, p. 114 (1827); Hist. Nat. Dipt. Vol. 1, p. 343 (1834); Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt. p. 20 (1840); Isis, Vol. 7, p. 546 (1840); Zetterstedt, Dipt. Sc. Vol. 1, p. 336 (1842); Boitard, Man. Ent. Vol. 3, p. 321 (1843); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 159 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 114 (1862); Siebke, Cat. Dipt. Norv. p. 26 (1877); Neuhaus, Dipt. March. p. 70 (1886); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 166 (1892); Grünberg, Süßwasserf. Deutschl. 2a, Vol. 1, p. 155 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 176, f. 66 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 46 (1913).
73. *H. geniculata*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, C. Europe. Vol. 1, p. 53 (1840).
74. *H. gracilis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 205 : Cent. 3, n° 44 Pennsylvania. (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 266 (1902).
75. *H. griseola*, Zetterstedt, Fauna Ins. Lappon, p. 557 (1838); Dipt. Scand. C. & N. Europe. Vol. 1, p. 350 (1842); Vol. 7, p. 3016 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 160 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 152 (1864); ibidem, Vol. 14, p. 3880 (1866); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 155 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 91 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 47 (1913).
fuscipes, Bonsdorff (not. Fabricius), Finl. tvåv. Ins. Dipt. Vol. 1, p. 159 (1861); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 126, var. (1887).
nigritarsis, Zetterstedt, Fauna, Ins. Lappon, p. 557 (1838); Dipt. Scand. Vol. 1, p. 351 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 161 (1861); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 380, 384, 399 (1866).
platyura, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 42 (1873); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 179 (1892).
76. *H. * Heerii*, Meunier, Verh. Akad. Wetensch. Amsterdam, Sect. 2, Aix, Lower Oligocene. Vol. 18 (5), p. 12, f. 12 (1916).

77. *H. heterogastra*, Nowicki, Verh. Naturf. Ver. Brünn. Vol. 6, p. 84 (1868); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 172 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 91 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 25 (1912).
abdominalis, Scholz (not Zetterstedt), Zeitschr. Ent. Breslau, Vol. 5, p. 19, 50 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 114 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 128 (1887).
?*albiventris*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 53 (1840); Wood, Ent. Mo. Mag. Vol. 49, p. 14 (1913).
78. *H. hirsuta*, Becker, Mém. Acad. Sc. Petrograd, Vol. 28 (7), p. 59 (1915). N. W. Siberia.
79. *H. hirta*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 141 (1892); C. Europe.
Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892).
80. *H. hirtula*, Zetterstedt, Fauna Ins. Lappon, p. 556 (1838); Dipt. Scand. Vol. 1, p. 348 (1842); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 179 (1892).
81. *H. Hudsoni*, Hutton, Trans. New Zeal. Inst. Vol. 33, p. 30 [1901] (*Empis*). New Zealand.
82. *H. hystrix*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 1112 (1892); C. Europe.
Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892).
83. *H. infans*, Zetterstedt, Dipt. Scand. Vol. 1, p. 346 (1842); ibidem, Vol. 8, N. Europe.
p. 3016 (1849); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 399 (1866); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 180 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 47 (1913).
infuscata, Brullé, Expéd. Morée, Zool. Vol. 3, Pt. 1, p. 300 (1832); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 181 (1892).
84. *H. interstincta*, Fallen, Empid. Suct. p. 24 [1816] (*Empis*); Zetterstedt, N. & C. Europe.
Fauna Ins. Lappon. p. 555 (1838); Dipt. Scand. Vol. 1, p. 343 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 490 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3016 (1849); Walker, Ins. Brit. Dipt. Vol. 1, p. 100 (1851); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Nylander, ibidem, Vol. 4, p. 247 (1858); Bonsdorff, Finl. Tväv. Ins. Dipt. Vol. 1, p. 160 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 152 (1864); ibidem, Vol. 14, p. 399 (1866); Cat. Dipt. Norv. p. 27 (1877); Beling, Arch. Naturg. Berlin, Vol. 48, Pt. 1, p. 218 (1882); Neuhaus, Dipt. March. p. 71 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 127 (1887); Girschner, Ent. Nachr. Vol. 15, p. 220 (1889); Mik, Wien. Ent. Zeit. Vol. 11, p. 78 (1892); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 131 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1892); Mik, Wien. Ent. Zeit. Vol. 13, p. 283 (1894); Lundbeck, Dipt. Danica, Vol. 3, p. 160, f. 58 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 54 (1913).
modesta, Meigen, Syst. Beschr. Vol. 3, p. 10 (1822).
pilosa, Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 160 (1861).
85. *H. irritans*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 378 (1909). Chile.
86. *H. Johnsoni*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 393 (1895); Alabama.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 268 (1902).
87. *H. lacteipennis*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 137 (1892); C. Europe.
88. *H. lasiochira*, ibidem. Vol. 42, p. 110 (1892); C. Europe.
89. *H. lasiopyga*, Lundbeck, Dipt. Danica, Vol. 3, p. 178, f. 68, 69 (1910). Denmark.
90. *H. laureæ*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 42 (1908). Canary Islands.
91. *H. leucoptera*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 205 : Cent. 3, n° 43 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 395 (1895); Florida.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 266 (1902).

92. *H.* ? litigiosa*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 93, 120, pl. 10, f. 14, 15, pl. 11, f. 1 (1908).
93. *H. litorea*, Fallen, Empid. Suec. p. 24 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 8 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 34 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 351 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 490 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3017 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 101 1851; ? Walker, Ins. Saunders. Dipt. Vol. 1, p. 204 (1852); Schiner, Fauna Dipt. Austr. Vol. 1, p. 113, 114 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 129 (1887); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 161 (1892); Mitteil. Naturw. Ver. Steiermark. Graz, Vol. 29, p. 91 (1892); Grünberg, Süsswasserf. Deutschl. 2a, Vol. 1, p. 155 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 174, f. 65 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 47 (1913).
univittata, Meigen, Syst. Beschr. Vol. 3, p. 9 (1822).
95. *H. longesetosa*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, Austria, p. 67 (1909).
96. *H. longevittata*, Zetterstedt, Dipt. Scand. Vol. 1, p. 358 [1842] (*longivittata*); Strobl, Verg. Zool.-bot. Ges. Wien, Vol. 42, p. 118 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892); ibidem, Vol. 46, p. 68 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 56 (1913).
var. *andermattensis*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 119 (1892). C. Europe.
var. *major*, Strobl (1909, not 1892), Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 68 (1909). C. Europe.
var. *styriaca*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 68 (1909). C. Europe.
97. *H. longicornis*, Strobl, Wien. Ent. Zeit. Vol. 13, p. 59 (1894); Becker, C. Europe.
ibidem, Vol. 13, p. 156 (1894) nov. gen. ?
98. *H. ? longirostris*, Macquart, Mém. Soc. Sc. Lille, p. 164 [1823] (*Empis*); France. Dipt. N. France, Vol. 3, p. 115 (1827); Meigen, Syst. Beschr. Vol. 7, p. 81 (1838); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 181 [1892] (? *Empis*).
99. *H. lucidifrons*, Becker, Bull. Mus. Hist. Nat. Paris, p. 116 (1909); E. Africa.
Ann. Soc. Ent. France, Vol. 79, p. 25 (1910).
100. *H. lugubris*, Zetterstedt, Svenska Vet. Akad. Handl. 1819, p. 81 [1819] (*Empis*); Fallen, Dipt. Suec. Suppl. 8 [1826] (*Empis*); Zetterstedt, Dipt. Scand. Vol. 1, p. 341 (1842); ibidem, Vol. 8, p. 3016 (1849); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 160 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Engel, Ent. Nachr. Berlin, Vol. 12, p. 45 (1886); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 127 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1892); Grünberg, Süsswasserf. Deutschl. 2a, Vol. 1, p. 155 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912).
trigramma, Meigen, Syst. Beschr. Vol. 6, p. 337 (1830); Schiner, Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Neuhaus, Dipt. March. p. 71 (1886).
101. *H. Lundbecki*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 54 (1913); Ent. Tidskr. 1919, p. 80 (1919). Sweden, Denmark, England.
pilipes, Zetterstedt, Dipt. Scand. Vol. 1, p. 346 (1842); ibidem, Vol. 8, p. 3016 (1849); ibidem, Vol. 12, p. 4606 (1855); Lundbeck, Dipt. Dan. Vol. 3, p. 171, f. 64 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).

102. *H. lurida*, Fallen, Empid. Suec. p. 22 [1816] (*Empis*); Meigen, Syst. C. & N. Europe. Beschr. Vol. 3, p. 8 (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 336 (1842); ibidem, Vol. 8, p. 3014 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 50 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 100 (1851); Pipping, Not Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 159 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 114 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 399 (1866); Brauer, Denkschr. Akad. Wiss. Wien, Vol. 47, p. 44, pl. 4, f. 77, 79 (1883); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 171 (1892); Grünberg, Süßwasserf. Deutschl. 2a, Vol. 1, p. 155 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 177, f. 67 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 46 (1913).
103. *H. lutea*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 18 : Cent. 4, No. 33 E. United States. (1863); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 267 (1902).
104. *H. macroptera*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 18 : Cent. 4, No. 32 E. United States. (1863); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 267 (1902); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1905).
105. *H. maluinensis*, Enderlein, Svenska Vet. Akad. Handl. Vol. 48 (3), Falkland Islands. p. 48, f. 7 (1912).
106. *H. manicata*, Meigen, Syst. Beschr. Vol. 3, p. 5 (1822); Macquart, Dipt. C. & N. Europe. N. France, Vol. 3, p. 112 (1827); Hist. Nat. Dipt. Vol. 1, p. 341 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 348 (1842); ibidem, Vol. 8, p. 3016 (1849); Walker, Ins. Brit. Dipt. Vol. 1, p. 99 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 114 (1862); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 163 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910).
pallens, Macquart, Mém. Soc. Sc. Lille, p. 163 [1823] (*Empis*).
squalens, Zetterstedt, Fauna Ins. Lappon, p. 556 (1838); Dipt. Sc. Vol. 1, p. 349 (1842); ibidem, Vol. 12, p. 4606 (1855).
107. *H. marginipennis*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 176 Spain. (1909).
108. *H. matrona*, Haliday, Ent. Mag. London, Vol. 1, p. 158 (1833); Walker, C. Europe. Ins. Brit. Dipt. Vol. 1, p. 98 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 113 (1862); Beling, Arch. Naturg. Berlin, Vol. 48, Pt. 1, p. 221 (1882); ?Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 168 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 91 (1892); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912).
109. *H. matroniformis*, Strobl, Wien. Ent. Zeit. Vol. 12, p. 40 (1893); S. Europe. Glasnik, Zem. Mus. Bosn. Herceg. Sarajevo, Vol. 14, p. 471 (1902); Mitth. Bosn. Herceg. Sarajevo, Vol. 9, p. 530 (1904).
110. *H. maura*, Fabricius, Spec. Ins. Vol. 2, p. 471 [1781] (*Empis*); Mant. Europe. Ins. Vol 2, p. 364 [1787] (*Empis*); Gmelin, Syst. Nat. Vol. 5, p. 2889 [1790] (*Empis*); Olivier, Encycl. Méth. Vol. 6, p. 387 [1791] (*Empis*); Fabricius, Ent. Syst. Vol. 4, p. 404 [1794] (*Empis*); Schellenberg, Gatt. Fliegen, p. 88, 89, pl. 30, f. 4 [1803] (*Empis*); Meigen, Classif. Beschr. Ent. Zweifl. Ins. Vol. 1, p. 222 [1804] (*Empis*); Fabricius, Syst. Antl. p. 139 [1805] (*Empis*); Fallen, Empid. Suec. p. 23 [1816] (*Empis*); Macquart, Mém. Soc. Sc. Lille, 1823, p. 162 [1823] (*Empis*); Zetterstedt, Fauna Ins. Lappon, p. 555 (1838); Dipt. Scand. Vol. 1, p. 341 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 490 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3015 (1849);

Walker, Ins. Brit. Dipt. Vol. 1, p. 98 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Beling, Arch. Naturg. Berlin, Vol. 48, p. 220 (1882); Neuhaus, Dipt. March. p. 71 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 128 (1887); Girschner, Ent. Nachr. Berlin, Vol. 15, p. 220 (1889); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 126 (1892); Mik. Wien. Ent. Zeit. Vol. 13, p. 283 (1894); Howlett, Ent. Mag. London (2), Vol. 13, p. 231 (1907); Grünberg, Süsswasserf. Deutschl. 2a, Vol. 1, p. 155, f. 199 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 159, f. 46, 57 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 55 (1913). — **Pl. 6, Fig. 55.**

crassipes, Fourcroy, Ent. Paris, Vol. 2, p. 465 [1785] (*Asilus*).

globulipes, Meigen, Syst. Beschr. Vol. 3, p. 3 (1822); Macquart, Dipt. N. France, Vol. 3, p. 110 (1827); Hist. Nat. Dipt. Vol. 1, p. 340 (1834); Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt. p. 20, f. 15 (1840); Isis, Vol. 7, p. 546, f. 15 (1840); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 [1862] (*globuliceps*); Leunis, Syn. Zool. Vol. 2, p. 402 (1886).

? *simplex*, Wiedemann, Zool. Mag. Vol. 1, p. 70 [1817] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 28 [1822] (*Empis*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 [1862] (*Empis*); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 58 (1867).

111. *H. melanochira*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 474 (1912). Formosa.
112. *H. migrata*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 491 (1849); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 271 (1902). Hudson Bay.
113. *H. minuta*, Zetterstedt, Dipt. Scand. Vol. 1, p. 359 (1842); ibidem, C. & N. Europe. Vol. 8, p. 3018 (1849); Loew, Nat. Hist. Rev. London, Vol. 3, p. 86 [1856] (*Ragas*); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 120 (1892); Meijere, Tijdschr. Ent. Vol. 1, p. 157, pl. 4, f. 8 [1907] (*Ragas*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 68 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 182, f. 70 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 65 [1910] (*Ragas*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 46 [1913] (*Ragas*).
nudinervia, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 121 (1892).
114. *H. mollicella*, White, Proc. Roy. Soc. Tasmania, 1916, p. 225, f. 41e Tasmania. (1917).
115. *H. mutabilis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 204 : Cent. 3, C. United States. No. 40 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 265 (1902).
116. *H. nigrina*, Fallen, Empid. Suec. p. 24 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 4, pl. 22, f. 5 (1822); Macquart, Dipt. N. France, Vol. 3, p. 112 (1827); Hist. Nat. Dipt. Vol. 1, p. 341 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 354 (1842); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 490 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3017 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 99 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 4270 (1852); Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 384 (1866); Neuhaus, Dipt. March. p. 70 (1886); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 111 (1892); Lundbeck, Dipt. Danica, Vol. 3 p. 156, f. 55 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).
117. *H. nigriventris*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 203 : Cent. 3, No. 38 E. United States. (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 265 (1902).
118. *H. nimia*, White, Proc. Roy. Soc. Tasmania, 1916, p. 223, f. 41c (1917). Tasmania.

119. *H. nitidula*, Zetterstedt, Fauna Ins. Lappon, p. 556 (1838); Dipt. Scand. C. & N. Europe. Vol. 1, p. 355 (1842); ibidem, Vol. 8, p. 3017 (1849); ibidem, Vol. 13, p. 5002 (1859); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 162 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 153 (1864); Siebke, Cat. Dipt. Norv. p. 29 (1877); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 127 (1887); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 123, note (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 76, f. 7 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3) p. 55 (1913).
var. *femorella*, Zetterstedt, Dipt. Scand. Vol. 1, p. 355 (1842); ibidem, Vol. 8, Europe. p. 3018 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 153 (1864); ibidem, Vol. 14, p. 399 (1866); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 127 (1887); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 122 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).
pinetorum, Schiner (not Zetterstedt), Fauna Dipt. Austr. Vol. 1, p. 114 (1862).
120. *H. niveipennis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 352 (1842); ibidem, C. & N. Europe. Vol. 8, p. 3017 (1849); Schiner, Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 149 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 173 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910).
121. *H. Novakii*, Mik, Wien. Ent. Zeit. Vol. 11, p. 83 (1892); Strobl, Verh. S. Europe. Zool.-bot. Ges. Wien, Vol. 42, p. 158, note (1892).
Mikii, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 158 (1892).
122. *H. nubila*, White, Proc. Roy. Soc. Tasmania, 1916, p. 224, f. 41d (1917). Tasmania.
123. *H. nugax*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 273 (1902). California.
124. *H. obscuritarsis*, Zetterstedt, Dipt. Scand. Vol. 13, p. 4999 (1859); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 180 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 47 (1913).
abdominalis, Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 159 (1861).
125. *H. orientalis*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 474 (1912). Formosa.
126. *H. palmarum*, Strobl, Mém. Soc. Esp. Hist. Nat. Vol. 3, p. 309 (1906). Spain.
127. *H. paludosa*, Becker, Deutsche Ent. Zeitschr. 1910, p. 643 (1910). Corsica.
128. *H. pectinipes*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 116 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892). C. Europe.
129. *H. perplexa*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 379 (1909). Peru.
130. *H. perturbans*, Bezzi, ibidem, Vol. 91, p. 380 (1909). Chile, Peru.
131. *H. perversa*, Oldenberg, Arch. Naturg. Berlin, Vol. 81; A (9), p. 166 (1916). Germany.
132. *H. peshawarensis*, Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 360 (1920). N. W. India.
133. *H. philina*, Speiser, Kilimandjaro-Meru Exped. Vol. 10, p. 108 (1910). Africa.
134. *H. pilipes*, Zetterstedt, Fauna Ins. Lappon. p. 555 (1838); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 161 (1861); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 51, f. 8 (1913). N. Europe.
anomala, Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt., p. 19, f. 12, 13 (1840); Isis, Vol. 7, p. 554, f. 12, 13 (1840); Schiner, Fauna, Zool.-bot. Dipt. Austr. Vol. 1, p. 116 (1862); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 113, 178 (1892); Oldenberg, Arch. Naturg. Berlin, Vol. 80 (9), p. 171 (1916).
brevivittata, Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 161 (1861).

135. *H. pilosa*, Zetterstedt, Dipt. Scand. Vol. 1, p. 324 (1842); Walker, Ins. C. & N. Europe. Brit. Dipt. Vol. 1, p. 99 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Beling, Arch. Naturg. Vol. 48, p. 219 (1882); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 129 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Oldenberg, Arch. Naturg. Berlin, Vol. 81 (9), p. 169 (1916).
interstincta, Meigen (not Fallen), Syst. Beschr. Vol. 3, p. 6 (1822); ibidem, Vol. 7, p. 80 (1838).
? senilis, Panzer, Fauna Germ. liv. pl. 3 [1798] (*Bibio*); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 235 [1804] (*Empis*).
spinipes, Macquart, Dipt. N. France, Vol. 3, p. 112 (1827); Hist. Nat. Dipt. Vol. 1, p. 341 (1834).
136. *H. pilosopectinata*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 169 C. Europe. (1892).
137. *H. pinetorum*, Zetterstedt, Dipt. Sc. Vol. 8, p. 3017 (1849); Jaroschewsky, C. & N. Europe. Arb. Ges. Naturf. Univ. Kharkoff, Vol. 11, p. 356 (1877); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 115 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1893); Wien. Ent. Zeit. Vol. 18, p. 20 (1899); Poppius, Medd. Soc. Fenn. Helsingfors, Vol. 32, p. 107 (1906); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910). var. *major*, Strobl (1892, not 1909), Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 116 C. Europe. (1892).
138. *H. plebeia*, Walker, Trans. Ent. Soc. Lond. n. s. Vol. 4, p. 148 (1857); North America. Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 270 (1902).
139. *H. pruinosa*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 7 (1822); C. & S. Europe. Walker, List Dipt. Brit. Mus. Vol. 3, p. 490 (1849); Ins. Brit. Dipt. Vol. 1, p. 100 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 114 (1862); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 133 (1892); Mik, Wien. Ent. Zeit. Vol. 13, p. 283 (1894); Progr. Akad. Gym. Wien, p. 19 (1894).
vulnerata, Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Becher, Denkschr. Akad. Wiss. Wien, Vol. 45, p. 147, pl. 3, f. 13 (1882); Mik, in Beck, Fauna Bernstein, Vol. 2 (2), p. 59 (1885).
140. *H. pseudochorica*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 109 C. & N. Europe. (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 88 (1892); ibidem, Vol. 46, p. 67 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 155, f. 54 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 56 (1913).
141. *H. pseudocornicula*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 174 Spain. (1909).
142. *H. pseudosartrix*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 152 C. Europe. (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892). var. *galactoptera*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 69 C. Europe. (1909).
143. *H. pubipes*, Loew, Berl. Ent. Zeitschr. Vol. 18, p. 43 (1873); Strobl, C. Europe. Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 139 (1892); Mitteil. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892); Lundbeck, Dipt. Danica, Vol. 3, p. 164, f. 60, 47 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Verrall, Mag. London, Vol. 48, p. 24 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 49 (1913).
quadriserrata, Pipping (not Meigen), Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 159 (1861).
144. *H. pulchripes*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 47, Finland. f. 6 (1913).

145. *H. quadriclarata*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 21 (1899). Spain.
146. *H. quadrifaria*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 416 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 87 (1892); Wien, Ent. Zeit. Vol. 18, p. 20 (1899); Becker, Zeitsch. Hym. Dipt. Vol. 7, p. 123 (1907); Lundbeck, Dipt. Danica, Vol. 3, p. 153, f. 52 (1910); Collin, Ent. M. Mag. London (2), Vol. 24, p. 106 (1913); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 56 (1913). var. *longeciliata*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 309 (1906). Spain.
147. *H. quadrifilosa*, Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 30 (1900). Siberia.
148. *H. quadrivittata*, Meigen, Syst. Beschr. Vol. 3, p. 7 (1822); Macquart, Dipt. N. France, Vol. 3, p. 113 (1827); Hist. Nat. Dipt. Vol. 1, p. 342 (1834); Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt. p. 20 (1840); Isis, Vol. 7, p. 546 (1840); Boitard, Man. Ent. Vol. 3, p. 321 (1843); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 51 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 100 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Beling, Arch. Naturg. Berlin, Vol. 48, p. 221 (1882); Verh. Zool.-bot. Wien, Vol. 38, p. 2 (1888); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 142 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892); Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 412 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 264 (1902); Grünberg, Süßwasserf. Deutschl. 2a, Vol. 1, p. 156 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 165 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 76 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 27 (3), p. 49 (1913).
- intermedia*, Fallen, Empid. Succ. p. 23, part. [1816] (*Empis*).
quadrilineata, Macquart, Mém. Soc. Sc. Lille, p. 162 [1823] (*Empis*).
149. *H. recedens*, Haliday, in Walker, Ins. Brit. Dipt. Vol. 1, p. 101 (1851); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 179 (1892). England.
150. *H. rufipes*, Macquart, Mém. Soc. Sc. Lille, p. 161 [1823] (*Empis*); Dipt. N. France, Vol. 3, p. 114 (1827); Hist. Nat. Dipt. Vol. 1, p. 343 (1834); Meigen, Syst. Beschr. Vol. 7, p. 80 (1838). C. Europe.
- fulvipes*, Macquart, Hist. Nat. Dipt. Vol. 1, p. 342 (1834); Meigen, Syst. Beschr. Vol. 7, p. 81 (1838); Boitard, Man. Ent. Vol. 3, p. 321 (1843); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 181 (1892).
151. *H. sartor*, Becker, Berl. Ent. Zeitschr. Vol. 32, p. 7 (1888); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 38, Sitzgsber. p. 97 (1888); Wien. Ent. Zeit. Vol. 11, p. 78 (1892); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 151 (1892); Girschner, Ent. Nachr. Berlin, Vol. 20, p. 61, 241 (1894); Mik, ibidem, Vol. 20, p. 49, 151 (1894); Wien. Ent. Zeit. Vol. 13, p. 197, 273, pl. 2, f. 8, 13 (1894); Progr. Akad. Gymn. Wien, p. 11, f. 8, 13 (1894); Wasmann, Bericht Offenb. Ver. Naturk. Offenbach, Vol. 41, p. 436 (1895). C. Europe, Alps
- alpina*, Loew, in litt. Osten-Sacken, Ent. Mag. London, Vol. 14, p. 216 (1877); Ent. Nachr. Berlin, Vol. 12, p. 1 (1886).
- sartrix*, Handlirsch, Verh. Zool.-bot. Ges. Wien, Vol. 39, p. 623 (1889); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892).
152. *H. scrobiculata*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 41 (1873); Kowarz, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 457 (1873); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 127 (1887); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 130 (1892); Mitteil. Ver. Naturk. Steiermark, Graz, Vol. 29, p. 89 (1892); Becker, Deutsche Ent. Zeitschr. p. 643 (1910); Oldenberg, Arch. Naturg. Berlin, Vol. 81, A (9), p. 169 (1916). C. & S. Europe.
- cerulea*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 128 (1887); Oldenberg, Arch. Naturg. Berlin, Vol. 81, A (9), p. 170 (1916).

153. *H. seriata*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 82 : Cent. 5, No. 63 (1864); E. United States.
Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander,
Trans. Amer. Ent. Soc. Vol. 28, p. 268 (1902).
154. *H. simplicipes*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 119 (1892). C. Europe.
155. *H. Smithii*, Hutton, Trans. New. Zeal. Inst. Vol. 33, p. 30 [1901] New Zealand.
(*Empis*).
156. *H. spinimana*, Zetterstedt, Fauna Ins. Lappon, p. 556 (1838); Dipt. Scand. Vol. 1, p. 344 (1842); ibidem, Vol. 8, p. 3016 (1849); ibidem, Vol. 11, p. 4269 (1852); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 384 (1866); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910).
ciliipes, Schiner (not Meigen), Fauna Dipt. Austr. Vol. 1, p. 113 (1862).
var. *spinigera*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 169 (1892); C. & N. Europe.
Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 91 (1892).
157. *H. striaticollis*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 122 (1907). Algeria.
158. *H. strobliana*, Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 142 (1899). S. Europe.
159. *H. sublineata*, Brullé, Expéd. Morée (Zool.), Vol. 3, p. 300 (1832);
Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 181 (1892). Greece.
160. *H. sulcifarsis*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 123 (1892); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 89 (1892).
161. *H. tanychira*, Kowarz, in litt. Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 148 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 173 (1910);
Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 54 (1913).
pilipes, Strobl (not Zetterstedt), Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 147 (1892).
162. *H. tanyhrix*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 50, f. 7 (1913). Finland.
163. *H. tarda*, Meunier, Ann. Sc. Nat. (Zool.), Vol. 7, p. 93, 120, pl. 10, f. 16 (1908). Baltic Amber.
164. *H. tarsata*, Siebke, Nyt Mag. Naturvid. Vol. 12, p. 152 (1864); Cat. Dipt. Nov. p. 27 (1877). N. Europe.
165. *H. tenella*, Fallen, Empid. Suec. p. 25 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 9 (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 353 (1842); Walker, Ins. Brit. Dipt. Vol. 1, p. 101 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 115 (1862); Neuhaus, Dipt. March. Vol. 71 (1886); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 176 (1892); Grünberg, Süßwasserf. Deutschl. 2a, Vol. 1, p. 156 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 77 (1910).
166. *H. tenuinervis*, Zetterstedt, Fauna Dipt. Lappon, p. 557 (1838); Dipt. Scand. Vol. 1, p. 349 (1842); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 178 (1892); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 57 (1913). N. Europe.
167. *H. ternovensis*, Strobl, Glasnik. Zem. Mus. Bosn. Herceg. Vol. 10, p. 413 (1898); Mitteil. Bosn. Herceg. Vol. 7, p. 572 (1900). S. Europe.
168. *H. testacea*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 82 : Cent. 5, No. 64 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 268 (1902). New Hampshire.
169. *H. tetragramma*, Loew, Berl. Ent. Zeitschr. Vol. 18, p. 43 (1873); C. Europe.
Kowarz, Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 457 (1873); Strobl, ibidem, Vol. 42, p. 138 (1892); Lichtwardt, Zeitschr. Hym. Dipt. Vol. 5, p. 310 (1905).
170. *H. thoracica*, Macquart, Dipt. N. France, Vol. 3, p. 115 (1827); Hist. Nat. Dipt. Vol. 1, p. 343 (1834); Meigen, Syst. Beschr. Vol. 7, p. 81 (1838); Walker, List Dipt. Brit. Mus. Vol. 3, p. 491 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 52 (1851); Walker,

- Ins. Brit. Dipt. Vol. 1, p. 102 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark. Graz, Vol. 29, p. 92 (1892); ibidem, Vol. 46, p. 71 (1909); Boitard, Man. Ent. Vol. 3, p. 321 (1843).
? ferruginea, Roser, Correspondenzblad Landw. Ver. Würtemb. Stuttgart, Vol 1, p. 53 (1840).
? melanocephala, Macquart, Mém. Soc. Sc. Lille, 1823, p. 161 [1823] (*Empis*).
magica, Mik, Wien. Ent. Zeit, Vol. 6, p. 100 (1887); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 175 (1892).
171. *H. Tiefi*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 150 (1892); C. Europe.
 Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 90 (1892).
172. *H. transfuga*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 491 (1849); Boreal America.
 Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 411 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 270 (1902).
173. *H. tristis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 82 : Cent. 5, No. 62 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 267 (1902); Slosson, Ent. News. Philad. Vol. 14, p. 286 (1903). New Hampshire.
174. *H. trivittata*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 204 : Cent. 3, No. 39 (1862); Coquillett, Proc. U. S. Nat. Mus., Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 200, 265 (1902); Slosson, Ent. News. Philad. Vol. 14, p. 268 [1903] (*habits*); McAtee, Ent. News. Philad. Vol. 20, p. 360 [1909] (*habits*). United States.
175. *H. tyrolensis*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 121 (1892). C. Europe.
176. *H. umbrosa*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 202 : Cent. 3, No. 34 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 264 et 350 (1902). E. United States.
brachystoma, Coquillett, Rept. New Jersey Board Agric. Vol. 29, Suppl. p. 652 [1899] (*Empis*) no description; Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 350 (1902).
177. *H. uncicauda*, Bezz, Suppl. Ent. Berlin, Vol. 3, p. 72 (1914). Formosa.
178. *H. unicolor*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 203 : Cent. 3, No. 37 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 265 (1902). E. United States.
179. *H. velutina*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 203 : Cent. 3, No. 36 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 394 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 264 (1902). E. United States.
180. *H. Wheeleri*, Melander, Psyche, Vol. 9, p. 214 (1901); Trans. Amer. Ent. Soc. Vol. 28, p. 270, f. 95 (1902). Wyoming.

16. GENUS PHLEBOCTENA, BEZZI

Phleboctena, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 382 (1909).

Characters. — Yellow species, measuring six millimeters, with stout thorax and short abdomen. Eyes separated on the front a distance equal to the length of the first antennal joint, two ocellar bristles; antennæ elongate, almost bare, third joint long and compressed conical, terminal style two-thirds the length of the third joint, distinctly two-jointed, the basal joint quadrate, the long terminal joint tipped by a microscopic seta; proboscis thick, vertical, as long as the head; palpi long, thin, porrect and finely hairy. Disk of the thorax with very short hairs, notopleural, postalar, single dorsocentral and four scutellar bristles long, no strong humerals, pleuræ bare. Genitalia retracted, finely golden-hairy. Legs simple, with very short hairs, only the coxæ and posterior femora and tibiae bearing short black bristles. Wings long and broad, costa thinned beyond the third vein, basal bristle present, auxiliary

vein abbreviated, first vein distinctly and closely setulose on its entire length except at extreme base, third vein forked, the anterior branch almost parallel with the axis, the posterior branch bent backward to terminate at the wing-tip, discal cell large, as long as the first basal cell, anal vein obsolete, anal angle well developed, no alula.

Type species : *P. lutea*, Bezzi. Dr. Bezzi has forwarded the only known specimen which has served to check the preceding description and to make possible the accompanying illustration.

Geographical distribution.

1. *P. lutea*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 382, f. 7, 8 Chile.
(1909). — Pl. 2, Fig. 15.

17. GENUS HILAREMPIS, BEZZI

Hilarempis, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 443 (1905); Melander, Williston's Man. N. Amer. Dipt. Vol. 226 (1908); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 362 (1909); Kertész, Cat. Dipt. Vol. 6, p. 82 (1909).

Heterempis, Brethes, Ann. Mus. Hist. Nat. Buenos Aires, Vol. 19, p. 92 (1909); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 362, note (1909).

Hilaropus, White, Proc. R. Soc. Tasmania, 1916, p. 226 (1917).

Characters. — Opaque blackish species. Head globose, eyes of both male and female widely separated, face quadrate, hairy or bare, lower occiput hairy, upper occiput more bristly; antennæ shorter than the head, located near the middle, distinctly three-jointed, the basal joints not setose, the third joint conical with a short terminal style which is tipped by a small bristle; proboscis not longer than the head, rather strong but slender, vertical, palpi thickened, cylindrical, projecting forward but curving upward, setose beneath. Discal bristles of the thorax small, one humeral, three prominent notopleurals, the acrostichals minute, arranged in two approximate or several rows, dorsocentrals small, scutellum margined with a variable number of bristles ranging from six to fourteen; pleurae bare. Abdomen with long loose hair, base of the second segment marked with two rows of pittings; pygidium usually strongly compressed and thrown forward over the abdomen, penis hidden. Legs not bristly, front tibiae and metatarsi of the male sometimes shaggy, hind femora not thickened nor the hind tibiae shortened, front metatarsi of male often enlarged as in *Hilara*. Costa extending around the entire wing although sometimes thinner on the hind margin, anal angle broadly rounded, auxiliary vein straight, vanishing at the end, first vein swollen at the tip, upper branch of the third vein short, anal cell narrow, anal vein detached from the underside of the anal cell.

Type species : *H. nudifacies*, Bezzi (Pl. 6, Fig. 54), by present designation. The genus is subtropical in its distribution.

Geographical distribution.

1. *H. argentifera*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 365, 371 Chile.
(1909).
2. *H. argentula*, Becker, Mission Arc Méridien Amér. Sud, Vol. 10, p. 169 Ecuador.
(1910).
3. *H. argyrosoma*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 761 Chile.
[1865] (?*Hilara*); Bezzii, Ann. Mus. Hungar. Vol. 3, p. 458 (1905);
Nova Acta Akad. Naturf. Halle, Vol. 91, p. 363 (1909).
4. *H. barbatula*, Bezzi, ibidem, Vol. 91, p. 365, 371 (1909). Peru.

5. *H. Benhami*, Miller, Trans. New Zeal. Inst. Vol. 45, p. 198, f. 1, 5 [1913] New Zealand.
(*Hilara*).
6. *H. bicingulata*, Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 16, p. 364, Chile.
367 (1909).
7. *H. brachyrrhyncha*, Thomsen, Eugen. Resa, Dipt. p. 474 [1869] (*Hilara*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 364 (1909).
8. *H. brachystoma*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 755 Chile.
[1865] (*Empis*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 448 [1905]
(*Haplomera*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 364, 373 (1909).
9. *H. carinata*, Bezzī, ibidem, Vol. 91, p. 363, 369 (1909). Bolivia.
10. *H. ?cervina*, Loew, Oefv. Vet. Akad. Förh. Stockholm, Vol. 14, p. 370 S. Africa.
[1857] (*Hilara*); Dipterenf. Südafr. Vol. 1, p. 267 [1860] (*Hilara*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 443 (1905).
11. *H. cyanescens*, Bezzī, ibidem, Vol. 2, p. 342 [1904] (*Empis*); Vol. 3, p. 443 New South Wales.
(1905).
12. *H. dumicola*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 756 [1865] Chile.
(*Empis*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 364 (1909).
13. *H. echinata*, White, Proc. Roy. Soc. Tasmania, 1916, p. 230 [1917] Tasmania.
(*Hilaropus*).
14. *H. elegans*, Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 364, 368 Peru.
(1909).
15. *H. fulva*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 765 [1865] Chile.
(*Empis*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905).
Philippii, Bezzī (new name for *Empis fulva* Philippi, not Macquart nor Walker), Nova Acta Akad. Naturf. Halle, Vol. 91, p. 363 (1909).
16. *H. fulvipes*, Hutton, Trans. New Zeal. Inst. Vol. 33, p. 31 [1901] (*Hilara*). New Zealand.
Huttoni, Bezzī (new name for *Hilara fulvipes* Hutton, not Macquart), Ann. Mus. Hungar. Vol. 2, p. 360 (1904).
17. *H. griseiventris*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 761 Chile.
[1865] (*Hilara*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 365 (1909).
18. *H. gymnaspis*, Bezzī, ibidem, Vol. 91, p. 365, 370 (1909). Peru.
19. *H. ?heterogastra*, Loew, Vet. Akad. Förh. Stockholm, Vol. 14, p. 370 S. Africa.
[1858] (*Hilara*); Dipterenf. Südafr. Vol. 1, p. 268 [1860] (*Hilara*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 443 (1905).
20. *H. hilaræformis*, Bezzī, ibidem, Vol. 2, p. 240 [1904] (*Empis*); ibidem, Australia.
Vol. 3, p. 443 (1905).
21. *H. holosericea*, Thomsen, Eugen. Resa, Dipt. p. 473 [1869] (*Hilara*); Bezzī, Patagonia.
Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 364 (1909).
22. *H. kaiteriensis*, Miller, Trans. New Zeal. Inst. Vol. 45, p. 200, f. 6-8 New Zealand.
[1913] (*Hilara*).
23. *H. laticornis*, Bigot, Mission Sc. Cap Horn, Zool. Vol. 6, p. 20 [1888] Cape Horn.
(*Hilara*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 364 (1909).
24. *H. macrocera*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 128 [1889] Chile.
(*Hilara*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 365 (1909).
25. *H. magellanica*, Bigot, Mission Sc. Cap Horn, Zool. Vol. 6, p. 20 [1888] Cape Horn.
(*Hilara*); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 364 (1909); Enderlein, Vet. Akad. Handl. Stockholm, Vol. 47 (3) 103 (1912).

26. *H. mendozana*, Brèthes, An. Mus. Hist. Nat. Buenos Aires, Vol. 19, p. 92, Argentina.
fig. [1909] (*Heterempis*).
 27. *H. nigrimana*, White, Proc. Roy. Soc. Tasmania, 1916, p. 228 [1917] Tasmania.
(*Hilaropus*).
 28. *H. nudifacies*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 446 (1905); Nova Peru.
Acta Akad. Naturf. Halle, Vol. 91, p. 364 (1909). — Pl. 6, Fig. 54.
 29. *H. ochracea*, Bigot, Mission Sc. Cap Horn, Zool. Vol. 6, p. 22, pl. 3, f. 4 Cape Horn.
[1888] (*Heleodromia*); Ann. Soc. Ent. France (6), Vol. 8, Bull. p. XXX
[1888] (*Hilara*); Bull. Soc. Zool. France, Vol. 13, p. 101 [1888]
(*Hilara*); Wien. Ent. Zeit. Vol. 7, p. 109 [1888] (*Hilara*); Ann. Soc.
Ent. France (6), Vol. 9, p. 113 [1889] (*Hilara*); Bezzi, Ann. Mus.
Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad. Naturf. Halle,
Vol. 91, p. 363 (1909).
 30. *H. pallida*, Philippi, Zool.-bot. Ges. Wien, Vol. 15, p. 761 [1865] Chile.
(? *Hilara*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 458 (1905); Nova
Acta Akad. Naturf. Halle, Vol. 91, p. 363 (1909).
 31. *H. pallidifurca*, White, Proc. Roy. Soc. Tasmania, 1916, p. 227 [1917] Tasmania.
(*Hilaropus*).
 32. *H. peregrina*, White, ibidem, p. 228, f. 43 [1917] (*Hilaropus*). Tasmania.
 33. *H. Philpotti*, Miller, Trans. New Zeal. Inst. Vol. 45, p. 202, f. 10 [1913] New Zealand.
(*Hilara*).
 34. *H. polychæta*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 445 (1905); Nova Acta Peru.
Akad. Naturf. Halle, Vol. 91, p. 365 (1909).
 35. *H. quadrifaria*, Becker, Mission Arc Méridien Amér. Sud. Vol. 10, p. 169 Ecuador.
(1919).
 36. *H. ? sordida*, Loew, Oefv. Vet. Akad. Förh. Stockholm, Vol. 15, p. 340 S. Africa.
[1858] (*Hilara*); Dipterent. Südafr. Vol. 1, p. 267, pl. 2, f. 51 [1860]
(*Hilara*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 443 (1905).
 37. *H. spinosa*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 363, 365 Chile.
(1909).
 38. *H. spinulosa*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 757 [1865] Chile.
(*Empis*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 444 (1905); Nova
Acta Akad. Naturf. Halle, Vol. 91, p. 363, 365 (1909).
 39. *H. tephrodes*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 755 [1865] Chile.
(*Empis*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova
Acta Akad. Naturf. Halle, Vol. 91, p. 365 (1909).
 40. *H. vanellus*, Schiner, Novara Reise, Dipt. p. 206 [1868] (*Hilara*); Bezzi, Colombia.
Ann. Mus. Hungar. Vol. 3, p. 444, 458 (1905); Nova Acta Akad.
Naturf. Halle, Vol. 91, p. 364 (1909).
 41. *H. xanthocera*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 444 (1905); Nova Peru.
Acta Akad. Naturf. Halle, Vol. 91, p. 364, 367 (1909).

18. GENUS HAPLOMERA, MACQUART

Haplomera, Macquart, Dipt. Exot. Vol. 1, 2, p. 279 or 163 [1838] (*Aplomera*); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 561 [1857] (*Aplomera*); Scudder, Nomencl. Univ. Index, p. 141 (1884); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 119 [1889] (*Aplomera*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448 (1905); Melander, Williston's Man. N. Amer. Dipt. p. 226 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 372 (1909); Kertész, Cat. Dipt. Vol. 6, p. 83 (1909).

Anodontina, Macquart, Dipt. Exot. Vol. 1, Pt. 2, pl. 13, f. 4 (1838) figure only.

Characters. — Opaque blackish species with robust hind legs. Head globose, eyes of both sexes widely separated, face vertically oblong, with sparse lateral hairs, occiput with few hairs; antennæ as long as the head or longer, located above the middle, basal joint with short fine setæ, third joint much lengthened, narrow, gradually tapering, ending in a short style which is tipped with a small bristle; proboscis about as long as the head, vertical, palpi porrect, rather long, cylindrical, curving upward, pilose and hairy. Bristles of the thorax weak, no humeral, about three notopleurals, acrostichal setulæ very short, biseriate but the rows distant, dorsocentral setulæ numerous and short, lateral setulæ sparse but evident, scutellum with several variable bristles; pleuræ bare. Abdomen of female often blunt and with the terminal segments retracted, pits of the second segment small; epipygium with compressed ventral keel, all its valves acute. Legs bare of bristles or nearly so, front coxæ rather lengthened, hind femora lengthened and thickened, sometimes thorny beneath, hind tibiæ shortened, sometimes geniculate next to the knee. Costa enclosing the entire wing, anal angle broadly rounded, not prominent, no basal bristle, auxiliary vein evanescent at the end, fork of the third vein short and oblique, anal vein reaching halfway along the narrow anal cell, anal vein rather more distinct than usual, almost attaining the margin, detached from the anal cell, no alula.

Type species : *H. Gayi*, Macquart, the original species.

Geographical distribution. — The distribution of *Haplomera* is subtropical, all of the species known coming from South America.

1. *H. argentata*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 758 [1865] Chile.
(*Pachymeria*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448 (1905);
Nova Acta Akad. Naturf. Halle, Vol. 91, p. 373 (1909).
2. *H. brachygastera*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 759 Chile.
[1865] (*Pachymeria*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448
(1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 374 (1909).
3. *H. chilensis*, Bezzi, ibidem, Vol. 91, p. 374 (1909). Chile.
4. *H. fulvipes*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 760 [1865] Chile.
(*Pachymeria*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448 (1905);
Nova Acta Akad. Naturf. Halle, Vol. 91, p. 374 (1909).
5. *H. Gayi*, Macquart, Dipt. Exot. Vol. 1, p. 279 (*Aplomera*), pl. 13, f. 4 Chile.
[1838] (*Anodontina*); Blanchard, Gay, Hist. Chile (Zool.), Vol. 7,
p. 374 [1852] (*Aplomera*); Philippi, Verh. Zool.-bot. Ges. Wien,
Vol. 15, p. 758 [1865] (*Aplomera*); Bezzi, Ann. Mus. Hungar. Vol. 3,
p. 458 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 373 (1909).
6. *H. gymnopoda*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448 (1905); Nova Peru.
Acta Akad. Naturf. Halle, Vol. 91, p. 374 (1909). — **Pl. 6, Fig. 58.**
7. *H. modesta*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 759 [1865] Chili.
(*Pachymeria*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448 (1905); Nova
Acta Akad. Naturf. Halle, Vol. 91, p. 373 (1909).
8. *H. notogramma*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 449 (1905); Nova Bolivia.
Acta Akad. Naturf. Halle, Vol. 91, p. 374 (1909).
9. *H. nudipes*, Macquart, Dipt. Exot. Vol. 1, p. 277, pl. 13, f. 2 [1838] (*Empis*); Chile.
Blanchard, Gay, Hist. Chile (Zool.), Vol. 7, p. 373 [1852] (*Empis*);
Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 753 [1865] (*Empis*);
Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 373 (1909).
10. *H. obscurata*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 759 [1865] Chile.
(*Pachymeria*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448 (1905); Nova
Acta Akad. Naturf. Halle, Vol. 91, p. 373 (1909).
11. *H. obscuripennis*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 759 Chile.
[1865] (*Pachymeria*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 448
(1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 373 (1909).

12. *H. rubripes*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 760 [1865] Chile.
(Pachymeria); Bezz, Ann. Mus. Hungar. Vol. 3, p. 448 (1905);
Nova Acta Akad. Naturf. Halle, Vol. 91, p. 373 (1909).
13. *H. Schrottkyi*, Bezz, Wien. Ent. Zeit. Vol. 28, p. 319, fig. (1909). Paraguay.
14. *H. Verasi*, Brèthes, Rev. Chilena, Hist. Nat. Vol. 20, p. 79 (1916). Chile.

19. GENUS DEUTERAGONISTA, PHILIPPI

Deuteragonista, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 770 (1865); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 121 (1889).

Amictoides, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 380 (1910).

Characters. — Short robust reddish species, measuring about five millimeters in length, resembling the bombyliid genus *Amictus* in appearance. Eyes of the male contiguous half-way between the antennæ and the ocelli, ocellar triangle large; basal joints of the antennæ nearly bare, the third joint long and conical, about three times as long as broad, the apical style half the length of the third joint, distinctly two-segmented and with short end-bristle, the basal segment about one-fourth as long as the apical; proboscis vertical, as long as the head, palpi curved and with a few fine hairs. Thoracic bristles black, no strong humeral, acrostichals short and arranged in several rows, anterior dorsocentrals short, hair-like and scattered, the middle ones longer and seriate, the posterior three long and bristle-like, notopleural bristles strong, two long and two short scutellars; pleuræ bare. Abdomen short, pygidium small, erect and closed, middle valve large, central piece compressed below. Legs simple, hairs short and fine, tibæ with rather distinct extensor bristles, front metatarsi of the male somewhat thickened. Costa greatly thinned beyond the third vein, basal bristle present, auxiliary vein stopping before the costa, no stigma, fork of third vein moderately short, discal cell not large, as long as first basal cell, anal vein obsolete, anal angle well developed, axillary incision distinct, no alula.

Type species: *D. bicolor*, Philippi. This genus was located near *Thereva* by Philippi. Bezz erected the genus *Amictoides* on *Hilara* (?) *breviventris* Philippi, stating in a footnote that it differed only in the structure of the antennal style. Dr. Bezz has loaned his single specimen of *breviventris*, which has furnished the basis for the preceding description and the accompanying illustration. This specimen has the style exactly as described by Philippi for *Deuteragonista*, and agrees in all points with the rest of Philippi's description of the genus. Philippi separated widely the species *bicolor* and *breviventris* in his paper on Chilean diptera, and it may be that the two forms are entirely distinct. In the meantime, as we have no specimens of *bicolor* for comparison, we have no basis for segregating the two.

Geographical distribution.

1. *D. bicolor*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 771, f. 49 Chile.
 (1865).
2. *D. breviventris*, Philippi, ibidem. Vol. 15, p. 762 [1865] (? *Hilara*); Bezz, Chile.
 Ann. Mus. Hungar. Vol. 3, p. 444 [1905] (*Apalocnemis*); Nova Acta
 Akad. Naturf. Halle, Vol. 91, p. 380 [1909] (*Amictoides*). — Pl. 2,
 Fig. 14.

20. GENUS TOREUS, MELANDER

Toreus, Melander, Ent. News Philad. Vol. 17, p. 376 (1906); Williston's N. Amer. Dipt. Man. p. 226 (1908); Kertész, Cat. Dipt. Vol. 6, p. 80 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 615 (1910).

Characters. — Rather large, brownish, entirely pollinose bristleless species with yellow legs. Eyes broadly separated, bare, the lower anterior facets of the male large; antennæ inserted below the middle of the head, nearly bare, the basal joints subequal, the third joint elongate, cylindrical at the bases and bluntly conical at the apex, terminated by a thickened two-jointed style the outer segment of which is short and bristle-like; proboscis three times as long as the head, rigid, directed somewhat forward, the labrum three-fifths as long as the labella, the two-jointed palpi short, narrow, flattened and incumbent; ocelli large, not raised, ocellar and vertical hairs reduced to microscopic size. Thorax moderately stout, entirely without bristles, no metapleural setæ. Abdomen robust, cylindrical, twice as long as the thorax, nearly bare, no pittings, six segments visible anterior to the epipygium, the seventh sternite forming a flattened peduncle bearing the pygidium, which comprises a pair of upright pointed lateral valves, a pair of flattened dorsal decussating filaments, a pair of basal long slender diverging filaments and a hooked downward-pointed penis. Legs entirely devoid of bristles, simple except that the hind femora and tibiae of the male are bent, pulvilli small. Wings rather broad, anal angle broadly rounded, costa extending around the entire wing, no basal bristle, auxiliary vein distinct except at the tip, straight, the other veins reaching the margin, third vein forked, basal cells long, equal in length, discal cell rather narrow, blunt, sections of the fourth vein proportioned $0.8 : 0.2 : 1 : 1.2$, of the fifth vein $1.1 : 0.3 : 1 : 1$, anal crossvein abruptly reflexed and fused with the underside of the anal cell, anal vein represented by a faint fold, no alula, alular fringe very weak, calypteres with a straight edge and with weak hairs.

Type species : *T. neomexicanus*, by original designation.

Geographical distribution.

1. *T. neomexicanus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 352 [1902] New Mexico.
(*Empis*); Ent. News, Philad. Vol. 17, p. 377, fig. (1906).

21. GENUS TENONTOMYIA, WHITE

Tenontomyia, White, Papers, Proc. Roy. Soc. Tasmania, 1916, p. 236 (1917).

Characters. — Head a little narrower than the thorax; eyes separated in both sexes; proboscis thick, a little shorter than height of the head and bearing long hairs, palpi not distinguishable; antennæ with basal joints extremely small, the third expanded, nearly oval in shape and drawn out at the apex into a narrow point, from which springs a long aristiform style which is about twice as long as the three antennal joints together. Thorax considerably arched, devoid of pubescence, two rows of fairly long dorsocentrals, lateral bristles long, four scutellars. Abdomen with a few short lateral bristles on hind margins; genitalia of male narrow but lengthened. Legs long, particularly the hind pair, and very slender, all joints simple and practically bare. Wings of medium size, auxiliary vein straight and apparently becoming coalescent with the first vein; third vein with perpendicular fork which closes the first submarginal cell before the apex of the marginal cell, discal cell somewhat triangular, with three complete posterior veins, stigma weak.

Genotype : *T. gracilipes*, White, the only known species. This is a brownish insect 5.5 mm. in length that has been taken a number of times in Tasmania. White described the third vein as simple, designating the anterior fork as a crossvein. The conformation suggests *Blepharoprocta* or *Empis clausa*, where the submarginal cell is closed by what is unquestionably the anterior fork of the third vein.

Geographical distribution.

1. *T. gracilipes*, White, Papers, Proc. Roy. Soc. Tasmania, 1916, p. 237, Tasmania.
f. 45 (1917). — PI. 8, Fig. 83.

22. GENUS ATRICHOPLAURA, BEZZI

Atrichoplaura, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 357 (1910).

Characters. — Slender or robust blackish species, two to four millimeters in length. Head globular, the antennæ inserted near the middle of the head, eyes of both sexes widely separated, bare, the facets not enlarged, marginal incision at the antennæ deep, ocellar triangle not elevated, ocellar bristles long and divergent, several diverging fronto-orbitals, face quadrate, sometimes bearing hairs; antennæ rather shorter than the head, basal joints subequal, more or less globular, not setose, the third joint narrowly conical, with a thickened terminal style shorter than the third joint and not geniculate at the short basal section; proboscis about as long as the head, projecting obliquely forward, labrum shorter than the labium, the labellar flaps broad, palpi long, cylindrical, up-turned, with a few long hairs. Thorax small, bristles weak or strong, one humeral and one posthumeral, the anterior dorsocentrals weak and numerous, several scutellars, the acrostichals weak and biserrate; pleuræ entirely bare. Abdomen typically slender, twice as long as the thorax, with weak hairs; pygidium typically rather small, closed, no projecting valves, penis filiform; abdomen of the female tapering. Legs simple, of the female without bristles, of the male the anterior tibiæ sometimes densely pubescent. Wings long and narrow or short and broad, the costa encompassing the entire margin, auxiliary vein straight, distinct, not meeting the costa, third vein simple, anal vein obsolescent, no alula, typically the anal angle broadly rounded and the anal crossvein abruptly and greatly reflexed, parallel with the axis of the wing.

Type species: *A. Schneusei* (Pl. 6, Fig. 60), by Bezzii's designation. The genus *Atrichoplaura* is a composite which ultimately will probably be subdivided. The species *nitida* has the habitus of *Hilara*, *crassa* resembles *Schistostoma*, while *Schneusei* and *hirtipes* are slender insects with long narrow wings and more or less hairy face. The genus appears to be restricted to South America.

Geographical distribution.

1. *A. argyiventris*, Becker, Mission Arc Méridien Amér. Sud, Vol. 10, p. 171 Ecuador.
(1919).
2. *A. crassa*, Bezzii, Nova Acta Acad. Naturf. Halle, Vol. 91, p. 359 (1910). Chile.
3. *A. hirtipes*, Bezzii, ibidem, Vol. 91, p. 361 (1910). Peru.
4. *A. nitida*, Bezzii, ibidem, Vol. 91, p. 358. Peru.
5. *A. Schneusei*, Bezzii, ibidem, Vol. 91, p. 362 (1910). — Pl. 6, Fig. 60. Peru.
6. *A. tephrodes*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 760 [1865] Chile.
(*Rhamphomyia*); Bezzii, Nova Acta Acad. Naturf. Halle, Vol. 91,
p. 323 [1910] (? *Rhamphomyia*).

23. GENUS OPEATOCERATA, NOV. GEN.

Characters. — Small, polished yellow species suggestive of *Leptopeza*, measuring about three millimeters in length. Antennæ inserted below the middle of the head, joints small, slightly pubescent, the third joint short-ovate, tipped with a lengthened two-piece arista three times the length of the third joint, of which the basal part is two-thirds the length of the third antennal joint and appears like a prong-like extension of that joint; proboscis inflexed, slender, slightly longer than the head-height. Thorax strongly convex, bristles very delicate and yellow, no pubescence, dorsocentrals uniserial, no acrostichals, humeral, posthumeral, notopleural or presutural setæ, two supra-alars, two parallel scutellars; propleuræ bare, a row of long delicate metapleural hairs, prothoracic spiracle concolorous,

metathoracic spiracle brown. Abdomen (♀) with nine segments, the eighth lengthened and compressed, styles long and narrow, abdominal hairs very sparse. Legs long, slender, hairy, hind tibiæ and metatarsi with long delicate extensor setæ. Wings more or less cuneiform in outline, costa stopping at apex, first vein ending near basal two-fifths of wing, auxiliary vein straight, evanescent apically, stigma strong, third vein ending at wing-tip, its anterior branch erect, submarginal cells wide, basal cells much shorter than the discal, fourth and intercalary veins evanescent apically, anal crossvein abruptly reflexed, in line with the outward continuation of the cubitus, only the base of the anal vein indicated, axillary incision weak, shallow; calypteres with a few long hairs.

Genotype : *Empis rubida*, Wheeler & Melander. This species was originally described from Mexico. Bezzī has recorded it from Bolivia. Pablo Schild has taken it in Costa Rica. The peculiar antennæ can well remove this form from the complex genus *Empis*. The genus is a likely ancestor to *Lamprempis*.

Geographical distribution.

1. *O. rubida*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, p. 368 Mexico, Bolivia.
 [1901] (*Empis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 302,
 f. 121 [1902] (*Empis*); Bezzī, Nova Acta Akad. Naturf. Halle,
 Vol. 91, p. 343, 348 [1909] (*Empis*); Deut. Ent. Zeitschr. Beiheft,
 p. 89 [1909] (*Empis*).

24. GENUS LAMPREMPIS, WHEELER & MELANDER

Lamprempis, Wheeler & Melander, Biol. Centr. Dipt. Suppl. p. 366 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 278 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252, 263 (1903); Bezzī, Ann. Mus. Hungar. Vol. 3, p. 434 (1905); Melander, Williston, N. Amer. Dipt. Man. p. 225 (1908); Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 340 (1909); Kertesz, Cat. Dipt. Vol. 6, p. 38 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 557 (1910).

Characters. — Brilliant metallic, blue or green or polished black species with elongate antennæ and erect fork of the third vein. Head longer than broad, placed low down on the thorax, eyes of the male contiguous above the antennæ, the upper facets enlarged, eyes bare, marginal excision at the antennæ very shallow and broad; face with parallel sides, usually shining, of the male rather narrow, of the female moderate, no cheeks; antennæ inserted well above the middle of the head; porrect, the first joint elongate cylindrical, more than twice as long as the second, and abundantly hairy; proboscis longer than the head, vertical, slender, the lobes of the labella long and very narrow; ocellar triangle hairy, raised in the male, occipital hairs abundant. Thorax very greatly hunchbacked, especially in front, devoid of strong bristles, but densely pilose, several weak but long notopleurals and scutellars; pleuræ partly pollinose, with a series of many long strong metapleural hairs, pectus with some long hairs. Abdomen less abundantly furnished with hairs; pygidium of moderate size, lateral valves broad at the apex, penis short and thick; abdomen of the female flattened, the terminal segments retractile into the fifth. Legs rather stout, strongly hairy, the hind pair largest, front metatarsi frequently swollen, hind femora, tibiæ and metatarsi of the male sometimes furnished with apophyses or groups of hairs, or they may become pennate with flat scales, posterior legs of the female often strongly compressed and feathered with scale-like hairs. Wings sometimes pictured, usually clear hyaline or with brownish tinge, rather pointed, the costal edge straight, costal hairs microscopic, no basal bristle, costa stopping at the third vein, auxiliary vein straight and distinct, almost attaining the costa, third

vein abruptly forked, fourth vein usually weak and sinuous, the first posterior cell widened before its apex, discal cell small, no longer than the equal basals, anal cell short, the crossvein abruptly reflexed and fusing with the under side of the anal cell, the anal vein faint but almost complete, alula distinct but not margined, finely fringed, base of the second anal vein very strong; calypteres with short fringe.

Type species: *L. chichimeca*, Wheeler & Melander (Pl. 2, Fig. 13), so designated by Coquillett in 1903. *Lamprempis* was first proposed as a subgenus of *Empis*, but the species are so highly specialized and distinctive that the group deserves generic rank. Bezzi (1909) has given a table of the known species, all of which are restricted to tropical and subtropical America.

Geographical distribution.

1. *L. benigna*, Osten-Sacken, Biol. Centr. Amer. Dipt. Vol. 1, p. 215 [1887] Mexico. (*Empis*); Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. p. 367 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 281 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 341, 342 (1909).
2. *L. calopoda*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 436 (1905); Nova Acta Peru. Akad. Naturf. Halle, Vol. 91, p. 341 (1909).
3. *L. chichimeca*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. Mexico. p. 368 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 280, f. 100-102 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 341 (1909).
4. *L. Columbi*, Schiner, Novara Reise, Dipt. p. 205 [1868] (*Empis*); Bezzi, Colombia. Ann. Mus. Hungar. Vol. 3, p. 434, 457 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 340 (1909).
5. *L. cyanea*, Bellardi, Ditt. Messic. Vol. 2, p. 98 [1861] (*Empis*); Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. p. 367 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 279 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 341 (1909).
6. *L. diaphorina*, Osten-Sacken, Biol. Centr. Amer. Dipt. Vol. 1, p. 215 [1887] Mexico. (*Empis*); Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. Vol. 1, p. 367 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 281 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 341 (1909).
7. *L. dolichopodina*, Schiner, Novara Reise, Dipt. p. 205 [1868] (*Empis*); South America. Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 341 (1909).
8. *L. gemmea*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 438 (1905); Nova Acta Bolivia. Akad. Naturf. Halle, Vol. 91, p. 342 (1909).
9. *L. setigera*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 272 (1903); Cuba. Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 341 (1909).
10. *L. suavis*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 33 : Cent. 8, No. 56 [1869] Mexico, Bolivia. (*Empis*); Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. p. 368 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 280 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 342 (1909).
11. *L. superba*, Loew, Wien. Ent. Monatschr. Vol. 5, p. 36 [1861] (*Empis*); Cuba. Berl. Ent. Zeitschr. Vol. 13, p. 34 : Cent. 8, No. 57 [1869] (*Empis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 279 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, p. 91, p. 340 (1909).

12. *L. tuberifera*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 435, f. 1 (1905); Nova Peru. Acta Akad. Naturf. Halle, Vol. 91, p. 340 (1909).
13. *L. violacea*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 32 : Cent. 8, No. 55 Mexico. [1869] (*Empis*); Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. p. 367 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 278, pl. 8, f. 103, 104 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 340 (1909).
14. *L. viridis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 395 [1895] Jamaica. (*Hilara*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 269 [1902] (*Hilara*); Cockerell, Ent. News Philad. Vol. 36, p. 49 (1903); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 434, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 341 (1909).

25. GENUS PORPHYROCHROA, NOV. GEN.

Characters. — Metallic steel-blue slender species, with long yellow legs. Eyes large, bare, facets uniform, the emargination at antennæ short and narrow; front very narrow, sides parallel, vertex filled by the ocellar triangle, face almost obliterated, only a linear indication at the oral edge; antennæ located well up on the head, the front being half as long as the face, long and slender, basal joints shining, the first joint cylindrical with apical setulæ, the second joint pyriform and loosely setulose, the third joint velvety, long conical, narrow, only a little tapering, indistinctly ending in a long thick cylindrical style which is nearly as long as the remainder of the antenna, its outer third a more slender segment; proboscis slender, pointed, rigid, thick only at extreme base where the labella is developed into lateral flaps; palpi minute, visible only as a black point; two erect parallel proclinate vertical bristles, one pair of proclinate ocellar, two frontal bristles arranged one before the other. Thorax not pubescent, five reclinate dorsocentral bristles then an interruption to the proclinate prescutellar pair, no acrostichals, one humeral, two or three supraalar setulæ, four scutellar bristles, metapleuræ with a few setulæ. Abdomen depressed, sparsely hairy; pygidium terminal, penis thick but visible at base only, middle valves large. Legs simple, slender, hairy, middle tibiæ with two apical bristles, hind tibiæ with terminal pecten. Wings with strong anal angle, no costal bristle, costa abruptly interrupted beyond third vein, a crease in the membrane beneath the humeral crossvein, auxiliary vein almost contiguous with first vein, evanescent at middle of costal cell, anal crossvein recurved, forming an angle of 140 degrees with anal vein which passes into a fold beyond anal cell, marginal cilia of two lengths, some shorter and some longer than the anterior crossvein which is located well toward the base of the complete discal cell, three complete veins issuing from the discal cell of which the anterior two are weak; calypteres with long cilia.

Type species: *P. palliata*, Coquillett. This was originally described as belonging to *Sciodromia*. Later Coquillett referred the species to *Microphorus*. The genus is much more closely related to *Rhamphomyia* than to either of these, but differs in the high-placed antennæ with lengthened first joint and arista. Other tropical species of *Rhamphomyia* resemble *palliata* in possessing metallic coloration. The description and the figure of the genus were obtained from the type specimen in the U. S. National Museum.

Geographical distribution.

1. *P. palliata*, Coquillett, Jour. New York Ent. Soc. Vol. 10, p. 140 [1902] Mexico. (*Sciodromia*); Proc. Ent. Soc. Wash. Vol. 5, p. 264 [1903] (*Microphorus*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 460 [1905] (*Microphorus*). — Pl. 2, Fig. 12.

26. GENUS EMPIS, LINNÆUS

Empis, Linnæus, Syst. Nat. Ed. 10, p. 603 (1758); Fauna Suec. p. 466 (1763); Latreille, Consid. Gen. p. 443 (1810); Meigen, Syst. Beschr. Vol. 3, p. 15 (1822); Macquart, Mém. Soc. Sc. Lille, 1823, p. 156 (1823); Curtis, Brit. Ent. Vol. 8, p. 18 (1824); Macquart, Dipt. N. France, Vol. 3, p. 115 (1827); Hist. Nat. Dipt. Vol. 1, p. 327 (1834); Zetterstedt, Fauna Ins. Lappon. p. 560 (1838); Westwood, Gen. Syn. p. 131 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 367 (1842); Boitard, Man. Ent. Vol. 3, p. 317 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 89 (1852); Rondani, Dipt. Ital. Vol. 1, p. 151 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 561 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 102 (1862); Liroy, Atti Inst. Ven. Sc. Venezia, 1864, p. 599 (1864); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 1-62, 157-175 (1867), Vol. 12, p. 231-240, 387-393 (1868); Beling, Arch. Naturg. Berlin, Vol. 48, p. 240 (1882); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 120 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389, 397 (1895); Williston, Man. N. Amer. Dipt. p. 74 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 282 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 249, 263 (1903); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 438 (1905); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 209-216, 297-304 (1906), Vol. 7, p. 25-32, 155-160 (1907); Melander, Williston's N. Amer. Dipt. Man. p. 226 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 342 (1909); Kertész. Cat. Dipt. Vol. 6, p. 40 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 537 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 79, 128 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 60 (1910); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 467 (1912); White, Proc. Roy. Soc. Tasmania, 1916, p. 231 (1917); Brunetti, Fauna Brit. Ind. Dip. Vol. 1, p. 347 (1920).

Anacrostichus, Bezzi, Deutsche Ent. Zeitschr. 1909, Beiheft, p. 93 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 96 (1910).

Argyrandrus, Bezzi, Deutsche Ent. Zeitschr. 1909, Beiheft, p. 100 (1909).

Coptophlebia, Bezzi, ibidem, p. 100 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 102 (1910).

Dionaea, Meigen, Nouv. Classif. p. 24 (1800); Hendel, Verh. Zool-bot. Ges. Wien, Vol. 58, p. 54 (1908).

Enoplempis, Bigot, Ann. Soc. Ent. France (5), Vol. 10, Bull. p. 47 (1880); Williston, Man. N. Amer. Dipt. p. 81 (1888); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 119 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 249 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 538 (1910).

Eriogaster, Macquart, not Germar, 1811 (*Lepidoptera*); Dipt. Exot. Vol. 1, Pt. 2, p. 162 or 278 (1838); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 561 (1857); (6), Vol. 9, p. 120 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 249 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 539 (1910).

Haplomerinx, Bezzi, Deutsche Ent. Zeitschr. 1909, Beiheft, p. 86 (1909).

Lissempis, Bezzi, ibidem, p. 99 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 101 (1910).

Niconia, Walker, Ins. Saunders. Dipt. Vol. 1, pl. 6, f. 1 (1852); Trans. Ent. Soc. Lond. Vol. 4, p. 147 (1857).

Pachymeria, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 262 (1829); Curtis, Brit. Ent. Vol. 8, p. 18, 4; Meigen, Syst. Beschr. Vol. 7, p. 88 (1838); Westwood, Gen. Syn. p. 131 (1840); Rondani, Dipt. Ital. Vol. 1, 151 (1856); Schiner, Fauna Dipt. Austr. Vol. 1, p. 110 (1862); Loew, Wien. Ent. Monatschr. Vol. 8, p. 353-366 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 119 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389 (1895); Williston, Man. N. Amer. Dipt. p. 74 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 330 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 254, 263 (1903); Melander, Williston N. Amer.

Dipt. Man. p. 226 (1908); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 582 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 98 (1910).

Pachymerina, Macquart, Hist. Nat. Dipt. Vol. 1, p. 333 (1834), Vol. 2, p. 657 (1835); Boitard, Man. Ent. Vol. 3, p. 319 (1843); Bigot, Ann. Soc. Ent. France (3). Vol. 5, p. 561 (1857); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 254 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 582 (1910).

Platyptera, Meigen, Illiger Mag. Ins. Vol. 2, p. 269 (1803); Curtis, Brit. Ent. Vol. 8, p. 18, 2 (1824).

Platypterygia, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 263 (1829); Westwood, Gen. Syn. p. 131 (1840); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 256 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 592 (1910).

Polyblepharis, Bezzi, Deutsche Ent. Zeitschr. 1909, Beiheft, p. 95 (1909).

Pterempis, Bezzi, ibidem, p. 87 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 108 (1910).

Xanthempis, Bezzi, Deutsche Ent. Zeitschr. 1909, Beiheft, p. 88 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 85 (1910).

Characters. — A dominant and variable genus whose species range from three to ten millimeters in length. More or less slender, sometimes abundantly hairy, sometimes nearly bare, usually black though often cinereous or yellow, pollinose or shining. Head narrower than the thorax, globular, or in *Xanthempis* pear-shaped, due to a conical development of the occiput; eyes contiguous on the front, or subcontiguous or fully separated, when contiguous the upper facets usually larger than those below, when separated the facets uniform, eyes of female always separated and with uniform facets; face usually broad, the lower edge somewhat ridged at the epistome, in *Lissempis* the face is narrow; antennæ generally longer than the head, inserted at or a little above the middle, distinctly three-jointed, the first joint usually longer than the second and both more or less cylindrical, the third joint lengthened, compressed conical, with a terminal two-jointed style whose apical part is longer than the basal segment and tapers; proboscis at least as long as the head, usually much longer, strong, vertical or slightly inflexed, in *Acallomyia* slender and obliquely projecting, labium of variable length, palpi cylindrical, one-jointed, short, extending rather upward; occipital hairs biseriate or abundant. Thorax rather large, not greatly convex, generally vittate to mark the acrostichal and dorsocentral spaces, prothorax of *Xanthempis* lobosely broadened, chaetotaxy very variable, sometimes the thorax is abundantly furnished with fine hairs, sometimes the dorsocentrals and acrostichals form definite single rows, sometimes the acrostichals are entirely absent, humeral, posthumeral, notopleural and supraalar bristles variable but usually present; metapleuræ always with some hairs or bristles, pectus with or without hairs. Abdomen rather slender and long, tapering in the female but in the male generally with quite prominent pygidium, which comprises two pairs of lateral valves a ventral piece and a long and slender or short and thick penis; in some species the last ventral segment of the male is pronged. Legs of the female often ornate with scales, sometimes the males have apophyses about the hind knees or on the hind trochanters, otherwise the legs are more or less hairy and bristly. Wings generally narrow but in some females (e. g. *E. borealis*) unusually broad, costa stopping at the end of the wing, basal bristle present or absent, auxiliary vein straight and incomplete, not reaching the costa, third vein forked, discal cell usually rather small, emitting three veins, of which the first may be shortened (*Coptophlebia*), anal cell short and narrow, the anal crossvein abruptly reflexed, the anal vein separate from the anal cell, anal angle of the wing usually more or less rectangular, sometimes obtuse, sometimes strong, auxillary incision distinct, alula small or absent, fringe of the calypteres variable.

Type species : *E. pennipes*, Linnæus (Pl. 5, Fig. 48, 49), so designated by Latreille, 1810. Curtis, Westwood and Rondani selected *E. tessellata*, Fabricius, 1794, as the type, and this course unfortunately has been usually followed by European dipterists. Bezzi has erected the subgenus

Pterempis for the species about *pennipes*, but for this restricted group the name *Empis* should be restored. Bezzi's subgenus *Empis* should be called *Platyptera*, Meigen, since the species *borealis*, the type of *Platyptera*, occurs there.

Empis is the dominant genus of the family, particularly in the Old World with nearly five hundred species described it has become very complex. Loew and Kuntze, in a series of papers, have subdivided the genus into groups of species centering about a few typical forms. Bezzi, 1909, effected a different separation based mainly on chætotaxy, and designated his groups as subgenera. These subgenera are differentiated in the previous table of genera. Their limits are far from definite but in the main they appear to be natural groupings. Holopticism appears to have developed in several lines of *Empis* and does not seem to be as ancient a character as Bezzi's definitions would indicate. That is to say, the presence of holopticism in *Empis* does not of itself bespeak relationship.

The mating habits of several species of *Empis* have been studied, particularly by Aldrich and by Hamm. Quite often the males have the habit of presenting a small insect to the female who sucks its juices during copulation. In the American species *aerobatica* the male elaborates a large frothy balloon around his prey and flies with this to induce the female to mate. Species of the subgenus *Xanthempis* do not mate in the air. The species of *Empis*, s. str. swarm in shady woods. It is particularly in this group that the legs of the females are pennate with scale-like hairs.

The earliest stages of several species of *Empis* are known. Beling has described the larvæ of *astiva*, *decora*, *nodosa*, *tessellata*, and *trigramma* and Kieffer of *meridionalis*. The larvæ live in rich moist earth, in which they pupate following hibernation. They are cylindrical, tapering anteriorly and consist of twelve segments. Ventrally they show transverse swellings. The pupæ possess bands of abdominal spines.

Empis minuta, Fabr. Described in 1787, belongs to *Phora*, as discussed by Lundbeck, Dipt. Danica, Vol. 3, p. 6, 189 (1910).

SUBGENUS ACALLOMYIA, NOV. SUBGEN.

Characters. — Slender, blackish species with long simple legs. Occipital hairs long and straggling; eyes of male contiguous along the front, the upper facets large; face narrowed; basal antennal joints short, third joint long and tapering; proboscis very slender, projecting obliquely forward, palpi without setæ. Prothorax small, mesonotum gibbose, bare of pubescence, acrostichals sparse but biserrate, dorsocentrals long, sparse, biserrate, lateral bristles long and hair-like, one humeral, one posthumeral, one presutural, several small notopleural, four scutellar, no propleural, metapleural setæ bunched. Abdominal hairs sparse, those before the incisures long and delicate, pygidium small. Wings normal, third vein curving to end at apex, discal cell not pointed, anal vein incomplete.

Type species: *Iteaphila peregrina*, Melander. *Empis brunnea*, Coquillett is a closely related but distinct species.

SUBGENUS ANACROSTICHUS, BEZZI

Characters. — Black, shining or more or less pollinose species of moderate size, with rather fine bristles and few hairs. Sex differences slight. Eyes separated, facets uniform, front of male sometimes narrower than of female, either parallel-sided or centrally contracted; head globose occipital setulæ not clearly biserrate; antennæ relatively short. Prothorax not bilobate; notal hairs undeveloped except a few laterally; no acrostichals, dorsocentrals almost uniserrate, the three hind ones stronger, one humeral bristle and many humeral hairs, one posthumeral, one or two notopleural, two or three presutural, one or two supra-alar; propleural hairs evident, metapleural long, fine and many, four or six scutellars, the inner pair cruciate or convergent. Abdomen bristleless, in the male with rather long

fine pubescence, last sternites of male sometimes deformed, pygidium robust, short, penis thick and hidden, ovipositor suddenly narrowed, styles long and slender. Legs rather stout, hind femora strong, flexor femoral setulae present. Wings with complete neuration, no basal seta, third vein ending a little before wing-tip, discal cell rather pointed, anal vein reaching margin, axillar incision rectangular or acute.

Type species : *Empis nitida*, Meigen, by Bezzi's designation.

The following species are to be located in this subgenus : *E. bistortæ*, Meig., *humilis*, Coq., *monticola*, Lw., *scoparia*, Coq., *tersa*, Coq., *tumida*, Meig. and *varipes*, Lw.

SUBGENUS ARGYRANDRUS, BEZZI

Characters. — Rather small, yellow, bare species with few bristles, the males with silvery pruinosity. Head round, eyes of male broadly contiguous, the upper facets large, of female narrowly separated, front of female sparsely hairy; first antennal joint not lengthened, third rather elongate; proboscis a little longer than the head. Prothorax very small, notum bare except for the bristles, no acrostichals, seven or eight strong uniseriate dorsocentrals, one humeral, one posthumeral, no notopleural, two presutural, one supra-alar; no propleural and a few metapleural black hairs, four scutellars, the middle ones crossed. Abdomen of male with fine bristles before the incisures, lacking in the female; pygidium small, closed, styles of female long. Legs long, slender, rather bare and with few setæ, simple in both sexes. Wings with complete neuration, rather broad, similar in both sexes, axillar incision rectangular but not deep, discal cell large, rather blunt, third vein straight, ending at apex, cilia of calypteres long and pale.

Type species : *Empis dispar*, Scholtz, by Bezzi's designation. *E. lamellicornis*, Becker belongs here. The American species *captus*, Coquillett, and *vaginifer*, Melander may be assigned here, but they are different from the European species.

SUBGENUS COPTOPHLEBIA, BEZZI

Characters. — Small, black or grayish, pubescent and setose species, sometimes differing greatly in the sexes. Head globose, occiput little setose; proboscis long, palpi small; eyes of male broadly contiguous, the upper facets larger than the lower, of female widely separated; antennæ rather short, basal joints about equal. Prothorax small; notum bare of hairs except at sides, acrostichals biserrate, rarely absent, central dersocentrals biserrate, the posterior three or four stronger and uniseriate; one longer and several shorter humerals, one each of posthumeral, notopleural, supra-alar and presutural bristles, two cruciate scutellars; propleural and metapleural setæ numerous. Pygidium small. Hind legs lengthened, usually feathered in female, front and hind metatarsi of male more or less thickened. Wings with acute axillar incision and well developed anal lobe, discal cell blunt, fourth and usually anal vein shortened, second submarginal cell large; calypteres long-ciliate.

Type species: *Empis hyalipennis*, Fallen, by Bezzi's designation. The species are more abundant in America and the Orient than in Europe, the following being referable to this subgenus: *E. abbreviata* Lw., *albinervis*, Meig., *albipennis*, Meig., *asema*, Mel., *ceylonica*, Bez., *clausa*, Coq., *confluens*, Beck., *coracina*, Bez., *corycrica*, Bez., *distans*, Lw., *dolorosa*, Wh.-Mel., *florisomma*, Lw., *hirticrus*, n. sp. (*hirtipes*, Coq.), *Hoffmannseggii*, Lw., *hystrichopyga*, Bez., *inclinata*, Bez., *Jacobsoni*, Meij., *labiata*, Lw., *leptomorion*, Bez., *melaena*, Bez., *papuana*, Bez., *patagiata*, Bez., *Pavesii*, Bez., *pilimana*, Lw., *plorans*, Bez., *sauteriana*, Bez., *squamipes*, Coq., *spiloptera*, Wied., *tenuinervis*, Bez., *totipennis*, Bell., *velutina*, Bez. and *volucris*, Meig.

SUBGENUS EMPIS, LINNÆUS, S. STR.

Characters. — Moderately small blackish or cinereous species, showing considerable differences in the sexes. Eyes of male broadly contiguous with the upper facets enlarged. Antennæ shortened, with basal joints about equal. Acrostichals biserrate, dorsocentrals biserrate, rarely pluriseriate, one posthumeral, several lateral bristles and usually lateral pubescence, two or four scutellars; metapleural hairs numerous. Abdomen bristleless. Legs with bristles, male metatarsi more or less thickened, middle tibiæ with fine short erect pubescence along front side; legs of female more or less compressed and variously pinnate. Wings with anal lobe well developed, axillary incision acute or rectangular, neuration complete, the anal vein more or less shortened, discal cell rather blunt apically.

Type species : *Empis pennipes*, Linnæus (Pl. 5, Fig. 48, 49), as discussed in a preceding paragraph. *Pterempis*, Bezzi is a synonym. The subgeneric position of *Niconia*, Walker, founded on the Brazilian *Empis pentaphora*, Wiedemann is not known.

This subgenus includes species that hover about herbage, in grassy fields and in shady woods. The females have characteristic feathered legs, such as occur also in *Coptophlebia*. The species are confined to Europe, and among others include: *E. aestiva*, Lw., *alpicola*, Str., *brunnipennis*, Meig., *caudatula*, Lw., *chioptera*, Meig., *ciliata*, Fab., *ciliatopennata*, Str., *cincinnatula*, Lw., *cinerea*, Zett., *dasyprocta*, Lw., *ghigiana*, Bez., *gymnopoda*, Bez., *hyalogyne*, Bez., *lepidopus*, Meig., *malleola*, Beck., *melanotricha*, Lw., *nigricoma*, Lw., *nigrithibialis*, Str., *pennaria*, Fall., *pilosa*, Lw., *plumipes*, Zett., *procera*, Lw., *prodromus*, Lw., *pseudomalleola*, Str., *pusio*, Egg., *rufiventris*, Meig., *saura*, Lw., *scopulifera*, Bez., *serotina*, Lw., *setosa*, Lw., *subpennata*, Macq., *tansphyra*, Lw. and *vernalis*, Meig.

SUBGENUS ENOPLEMPIS, BIGOT

Characters. — Species of moderate size, usually black and heavily grayish pollinose, greatly differing in leg structure in the sexes. Head globose, occiput with two rows of setæ; proboscis one and one-half to two times the head-height, palpi sparsely hairy; eyes separated, facets uniformly small; face with parallel sides; antennæ long, the second joint about half the length of the first, style about one-third the last joint. Prothorax not lobose; notum bare of hairs; acrostichals and dorsocentrals small and uniserrate, scutellars not cruciate, one humeral, one or two presutural; no propleural, a row of metapleural setulæ. Abdomen scarcely hairy, pits prominent, pygidium small, closed. Legs robust, males usually with apophyses about the hind knees or on the hind trochanters, bristles reduced, tarsi short-spinose beneath. Wings narrow, axillar incision rectangular, anal lobe small, discal cell not blunt.

Type species : *Enoplempis mira*, Bigot, the original unique species. This group is well represented in North America, the following species belonging here: *E. aripes*, Mel., *aerobatica*, Mel., *arthritica*, Mel., *Bigoti*, Mel., *cacuminifer*, Mel., *canaster*, Mel., *clauda*, Coq., *dolabraria*, Mel., *enodis*, Mel., *fulcata*, Mel., *gladiator*, Mel., *longipes*, Lw., *loripedis*, Coq., *manca*, Coq., *metapleuralis*, Bez., *mira*, Big., *mixopolia*, Mel., *nodipes*, Mel., *nuda*, Lw., *podagra*, Mel., *poplitea*, Lw., *stenoptera*, Lw., *teres*, Mel. and *valentis*, Coq.

SUBGENUS HAPLOMERINX, BEZZI

Characters. — Body more or less yellowish, without white pruinosity in the male. Eyes of male contiguous, the upper facets distinctly larger than the lower. No acrostichals, dorsocentrals uniserrate, one or two presuturals, scutellar bristles not cruciate. Axillar incision of wings obtuse, discal cell pointed.

This brief diagnosis is combined from the characters given in the identification key in Dr. Bezzi's paper. No type was designated and no species have been assigned to this subgenus.

SUBGENUS LISSEMPIS, BEZZI

Characters. — Small black shining nearly bare species with little difference between the sexes. Head round, occiput with few bristles; eyes with small uniform facets, narrowly separated on front and face, the face narrower than in the other groups; first antennal joint rather long, third narrow; proboscis little longer than the head, palpi small, black, nearly bare. Prothorax small, not lobate; bristles reduced, yellow, dorsocentrals and acrostichals fine and usually biseriate, no humeral, notopleural or posthumeral, two or four scutellars, the middle ones not crossed; propleural and metapleural hairs numerous. Abdomen bristleless, with rather long hairs, pygidium small, open, ovipositor long, styles long and narrow. Legs hairy and setose, especially in male, hind legs lengthened, with thickened metatarsi, legs of female simple. Wings more or less narrowed, alike in both sexes, second submarginal cell rather large, third vein ending at apex, discal cell rather blunt, anal vein small or wanting, axillar incision wide and shallow.

Type species : *Empis nigritarsis*, Meigen, by Bezzi's designation. Additional species referable to this subgenus are: *E. crassipes*, Meig., *cuneipennis*, Bez., *liosoma*, Bez. and *nitidissima*, Str.

SUBGENUS PACHYMERIA, STEPHENS

Characters. — Generally black, robust species of moderate size, sometimes with yellowish abdomen; sexes slightly different in color and leg structure. Dichoptic, facets small and uniform, front of male with the sides parallel or centrally convergent; head globose, occiput hairy and setose; proboscis long, palpi variable, short or long, setose, yellow or black; antennæ rather short, third joint usually strongly narrowed apically. Thorax bristly and pubescent, acrostichals in two or four rows, dorsocentrals bi- or pluriseriate, sometimes not distinct from the hairs, humerals, posthumerals and notopleurals sometimes not distinct, sometimes numerous, two or more presuturals, one or more supralars, four to eight scutellars; propleural and metapleural hairs abundant. Abdomen without bristles, pygidium compressed, penis sometimes long and free, ovipositor short. Legs strong, short, hind femora more or less thickened, of female sometimes fringed. Wings of female sometimes a little broader than of male, neuration complete, third vein straight, ending slightly before apex, second submarginal cell rather small, discal cell pointed, axillar incision sharp and usually deep.

Type species : *Empis femorata*, Fabricius (Pl. 5, Fig. 47), the single original species. *Pachymerina*, Macquart is synonymous, its genotype being also *E. femorata*, Fab., so designated by Coquillett in 1903. The following species belong to this group: *E. brevis*, Lw., *contigua*, Lw., *Erberi*, Lw., *grisea*, Fall., *Johnsoni*, Mel., *mediterranea*, Lw., *obscuripes*, Lw., *opaca*, Meig., *otiosa*, Coq., *palparis*, Egg., *picena*, Bez., *ptilocnemis*, Lw., *pudica*, Lw., *ruficornis*, Lw., *specularis*, Bez., *subclavata*, Lw., and *trianguligera*, Str.

SUBGENUS PLATYPTERA, MEIGEN

Characters. — Middle-sized to large, generally blackish or brownish, pollinose species. Head globose, occiput hairy, eyes of male contiguous, upper facets somewhat larger than the lower ones; first antennal joint usually twice as long as the second and hairy; palpi hairy. Prothorax small, mesonotum hairy, dorsocentrals pluriseriate, rarely uniseriate, acrostichals bi- or pluriseriate, usually several humeral, posthumeral, notopleural and supra-alar setæ to be distinguished among the lateral hairs; pectus hairy, metapleural setæ bunched. Abdomen more or less hairy and long-setose on the margins;

pygidium large or rather small. Legs setose, metatarsi not swollen, female legs rarely pannate. Wings of female of *E. borealis* remarkably broad, but in the other species normal, neuration usually complete, discal cell more or less pointed, anal lobe full, axillary incision deep and sharp.

This genus originally included two species, *Empis borealis*, Linn., and *platyptera*, Panz. Curtis in 1824 designated *borealis* as type, unmindful of the rule in present-day nomenclatorial procedure that gives preference to a tautonymic species. Accepting Curtis' selection safeguards much confusion for otherwise the host of species described as *Rhamphomyia* would require a change of name to the earlier described *Platyptera*. Hendel would recognize in *Dionaea* the genus *Platyptera*, and acceptance of this conclusion would still further imperil the nomenclatorial stability of the dominant genus *Rhamphomyia*. *Eriogaster*, Macquart founded on *Empis laniventris*, Eschscholz is subgenerically equivalent to the species grouped about *E. borealis*. *Platypterygia*, Stephens was erected for *E. borealis*. This is the group for which most European dipterists reserve the name *Empis*, s. str., based on *E. tessellata*.

The following species belong to the subgenus *Platyptera*: *E. alampra*, Lw., *Aldrichii*, Mel., *borealis*, Linn., *brachysoma*, Coq., *brevicornis*, Lw., *cæligena*, Mel., *calcarata*, Bez., *cognata*, Egg., *cothurnata*, Brullé, *crassa*, Now., *dedecor*, Lw., *discolor*, Lw., *divergens*, Lw., *erosa*, Lw., *fallax*, Egg., *fraterna*, Lw., *fumida*, Coq., *infumata*, Coq., *laevigata*, Lw., *laniventris*, Eschsch., *livida*, Linn., *lucida*, Zett., *maculipes*, Zett., *meridionalis*, Meig., *morio*, Fab., *nigricans*, Meig., *nitidiventris*, Lw., *obesa*, Lw., *opaca*, Meig., *plebeja*, Lw., *polita*, Macq., *ravida*, Coq., *scatophagina*, Mel., *serena*, Pok., *strigata*, Lw., *tessellata*, Fab., *variegata*, Meig., and *virgata*, Coq.

SUBGENUS POLYBLEPHARIS, BEZZI

Characters. — Rather large, cinereous, heavily bristly species, with sex differences in color and size. Both sexes dichoptic, facets small and uniform; sides of front parallel; head globular, with strong occipital bristles; proboscis long, palpi large, with many long bristles; first antennal joint short, third not long, excavated below the tip. Thorax strongly bristly and sometimes also hairy; acrostichals dense, more or less definitely biseriate, central dorsocentrals in two or more rows, the posterior three to seven strong and uniserial, several each of humeral, posthumeral, notopleural, presutural and supra-alar bristles, four to eight scutellars; propleural and metapleural bristles very strong. Abdomen hairy and with very strong lateral bristles before the incisures, pygidium small and closed, female abdomen blunt, ovipositor short. Legs stout, strongly bristly and hairy, the hind coxae and hind femora greatly thickened, especially in the male, hind femora more or less curved. Wings alike in both sexes, unusually narrow and pointed, neuration complete and light colored, no basal seta, third vein anteriorly located and ending much before apex, second submarginal cell very small, discal cell small, rather pointed, axillary incision acute and rather deep.

Type species: *Empis albicans*, Meigen, by Bezz's designation. The other recorded species are *curvipes*, Lw., *eumera*, Lw., and *phænomeris*, Lw., all restricted to Southeastern Europe and Asia Minor. The American species *frontalis*, Coq., and *spectabilis*, Lw., may be assigned here, although probably they are phylogenetically distinct.

SUBGENUS PYRREMPIS, NOV. SUBGEN.

Characters. — Full-sized, yellow, setose species. Head globular, occipital setæ biseriate and strong; eyes of male narrowly separated above the middle of the front, facets uniform; antennæ setulose at base, the second joint much shorter than the first; proboscis twice the head-height, palpi with few hairs. Prothorax not lobate, but margined with stiff setæ; dorsocentrals uniserial, acrostichals wanting or the anterior ones weakly indicated, humeral, posthumeral, presutural, two notopleural, and one or two supra-alar bristles present; propleural setæ present, metapleural setæ bunched. Abdomen shining,

with long thin setæ before the incisures, pygidium large or small. Legs setose, normal. Wings normal, anal angle full, axillar incision deep and acute, costa with basal spine.

Type species: *Empis rufescens*, Loew. *E. tridentata*, Coq. also belongs here. Both species inhabit North America.

SUBGENUS XANTHEMPIS, BEZZI

Characters. — Yellowish, bare, middle-sized species not showing sexual dimorphism. Eyes of both sexes separated, facets uniform, front of male parallel-sided; head piriform, the occiput projecting, its bristles in two regular series; first antennal joint usually lengthened, third narrow and long. Prothorax neck-like, bilobed and setigerous above; notal bristles small, no pubescence, dorso-centrals uniseriate, no acrostichals, typically one humeral, no notopleural, one posthumeral, one supra-alar and two or four scutellar bristles, the inner ones cruciate; metapleural setulae fine and moderately few. Abdomen sparingly hairy and not bristly, pygidium large, with large free penis, styles of ovipositor rather long. Legs not stout, simple, slightly hairy. Wings with strong veins, third vein curving back and ending at wing-tip, discal cell blunt, anal angle obtuse.

Xanthempis is regarded as ancestral to the other divisions of *Empis*, as is shown by the yellow, bare body with reduced bristles, the dichoptic males and lack of sexual dimorphism. The species inhabit shady places in woods and thickets during early summer and do not hover in the aerial dance characteristic of other species. Nearly all the known species are European, although two of the following list are American.

Type species: *Empis stercorea*, Linnaeus, by Bezzi's designation. Additional species are the following: *E. æmula*, Lw., *æqualis*, Lw., *albifrons*, Bez., *caucasica*, Bez., *concolor*, Verr., *digramma*, Meig., *feruginea*, Meig., *Kuntzei*, Beck., *læta*, Lw., *loewiana*, Bez., *lutea*, Meig., *pallida*, Lw., *parvula*, Egg., *pitto-procta*, Lw., *pæcilioptera*, Lw., *punctata*, Meig., *scutellata*, Curt., *seminigra*, Lw., *styriaca*, Strobl, *testacea*, Fabr., *testiculata*, Bez., *trigramma*, Meig., and *univittata*, Lw.

Geographical distribution.

1. *E. abbreviata*, Loew, Berlin. Ent. Zeitschr. Vol. 13, p. 84 (1869); Strobl, C. Europe.
Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 69 (1892);
Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 303 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 102 [1909] (*Coptophlebia*).
2. *E. abbrevinervis*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 328 (1911). Java.
3. *E. abcirus*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 494 (1849); Melander, Georgia.
Trans. Amer. Ent. Soc. Vol. 28, p. 299 (1902).
4. *E. abrupta*, Thomson, Eugen. Resa, Dipt. p. 473 (1870). Cape of Good Hope.
5. *E. adusta*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 266 (1869); Kuntze, Corfu.
Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906).
6. *E. æmula*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 228 (1873); Kuntze, C. Europe.
Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi, Deutsche Ent.
Zeitschr. Beiheft, p. 90 [1909] (*Xanthempis*).
7. *E. æqualis*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 13, 20 (1867); Strobl, C. Europe.
Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 60 (1892);
Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi,
Deutsche Ent. Zeitschr. Beiheft, p. 89 [1909] (*Xanthempis*).
? *pallens*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1,
p. 53 (1840); Bezzi, Deutsch. Ent. Zeitschr. Beiheft, p. 89 (1909).
8. *E. æripes*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 328 (1902). — N. W. United States.
Pl. 9, Fig. 141.

9. *E. aerobatica*, Aldrich & Turley, Amer. Natur. Vol. 33, p. 802 [1899] (*Empis*, sp. habits); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 323 (1902); Slosson, Ent. News Philad. Vol. 14, p. 268 (1903); McAtee, Ent. News Philad. Vol. 20, p. 359 (1909). W. United States.
10. *E. aestiva*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 39, 54 (1867); Beling, Arch. Naturg. Berlin, Vol. 48, p. 211 (1882); Strobl, Verh. Siebenb. Ver. Nat. Hermannstadt, Vol. 46, p. 22 (1897); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 302 (1906); Lundbeck, Dipt. Danica, Vol. 3, p. 125, f. 41 [1910] (*Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 64 (1910).
- vulgaris*, Zetterstedt (not Meigen), Dipt. Scand. Vol. 1, p. 387, 21 (1842); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 53 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 109 (1862); Siebke, Enum. Ins. Norv. Vol. 4, p. 33, 14 (1877); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 399 (1866); Lundbeck, Dipt. Danica, Vol. 3, p. 121, 127 (1910).
11. ? *E. aestiva*, Scopoli, Ent. Carn. p. 365 [1763] (*Asilus*); Loew, Linn. Ent. Berlin, Vol. 4, p. 87, note (1849); Schiner, Verh. Zool.-bot. Ges. Wien, Vol. 6, p. 420 (1856) ? gen. C. Europe.
- . *E. affinis*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 263 (1829), no description.
12. *E. agasthus*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 496 (1849); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895). Canada.
13. *E. alampra*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 37 (1872); Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 137 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 27 (1907). C. & S. Europe.
14. *E. albicans*, Meigen, Syst. Beschr. Vol. 3, p. 20 (1822); Schiner, Fauna Dipt. Austr. Vol. 1, p. 105 (1862); Loew, Berl. Ent. Zeitschr. Vol. 12, p. 170, 387, 393 (1868); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 32 (1907); Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 96 [1909] (*Polyblepharis*). C. & S. Europe.
15. *E. albicincta*, Loew, Oefv. Vet. Akad. Förh. Stockholm, Vol. 15, p. 340 (1858); Dipterent. Südafr. p. 264, pl. 2, f. 47 (1860). Cape of Good Hope.
16. *E. albidiseta*, Becker, Zeitschr. Hym. Dipt. 1907, p. 229 (1907). Tunis.
17. *E. albifrons*, Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 90, 91 (1909). C. Europe.
- nana*, Loew (not Macquart), Berl. Ent. Zeitschr. Vol. 11, p. 18, 21, 10 (1867); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 60 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906).
18. *E. albinervis*, Meigen, Syst. Beschr. Vol. 3, p. 26 (1822); Walker, Ins. Brit. Vol. 1, p. 94 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 106 (1862); Strobl, Progr. Seitenstetten, Vol. 14, p. 9 (1880); Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 74 (1892); Glasnik Zem. Mus. Bosn. Herceg. Sarajevo, Vol. 14, p. 470 (1902); Mitt. Bosn. Herceg. Sarajevo, Vol. 9, p. 529 (1904); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 28 (1907); Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 101 [1909] (*Coptophlebia*); Lundbeck, Dipt. Dan. Vol. 3, p. 105 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910). Europe.
- albipennis*, Zetterstedt (not Meigen), Dipt. Scand. Vol. 1, p. 386 (1842).
- hyalipennis*, Schiner, part. Fauna Dipt. Austr. Vol. 1, p. 106 (1862).
- var. *impennis*, Strobl, Glasnik Mus. Bosn. Herceg. Sarajevo, Vol. 14, p. 470 (1902); Mitteil. Bosn. Herceg. Sarajevo, Vol. 9, p. 529 (1904); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 66 (1909). C. Europe.
- var. *pennata*, Strobl (not Schrank), Glasnik Mus. Bosn. Herceg. Sarajevo, Vol. 14, p. 470 (1902); Mitt. Bosn. Herceg. Sarajevo, Vol. 9, p. 529 (1904). C. Europe.
19. *E. albipennis*, Meigen (not Zetterstedt), Syst. Beschr. Vol. 7, p. 84, 60 (1838); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 28 (1907); Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 101, note [1909] (*Coptophlebia*). Spain.

20. *E. albohalteralis*, Brunetti, Fauna Brit. Ind. Brachyc. Vol. 1, p. 350 (1920). Baluchistan.
21. *E. Aldrichii*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 309, f. 110 (1902). N. W. United States.
22. *E. algaebrasensis*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 172 (1909). Spain.
23. *E. algira*, Macquart, Dipt. Exot. Vol. 1, Pt. 2, p. 159 (1839); Explor. Scien. Algérie, Zool. Vol. 3, p. 443 (1849); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906); Becker, ibidem, Vol. 7, p. 125 (1907). Algeria.
24. *E. alpicola*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 71 (1892), Vol. 34, p. 204 (1898); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 303 (1906). Alps.
25. *E. alpina*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 24 (1867); Pokorný, Verh. Zool.-bot. Ges. Wien, Vol. 37, p. 393 (1887); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906). Alps.
26. *E. ambigua*, Bezzi, Ann. Mus. Hungar., Vol. 3, p. 439 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909). Peru.
27. *E. amplitarsis*, Brunetti, Fauna Brit. Ind. Brachyc. Vol. 1, p. 349, f. 29 (1920). Bombay.
28. *E. amyntis*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 493 (1849); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 398 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 299 (1902); Aldrich, Cat. Dipt. N. Amer. p. 321 [1905] (*amystis*). New York.
29. *E. angustipennis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 351 (1909). Peru.
30. *E. annulata*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 758 [1865] (*Pachymeria*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 457 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909). Chile.
31. *E. annulipes*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, p. 369 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 289, f. 115 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909). Mexico.
32. *E. antarctica*, Walker, Trans. Linn. Soc. Lond. (Zool.), Vol. 17, p. 341, 26 (1837); List Dipt. Brit. Mus. Vol. 3, p. 495 (1849); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909). Magellan Straits.
33. *E. anthracina*, Bigot, Mission Scient. Cap Horn, Dipt. p. 19 (1888); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, 345 (1909). Cape Horn.
34. *E. Apfelbecki*, Strobl, Glasnik. Zem. Bosn. Herceg. Sarajevo, Vol. 10, p. 407 (1898); Wiss. Mitt. Bosn. Herceg. Sarajevo, Vol. 7, p. 567 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 304 (1906). S. Europe.
35. *E. apicalis*, Loew, Berl. Ent. Zeitschr. Vol. 9, p. 237 (1865); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906). Imeretia.
- . *E. aprica*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 262 (1829), no description.
36. *E. aquilus*, White, Proc. Roy. Soc. Tasmania, 1916, p. 234 (1917). Tasmania.
37. *E. ardesiaca*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 30 (1822); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 58, 165 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 25 (1907). Spain.
38. *E. argyrozona*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 754 (1865); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909). Chile.
39. *E. armipes*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 323 : Cent. 1, No. 32 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 300, 353 (1902). E. United States.
40. *E. arthritica*, Melander, ibidem, Vol. 28, p. 318, pl. 9, f. 135, 144 (1902). E. United States.
41. *E. asema*, Melander, ibidem, Vol. 28, p. 294, pl. 9, f. 130 (1902); Tucker, Kansas Univ. Sc. Bull. Vol. 4, p. 97 (1907). S. United States.

42. *E. assimilis*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 79 C. Europe.
(1893); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 29 (1907).
43. *E. atra*, Wiedemann, Aussereurop. Zweifl. Ins. Pt. 2, p. 1 (1829); Schiner, Brazil.
Novara Reise, Dipt. p. 203 (1868); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 344 (1909).
penthophora, Wiedemann, Aussereurop. Zweifl. Ins. Pt. 2, p. 4, 6 (1830);
Walker, Ins. Saunders, Dipt. Vol. 1, pl. 6, f. 1 [1852] (*Niconia*);
Trans. Ent. Soc. Lond. Vol. 4, p. 147 [1857] (*Niconia*).
44. *E. atrifemur*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, Mexico,
p. 370 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 290,
f. 116, 117 [1902] (*antifemor*); Bezzii, Nova Acta Akad. Naturf. Halle,
Vol. 91, p. 346 (1909).
45. *E. avida*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 405 (1895); Illinois.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 307 (1902).
46. *E. asteca*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, p. 369 Mexico.
(1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 291, f. 119
(1902); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909).
47. *E. barbata*, Macquart, Mém. Soc. Sc. Lille, p. 164 (1823). France.
48. *E. basalis*, Loew, Bes. Eur. Dipt. Vol. 3, p. 223 (1873); Kuntze, Zeitschr.
Hym. Dip. Vol. 6, p. 299 (1906). S. Russia.
49. *E. basilaris*, Becker, Mitteil. Zool. Mus. Berl. Vol. 4, p. 44 (1908). Canary Islands.
50. *E. bellatorius*, White, Proc. Roy. Soc. Tasmania, 1916, p. 232 (1917). Tasmania.
51. *E. bicolor*, Bellardi, Mem. Accad. Sc. Torino (2), Vol. 21, p. 198, Saggio,
Vol. 2, p. 98 (1862); Wheeler & Melander, Biol. Centr. Amer.
Dipt. Vol. 1, p. 370 (1901); Melander, Trans. Amer. Ent. Soc.
p. 291, f. 112, 113 (1902); Bezzii, Nova Acta Akad. Naturf. Halle,
Vol. 91, p. 344, 345 (1909). Mexico.
52. *E. bifasciata*, Olivier, Encycl. Méthod. Vol. 6, p. 390 (1791). France.
53. *E. Bigoti*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 319 (1902);
Coquillett, Proc. Ent. Soc. Wash. Vol. 5 p. 365 (1903). California.
- cinerea*, Bigot (not Zetterstedt), Ann. Soc. Ent. France (6), Vol. 2, Bull.
(9), 91, p. 112 [1881] (*Enoplectus*); Coquillett, Proc. U. S. Nat. Mus.
Vol. 18, p. 388, 397 (1895).
54. *E. bistortæ*, Meigen, Syst. Beschr. Vol. 3, p. 29 (1822); Curtis, Brit. Ent.
Vol. 8, p. 18 (1824); Macquart, Hist. Nat. Dipt. Vol. 1, 329 [1834];
(*bistorta*); Schummel, Uebers. Schles. Ges. Vat. Cultur. Breslau,
p. 189 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 496 (1849);
Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19) p. 53 (1851); Schiner,
Fauna Dipt. Austr. Vol. 1, p. 104 (1862); Fr. Stein, Stettin. Ent.
Zeit Vol. 34, p. 242 (1873); Strobl, Mitteil. Nat. Ver. Steiermark,
Graz, Vol. 29, p. 84 (1892); Bezzii, Bull. Soc. Ent. Ital. Vol. 30,
p. 139 (1899); Czizek, Zeitschr. Mähn. Landesmus. Brünn, Vol. 7,
p. 165 (1907); Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 94
[1909] (*Anacrostichus*).
truncata, Loew (not Meigen), Berl. Ent. Zeitschr. Vol. 12, p. 235, 240, 3 (1868).
55. *E. bistortæ*, Walker, (not Meigen), Ins. Brit. Vol. 1, p. 92 (1851). England.
56. *E. bivittata*, Wiedemann, Anal. Ent. p. 28 (1824); Aussereurop. Zweifl. Ins.
Pt. 2, p. 3 (1824); Loew, Oefv. Vet. Akad. Förh, Stockholm, Vol. 14,
p. 369 (1857); Dipterent. Südafri. p. 263 (1860).
57. *E. borealis*, Linneus, Syst. Nat. ed. 10, p. 601 (1758); Fauna Suec. p. 1895 Europe, Siberia.
(1761); Syst. Nat. ed. 12, Vol. 2, p. 1003 (1767); Fabricius, Syst.
Ent. p. 801 (1775); De Geer, Mém. Hist. Ins. Vol. 6, p. 255, pl. 14,
f. 17-19 (1776); Sulzer, Abgek. Gesch. Ins. 221, pl. 28, f. 15
(1776); Müller, Zool. Dan. Prödr. p. 182 (1776); Fabricius, Spec.

Ins. Vol. 2, p. 471 (1781); Mant. Ins. Vol. 2, p. 364 (1787); De Geer, Mém. Ins. (ed. Götze), Vol. 6, p. 102, pl. 14, f. 17 (1782); Herbst, Gem. Naturg. Vol 8, p. 117, pl. 344, f. 1 (1787); Gmelin, Syst. Nat. Vol. 5, p. 2889 (1788); Retzius, Gen. Sp. Ins. p. 190 (1783); Olivier, Encycl. Méthod. Vol. 6, p. 387 (1791); Fabricius, Ent. Syst. Vol. 4, p. 403, (1794); Schrank, Fauna Boica, Vol. 3, p. 170 (1803); Latreille, Hist. Nat. Crust. Ins. Vol. 14, p. 311 [1804] (*Platyptera*); Dict. Hist. Nat. Vol. 24, p. 191 [1804] (*Platyptera*); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 219, pl. 11, f. 25, 26 (1804); Fabricius, Syst. Antl. p. 137 (1805); Fallen, Empid. Suec. p. 16 (1815); Billberg, Enum. Ins. p. 120 (1820); Meigen, Syst. Beschr. Vol. 3, p. 36 (1822); Macquart, Mém. Soc. Sc. Lille (1823) p. 158 (1823); Curtis, Brit. Ent. Vol. 1, p. 18 (1824); Macquart, Dipt. N. France, Vol. 3, p. 122 (1827); Boitard, Man. Ent. Vol. 2, p. 369 [1828] (*Platyptera*); Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 263 [1829] (*Platypterygia*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 328 (1834); Zetterstedt, Fauna Ins. Lappon. p. 560 (1838); Loew, Be- merk. Posen Gegend Art. Zweifl. Ins. (1840), p. 20 : Isis, Vol. 7, p. 546 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 370 (1842); Gimberthal, Bull. Soc. Nat. Mosc. Vol. 20 (2), p. 163 (1847); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 497 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3123 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 52 (1851); Walker, Ins. Brit. Vol. 1, p. 90 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 427 (1852); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. Ins. Vol. 1, p. 164 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 105 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 122 (1887); Strobl, Mitteil. Steiermark, Graz, Vol. 29, p. 63 (1892), Vol. 34. p. 203 (1898); Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9) p. 28 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906); Howlett, Ent. M. Mag. London, Vol. 43, p. 229 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 137, f. 45 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 43 (1913).

58. *E. brachysoma*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 409 (1900); Alaska.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 308 (1902).
59. *E. brevicornis*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 263 (1869); Verrall, C. Europe.
Ent. M. Mag. London, Vol. 30, p. 140 (1894); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906); Lundbeck, Dipt. Dan. Vol. 3, p. 131, f. 43 (1910).
60. *E. brevipennata*, Macquart, Dipt. France, Vol. 3, p. 126 (1827); Hist. Nat. Dipt. Vol. 1, p. 332 (1834); Meigen, Syst. Beschr. Vol. 7, p. 83 [1838] (*brevipennis*); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Schiner, Fauna, Dipt. Austr. Vol. 1, p. 109 [1862] (*brevipennis*).
61. *E. brevirostris*, Macquart, Dipt. Exot. 4th. Suppl. 400, p. 9, pl. 9, f. 7, Tasmania.
(1847); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 343, note (1904).
62. *E. brevis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 196 : Cent. 3, No. 22 [1862] E. United States.
(*Pachymeria*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 331 [1902] (*Pachymeria*).
63. *E. Brouni*, Hutton, Trans. New. Zeal. Inst. Vol. 33, p. 30 (1901). New Zealand.
64. *E. brunnea*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 270 (1903). California.
65. *E. brunnipennis*, Meigen, Syst. Beschr. Vol. 3, p. 31 (1822); Curtis, Brit. Ent. Vol. 8, p. 18 (1824); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 53 (1851); Walker, Ins. Brit. Vol. 1, p. 92 (1851); Schiner,

- Fauna Dipt. Austr. Vol. 1, p. 109 (1862); Strobl, Wien. Ent. Zeit. Vol. 12, p. 39 (1893); Glasnik Zem. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 410 (1898); Wiss. Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 570 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 32 (1907).
66. *E. brunneipes*, Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 224 (1804). C. Europe.
67. *E. *bulbirostris*, Meunier, Loew, Bernsteinfauna, p. 41 [1850] (*unnamed*); Meunier, Miscell. Ent. Vol. 7, p. 178 (1899). Lower Oligocene, Baltic Amber.
68. *E. bullata*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 442 (1905); Nova Acta, Akad. Naturf. Halle, Vol. 91, p. 347 (1909). Bolivia.
69. *E. cacuminifer*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 304, pl. 9, f. 124 (1902). United States.
70. *E. cæligena*, Melander, ibidem, Vol. 28, p. 314, f. 107 (1902). Alabama.
71. *E. calcarata*, Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 131 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906). Italy.
72. *E. canaster*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 326, f. 139, 140 (1902). W. United States.
73. *E. candida*, Rossi, Fauna Etrusca, Vol. 2, p. 335 (1790), ed. 2, p. 505 Italy. (1807).
74. *E. candidata*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 226 (1873); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 299 (1906). S. Russia.
75. *E. captus*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 405 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 310 (1902). S. E. United States.
76. *E. carbonaria*, Brunetti, Rec. Indian Mus. Vol. 9, p. 27 (1913); Fauna Brit. India Dipt. Vol. 1, p. 351, f. 30 (1920). Ceylon.
77. *E. ?*carbonum*, Germar, Fauna Ins. Vol. 19, p. 21, pl. 21 (1837); Verz. Samml. Bayr. p. 71 (1840); Giebel, Deutschl. Petref. p. 642 (1852); Ins. Vorwelt. p. 207 (1856); Handlirsch, Fossil. Ins. p. 1017 [genus?] (1908). Upper Miocene, Bavaria.
78. *E. caucasica*, Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 90, 92, [1909] (Xanthempis). Caucasus.
79. *E. caudatula*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 33, 53 (1867); Strobl, Mitteil. Natur.-Ver. Steiermark, Graz, Vol. 29, p. 65 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906); Lundbeck, Dipt. Dan. Vol. 3, p. 124 [1910] (*Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 64 (1910). C. Europe.
- morosa*, Macquart (not Meigen), Mém. Soc. Sc. Lille, 1827, p. 126, 17 (1827); Hist. Nat. Dipt. p. 332, 22 (1834); Boitard, Man. Ent. p. 318 (1843); Scholtz, Zeitschr. Ent. Breslau, Vol. 5, p. 53 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 109 (1862).
80. *E. centralis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 26 (1913); Fauna Brit. India, Dipt. Vol. 1, p. 353 (1920). W. Himalayas.
81. *E. ceylonica*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 343 (1904), Vol. 10, p. 467 [1912] (*Coptophlebia*); Brunetti, Fauna Brit. India, Dipt. Vol. 1, p. 352 (1920). Ceylon.
82. *E. chioptera*, Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 221 [1804] (not 1822); Fallen, Empid. Suec. p. 21 (1815); Curtis, Brit. Ent. Vol. 8, p. 18 (1824); Macquart, Dipt. N. France, Vol. 3, p. 128 (1827); Brullé, Expéd. Morée, Vol. 3, p. 128 (1832); Macquart, Hist. Nat. Dipt. Vol. 1, p. 330 (1834); Zetterstedt, Fauna Ins. Lappon, p. 561 (1838); Dipt. Scand. Vol. 1, p. 376 (1842); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 495 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3024 (1849); Scholtz, Zeitschr. Ent. Breslau, Vol. 5 (19).

p. 52 (1851); Walker, Ins. Brit. Vol. 1, p. 92 (1851); Bonsdorff, Finl. Ins. Vol. 1, p. 165 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 106 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 43, 54 (1867); Glover, Manuscr. Notes, p. 21, pl. 11, f. 6 (1874); Leunis, Synops. Zool. Vol. 2, p. 402 (1886); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 70 (1892); Glasnik Mus. Bosn. Herceg. Sarajevo, Vol. 14, p. 470 (1902); Mitteil. Bosn. Herceg. Sarajevo, Vol. 9, p. 529 (1904); Wesché, Trans. Linn. Soc. Lond. [Zool.] (2), Vol. 9, p. 4, pl. 8, f. 2 (1904); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 304 (1906); Czizek, Zeitschr. Mähr. Landesmus. Brünn, Vol. 7, p. 165 (1907); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 65 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 118, f. 39, 40 [1910] (*Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 64 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 43 [1913] (*Pterempis*).

alpicola, Strobl, part. Mitteil. Natur. Ver. Steiermark, Graz, Vol. 29, p. 71 (1893).

? *crassipes*, Schrank [1781] (not Meigen), Ins. Austr. p. 484 (1781); Gmelin, Syst. Nat. Vol. 5, p. 2890 (1790); Schrank, Fauna Boica, Vol. 3, p. 172, 2576 (1803); Meigen, Classif. Beschr. Zweifl. Ins. Vol. 1, p. 234 (1804).

var. *sicula*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 45, 55 (1867); Strobl, Wien. Ent. Zeit. Vol. 18, p. 17, 25 (1899); Mitteil. Bosn. Herceg. Sarajevo, Vol. 9, p. 529 (1904); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 304 (1906). S. Europe.

83. *E. chiragra*, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 353 (1909). Peru.

84. *E. ciliata*, Fabricius, Mant. Ins. Vol. 2, p. 365 (1787); Schaeffer, Icon. Ins. pl. 192, f. 3 (1779); Gmelin, Syst. Nat. Vol. 5, p. 2890 (1788); Olivier, Encycl. Méthod. Vol. 6, p. 388 (1791); Fabricius, Ent. Syst. Vol. 4, p. 405 (1794); Syst. Antl. p. 140 (1805); Meigen, Syst. Beschr. Vol. 3, p. 20 (1822); Macquart, Mém. Soc. Sc. Lille, p. 160 (1823); Dipt. N. France, Vol. 3, p. 122, pl. 3, f. 6 (1827); Hist. Nat. Dipt. Vol. 1, p. 330, 15 (1834); Zetterstedt, Dipt. Scand. Vol. 12, p. 4611, note (1855); Schiner, Fauna Dipt. Austr. Vol. 1, p. 104 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 2, 7 (1867); Giard, Bull. Scient. Dept. Nord, France, Lille, Vol. 5, p. 192 (1873); Kunckel d'Herculais, Rech. Dipt. Vol. 2, pl. 20, f. 6 (1881); Adolph, Nova Acta Akad. Naturf. Halle, Vol. 47, p. 271, pl. 24, f. 11 (1885); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 297 (1906).

aurata, Villers, Ent. Linn. Vol. 3, p. 571 (1789).

boja, Schrank, Fauna Boica, Vol. 3, p. 170 (1803).

pennata, Panzer (not Schrank), Fauna Germ. 91, 22 (1804); Fabricius, Syst. Antl. p. 140, 11 (1805); Macquart, Mém. Soc. Sc. Lille, p. 160 (1823).

pennipes, Meigen (not Linnæus), Classif. Beschr. Ent. Zweifl. Ins. Vol. 1, p. 219, pl. 11, f. 27 (1804); Fallen, Empid. Suec. p. 19 (1816).

85. *E. ciliatopennata*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 80 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 26 (1907). C. & S. Europe.

var. *cantabrica*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 20 (1899). Spain.

86. *E. cincinnatula*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 31, 52 (1867); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 65 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906). C. Europe.

87. *E. cinerea*, Müller, Zool. Dan. Prodr. p. 182 (1776). Denmark.

88. *E. cinerea*, Zetterstedt, Dipt. Scand. Vol. 12, p. 4609, 7-8 (1855); Frey, Medd. Soc. Sc. Fenn. Helsingfors, Vol. 32, p. 108 (1906); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 31 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 127 [1910] (? *Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 40, pl. 1, f. 4 [1913] (*Pterempis*). N. Europe.

89. *E. cingulata*, Gimmerthal, Bull. Soc. Nat. Moscou, Vol. 7, p. 113 (1834). Russia.
 90. *E. clauda*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 407 (1900); Alaska.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 321 (1902).
 91. *E. clauda*, Schrank, Fauna Boica, Vol. 3, 171 (1803). C. Europe.
 92. *E. clausa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 401 (1895); W. United States.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 293, f. 129 (1902);
 Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 88 [1909] (*Coptophlebia*).
 93. *E. cognata*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 340 (1860); C. & S. Europe.
 Schiner, Fauna Dipt. Austr. Vol. 1, p. 105 (1862); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906).
 —. *E. cognata*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 263 (1829), no description.
 94. *E. colonica*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 498 (1849); Nova Scotia.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 300 (1902).
 95. *E. comantis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 402 (1895); California.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 308 (1902).
 96. *E. completa*, Loew, Dipterent. Südafr. p. 266 (1857); Oefv. Vet. Akad. Caffraria.
 Förh. Stockholm, Vol. 14, p. 370 (1857); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 343, note 4 (1904).
 97. *E. compta*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 405 (1895); C. United States.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 306 (1902).
 98. *E. concisa*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 70 (1869); Kuntze, Ephesus, Asia Minor.
 Zeitschr. Hym. Dipt. Vol. 6, p. 298 (1906).
 99. *E. concolor*, Verrall, Ent. Mag. London, Vol. 8, p. 283 (1872); Kuntze, W. Europe.
 Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 90 [1909] (*Xanthempis*).
 occipitalis, Bezzi, Deutsche Ent. Zeitschr. Suppl. p. 91 [1909] (*Xanthempis*).
 100. *E. confusa*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 227 (1907); Bezzi, N. Africa.
 Deutsche Ent. Zeitschr. Beiheft, p. 101, 102 [1909] (*Coptophlebia*).
 101. *E. connexa*, Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9) p. 28, Siberia.
 pl. 2, f. 29, 30 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 156 (1907).
 102. *E. contigua*, Loew, Wien. Ent. Monatschr. Vol. 8, p. 360 [1864] Greece.
 (*Pachymeria*); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 157 (1907);
 Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 96, note, & 98, note [1909] (*Pachymeria*).
 103. *E. coracina*, Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 102, 103 [1909] France.
 (*Coptophlebia*).
 104. *E. corcyrica*, Bezzi, ibidem, p. 102, 103 [1909] (*Coptophlebia*). Corfu.
 105. *E. corvina*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 90 (1869); Kuntze, C. Europe.
 Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906).
 106. *E. cothurnata*, Brullé, Expéd. Morée, Vol. 3, p. 298, pl. 46, f. 8 (1834); Greece.
 Loew, Wien. Ent. Monatschr. Vol. 8, p. 256 (1864); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906).
 morio, Schiner, part. (not Fabricius) Fauna Dipt. Vol. 1, p. 109 (1862).
 107. *E. cotoxanthus*, Blanchard, Hist. Fis. Polit. Chile (Zool.) Vol. 7, p. 372, Chile.
 pl. 3, f. 3 (1852); Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 753 (1865); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 343 (1909).
 108. *E. coxalis*, Thomson, Eugen. Resa, Ins. p. 471 (1858); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909). Patagonia.
 109. *E. crassa*, Nowicki, Verh. Naturf. Ver. Brünn. Vol. 6, p. 82 (1868); C. Europe.
 Loew, Berl. Ent. Zeitschr. Vol. 13, p. 77 (1869); Strobl, Mitteil.

- Naturw. Ver. Steiermark, Graz, Vol. 29, p. 64 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 299 (1906).
grandis, Loew, in litt.
110. *E. crassifila*, Loew, Oefv. Vet. Akad. Forhandl. Stockholm, Vol. 14, Cape of Good Hope. p. 369 (1857); Dipterenf. Südafr. p. 264, pl. 2, f. 48 (1860).
111. *E. crassipes*, Meigen (not Schrank), Syst. Beschr. Vol. 3, p. 33 (1822); C. Europe. Loew, Berl. Ent. Zeitschr. Vol. 11, p. 59 (1857); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 99, note [1909] (*Lissemphis*).
112. *E. crassitarsata*, Macquart, Mém. Soc. Sc. Lille, p. 162 (1823). France.
113. *E. cuneipennis*, Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 126 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906); Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 99 [1909] (*Lissemphis*).
114. *E. curta*, Loew. Beschr. Eur. Dipt. Vol. 1, p. 261 (1869); Kuntze, S. Russia. Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906).
115. *E. curvipes*, Loew, Berl. Ent. Zeitschr. Vol. 12, p. 391, 393 (1868); S. Russia. Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 32 (1907); Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 96 [1909] (*Polyblepharis*).
116. *E. cylindrica*, Fabricius, Ent. Syst. Vol. 4, p. 403 (1794); Coquebert, Barbary. Illustr. Icon. Ins. p. 121, pl. 37, f. 8 (1804); Fabricius, Syst. Antl. p. 138 (1805); Wiedemann, Aussereurop. Zweifl. Ins. Pt. 2, p. 5 (1830).
117. *E. dasychira*, Mik, Jährbed. Akad. Gymn. Wien (1878); Dipt. Untersuch. C. Europe. p. 24, pl. 1, f. 21 (1878); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 73 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 28 (1907).
118. *E. dasynota*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 73 (1869); Kuntze, Greece. Zeitschr. Hym. Dipt. Vol. 6, p. 290 (1905).
119. *E. dasypoda*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 344 (1860); Sicily. Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 28 (1907).
120. *E. dasyprocta*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 46, 55 (1867); C. Europe. Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 303 (1906).
anfractuosa, Mik, Wien. Ent. Zeit. Vol. 3, p. 4 (1884); in Beck, Fauna Hernstein, Vol. 2 (2), p. 57, f. 5-7 (1885); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 24, p. 203 (1898).
chioptera, Meigen, Syst. Beschr. Vol. 3, p. 27, pl. 22, f. 19 (1822).
?rapida, Meigen, Syst. Beschr. Vol. 7, p. 86, 64 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); ?Loew, Berl. Ent. Zeitschr. Vol. 11, p. 61, 18 (1867); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 70 (1893).
121. *E. dasythrix*, Meijere, Tijdschr. v. Ent. Vol. 61, p. 132, pl. 8, f. 1, 2 Belgium. (1918).
122. *E. decora*, Meigen, Syst. Beschr. Vol. 3, p. 22, pl. 22, f. 18 (1822); C. & S. Europe. Curtis, Brit. Ent. Vol. 8, p. 18 (1824); Macquart, Dipt. N. France, Vol. 3, p. 124 (1827); Hist. Nat. Dipt. Vol. 1, p. 331 (1834); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Walker, Ins. Brit. Vol. 1, p. 92 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 125 (1887); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 76 (1893); Bezzii, Bul. Soc. Ent. Ital. Vol. 30, p. 135 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 29 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 112 [1910] (*Pteremphis*).
var. genualis, Strobl, Wien. Ent. Zeit. Vol. 12, p. 39 (1893); Glasnik Zem. S. Europe. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 408 (1898); Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 135 (1899); Strobl, Wien. Ent. Zeit. Vol. 38, p. 19 (1899); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 569 (1900).

123. *E. decoripes*, Röder, Ent. Nachr. Berlin, p. 203 (1894); Kuntze, Zeitschr. Asia Minor. Hym. Dipt. Vol. 6, p. 299 (1906).
124. *E. dedecor*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 65 (1869); Kuntze, S. E. Europe. Zeitschr. Hym. Dipt. Vol. 6, p. 297 (1906); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 173 (1909).
125. *E. depilis*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 231 (1873); Kuntze, S. Russia. Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906).
126. *E. digramma*, Meigen, Syst. Beschr. Vol. 7, p. 87 (1838); Bezzi, Europe. Deutsche Ent. Zeitschr. Beiheft, p. 89 [1909] (*Xanthempis*).
bilineata, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 14, 20 (1867); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 60 (1892); Elliot, Trans. Ent. Soc. Lond. 1896, p. 117, 128 (1896); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Lundbeck, Dipt. Dan. Vol. 3, p. 86, f. 24, 25 [1910] (*Xanthempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910).
punctata, Fabricius (not Meigen), Syst. Antl. p. 142, 22 (1805); Fallen, part. Dipt. Suec. Emp. p. 19, 6 (1815); Meigen, Syst. Beschr. Vol. 6, p. 338, 51 (1830); Schiner, Fauna Austr. Dipt. Vol. 1, p. 107 (1862); Müller, Kosmos, Vol. 9, p. 416 (1881); Ent. Nachr. Bul. Vol. 8, p. 116 (1882); Mik, Wien. Ent. Zeit. Vol. 1, p. 203 (1882).
testacea, Zetterstedt (not Fabricius), Dipt. Scand. Vol. 1, p. 378, 11 (1842), Vol. 8, p. 3024 (1849).
127. *E. dimidiata*, Meigen, Gistl. Faunus, Vol. 2, p. 58 (1835); Bezzi, C. & S. Europe. Deutsche Ent. Zeitschr. Beiheft, p. 91 (1909).
mesogramma, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 22 (1867).
monogramma, Meigen, Syst. Beschr. Vol. 7, p. 87 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 62 (1892); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 129 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906); Bezzi, Ann. Mus. Hungar. Vol. 6, p. 394 (1908).
128. *E. discolor*, Loew, Neue Beitr. Dipt. Vol. 4, p. 34 (1856); Schiner, C. Europe, Alps. Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Pokorny, Verh. Zool.-bot. Ges. Wien. Vol. 37, p. 394 (1887); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 62 (1892); Kuntze, Zeitschr., Hym. Dipt. Vol. 6, p. 213 (1906); Bezzi, Ann. Mus. Hungar. Vol. 6, p. 394 (1908).
129. *E. dispar*, Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 53 (1851); C. Europe, Alps. Schiner, Fauna Dipt. Austr. Vol. 1, p. 109 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 22 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906); Bezzi, Ann. Mus. Hungar. Vol. 6, p. 394 (1908); Deutsche Ent. Zeitschr. Beiheft, p. 100 [1909] (*Argyrandrus*).
130. *E. distans*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 32 : Cent. 8, No. 54 (1869); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 295, f. 127 (1902); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 101 [1909] (*Coptophlebia*). S. & E. United States.
131. *E. divergens*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 72 (1869); Kuntze, Greece. Zeitschr. Hym. Dipt. Vol. 6, p. 298 (1906).
132. *E. divisa*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 257 (1869); Kuntze, Greece. Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906); Bezzi, Ann. Mus. Hungar. Vol. 6, p. 394 (1908).
133. *E. dolabraria*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 325 (1902); California. Aldrich, Cat. Dipt. N. Amer. p. 322 [1905] (*dolabraria*). var. *disconvenita*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 326 (1902). California.
134. *E. dolorosa*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, Mexico. p. 370 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 295, f. 128 (1902); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 101 (1909); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909).

135. *E. dumetorum*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 756 Chile.
 (1865); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909).
136. *E. Dusmetii*, Strobl, Verh Zool.-bot Ges. Wien, Vol. 59, p. 173 (1909). Spain.
137. *E. elegans*, Brunetti, Rec. Indian Mus. Vol. 9, p. 26 (1913); Fauna Brit. W. Himalayas.
 India Dipt. Vol. 1, p. 356 (1920).
138. *E. elongata*, Meigen, Syst. Beschr. Vol. 7, p. 84 (1838); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 61 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 297 (1906).
139. *E. enodis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 303, pl. 9, Illinois.
 f. 125 (1902).
140. *E. Erberi*, Nowick, Beitr. Ins. Faune Galiziens, p. 5 [1873] (*Pachymeria*); C. & S. Europe.
 Loew, Beschr. Eur. Dipt. Vol. 3, p. 217 [1873] (*Pachymeria*); Bezzi, Bull. Soc. Ent. Ital. Vol. 31, p. 81 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 158 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98 [1909] (*Pachymeria*); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912).
141. *E. erosa*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 260 (1869); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906). S. Europe.
142. *E. eudamides*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 493 (1849); North America.
 Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 299 [1902] (*eudamides*).
143. *E. eumera*, Loew, Berl. Ent. Zeitschr. Vol. 12, p. 388, 393 (1868); S. Russia.
 Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 32 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 95, 96 [1909] (*Polyblepharis*).
phenomeris, Loew, Berl. Ent. Zeitschr. Vol. 12, p. 172 [♀] (1868).
144. *E. eupæza*, Loew, Zeitschr. Geg. Naturw. Berlin, Vol. 43 p. 417 (1874); Transcasplia.
 Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 27 (1907).
145. *E. Eversmanni*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 227 (1873); Russia.
 Jaroschewsky, Arb. Gess. Naturf. Univ. Kharkow, Vol. 11, p. 353 (1877), Vol. 12, p. 9 (1880); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906); Bezz, Deutsche Ent. Zeitschr. Beiheft, p. 91, note (1909).
146. *E. exilis*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 269 (1903). Missouri.
147. *E. exotica*, Wiedemann, Anal. Ent. p. 28 (1824); Aussereurop. Zweifl. Ins. Pt. 2, p. 2 (1830). Cape of Good Hope.
148. *E. falcata*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 326, pl. 9, W. United States.
 f. 137 (1902).
149. *E. fallax*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 340 (1860); C. & S. Europe.
 Schiner, Fauna Dipt. Austr. Vol. 1, p. 105 (1862); Strobl, Glasnik Mus. Bosn. Herceg. Sarajevo, Vol. 14, p. 470 (1902); Mitth. Bosn. Herceg. Sarajevo, Vol. 9, p. 529 (1904); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906).
- . *E. fasciata*, Schummel, Arbeiten Schles. Ges. Vaterl. Kult. Breslau, 1832, p. 70 (1832), no description.
150. *E. fasciculata*, Strobl, Jahrb. Mus. Karnten, Klagenfurt, Vol. 26, p. 200 C. Europe.
 (1901); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 29 (1907).
151. *E. femorata*, Fabricius, Ent. Syst. Suppl. 568 (1798); Coquebert, C. & S. Europe.
 Illustr. Icon. pl. 27, f. 8 (1804); Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 223 (1804); Fabricius, Syst. Antl. p. 140 (1805); Latreille, Gen. Crust. Ins. Vol. 4, p. 303 (1809); Meigen, Syst. Beschr. Vol. 3, p. 40, pl. 22, f. 20 (1822); Macquart, Mém. Soc. Sc. Lille, p. 160 (1823); Curtis, Brit. Ent. Vol. 8, p. 18 [1824] (*Pachymeria*); Macquart, Dipt. N. France, Vol. 3, p. 128 (1827); Hist. Nat. Dipt. Vol. 1, p. 333, pl. 8, f. 2 (1834); Meigen, Syst.

- Beschr. Vol. 7, p. 89, 67, f. 13-17 [1838] (*Pachymeria*); Blanchard, Hist. Nat. Ins. Vol. 3, p. 581 [1840] (*Pachymeria*); Boitard, Man. Ent. Vol. 3, p. 319 [1843] (*Pachymeria*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 110 [1862] (*Pachymeria*); Loew, Wien. Ent. Monatschr. Vol. 8, p. 362 [1864] (*Pachymeria*); Leunis, Synops. Zool. Vol. 2, p. 402 [1886] (*Pachymeria*); Neuhaus, Dipt. March. p. 72 [1886] (*Pachymeria*); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 158 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98 [1909] (*Pachymeria*); Hamm, Ent. Mag. London, Vol. 45, p. 160 [1909] (*Pachymeria*) (mating); Kuntze, Deutsche Ent. Zeitschr. p. 548 (1913).
- ? *confidens*, Harris, Engl. Ins. p. 151, pl. 44, f. 7 (1782).
- ? *dubia*, Schiner, Verh. Zool.-bot. Ges. Wien, Vol. 6, p. 421 [1856] (*Pachymeria*); *quinquevittata*, Macquart, Mém. Soc. Sc. Lille, 1827, p. 129, 23 [1827] (*Pachymeria*); Hist. Nat. Dipt. Vol. 1, p. 334 [1834] (*Pachymeria*), Vol. 2, p. 657 [1835] (*Pachymeria*); Meigen, Syst. Beschr. Vol. 7, p. 89 [1838] (*Pachymeria*).
- ruralis*, Meigen, Syst. Beschr. Vol. 3, p. 40, 46 (1822); Curtis, Brit. Ent. Vol. 8, p. 18, 4 [1824] (*Pachymeria*); Meigen, Syst. Beschr. Vol. 7, p. 89, 4 [1838] (*Pachymeria*).
- sericea*, Olivier, Encycl. Méth. Vol. 6, p. 389 (1791).
152. *E. filata*, Loew, Berl. Ent. Zeitsch. Vol. 17, p. 39 (1873); Kuntze, C. Europe. Zeitschr. Hym. Dipt. Vol. 6, p. 304 (1906).
153. *E. fimbria*, Walker, Ins. Saunders. Dipt. p. 204 (1852); Bezzi, Nova Act. Akad. Naturf. Halle, Vol. 41, p. 345 (1909).
154. *E. fumana*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 341 (1860); Schiner, Fauna Dipt. Austr. Vol. 1, p. 104 (1862); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 300 (1906).
155. *E. flabilis*, White, Proc. Roy. Soc. Tasmania, 1916, p. 235 (1917). Tasmania.
156. *E. flava*, Müller, Fauna Ins. Fridrichsdal, p. 87 (1764); Zool. Dan. Prodr. p. 182 (1776); Villers, Ent. Linn. Vol. 3, p. 571 (1789). N. Europe.
157. *E. flavicans*, Olivier, Encycl. Méthod. Vol. 6, p. 388 (1791). W. Europe.
158. *E. flavinervis*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 755 (1865); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909). Chile.
159. *E. flavitarsis*, Roser, Correspondenzbl. Landw. Ver. Württ. Stuttgart, Vol. 1, p. 53 (1840); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 97 (1909). C. Europe.
- leucopeta*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 40 (1873); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 157 (1907).
160. *E. flavobasalis*, Matsumura, Addit. Vol. 2, p. 353, pl. 21, f. 16 (1916). Japan.
161. *E. florisomma*, Loew, Neue Beitr. Dipt. Vol. 4, p. 35 (1856); Schiner, Fauna Dipt. Austr. Vol. 1, p. 106 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 26, 52 (1867); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 65 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 300 (1906). C. Europe.
162. *E. *florissantana*, Cockerell, Proc. Acad. Nat. Sc. Philad. 1914, p. 645 Florissant, Miocene. (1914).
163. *E. fraterna*, Loew, Berl. Ent. Zeitschr. Vol. 9, p. 239 (1865); Kuntze, S. Europe. Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906).
164. *E. Frauscheri*, Strobl, Jahrb. Mus. Karnten, Klagenfurt, Vol. 26, p. 198 C. Europe. (1901); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906).
165. *E. frontalis*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 271 (1903). Alaska.
166. *E. fulva*, Macquart, Mém. Soc. Sc. Lille, p. 160 (1823). France.
167. *E. fulvipes*, Wiedeman-Meigen, Syst. Beschr. p. 31 (1822); Strobl, Wien. Ent. Zeit. Vol. 18, p. 14 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 216 (1906); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 173 (1909). Portugal.

168. *E. fumida*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 409 (1900); Alaska.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 313 (1902).
169. *E. fumosa*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 5, 8 (1867); Kuntze, C. Europe.
Zeitschr. Hym. Dipt. Vol. 6, p. 298 (1906).
170. *E. funebris*, Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 226 C. Europe.
(1804); Syst. Beschr. Vol. 3, p. 18 (1822); Macquart, Hist. Nat.
Dipt. Vol. 1, p. 328 (1834); Schiner, Fauna Dipt. Austr. Vol. 1,
p. 104 (1862); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 216 (1906).
171. *E. funesta*, Meigen, Syst. Beschr. Vol. 7, p. 84 (1838); Schiner, Fauna C. Europe.
Dipt. Austr. Vol. 1, p. 109 (1862); Loew, Berl. Ent. Zeitschr.
Vol. 11, p. 61 (1867).
172. *E. fuscipes*, Brullé (not Gmelin), Expéd. Morée, Vol. 3, p. 299 (1834); Greece.
Loew, Berl. Ent. Zeitschr. Vol. 11, p. 62 (1867).
173. *E. fuscipes*, Gmelin, Syst. Nat. Vol. 5, p. 2891 (1790); Olivier, Encycl. Europe.
Méthod. Vol. 6, p. 391 (1791).
174. *E. Gaigeri*, Gercke, Wien. Ent. Zeit. Vol. 5, p. 163, pl. 2, f. 3-4 (1886); S. Europe.
Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906).
175. *E. geniculata*, Macquart (not Zetterstedt or Kirby; see *Iteaphila luctuosa* Algeria.
and *Macquarti*), Explor. Algérie (Zool.), Vol. 3, p. 443, pl. 3, f. 4
(1849).
Macquarti, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 127 (1907).
176. *E. ghigiana*, Bezzì, Atti Soc. Ital. Sc. Nat. Milano, Vol. 60, p. 435 [1921] Cirenaica.
(*Pterempsis*).
177. *E. gibbipes*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 308 (1906). Spain.
178. *E. gibbosa*, Gmelin, Syst. Nat. Vol. 5, p. 2891 (1790); Olivier, Encycl. Europe.
Méthod. Vol. 6, p. 391 (1791).
179. *E. gladiator*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 316, pl. 9, Kansas.
f. 134 (1902).
180. *E. gracilipes*, Philippi, Verh. Zool.-bot. Ges. Wien. Vol. 15, p. 755 W. South America.
(1865); Bezzì, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347, 356
(1909).
181. *E. gracilis*, Curtis, Brit. Ent. Vol. 8, p. 18 (1824). England.
182. *E. gracilitarsis*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 18 (1899); Kuntze, Spain.
Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906).
183. *E. gravipes*, Loew, Neue Beitr. Dipt. Vol. 4, p. 34 (1856); Schiner, Fauna Europe, Siberia.
Dipt. Austr. Vol. 1, p. 105 (1862); Strobl, Mitteil. Nat. Ver.
Steiermark, Graz, Vol. 29, p. 63 (1893); Kuntze, Zeitschr. Hym.
Dipt. Vol. 6, p. 214 (1906).
184. *E. gravis*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 30 (1822); Lusitania.
Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 29 (1907).
185. *E. grisea*, Fallen, part. Empid. Suec. p. 22, part (1815); Zetterstedt, Europe.
Dipt. Scand. Vol. 1, p. 373 (1842); Scholz, Zeitschr. Ent. Breslau,
Vol. 5 (19) p. 52 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 5007
(1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Loew,
Wien. Ent. Monatschr. Vol. 8, p. 355 (1864); Mik, Wien. Ent. Zeit.
Vol. 1, p. 180 (1882); ibidem, Vol. 9, p. 294 (1890); Strobl, Mitteil.
Nat. Ver. Steiermark, Graz, Vol. 29, p. 85 [1893] (*Pachymeria*); Bezzì,
Bull. Soc. Ent. Ital. Vol. 30, p. 140 (1899); Kuntze, Zeitschr. Hym.
Dipt. Vol. 7, p. 157 (1907); Bezzì, Deutsche Ent. Zeitschr. Beiheft,
p. 98 [1909] (*Pachymeria*); Lundbeck, Dipt. Dan. Vol. 3, p. 99, f. 34
[1910] (*Pachymeria*); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910).
flavipes, Zetterstedt, Dipt. Scand. Vol. 1, p. 374 (1842) label name.
- var. *nigriventris*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 85 Europe.
[1893] (*Pachymeria*).

- var. *nodosa*, Beling, Arch. Naturg. Berlin, Vol. 48, p. 208 (1862). C. Europe.
 var. *pilipes*, Meigen, Syst. Beschr. Vol. 3, p. 31 (1822); Curtis, Brit. Ent. Vol. 8, Europe.
 p. 18 (1824); Walker, List Dipt. Brit. Mus. Vol. 3, 496 (1849); Ins.
 Brit. Vol. 1, p. 94 (1851).
 var. *rufiventris*, Strobl (not Meigen), Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, C. Europe.
 p. 85 [1892] (*Pachymeria*).
 186. *E. griseonigra*, Brunetti, Rec. Indian Mus. Vol. 9, p. 25 (1913); Fauna W. Himalayas.
 Brit. India Dipt. Vol. 1, p. 353 (1920).
 187. *E. gulosa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 408 (1895); Illinois.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 297 (1902).
 188. *E. gymnopoda*, Bezzii, Ann. Mus. Hungar. Vol. 6, p. 395 (1908). C. Europe.
 189. *E. Haemis*, Loew, Wien. Ent. Monatschr. Vol. 6, p. 168 (1862); Kuntze, Bulgaria.
 Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906).
 190. *E. haemorrhica*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 67 (1869); Greece.
 Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 298 (1906).
 191. *E. helophila*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 28, 52 (1867); Mik, C. Europe.
 Dipt. Hernst. p. 59 (1885); Becker, Berl. Ent. Zeitschr. Vol. 33,
 p. 169 (1889); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906).
 chioptera, Becker (not Meigen), Berl. Ent. Zeitschr. Vol. 31, p. 123 (1887).
 192. *E. heteroptera*, Macquart, Dipt. Exot. Vol. 1, p. 276, pl. 13, f. 3 (1838). Cape of Good Hope.
 193. *E. hirsuta*, Becker, Mém. Acad. Sc. Petrograd, Vol. 28, No. 7, p. 59 Karskaja Tundra.
 (1915).
 194. *E. hirta*, Loew, Berl. Ent. Zeitschr. Vol. 9, p. 240 (1865); ibidem, Transcaucasia.
 Vol. 11, p. 4, 8 (1867).
 195. *E. hirticrus*, new name for *hirtipes*, Coquillett (not Wiedemann), Proc. New Mexico.
 Wash. Ent. Soc. Vol. 5, p. 270 (1903).
 tenebrosa, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 270 (1903).
 196. *E. hirtipes*, Wiedemann (not Coquillett), Anal. Ent. p. 28 (1824); Aus- Cape of Good Hope.
 sereeurop. Zweifl. Ins. Pt. 2, p. 3 (1830).
 197. *E. Hoffmannseggii*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 83, 84 (1869); C. & S. Europe.
 Becker, Berl. Ent. Zeitschr., Vol. 31, p. 123 (1887); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 302 (1906); Bezzii, Deutsche Ent.
 Zeitschr. Beiheft, p. 102, note [1909] (*Coptophlebia*).
 grissa, Wiedemann-Meigen (not Fallen), Syst. Beschr. Vol. 3, p. 30 (1822);
 Zetterstedt, Dipt. Scand. Vol. 1, p. 374, note (1842); Loew, Berl. Ent.
 Zeitschr. Vol. 11, p. 58, 164 (1867).
 198. *E. holosericea*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 495 (1849). France.
 199. *E. humilis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 403 (1895); Illinois.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 297 (1902).
 200. *E. hyalipennis*, Fallen, Empid. Suec. p. 21 (1815); Meigen, Syst. Beschr. Europe.
 Vol. 3, p. 24 (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 385 (1842);
 Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 53 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 106, part (1862); Malloch, Ent. Mag. Lond. Vol. 42, p. 257 (1906); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 28 (1907); Bezzii, Deutsche Ent. Zeitschr. Beiheft, 1909, p. 101 [1909] (*Coptophlebia*); Lundbeck, Dipt. Dan. Vol. 3, p. 103 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 40 [1913] (*Coptophlebia*).
 proxima, Meigen, Syst. Beschr. Vol. 7, p. 85 (1838).
 201. *E. hyalogyne*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 468, 473 [1912] Formosa.
 (*Pterempis*); Suppl. Ent. Berlin, Vol. 3, p. 72 (1914).
 202. *E. hystrichopyga*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 467, 469 [1912] Formosa.
 (*Coptophlebia*).
 203. *E. hystrix*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 47, 55 (1867); Kuntze, S. Europe.
 Zeitschr. Hym. Dipt. Vol. 7, p. 25 (1907).

204. *E. inclinata*, Bezzia, Ann. Mus. Hungar. Vol. 10, p. 468, 470 [1912] Formosa.
(*Coptophlebia*).
205. *E. incompleta*, Macquart, Dipt. Exot. Suppl. Vol. 1, p. 96, pl. 9, f. 1 (1846). Caffraria.
206. *E. inconspicua*, Brunetti, Rec. Indian Mus. Vol. 9, p. 28 (1913); ibidem, N. India.
Fauna Brit. India, Dipt. Vol. 1, p. 357 (1920).
207. *E. induta*, Bezzia, Nova Acta Akad. Nat. Halle, Vol. 91, p. 343, 348 (1909). Peru.
208. *E. infumata*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 409 (1900); Alaska.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 814 (1902).
209. *E. inornata*, Loew, Oefv. Vet. Akad. Förh. Stockholm, Vol. 14, Vol. 370 Cape of Good Hope.
(1857); Dipterent. Südafr. Vol. 2, p. 266 (1860).
210. *E. *inscita*, Meunier, Ann. Sc. Nat. (Zool.), Vol. 7, p. 93, 121, pl. 11, f. 5-6 (1908). Baltic Amber.
211. *E. intercepta*, new name. Corsica.
interrupta, Becker (not Macquart), Deutsche Ent. Zeitschr. p. 641 (1910).
212. *E. interrupta*, Macquart, Mém. Soc. Sc. Lille, p. 163 (1823). France.
213. *E. Jacobsoni*, Meijere, Tijdschr. v. Ent. Vol. 1, p. 251, pl. 6, f. 18 (1907); Java.
Bezzia, Ann. Mus. Hungar. Vol. 10, p. 468 [1912] (*Coptophlebia*).
214. *E. Johnsoni*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 303, f. 123 Pennsylvania.
(1902).
215. *E. Kerteszi*, Bezzia, Termes. Füzet. Vol. 23, p. 251 (1900); Kuntze, C. Europe.
Zeitschr. Hym. Dipt. Vol. 6, p. 300 (1906).
216. *E. Kuntzei*, Becker-Bezzia, Deutsche Ent. Zeitschr. 1909, Suppl. p. 90, 91 Corsica.
[1909] (*Xanthempis*); Becker, Deutsche Ent. Zeitschr. 1910, p. 641 (1910).
217. *E. labiata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 323; Cent. 1, No. 33 E. United States.
(1861); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 294 (1902).
218. *E. leta*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 82 (1869); Jahrb. Krakau, C. Europe.
Vol. 41, p. 13 (1870); Strobl. Mitteil. Nat. Ver. Steiermark, Graz,
Vol. 29, p. 60 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211
(1906); Bezzia, Deutsche Ent. Zeitschr. 1909, Beiheft, p. 89 [1909]
(*Xanthempis*).
219. *E. levigata*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 75: Cent. 5, No. 49 New Hampshire.
(1864); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 302 (1902).
220. *E. levius*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 40 (1873); Kuntze, C. Europe.
Zeitschr. Hym. Dipt. Vol. 6, p. 302 (1906).
221. *E. lamellicornis*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 124 (1887); Europe
Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 177 (1892);
Wien, Ent. Zeit. Vol. 18, p. 20 (1899); Mem. Soc. Esp. His. Nat.
Vol. 3, p. 309 (1906); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 30
(1907); Lundbeck, Dipt. Dan. Vol. 3, p. 116 (1910).
222. *E. Landbecki*, Philippi, Verh. Zool-bot. Ges. Wien, Vol. 15, p. 754 Chile.
(1865); Bezzia, Nova Acta Akad. Nat. Halle, Vol. 91, p. 346 (1909).
223. *E. laniventris*, Eschscholz, Entomgr. Vol. 1, p. 113 (1822); Wiedemann, Alaska, Bering Island,
Aussereurop. Zweifl. Ins. Pt. 2, p. 6 (1830); Macquart, Dipt. Washington.
Exot. Vol. 1, Pt. 2, p. 162 [1838] (*Eriogaster*); Coquillett, Dipt. Commander Isl. p. 343 (1899); Proc. Wash. Acad. Sc. Vol. 2, p. 408
(1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 300 (1902);
Bezzia, Deutsche Ent. Zeitschr. Beiheft, p. 88 (1909).
224. *E. lasionota*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 264 (1869); Kuntze, C. & S. Europe.
Zeitschr. Hym. Dipt. Vol. 6, p. 300 (1906).
225. *E. lepidopus*, Meigen (not Walker), Syst. Beschr. Vol. 3, p. 23 (1822); C. Europe.
Curtis, Brit. Ent. Vol. 8, p. 18, 4 (1824); Meigen, Syst. Beschr. Vol. 7, p. 88 (1838); Walker, List Dipt. Brit. Mus. Vol. 3, p. 495
(1849); Bezzia, Bull. Soc. Ent. Ital. Vol. 30, p. 136 (1899); Villeneuve,

- Bull. Soc. Ent. France, p. 212 (1903); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 298, 299 (1906).
- atra*, Macquart (not Wiedemann), Dipt. N. France, Vol. 3, p. 124 (1827); Hist. Nat. Dipt. Vol. 1, p. 331 (1834); Meigen, Syst. Beschr. Vol. 7, p. 82 (1838); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 60 (1867).
- setigera*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 80 (1869); Strobl, Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 570 (1900).
226. *E. lepidopus*, Walker (not Meigen), Ins. Brit. Vol. 1, p. 93 (1851). England.
227. *E. leptogastra*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 17 : Cent. 3, No. 30 (1863); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 300 (1902).
228. *E. leptomorion*, Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 101, 102 [1909] C. Europe, Alps. (*Coptophlebia*).
229. *E. leucoptera*, Gmelin, Syst. Nat. Vol. 5, p. 2891 (1790); Olivier, Encycl. Europe. Méthod. Vol. 6, p. 392 (1791).
230. *E. leucoptera*, Macquart (not Meigen), Mém. Soc. Sc. Lille, p. 161 (1823). France.
231. *E. leucostigma*, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347, 356 (1909). Peru.
232. *E. levicula*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 406 (1896); Illinois. Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 307 (1902).
233. *E. lineata*, Meigen (not Fabricius or Villers), Classif. Beschr. Eur. Zweifl. C. Europe. Ins. Vol. 1, p. 224 (1804); Syst. Beschr. Vol. 3, p. 29 (1822); Schiner, Fauna Dipt. Austr. Vol. 1, p. 109 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 58 (1867).
234. *E. lineata*, Villers, Ent. Linn. Vol. 3, p. 571, pl. 10, f. 20 (1789); Olivier Europe. Encycl. Méth. Vol. 6, p. 388 (1791).
235. *E. lioes*, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 344, 350 (1909). Peru.
236. *E. liosoma*, Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 99 [1909] (*Lissempis*). Tunis. *infumata*, Becker (not Coquillett), Zeitschr. Hym. Dipt. Vol. 7, p. 230 (1907).
237. *E. livida*, Linnaeus, Syst. Nat. Vol. 10, p. 604 (1758); Fauna Suec. Europe. p. 1897 (1761); Geoffroy, Hist. Ins. Vol. 4, p. 474 (1762); O. F. Müller, Ins. Friedrichsdal, p. 87 (1764); Linnaeus, Syst. Nat. Vol. 12 (2), p. 1003 (1767); Harris, Engl. Ins. p. 149, pl. 44, f. 1 (1776); Barbut, Genres Ins. Linn. p. 311, pl. 17, f. 3 (1781); De Geer, Hist. Ins. Vol. 6, p. 101, pl. 14, f. 14 (1782); Retzius, Gen. Sp. Ins. p. 190 (1783); Herbst, Gem. Naturg. Vol. 8, p. 118, pl. 344, f. 2 (1787); Gmelin, Syst. Nat. Vol. 5, p. 2889 (1788); Rossi, Fauna Etrusca, Vol. 2, p. 335 (1790); Schellenberg, Genres Mouch. p. 88, pl. 35, f. 1 (1803); Schrank, Fauna Boica Vol. 3, p. 171 (1803); Latreille, Dict. Hist. Nat. Vol. 24, p. 191 (1804); Hist. Crust. Ins. Vol. 14, p. 312, pl. 111, f. 1 (1804); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 225 (1804); Rossi, Fauna Etrusca (2), p. 505 (1807); Latreille, Gen. Crust. Ins. Vol. 4, p. 303 (1809); Fallen, Empid. Suec. p. 18 (1815); Lamarck, Anim. s. Vert. Vol. 3, p. 401 (1816); Billberg, Enumer. Ins. p. 120 (1820); Meigen, Syst. Beschr. Vol. 3, p. 35 (1822); Macquart, Mém. Soc. Sc. Lille, 1823, p. 159 (1823); Curtis, Brit. Ent. Vol. 8, p. 18, 2 (1824); Macquart, Dipt. N. France, Vol. 3, p. 119, pl. 3, f. 7 (1827); Boitard, Man. Ent. Vol. 2, p. 369 (1828); Macquart, Hist. Nat. Dipt. Vol. 1, p. 329 (1834); Zetterstedt, Fauna Ins. Lappon. p. 560 (1838); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. 1840, p. 20, pl. 1, f. 19 : Isis, Vol. 7, p. 547 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 372 (1842); Boitard, Man. Ent. Vol. 3, p. 317 (1843); Curtis, Jour. Agric. Soc. Lond. Vol. 6, p. 2 (1845); Walker,

List Dipt. Brit. Mus. Vol. 3, p. 492 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3023 (1849); L. Dufour, Mém. Acad. Sc. Paris, Vol. 11, pl. 5, f. 61 (1850); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19); p. 52 (1851); Walker, Ins. Brit. Vol. 1, p. 90, pl. 3, f. 1 (1851); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. Ins. Vol. 1, p. 165 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 104 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 399 (1866); Menzbier, Bull. Soc. Nat. Moscou, Vol. 55, p. 1, pl. 3, f. 7, 13 (1880); Becher, Denkschr. Akad. Wiss. Wien, Vol. 45, p. 147, pl. 3, f. 11 (1882); Leunis, Synops. Zool. Vol. 2, p. 402 (1886); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 63 (1892); Elliot, Trans. Ent. Soc. Lond. 1896, p. 117, 128 (1896); Osten-Sacken, Berl. Ent. Zeitsch., Vol. 41, p. 280 (1896); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906); Hamm, Ent. M. Mag. London, Vol. 44, p. 181, 184 (1908); Lundbeck, Dipt. Dan. Vol. 3, p. 133 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 44 (1913).

? *constans*, Harris, Engl. Ins. p. 150, pl. 44, f. 2 (1782).

fugeo, Harris, ibidem, p. 150, pl. 44, f. 2 (1782).

lineata, Fabricius, Syst. Antl. p. 141 (1805).

melanopa, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 263 (1829), no description.

reticulata, Fourcroy, Ent. Paris, Vol. 2, p. 465 [1785] (*Asilus*).

238. *E. lobalis*, Thomson, Eugen. Resa Ins. p. 472 (1868); Beazzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909). Patagonia.

239. *E. loewiana*, Beazzi, Deutsche Ent. Zeitschr. Beiheft, 1909, p. 90, 91 C. Europe. [1909] (*Xanthempis*).

dimidiata, Loew (not Meigen), Berl. Ent. Zeitschr. Vol. 11, p. 12, 19 (1867);

Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 59 (1892);

Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906).

parvula, Strobl (not Egger), Progr. Seitenst. Vol. 14, p. 58 (1880).

240. *E. longicornis*, Macquart, Mém. Soc. Sc. Lille, 1823, p. 159 (1823). France.

241. *E. longimana*, Loew, Beschr. Eur. Dipt. Vol. 2, p. 238 (1871); Kuntze, Siberia. Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906).

242. *E. longipennis*, Loew, Berl. Ent. Zeitschr. Vol. 12, p. 239, 240 (1868); W. Siberia. Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 157 (1907).

243. *E. longipes*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 76 : Cent. 5, No. 51 E. United States. (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 398 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 298, 353 (1902).

244. *E. loripedis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 400 (1895); C. United States. Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 316, f. 131 (1902).

245. *E. lucida*, Zetterstedt, Fauna Ins. Lappon. p. 561 (1838); Dipt. Scand. Europe; Siberia. Vol. 1, p. 373 (1842); ibidem, Vol. 7, p. 3024 (1849); Walker, Ins. Brit. Vol. 1, p. 91 (1851); Bonsdorff, Finl. Ins. Vol. 1, p. 165 (1861); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 153 (1864); Loew, Berl. Ent. Zeitschr., Vol. 12, p. 238, 240 (1868); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 122 (1887); Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 28 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 156 (1907); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 45 (1913).

246. *E. lucidilabris*, Beazzi, Ann. Mus. Nat. Hungar. 1905, p. 440 (1905); Peru. Nova Acta Akad. Naturf. Halle, Vol. 91, p. 344, 349 (1909).

247. *E. lugens*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 760 [1865] Chile. (*Hilara*); Beazzi, Ann. Mus. Hungar. 1905, p. 444 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909).

248. *E. lugubris*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 75 (1869); Kuntze, Rhodes.
Zeitschr. Hym. Dipt. Vol. 6, p. 297, 299 (1906).
249. *E. lutea*, Meigen. Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 228 (1804); C. & S. Europe.
Syst. Beschr. Vol. 3, p. 37 (1822); Curtis, Brit. Ent. Vol. 8, p. 18 (1824); Macquart, Dipt. N. France, Vol. 3, p. 118 (1827); Hist. Nat. Dipt. Vol. 1, p. 330 (1834); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. 1840, 20 : Isis, Vol. 7, p. 547 (1840); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Gimmerthal, Bull. Soc. Nat. Mosc. Vol. 20, 2, p. 163 (1847); Walker, List, Vol. 3, p. 497 (1849); Ins. Brit. Vol. 1, p. 96 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 17, 21 (1867); Verrall, Ent. Mag. Lond. Vol. 8, p. 283 (1872); Strobl, Mitteil. Nat. Ver. Steiermark, Graz. Vol. 29, p. 60 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 91 [1909] (*Xanthempis*); Verrall, Brit. Flies, Vol. 5, p. 9, f. 32 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 95, f. 32 [1910] (*Xanthempis*).
certa, Harris, Engl. Ins. p. 150, pl. 44, f. 6 (1782).
ochracea, Curtis, Brit. Ent. Vol. 8, p. 18 (1834).
- var. *ministriata*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 125 (1887); Strobl, Mitteil. Alps.
Nat. Ver. Steiermark, Graz, Vol. 29, p. 60 (1893); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 92 [1909] (*Xanthempis*).
250. *E. * macilenta*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 93, 121, pl. 11, Baltic Amber.
f. 2, 3 [1908] (*Hilara*).
251. *E. macra*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 24 (1867); Kuntze, Greece.
Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906).
252. *E. macropalpa*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 344 (1860); S. Europe.
Loew, Berl. Ent. Zeitschr. Vol. 11, p. 10 (1867); Kuntze,
Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906).
253. *E. * macrophtalma*, Förster, Abh. Geol. Spezialk. Elsass, Vol. 3, p. 482, Elsass, Middle
pl. 14, f. 29 (1891). Oligocene.
254. *E. macropus*, Loew, Ofv. Vet. Akad. Förh. Stockholm, Vol. 14, p. 369 (1857); Dipterens. Südafr. p. 265, pl. 2, f. 49 (1860). Caffraria.
255. *E. macrorrhyncha*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 757. Chile.
pl. 28, f. 47 (1865); Bezzi, Nova Acta Akad. Wiss. Halle, Vol. 91, p. 344 (1909).
256. *E. macrura*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 131 (1889); Argentina.
Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909).
257. *E. maculata*, Fabricius, Spec. Ins. Vol. 2, p. 472 (1781); C. & S. Europe.
Man. Ins. Vol. 2, p. 365 (1787); Gmelin, Syst. Nat. Vol. 5, p. 2890 (1788);
Olivier, Encycl. Méth. Vol. 6, p. 389 (1791); Fabricius, Ent. Syst. Vol. 4, p. 406 (1794); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 233 (1804); Fabricius, Syst. Antl. p. 141 (1805); Meigen, Syst. Beschr. Vol. 3, p. 33 (1822); Schiner, Verh. Zool.-bot. Ver. Wien, Vol. 6, p. 420 (1856); Loew, Wien. Ent. Monatschr. Vol. 6, p. 167 (1862); Silliman's Journ. Ent. Vol. 37, p. 323 (1864); Berl. Ent. Zeitschr. Vol. 9, p. 238 (1865); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906).
argyreata, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 341 (1860); Schiner,
Fauna Dipt. Austr. Vol. 1, p. 103 (1862); Becker, Denkschr. Akad. Wiss. Wien, Vol. 45, p. 147, pl. 3, f. 11, d, e (1882); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 62 (1892).
? *lineata*, Scopoli, Ent. Carn. p. 364 [1763] (*Asilus*).
- var. *affinis*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 339 (1860); Loew, Berl. Ent. Zeitschr. Vol. 9, p. 238 (1865). Italy.

- var. confusa*, Loew, *ibidem*, Vol. 9, p. 238 (1865); Mik, Beck. Fauna Bernstein, C. & S. Europe.
 Vol. 2 (2), p. 57 (1885); Strobl, Wien. Ent. Zeit. Vol. 18, p. 13 (1899);
 Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 64 (1909).
- maculata*, Macquart-Schiner, Dipt. N. France, Vol. 3, p. 120 (1827); Hist.
 Nat. Dipt. Vol. 1, p. 329 (1834); Schiner, Fauna Dipt. Austr. Vol. 1,
 p. 103 (1862).
258. *E. maculipes*, Zetterstedt, Dipt. Scand. Vol. 1, p. 374 (1842); *ibidem*, N. Europe.
 Vol. 8, p. 3024 (1849); *ibidem*, Vol. 11, p. 4271 (1852); Bonsdorff,
 Finl. Ins. Vol. 1, p. 165 (1861); Lundbeck, Dipt. Dan. Vol. 3, p. 101,
 note (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910); Frey,
Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 44 (1913).
- grisea*, Fallen, part. Empid. Suec. p. 22, part (1815).
- incompta*, Wahlberg, Zetterstedt, Dipt. Scand. Vol. 1, p. 374 (1842).
259. *E. mærens*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 34, 53 (1867); Strobl, Alps, Austria.
 Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 66 (1892); Kuntze,
Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906).
260. *E. *mala*, Meunier, Ann. Sc. Nat. (Zool.), Vol. 7, p. 93, 123, pl. 11, Baltic Amber.
 fig. 11, pl. 12, f. 1 (1908).
261. *E. *malefica*, Meunier, Ann. Sc. Nat. (Zool.), Vol. 7, p. 93, 122, pl. 11, Baltic Amber.
 fig. 7, 8 (1908).
262. *E. malleola*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 120 (1887); Strobl, Alps.
 Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 76 (1893); Kuntze,
Zeitschr. Hym. Dipt. Vol. 7, p. 30 (1907).
263. *E. manca*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 406 (1895); California.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 320 (1902); Coquillett, Proc. Wash. Ent. Soc. Vol. 5, p. 265 (1903).
264. *E. marginata*, Brunetti, Rec. Indian Mus. Vol. 13, p. 79 [1917] Simla District, India.
 (*Pachymeria*) Fauna Brit. India, Dipt. Vol. 1, p. 356 (1920).
265. *E. maura*, Macquart, Dipt. Exot. Vol. 1, Pt. 2, p. 160 (1838); Explor. Algér. Zool. Vol. 3, p. 443 (1849); Kuntze, *Zeitschr. Hym. Dipt.* Vol. 7, p. 32 (1907).
266. *E. mediocris*, Becker, *Zeitschr. Hym. Dipt.* Vol. 7, p. 228 (1907). Algeria.
267. *E. mediterranea*, Loew, Wien. Ent. Monatschr. Vol. 8, p. 364 [1864] Greece, Asia Minor.
 (*Pachymeria*); Kuntze, *Zeitschr. Hym. Dipt.* Vol. 7, p. 158 (1907);
 Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98 [1909] (*Pachymeria*).
268. *E. melæna*, Bezzi, Ann. Mus. Hungar. Vol. 6, p. 394 (1908); Deutsche C. & S. Europe.
Zeitschr. Beiheft, p. 101 [1909] (*Coptophlebia*).
269. *E. melanotricha*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 39 (1873); Strobl, C. & S. Europe.
 Glasnick Zem. Mus. Bosn. Herceg. Vol. 10, p. 410 (1898); Mitteil.
 Bosn. Herceg. Vol. 7, p. 570 (1900); Kuntze, *Zeitschr. Hym. Dipt.* Vol. 7, p. 27 (1907).
270. *E. *melia*, Heyden, Paleontologica, Vol. 17, p. 259, pl. 45, f. 27 (1870); Ratt, Upper Oligocene
 Scudder, Zittel's Handb. Paleont. Vol. 1, p. 807, f. 1074 (1885);
 Handlirsch, Foss. Ins. p. 1017 (1908) Genus?
271. *E. meridionalis*, Meigen, Syst. Beschr. Vol. 3, p. 33 (1822); Schiner, C. Europe.
 Fauna Dipt. Austr. Vol. 1, p. 103 (1862); Strobl, Progr. Seitenst.
 Vol. 14, p. 9 (1880); Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29,
 p. 62 (1892); Kieffer, Illustr. Zeitschr. Ent. Berlin, Vol. 5,
 p. 131-133 (1900); Kuntze, *Zeitschr. Hym. Dipt.* Vol. 6, p. 212
 (1906).
272. *E. metapleuralis*, Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 92 [1909] Sarepta.
 (*Enoplempis*).
273. *E. micans*, Schiner, Novara Reise, Dipt. p. 204 (1868); Bezzi, Nova Acta Akad. Nat. Halle, Vol. 91, p. 343 (1909). South America.

274. *E. micropyga*, Bezzii, Ann. Mus. Hungar. Vol. 3, p. 441 (1905); Nova Peru. Acta Akad. Nat. Halle, Vol. 91, p. 347 (1909).
275. *E. mikii*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 15 (1898); Kuntze, Spain. Zeitschr. Hym. Dipt. Vol. 6, p. 300 (1906).
276. *E. *miocenica*, Cockerell, Proc. Acad. Nat. Sc. Philad. 1914, p. 645 (1914). Florissant, Miocene.
277. *E. mira*, Bigot, Bull. Soc. Ent. France (5), Vol. 10, p. 47 [1880] California. (*Enoplempis*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 319 (1902).
278. *E. mixopolia*, Melander, ibidem, Vol. 28, p. 327 (1902). W. United States.
279. *E. modesta*, Meigen, Syst. Beschr. Vol. 7, p. 86 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Neuhaus, Dipt. March. p. 72 (1886); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 31 (1907). C. Europe.
280. *E. montezuma*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, Mexico. p. 369 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 290, f. 114 (1902); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909).
281. *E. monticola*, Loew, Berl. Ent. Zeitschr. Vol. 12, p. 233, 240 (1868); Alps. Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 156 (1907); Bezzii, Deutsche Ent. Zeitschr. Beiheft, 1909, p. 94 [1909] (*Anacrostichus*).
282. *E. *mordax*, Meunier, Ann. Sc. Nat. [Zool.] (9), Vol. 6, p. 94, 123, pl. 12, f. 2 (1908). Baltic Amber, Lower Oligocene.
283. *E. morenae*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 15 (1899); Kuntze, Spain. Zeitschr. Hym. Dipt. Vol. 6, p. 216 [1906] (*morena*).
284. *E. morio*, Fabricius, Ent. Syst. Vol. 4, p. 405 (1794); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 232 (1804); Fabricius, Syst. Antl. p. 141 (1805); Wiedemann, Aussereurop. Zweifl. Ins. Pt. 2, p. 2 (1830); Macquart, Dipt. Exot. Vol. 1, p. 2, 159 (1839); Explor. Algér. Zool. Vol. 3, p. 443 (1849); Loew, Wien. Ent. Monatschr. Vol. 8, p. 257 (1864); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 124 (1907); Bezzii, Revista S. Fiel, Vol. 8, p. 53 (1909). N. Africa; S. Europe.
- castellana*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 13 (1899).
- hispanica*, Loew, Rosenhauer : Thiere Andalus. p. 385 (1856); Wien. Ent. Monatschr. Vol. 8, p. 256 (1864); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906); Becker, Wien. Ent. Zeit. Vol. 30, p. 72 (1911); Villeneuve, Wien. Ent. Zeit. Vol. 31, p. 97 (1912).
285. *E. morosa*, Meigen, Syst. Beschr. Vol. 3, p. 26, ♂, not ♀ (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 387 (1842); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 57 (1867); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 122 (1887); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 65, note (1892). C. Europe.
286. *E. *morosella*, new name. Baltic Amber, Lower Oligocene.
- morosa*, Meunier (not Meigen or Macquart), Ann. Sc. Nat. (Zool.) (9), Vol. 7, p. 94, 125, pl. 12, f. 5, 6 (1908).
287. *E. ?mucronata*, Scopoli, Ent. Carn. p. 363 [1763] (*Asilus*); Olivier, Encycl. Méth. Vol. 6, p. 390 (1791); Schiner, Verh. Zool.-bot. Ges. Wien, Vol. 6, p. 420 [1856] (?genus).
288. *E. nepticula*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 259 (1869); Strobl, Wien. Ent. Zeit. Vol. 18, p. 14 (1899); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 567 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906). S. Europe.
289. *E. nigerrima*, Loew, Wien. Ent. Monatschr. Vol. 6, p. 171 (1862); Berl. Ent. Zeitschr. Vol. 11, p. 2, 7 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 297 (1906). S. Europe.
290. *E. nigra*, Villers, Ent. Linn. Vol. 3, p. 571 (1789); Olivier, Encycl. Europe. Méth. Vol. 6, p. 389 (1791).

- . *E. nigrescens*, Schummel, Veränd. Schles. Ges. Cult. Breslau, 1832, p. 70 (1832), no description.
291. *E. nigricans*, Macquart, Mém. Soc. Sc. Lille, 1823, p. 161 (1823). France.
292. *E. nigricans*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 224 (1804); Syst. Beschr. Vol. 3, p. 34 (1822); Macquart, Dipt. N. France, Vol. 3, p. 120 (1827); Hist. Nat. Dipt. Vol. 1, p. 328 (1834); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906); Lundbeck, Dipt. Dan. Vol. 3, p. 135, f. 44 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910).
- rustica*, Fallen, Empid. Suec. p. 18 (1816); Zetterstedt, Dipt. Scand. Vol. 1, p. 371 (1842); ibidem, Vol. 12, p. 4608 (1855); Schiner, Fauna Dipt. Austr. Vol. 1, p. 104 (1862); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 62 (1892).
293. *E. nigricomata*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 6, 8 (1867); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 64 (1892); ibidem, Vol. 34, p. 203 (1897); Strobl, Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 567 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 298 (1906); Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 308 (1906); Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 64 (1909).
294. *E. nigricrus*, Gmelin, Syst. Nat. Vol. 5, p. 2891 (1790); Olivier, Encycl. Méthod. Vol. 6, p. 391 (1791). Europe.
295. *E. nigrimana*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 225 (1907). Algeria.
296. *E. nigritarsis*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 221, (1804); Syst. Beschr. Vol. 3, p. 32, 31 (1822); Curtis, Brit. Ent. Vol. 8, p. 18, 3 (1824); Meigen, Syst. Beschr. Vol. 6, p. 337 (1830); Macquart, Hist. Nat. Dipt. Vol. 1, p. 329 (1834); Boitard, Man. Ent. Vol. 3, p. 317 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 496 (1849); Ins. Brit. Vol. 1, p. 95 (1851); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 59 (1867); Strobl, Glasnick Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 405 (1898); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 125 (1899); Strobl, Bos. Mitteil. Herceg. Sarajevo, Vol. 7, p. 566 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 99 [1909] (*Lissembris*); Lundbeck, Dipt. Dan. Vol. 3, p. 101, f. 35 [1910] (*Lissembris*).
- nitidicollis*, Curtis, Brit. Ent. Vol. 8, p. 18 (1824); Schiner, Fauna Dipt. Austr. Vol. 1, p. 109 (1862).
297. *E. nigritibialis*, Strobl, Glasnik Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 410 (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 570 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 31 (1907).
298. *E. nitida*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 223 (1804); Syst. Beschr. Vol. 3, p. 39 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 334 [1834] (*Pachymerina*); Meigen, Syst. Beschr. Vol. 7, p. 89 [1838] (*Pachymeria*); Boitard, Man. Ent. Vol. 3, p. 319 [1843] (*Pachymerina*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3027 [1849] (*Pachymeria*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 54 [1851] (*Pachymeria*); Bonsdorff, Finl. Ins. Vol. 1, p. 167 [1861] (*Pachymeria*); Loew, Berl. Ent. Zeitschr. Vol. 12, p. 232, 240, 1 (1868); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 86 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 156 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 94 [1909] (*Anacrostichus*); Lundbeck, Dipt. Dan. Vol. 3, p. 96, f. 33 [1910] (*Anacrostichus*); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 40 [1913] (*Anacrostichus*).

299. *E. nitidissima*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 82 C. Europe.
 (1892); ibidem, Vol. 34, p. 205 (1897); Kuntze, Zeitschr. Hym.
 Dipt. Vol. 7, p. 26 (1907).
300. *E. nitidiventris*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 38 (1873); Kuntze, C. Europe.
 Zeitschr. Hym. Dipt. Vol. 7, p. 30 (1907).
301. *E. nitidula*, Zetterstedt, Dipt. Scand. Vol. 13, p. 5008 (1859); Kuntze, N. Europe.
 Zeitschr. Hym. Dipt. Vol. 7, p. 29 (1907); Wahlgren, Ent. Tidskr.
 Vol. 31, p. 63 (1910).
302. *E. nodipes*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 324, pl. 9. New Mexico.
 f. 143 (1902).
303. *E. nuda*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 195 : Cent. 2, No. 20 Illinois.
 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 299 (1895);
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 305, f. 126 (1902).
304. *E. nuntia*, Meigen, Syst. Beschr. Vol. 7, p. 85, 62 (1838); Schiner, Fauna C. Europe.
 Dipt. Austr. Vol. 1, p. 108 (1862); Becker, Berl. Ent. Zeitschr. Vol.
 31, p. 123 (1887); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 31 (1907).
305. *E. obesa*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 321 : Cent. 1, No. 28 E. United States.
 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 400 (1895);
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 309, 353 (1902).
ravida, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 403 (1895); Melander,
 Trans. Amer. Ent. Soc. Vol. 28, p. 310 (1902); Bezzii, Deutsche Ent.
 Zeitschr. Beiheft, p. 88 (1909).
306. *E. obscuripes*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 218 [1873] (*Pachymeria*); Asia Minor.
 Kuntze, Zeit. Hym. Dipt. Vol. 7, p. 157 (1907); Bezzii, Deutsche Ent.
 Zeitschr. Beiheft, p. 98 [1909] (*Pachymeria*).
307. *E. ochropus*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 755 (1865); Chile.
 Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909).
308. *E. ollioides*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 493 (1849); Coquillett, Nova Scotia.
 Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895); Melander, Trans.
 Amer. Ent. Soc. Vol. 28, p. 299 (1902).
309. *E. opaca*, Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 226 (1804); C. & S. Europe.
 Fabricius, Syst. Antl. p. 138 (1805); Meigen, Syst. Beschr
 Vol. 3, p. 17, pl. 22, f. 17 (1822); Macquart, Mem. Soc. Sc. Lille,
 p. 159 (1823); Curtis, Brit. Ent. Vol. 8, p. 18, 2 (1824); Macquart,
 Dipt. N. France, p. 119 (1827); Hist. Nat. Dipt. Vol. 1, p. 328, 5
 (1834); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. 1840,
 p. 20, pl. 1, f. 18; Isis, Vol. 7, p. 546 (1840); Westwood, Introd.
 Classif. Ins. Vol. 2, p. 547 (1840); Boitard, Man. Ent. Vol. 3,
 p. 317 (1843); Walker, List Dipt. Brist. Mus. Vol. 3, p. 492 (1849);
 Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), 52 (1851); Walker, Ins.
 Brit. Vol. 1, p. 91, 4 (1851); Zetterstedt, Dipt. Scand. Vol. 12,
 p. 4608, note (1855); Schiner, Fauna Dipt. Austr. Vol. 1, p. 105
 (1862); Glover, Manuser. Notes, p. 21, pl. 11, f. 7 (1874); Neuhaus,
 Dipt. March. p. 72 (1886); Bezzii, Bull. Soc. Ent. Ital. Vol. 30,
 p. 129 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906);
 Hamm, Ent. Mag. London, Vol. 45, p. 132-134, 157 (1909); Lund-
 beck, Dipt. Danica, Vol. 3, p. 139 (1910).
rufipes, Fabricius, Syst. Antl. p. 138 (1805).
 var. *apteropus*, Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 130 (1899); Kuntze, Italy.
 Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906).
310. *E. otakouensis*, Miller, Trans. New Zeal. Inst. Vol. 42, p. 235, pl. 30, New Zealand.
 f. 1-6 (1909).
311. *E. otiosa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 407 (1895); E. United States.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 302, f. 122 (1902);
 Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 88 (1909).

312. *E. pachymera*, Macquart, Dipt. Exot. Vol. 1, p. 277 (1838); Blanchard, Chile.
 Gay, Hist. Chile, Zool. Vol. 7, p. 373 (1852); Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 753 [1865] (not *Pachymeria*); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909).
313. *E. pachymerina*, Schiner, Novara Reise Dipt. p. 205 (1868); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 344 (1909). South America.
314. *E. pachypodiata*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 132 (1889); Australia.
 Bezzi, Ann. Mus. Hungar. Vol. 2, p. 343, note (1904).
315. *E. pachystoma*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 757 (1865); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909). Chile.
316. *E. pallida*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 322 : Cent 1, No. 30 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 301 (1902). New York.
317. *E. pallipes*, Olivier, Encycl. Méthod. Vol. 6, p. 389 (1791). W. Europe.
318. *E. palparis*, Egger, Verh. Zool.-bot. Wien, Vol. 10, p. 345 [1860] (*Pachymeria*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 110 [1862] (*Pachymeria*); Loew, Wien. Ent. Monatschr. Vol. 8, p. 360 [1864] (*Pachymeria*); Verrall, Ent. Mag. London, Vol. 30 p. 141 [1894] (*Pachymeria*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 85 (1895); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 158 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98 [1909] (*Pachymeria*).
femorata, Walker (not Fabricius), List Dipt. Brit. Mus. Vol. 3, p. 498 (1849); Ins. Brit. Vol. 1, p. 96 (1851).
? scotica, Curtis, Brit. Ent. Vol. 8, p. 18, 4 [1824] (*Pachymeria*).
319. *E. ? palustris*, Scopoli, Ent. Carn. p. 369 (1763); Schiner, Verh. Zool.-bot. Ver. Wien, Vol. 6, p. 421 (1856) ? gen. C. Europe.
320. *E. papuana*, Bezzii, Ann. Mus. Hungar. Vol. 2, p. 344 (1904); ibidem, New Guinea. Vol. 10, p. 468 [1912] (*Coptophlebia*).
321. *E. parvula*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 343 (1860); C. Europe.
 Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 18, 21, 11 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 91 [1909] (*Xanthempis*).
322. *E. patagiata*, Bezzii, Suppl. Ent. Berlin, Vol. 3, p. 71 [1914] (*Coptophlebia*). Formosa.
323. *E. Pavesii*, Bezzii, Bull. Soc. Ent. Ital. Vol. 27, p. 50 (1895); ibidem, Italy. Vol. 30, p. 135 (1899) note; Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 303 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 102 [1909] (*Coptophlebia*).
324. *E. pegasus*, Osten-Sacken, Biol. Centr. Amer. Dipt. Vol. 1, p. 216 (1887); Panama.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 292 (1902); Bezzii, Nova Acta Akad. Nat. Halle, Vol. 91, p. 344 (1909).
325. *E. pellucida*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 408 (1900); Alaska.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 313 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903).
326. *E. pennaria*, Fallen, Empid. Suec. p. 20, part (1815); Meigen, Syst. Beschr. Vol. 3, p. 23 (1822); Curtis, Brit. Ins. Vol. 8, p. 18, 4 (1824); Macquart, Dipt. N. France, Vol. 3, p. 125 (1827); Hist. Nat. Dipt. Vol. 1, p. 331, 19 (1834); Zetterstedt, Fauna Ins. Lappon. p. 562 (1838); Dipt. Scand. Vol. 1, p. 383 (1842); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 495 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3025 (1849); Walker, Ins. Brit. Vol. 1, p. 93 (1851); Bonsdorff, Finl. Ins. Vol. 1, p. 166 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 106 (1862);

- Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 79 (1893); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 570 (1900); Jahrb. Landesmus. Kärnten, Klagenfurt, Vol. 47, p. 200 (1901); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 31 (1907).
var. *baldensis*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 20 (1899); Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 66 (1909).
var. *flaviventris*, Strobl, Glasnik Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 410 (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 570 (1900).
327. *E. pennipes*, Linnæus, Syst. Nat. Vol. 10, p. 604 (1758); Fauna Suec. p. 466 (1761); Scopoli, Ent. Carn. p. 365 [1763] (*Asilus*); Linnæus, Syst. Nat. Vol. 12, p. 2, 1003 (1767); Sulzer, Ins. pl. 21, f. 137 (1776); Schaeffer, Icon. Ins. pl. 192, f. 3 (1779); Fabricius, Spec. Ins. Vol. 2, p. 471 (1781); Mant. Ins. Vol. 2, p. 364 (1787); Gmelin, Syst. Nat. Vol. 5, p. 2889 (1788); Fabricius, Ent. Syst. Vol. 4, p. 404 (1794); Panzer, Fauna Germ. Vol. 74, p. 18 (1801); Schellenberg, Gehr. Mouches, p. 88, pl. 35, f. 2 (1803); Fabricius, Syst. Antl. p. 138 (1805); Latreille, Gen. Crust. Ins. Vol. 4, p. 303 (1809); Consid. Gén. 443 (1810); Lamarck, Anim. s. Vert. Vol. 3, p. 401 (1816); Billberg, Enumer. Ins. p. 120 (1820); Wood, Illiger, Linn. Gen. Ins. Vol. 2, p. 100, pl. 68 (1821); Meigen, Syst. Beschr. Vol. 3, p. 21 (1822); Macquart, Dipt. N. France, Vol. 3, p. 123 (1827); Boitard, Man. Ent. Vol. 2, p. 369 (1828); Macquart, Hist. Nat. Dipt. Vol. 1, p. 331 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 380 (1842); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 494 (1849); Ins. Brit. Vol. 1, p. 92, 11 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 5008 (1859); Bonsdorff, Finl. Ins. Vol. 1, p. 166, 9 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Leunis, Synop. Zool. Vol. 2, p. 402 (1886); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 74 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 26 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 109, f. 38 [1910] (*Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910); Frey, Acta Soc. Sc. Helsingfors, Vol. 37 (3), p. 42 [1913] (*Pterempis*). — Pl. 5, Figs. 48, 49.
ciliata, Meigen (not Fabricius), Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 220 (1804); Fallen, Empid Suec. p. 20, part. (1815).
longirostris, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 223 (1804).
pennata, Schrank, Ins. Austr. p. 987 (1781); Fauna Boica, Vol. 3, p. 170 (1803).
328. *E. penniventris*, Bezzi, Nova Acta Akad. Nat. Halle, Vol. 91, p. 352 Peru. (1909).
329. *E. *perdita*, Cockerell, Proc. U. S. Nat. Mus. Vol. 51, p. 92, f. (1916). Florissant, Miocene
330. *E. peregrina*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 331, f. 97 [1902] (*Iteaphila*). New Mexico.
331. *E. perpendicularis*, Loew, Ofv. Vet. Akad. Förhandl. Stockholm, Vol. 14, p. 370 (1857); Dipterenf. Südafr. p. 266, 5, pl. 2, f. 50 (1860). Caffraria.
332. *E. *personata*, Meunier, Ann. Sc. Nat. (Zool.) p. 7, 93, 122, pl. 11, f. 9, 10 Baltic Amber. (1908).
333. *E. petulans*, Becker, Deutsche Ent. Zeitschr. p. 641 (1910). Corsica.
334. *E. phænomeris*, Loew, Berl. Ent. Zeitschr. Vol. 12, p. 172, 388, 393 (1868); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 32 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 96 [1909] (*Polyblepharis*).
335. *E. picena*, Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 142 (1898); Kuntze, Italy. Zeitschr. Hym. Dipt. Vol. 7, p. 158 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98 [1909] (*Pachymeria*).

336. *E. picipes*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 226 C. Europe.
 (1804); Syst. Beschr. Vol. 3, p. 19 (1822); Zetterstedt, Dip. Scand. Vol. 1, p. 347 (1842); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Loew, Bericht Naturf. Ver. Augsburg, Vol. 20, p. 45 (1869); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906).
sulcipes, Meigen, Syst. Beschr. Vol. 3, p. 19 (1822); Curtis, Brit. Ent. Vol. 8, p. 18 (1824); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Loew, Bericht Naturf. Ver. Augsburg, Vol. 20, p. 45 (1869).
337. *E. pilicornis*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 3, 8 (1867); Kuntze, Spain.
 Zeitschr. Hym. Dipt. Vol. 6, p. 298 (1906).
338. *E. pilimana*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 86 (1869); Strobl, C. Europe.
 Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 68, note (1892); ibidem, Vol. 34, p. 203 (1897); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 303 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 102 [1909] (*Coptophlebia*); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 46, p. 65 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 64 (1910).
339. *E. pilitibia*, Macquart, Dipt. Exot. 6, Suppl. p. 87, 9 (1854). Cape of Good Hope.
340. *E. pilosa*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 9 (1867); Kuntze, C. Europe.
 Zeitschr. Hym. Dipt. Vol. 6, 299 (1906).
341. *E. pittoptera*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 229 (1873); Kuntze, Siberia.
 Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, 91, note (1909).
342. *E. platyptera*, Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 46, Austria.
 p. 64 (1909).
343. *E. plebeja*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 38 (1873); Kuntze, C. Europe.
 Zeitschr. Hym. Dipt. Vol. 7, p. 27 (1907).
344. *E. plorans*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 468 & 470 [1912] Formosa.
 (*Coptophlebia*).
345. *E. plumipes*, Zetterstedt, Dipt. Scand. Vol. 1, p. 382 (1842); Scholz, C. & N. Europe.
 Zeitschr. Ent. Breslau, Vol. 5 (19), p. 53 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 4271 (1852); Schiner, Fauna Dipt. Austr. Vol. 1, p. 109 (1862); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 77 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 129 (1907); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 42 [1913] (*Pterempis*).
ciliata, Fallen, part, Empid. Suec. p. 20, var. (1815).
346. *E. podagra*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 318 (1902). Idaho.
347. *E. podagraria*, Meigen, Syst. Beschr. Vol. 6, p. 338 (1830); Kuntze, C. Europe.
 Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906).
348. *E. poecila*, Philippi, Verh. Zool-bot. Ges. Wien, Vol. 15, p. 753 (1865); Chile.
 Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 344 & 350 (1909).
349. *E. poeciloptera*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 322 : Cent. 1, No. 31 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 298 (1902). New York.
350. *E. *Poepigia*, Giebel, Ins. Vorw. p. 207 (1856); Handlirsch, Foss. Ins. p. 1017, genus? (1908). Baltic Amber, Lower Oligocene.
351. *E. polita*, Macquart, Dipt. Exot. Vol. 1, p. 278 (1838); Blanchard, in. Gay Hist. Chile. Zool. Vol. 7, p. 373 (1852); Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 753 (1865); Bezzi, Nova Acta Akad. Naturf. Vol. 91, p. 355 (1909). Chile.
? collina, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 756 (1865); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909).
352. *E. poplitea*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 16 : Cent. 3, No. 29 (1863); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 399 (1895); W. North America.

- Proc. Wash. Acad. Sc. Vol. 2, p. 407 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 321, 353 (1902).
- serperastrorum*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 324, f. 136, 145 (1902).
353. *E. praecox*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 50, 56 (1867); Kuntze, Rhodes. Zeitschr. Hym. Dipt. Vol. 7, p. 25 (1907).
354. *E. præputiata*, Loew, Bes. Eur. Dipt. Vol. 3, p. 225 (1873); Kuntze, S. Russia. Zeitschr. Hym. Dipt. Vol. 6, p. 300 (1906).
355. *E. procera*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 37 (1873); Bezzi, Bull. C. & S. Europe. Soc. Ent. Ital. Vol. 30, p. 137 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 27 (1907).
356. *E. prodromus*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 42, 54, 13 (1867); C. & N. Europe. Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 70 (1892); Verrall, Ent. Mag. London, Vol. 30, p. 140 (1894); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 34, p. 203 (1897); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 304 (1906); Lundbeck, Dipt. Dan. Vol. 3, p. 120 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 64 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 43 [1913] (*Pterempsis*).
357. *E. pruinosa*, Wiedemann, Anal. Ent. p. 28 (1824); Aussereurop. Zweifl. Ins. Vol. 2, p. 4 (1830). Cape of Good Hope.
358. *E. pseudodecora*, Strobl, Glasnik Bosn. Herceg. Sarajevo, Vol. 10, p. 409 (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 569 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 27 (1907). Bosnia.
359. *E. pseudomalleola*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, C. Europe. p. 75 (1893); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 31 (1907).
360. *E. pteropoda*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 343 (1860); S. Europe. Schiner, Fauna Dipt. Austr. Vol. 1, p. 105 (1862); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906).
361. *E. ptilocnemis*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 222 [1873] (*Pachymeria*); Transcaspia. Zeitschr. Ges. Naturw. Berlin, Vol. 43, p. 417 [1874] (*Pachymeria*); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 158 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98 (1909) note (*Pachymeria*).
362. *E. ptilopoda*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 22 (1822); S. W. Europe. Brullé, Expéd. Morée, Zool. Vol. 3, 1, p. 299 (1832); Strobl, Wien. Ent. Zeit. Vol. 18, p. 18 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 31 (1907).
363. *E. pudica*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 324: Cent. 2, No. 35 N. America. [1861] (*Pachymeria*); Wien. Ent. Monatschr. Vol. 8, p. 364 [1864] (*Pachymeria*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 330 [1902] (*Pachymeria*). *tumida*, Loew (not Meigen), Wien. Ent. Monatschr. Vol. 8, p. 365 [1864] (*Pachymeria*).
364. *E. pulchripes*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 258 (1869); Kuntze, Greece. Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906).
365. *E. pulicaria*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 41, 54 (1867); C. Europe. Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 302 (1906).
366. *E. *pulvillata*, Loew, Bernsteinfauna, p. 41 (1850); Giebel, Ins. Vorwelt, Baltic Amber, p. 208 (1856); Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 94, 124 Lower Oligocene. (1908).
367. *E. punctata*, Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 227 C. & N. Europe. (1804); Syst. Beschr. Vol. 3, p. 38 (1822); Curtis, Brit. Ent. Vol. 8, p. 18, 3 (1824); Macquart, Dipt. N. France, Vol. 3, p. 118 (1827);

- Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 20 (1840); Isis, Vol. 7, p. 547 (1840); Berl. Ent. Zeitschr. Vol. 11, p. 15, 20 (1867); Verrall, Ent. Mag. London, Vol. 8, p. 282 (1872); Neuhaus, Dipt. March. p. 72 (1886); Kuntze, Zeitschr. Hym. Dipt. Vol. 6 p. 211 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft. p. 90 [1909] (*Xanthempis*); Hamm, Ent. Mag. London, Vol. 45, p. 160 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 88, f. 26 [1910] (*Xanthempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 39 [1913] (*Xanthempis*).
dorsalis, Curtis, Brit. Ent. Vol. 8, p. 18 (1824).
ignota, Meigen, Syst. Beschr. Vol. 6, p. 338 (1830); Curtis, Brit. Ent. Vol. 8, p. 18 (1834); Macquart, Hist. Nat. Dipt. Vol. 1, p. 329 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 380 (1842); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 497 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3025 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 52 (1851); Walker, Ins. Brit. Vol. 1, p. 95 (1851); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. Ins. Vol. 1, p. 166 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 104 (1862); Neuhaus, Dipt. March. p. 72 (1886).
testacea, Curtis (not Fabricius), Brit. Ent. Vol. 8, p. 18, 3 (1834).
368. *E. ? punctata*, Scopoli, Ent. Carn. p. 364 [1763] (*Asilus*); Olivier, Encycl. C. Europe.
Méthod. Vol. 6, p. 390 (1791); Schiner, Verh. Zool.-bot. Ges. Wien, Vol. 6, p. 420 (1856) ? gen.
369. *E. pusio*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 342 (1860); C. & S. Europe.
Schiner, Fauna Dipt. Austr. Vol. 1, p. 106 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 37 & 53 (1867); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 68 (1892); Glasnik Bosn. Herceg. Sarajevo, Vol. 10, p. 406 (1902); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 567 (1904); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 302 (1906).
simplicipes, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 37 & 53 (1867).
370. *E. quadrilineata*, Gmelin, Syst. Nat. Vol. 5, p. 2890 (1790). Europe.
371. *E. quadrivittata*, Arribalzaga, Nat. Hist. Argent. Vol. 1, p. 292 (1878); Argentina.
Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 343 (1909).
372. *E. raptoria*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 467, 468 (1912). Formosa.
373. *E. rava*, Loew, Wien, Ent. Monatschr. Vol. 6, p. 170 (1862); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906). S. Europe.
374. *E. reciproca*, Walker, Trans. Ent. Soc. Lond. n. s. Vol. 4, p. 147 (1857); N. America.
Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 399 (1895).
375. *E. rostrata*, Brunetti, Rec. Indian Mus. Vol. 9, p. 25 (1913); Fauna Brit. W. Himalayas.
India Dipt. Vol. 1, p. 355, f. 31 (1920).
376. *E. rufescens*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 76 : Cent. 5, No. 52 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 398 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 302, f. 120 (1902). E. United States.
377. *E. ruficornis*, Loew, Wien. Ent. Monatschr. Vol. 8, p. 365 [1864] (*Pachymeria*); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 157 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98, note [1909] (*Pachymeria*). Bessarabia.
378. *E. rufipes*, Gmelin, Syst. Nat. Vol. 5, p. 2890 (1790); Olivier, Encycl. Europe.
Méthod. Vol. 6, p. 391 (1791).
379. *E. ? rufipes*, Scopoli, Ent. Carn. p. 368 (1763); Schiner, Verh. Zool.-bot. Ver. Wien, Vol. 6, p. 421 (1856) ? genus. C. Europe.
380. *E. rufipes*, Wiedemann, Ausser. Eur. Zweifl. Ins. Vol. 2, p. 5 (1830); Russia.
Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 216 (1906).
381. *E. rufiventris*, Meigen, Syst. Beschr. Vol. 7, p. 86 (1838); Schiner, Fauna C. & N. Europe.
Dipt. Austr. Vol. 1, p. 108 (1862); Strobl, Mitteil. Naturf. Ver.

- Steiermark, Graz, Vol. 29, p. 77 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 29 (1907); Lundbeck, Dipt. Danica, Vol. 3, p. 111 [1910] (*Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910); Collin, Ent. M. Mag. London (2), Vol. 24 p. 106 (1913).
entralis, Zetterstedt, Dipt. Scand. Vol. 1, p. 381, 15 (1842); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 53 (1851).
382. *E. salicina*, Lioy, Atti Soc. Ital. Sc. Nat. Milano, Vol. 5, p. 380 (1864). Italy.
383. *E. sauteriana*, Bezzii, Suppl. Entom. Berlin, Vol. 2, p. 70 [1914] Formosa.
(*Coptophlebia*).
384. *E. scatophagina*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 351 (1902). Alaska.
385. *E. scaura*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 27, 52 (1867); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 65 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906).
386. *E. scoparia*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 269 (1903). New Hampshire.
387. *E. scopulifera*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 468, 472 [1912] (*Pterempis*). Formosa.
388. *E. scutellariae*, Bezzii, Ann. Mus. Hungar. Vol. 6, p. 393, 394 (1908). Italy.
389. *E. scutellata*, Curtis, Brit. Ent. Vol. 8, p. 183 (1824); Verrall, Ent. Mag. London, Vol. 8, p. 281 (1872); Hamm, Ent. Mag. London, Vol. 45, p. 160 (1909). England.
- testacea*, Walker (not Fabricius), Ins. Brit. Dipt. Vol. 1, p. 96 (1851).
390. *E. semicinerea*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 16, 24 (1867); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 123 (1887); Strobl, Mitteil. Natur. Ver. Steiermark, Graz, Vol. 29, p. 61 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906); Bezzii, Deutsche Ent. Zeitschr. Beiheft, p. 91 [1909] (*Xanthempis*).
391. *E. serena*, Pokorny, Verh. Zool.-bot. Ges. Wien, Vol. 37, p. 393 (1887); Alps. Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 212 (1906); Bezzii, Ann. Mus. Hungar. Vol. 6, p. 394 (1908).
392. *E. sericans*, Brullé, Expéd. Morée (Zool.), Vol. 3, p. 298 (1834). Greece.
393. *E. sericata*, White, Proc. Roy. Soc. Tasmania, 1916, p. 233, f. 44 (1917). Tasmania.
394. *E. serotina*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 48, 55 (1867); Strobl, Jahrb. Mus. Kärnten, Klagenfurt, Vol. 47, p. 199 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 25 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 122 [1910] (*Pterempis*).
? longirostris, Meigen, Syst. Beschr. Vol. 6, p. 338 (1830); Zetterstedt, Dipt. Scand. Vol. 1, p. 388, note (1842); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 59 (1867).
? obscura, Macquart, Dipt. N. France, Vol. 3, p. 126 (1827); Hist. Nat. Dipt. Vol. 1, p. 332 (1834); Meigen, Syst. Beschr. Vol. 7, p. 83 (1838); Zetterstedt, Dipt. Scand. Vol. 8, p. 3026 (1849); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 60 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 32 (1907).
obscurifrons, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 66 (1892); Strobl, ibidem, Vol. 46, p. 65 (1909).
395. *E. serrata*, Schrank, Fauna Boica, Vol. 3, p. 171 (1803). C. Europe.
396. *E. setosa*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 30, 52 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906). Island Chio.
397. *E. ?setosa*, Scopoli, Ent. Carn. p. 365 [1763] (*Asilus*); Olivier, Encycl. Méthod. Vol. 6, p. 390 (1791); Schiner, Verh. Zool.-bot. Ver. Wien, Vol. 6, p. 420 (1856) ? gen. C. Europe.
398. *E. sibillina*, Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 133 (1899); Kuntze, Italy. Zeitschr. Hym. Dipt. Vol. 6, p. 216 (1906).
399. *E. similis*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 45 (1908). Canary Islands.
400. *E. sordida*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 321 : Cent. 1, No. 29 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 397 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 298 (1902). E. United States.

401. *E. spectabilis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 196 : Cent. 2, No. 21 (1862); Glover, Manusc. Notes, p. 21, pl. 3, f. 22 (1874); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 398 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 311, f. 109 (1902); McAtee, Ent. News, Philad., Vol. 20, p. 359 (1909). E. United States.
402. *E. specularis*, Bezzi, Revista S. Fiel. Vol. 8, p. 55 [1909] (*Pachymeria*). Spain.
403. *E. spiloptera*, Wiedemann, Aussereurop. Zweifl. Ins. Vol. 2, p. 5 (1830); Loew, Berl. Ent. Zeitschr., Vol. 8, p. 99 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 398 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 292 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 344 (1909).
- picta*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 16 : Cent. 3, No. 28 (1863).
404. *E. spinifera*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346, 354 (1909). Peru.
405. *E. spirifera*, Bezzi, Revista S. Fiel. Vol. 8, p. 54 (1909). Spain.
406. *E. squamata*, Brunetti, Rec. Indian Mus. Vol. 9, p. 27 (1913); Faun. Brit. Ind. Dip. Vol. 1, p. 352, pl. 4, f. 11 (1920). Ceylon.
407. *E. squamipes*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 271 (1903) Mexico.
408. *E. stenoptera*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 75 : Cent. 5, No. 50 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 399 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 304, 353 (1902). New Hampshire.
409. *E. stercorea*, Linnæus, Fauna Suec. p. 467 (1761); Syst. Nat. Ed. 12, Europe.
 Pt. 2, p. 1004 (1767); Fabricius, Syst. Ent. p. 802 (1775); Spec. Ins. Vol. 2, p. 472 (1781); Schrank, Ins. Austr. p. 483 (1781); Fabricius, Mant. Ins. Vol. 2, p. 365 (1787); Gmelin, Syst. Nat. Vol. 5, p. 2890 (1788); Rossi, Fauna Etrusc. Vol. 2, p. 334 (1790); Olivier, Encycl. Method. Vol. 6, p. 389 (1791); Fabricius, Ent. Syst. Vol. 4, p. 406 (1794); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 228 (1804); Fabricius, Syst. Antl. p. 141 (1805); Rossi, Fauna Etrusc. (2), p. 505 (1807); Fallen, Empid Suec. p. 18 (1815); Wiedemann, Zool. Mag. Berlin, Vol. 1, p. 2 (1817); Billberg, Enumer. Ins. p. 120 (1820); Meigen, Syst. Beschr. Vol. 3, p. 36 (1822); Macquart, Mém. Soc. Sc. Lille, 1823, p. 159 (1823); Curtis, Brit. Ent. Vol. 8, p. 18, 3 (1824); Macquart, Hist. Nat. Dipt. Vol. 1, p. 330 (1834); Zetterstedt, Fauna Ins. Lappon. p. 560 (1838); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. 1840, p. 20 : Isis, Vol. 7, p. 547 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 377, part (1842); Boitard, Man. Ent. Vol. 3, p. 318 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 497 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3024 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 52 (1851); Walker, Ins. Brit. Vol. 1, p. 95 (1851); Zetterstedt, Dipt. Scand. Vol. 12, p. 4610 (1855); Schiner, Verh. Zool.-bot. Ges. Wien, Vol. 6, p. 720 (1856); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. Ins. Vol. 1, p. 166 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 153 (1864); ibidem, Vol. 14, p. 380 (1866); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 12, 19 (1867); Verrall, Ent. Mag. London, Vol. 8, p. 281 (1872); Jaroschewsky, Arb. Ges. Naturf. Univ. Kharkow, Vol. 11, p. 355 (1877); Beling, Arch. Naturg. Berlin, Vol. 48, p. 206, 16 (1882); Brauer, Denkschr. Wiss. Wien, Vol. 44, pl. 2, f. 7 (1882); Leunis, Synops. Zool. Vol. 2, p. 402 (1886); Strobl, Mitteil. Ver. Steiermark, Graz, Vol. 29, p. 59 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi, Deutsche Ent. Zeitschr. p. 90 [1909] (*Xanthempis*); Lundbeck, Dipt. Danica.

- Vol. 3, p. 91, f. 23, 28, 29 [1910] (*Xanthempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 62, f. 1 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), 39 [1913] (*Xanthempis*).
ferruginea, Scopoli, Ent. Carn. p. 364 [1763] (*Asilus*).
pertinax, Harris, Engl. Ins. p. 120, pl. 44, f. 5 (1782).
rapax, Pallas-Wiedemann, Zool. Mag. Berlin (1818), p. 25, no description.
stercoraria, Macquart, Dipt. N. France, Vol. 3, p. 117 (1827).
stigma, Meigen, Syst. Beschr. Vol. 7, p. 88 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862).
410. *E. *sticornis*, Loew, Bernsteina fauna, p. 41 [1850] (*unnamed*); Meunier, Miscell. Ent. Vol. 7, p. 178 (1899). Baltic Amber, Lower Oligocene.
411. *E. strigata*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 159 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906); Bezzi, Ann. Mus. Hungar. Vol. 6, p. 394 (1908). S. Russia.
412. *E. styriaca*, Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 61 (1892); ibidem, Vol. 34, p. 202 (1897); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 91 [1909] (*Xanthempis*). Alps.
413. *E. subciliata*, Loew, Beschr. Eur. Dipt. Vol. 2, p. 240 (1871); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 214 (1906). Siberia.
414. *E. subciliipes*, Brunetti, Rec. Indian Mus. Vol. 9, p. 24 (1913); Fauna Brit. India Dipt. Vol. 1, p. 354 (1920). W. Himalayas.
415. *E. subclavata*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 220 [1873] (*Pachymeria*); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 158 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98, note [1909] (*Pachymeria*). Greece.
416. *E. suberis*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 226 (1907); Deutsche Ent. Zeitschr. p. 641 (1910). Tunis, Corsica.
417. *E. subnitida*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 124 (1914). E. Africa.
418. *E. subpennata*, Macquart, Dipt. N. France, Vol. 3, p. 126 (1827); Hist. Nat. Dipt. Vol. 1, p. 332 (1834); Meigen, Syst. Beschr. Vol. 7, p. 83 (1838); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 60 (1867). C. Europe.
419. *E. surata*, Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 159 (1907). S. Europe.
420. *E. tanysphyra*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 38 (1873); Strobl, Glasnik Bosn. Herceg. Sarajevo, Vol. 10, p. 409 (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 569 (1900); Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 309 (1906); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 30 (1907). C. & S. Europe.
421. *E. tenebrosa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 404 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 311 (1902). Texas.
422. *E. tenuinervis*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 468, 471 (1912) (*Coptophlebia*). Formosa.
423. *E. tenuipes*, Loew, Berl. Ent. Zeitschr. Vol. 13, p. 92 (1869); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 304 (1906). C. Europe.
424. *E. tenuirostris*, Thomson, Eugen. Resa, Zool. p. 473 (1868); Bezzii, Ann. Mus. Hungar. Vol. 2, p. 341, note (1904). Australia.
425. *E. teres*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 315, f. 133, 142 (1902). Idaho.
426. *E. tersa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 404 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 311 (1902). North Carolina.
427. *E. tessellata*, Fabricius, Ent. Syst. Vol. 4, p. 405 (1794); Coquebert, Ill. Icon. p. 88, pl. 20, fig. 12 (1804); Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 227 (1804); Fabricius, Syst. Antl. p. 140 (1805); Latreille, Gen. Crust. Ins. Vol. 4, p. 303 (1809); Fallen, Empid. Suec. p. 17 (1815); Lamarck, Anim. S. Vert. Vol. 3, p. 402 (1816); Wiedemann, Zool. Mag. Vol. 1, p. 2, 25 (1817); Meigen, Europe, N. Africa.

Syst. Beschr. Vol. 3, p. 17 (1822); Curtis, Brit. Ent. p. 18, 2 (1824); Macquart, Dipt. N. Fr. Vol. 3, p. 121 (1827); Meigen, Syst. Beschr. Vol. 6, p. 337 (1830); Macquart, Hist. Nat. Dipt. Vol. 1, p. 327 (1834); Blanchard, Hist. Nat. Ins. Vol. 3, p. 581 (1840); Loew, Bemerk. Posen. Gegend. Art. Zweifl. Gatt. 1840, p. 20, pl. 1, f. 20: Isis, Vol. 7, p. 547 (1840); Zetterstedt, Dipt. Sc. Vol. 1, p. 368 (1842); Boitard, Man. Ent. Vol. 3, p. 317 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 492 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3023 (1849); Dahlbom, Vet. Akad. Handl. Stockholm, p. 161 (1850); Schötz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 52 (1851); Walker, Ins. Brit. Vol. 1, p. 90 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 4271 (1852); Nylander, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 118 [note] (1858); Bonsdorff, Finl. Ins. Vol. 1, p. 164 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 103 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 379, 399 (1866); Perty, Mitteil. Naturf. Ges. Bern. 1867, p. 306 (1867); Meinert, Trophi Dipt. p. 54, pl. 4, f. 11, 15 (1881); Beling, Arch. Naturg. Berlin, Vol. 47, p. 205, 15 (1882); Leunis, Synops. Zool. Vol. 2, p. 402 (1886); Neuhaus, Dipt. March. p. 71 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 123 (1887); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 63 (1892); Leonardi, Ins. Nov. Vol. 3, p. 362, f. 195 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 215 (1906); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 124 (1907); Hamm, Ent. Mag. Vol. 45, p. 158 (1909), sex relations; Lundbeck, Dipt. Danica, Vol. 3, p. 129, f. 42 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 44 (1913); Pearce, Typical Flies, p. 24, f. 79, 80 (1915).

rufipes, Stæger, Zetterstedt, Dipt. Scand. Vol. 1, p. 369 (1842), not described.

var. *atricipes*, Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 63 (1893). Europe.

var. *livida*, Fabricius (not Linnæus), Spec. Ins. Vol. 2, p. 471 (1781); Ent. Syst. Vol. 4, p. 404 (1794); Syst. Antl. p. 139 (1805). Europe.

var. *tipuloides*, Linnæus, Syst. Nat. Ed. 10, p. 606 [1758] (*Asilus*); Linnæus, Fauna Suec. p. 47 [1761] (*Asilus*); Müller, Fauna Friedrich. p. 88 [1764] (*Asilus*); Linnæus, Syst. Nat. Ed. 12, Vol. 2, p. 1008 [1767] (*Asilus*); Müller, Zool. Danica Prodr. p. 181 [1776] (*Asilus*); Gmelin, Syst. Nat. Vol. 5, p. 2899 [1790] (*Asilus*); Schrank, Fauna Boica, Vol. 3, p. 161 [1803] (*Asilus*); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 63 (1892); Verh. Zool-bot. Ges. Wien, Vol. 59, p. 172 (1909).

428. *E. testacea*, Fabricius, Syst. Antl. p. 141 (1805); Latreille, Gen. Crust. C. Europe.
Ins. Vol. 4, p. 303 (1809); Meigen, Syst. Beschr. Vol. 3, p. 37 (1822); Walker, List Dipt. Brit. Mus. Vol. 3, p. 497 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), 52 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 158 (1867); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 210 (1906); Bezzì, Deutsche Ent. Zeitschr. Beiheft, p. 90 [1909] (*Xanthempis*).
punctata, Fallen, part, Empid. Suec. p. 19, part (1815).

429. *E. testiculata*, Bezzì, Deutsche Ent. Zeitschr. Beiheft, p. 91, 92 [1909] Crete.
(*Xanthempis*).

430. *E. Thalhammeri*, Strobl, Glasnik Bosn. Herceg. Sarajevo, Vol. 10, p. 407 (1898); Mitteil. Bosn. Herceg. Vol. 7, p. 568 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 30 (1907). Bosnia.

431. *E. thermophila*, Wiedemann, Aussereur. Zweifl. Ins. Vol. 2, p. 4 (1830); Uruguay.
Bezzì, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909).

—. *E. thoracica*, Eversmann, Bull. Soc. Nat. Moscou, Vol. 7, p. 424 (1834), no description.

432. *E. * tibialis*, Loew, Bernsteinfauna, p. 41 [1850] (*unnamed*); Meunier, Miscell. Ent. Vol. 102, p. 178 (1899).
433. *E. totipennis*, Bellardi, Mem. Accad. Sc. Torino (2), Vol. 21, p. 199: Saggio, Vol. 2, p. 99 (1865); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 292 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909).
434. *E. trianguligera*, Strobl, Glasnik Bosn. Herceg. Sarajevo, Vol. 10, p. 411, (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 571 (1900); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 226 (1907); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 158 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 98, note [1909] (*Pachymeria*).
435. *E. tridentata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 23, p. 609 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 301, f. 132 (1902).
436. *E. trigramma*, Wiedemann-Meigen, Syst. Beschr. Vol. 2, p. 38, (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 379 (1842); ibidem, Vol. 8, p. 3025 (1849); Walker, Ins. Brit. Vol. 1, p. 95 (1851); Zetterstedt, Dipt. Scand. Vol. 12, p. 4612 (1855); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 17, 21, 158 (1867); Verrall, Ent. Mag. London, Vol. 8, p. 282 (1872); Beling, Arch. Naturg. Berlin, Vol. 41, p. 1, 39 (1880); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 60 (1892); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzi, Deutsche Ent. Zeitschr. p. 91 [1909] (*Xanthempis*); Verrall, Brit. Flies, Vol. 5, p. 27, f. 49 (1909); Hamm, Ent. Mag. Vol. 45, p. 159 (1909), mating habits; Lundbeck, Dipt. Danica, Vol. 3, p. 93, f. 30, 31 [1910] (*Xanthempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910).
- erruginea*, Meigen (not Scopoli), Syst. Beschr. Vol. 7, p. 88 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Kuntze, Zeitschr. Hym. Dip., Vol. 6, p. 211 (1906).
- punctata*, Fallen, Empid. Suec. p. 19, part (1815).
437. *E. trilineata*, Gmelin, Syst. Nat. Vol. 5, p. 2890 (1790); Olivier, Encycl. Méth. Vol. 6, p. 391 (1791).
438. *E. trilineata*, Pallas-Wiedemann, Zool. Mag. Vol. 1, p. 2, 25 (1817); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 216 (1906).
439. *E. tristis*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 35, 53 (1867); ibidem, Vol. 13, p. 91 (1869); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 301 (1906).
440. *E. * tritara*, new name.
- tristis*, Meunier, Ann. Sc. Nat. (Zool.) p. 7, 94, 124, pl. 12, f. 3, 4 (1908).
441. *E. trivittata*, Macquart, Dipt. N. France, Vol. 3, p. 125 (1827); Hist. Nat. Dipt. Vol. 1, p. 331 (1834); Meigen, Syst. Beschr. Vol. 7, p. 82 (1838); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 32 (1907).
442. *E. truncata*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 31 (1822); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 156 (1907).
443. *E. tumida*, Meigen, Syst. Beschr. Vol. 3, p. 39 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 334 [1834] (*Pachymerina*); Meigen, Syst. Beschr. Vol. 7, p. 89 [1838] (*Pachymeria*); Boitard, Ent. Man. Vol. 3, p. 319 [1843] (*Pachymerina*); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 59 [1867] (*Pachymeria*); Strobl, Progr. Seitenst. Vol. 14, p. 9 [1880] (*Pachymeria*); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 156 (1907); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 94, note [1909] (*Anacrostichus*).
444. *E. umbrina*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 41 (1822); Curtis, Brit. Ent. Vol. 8, p. 18 (1824).
- . *E. umbripennis*, Eversmann, Bull. Soc. Nat. Moscou, Vol. 7, p. 424 (1834) no description.
- C. Europe.
- Baltic Amber.
- France.
- Portugal.
- Great Britain.
- Mexico.
- S. Europe, N. Africa.
- E. United States.
- Europe.
- Russia.
- Grecian Archipelago.

445. *E. unicolor*, Brullé, Expéd. Morée (Zool.), Vol. 3, p. 297 (1834); S. Europe.
 Macquart, Hist. Nat. Dipt. Vol. 1, p. 327 (1834); Meigen, Syst. Beschr. Vol. 7, p. 82 (1838); Boitard, Ent. Man. Vol. 3, p. 317 (1843); Loew, Berl. Ent. Zeitschr. Vol. 13, p. 71, note (1869); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 298 (1906).
446. *E. unicolor*, Walker, Ins. Brit. Vol. 1, p. 91 (1851). England.
447. *E. univittata*, Loew, Berl. Ent. Zeitschr. Vol. 11, p. 12, 20 (1867); C. & N. Europe.
 Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 211 (1906); Bezzī, Deutsche Ent. Zeitschr. Beiheft, p. 90 [1909] (*Xanthempis*); Lundbeck, Dipt. Dan. Vol. 3, p. 90, f. 27 [1910] (*Xanthempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 62 (1910).
stercorea, Zetterstedt, var. b, Dipt. Scand. Vol. 1, p. 378 (1842).
448. *E. uruguayensis*, Arribálzaga, Natural. Argent. Buenos-Aires, Vol. 1, p. 293 (1879); Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909). Uruguay, Argentina.
449. *E. vaginifer*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 352 (1902). E. United States.
450. *E. valdiviana*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 755 (1865); Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 346 (1909). Chile.
451. *E. valentis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 402 (1895); California.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 320, f. 138 (1902).
452. *E. validis*, Adams, Kansas. Univ. Sc. Bull. Vol. 3, p. 158 (1905). Rhodesia.
453. *E. variabilis*, Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 291 (1857); Philippi, Verh. Zool.-bot. Ges. Wien. Vol. 15, p. 753 (1865); Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 347 (1909). Chile.
454. *E. varians*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 130 (1889); Chile.
 Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 343 (1909).
455. *E. variegata*, Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 225 (1804); Loew, Wien. Ent. Monatschr. Vol. 6, p. 168 (1862); Berl. Ent. Zeitschr. Vol. 9, p. 238 (1865); Mik, Dipt. Bernst. Vol. 2, p. 57 (1885); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 62, note (1893); Kuntze, Zeitschr. Hym. Dipt. Vol. 6, p. 213 (1906). C. Europe.
456. *E. varipes*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 324 : Cent. 1, No. 34 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 399 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 307 (1902). E. North America.
457. *E. velutina*, Bezzī, Ann. Mus. Hungar. Vol. 10, p. 468, 471 [1912] (Formosa).
Coptophlebia.
 var. *cineraria*, Bezzī, Suppl. Ent. Berlin, Vol. 3, p. 71 [1914] (*Coptophlebia*). Formosa.
458. *E. vernalis*, Meigen, Syst. Beschr. Vol. 3, p. 27 (1822); Curtis, Brit. Ent. Vol. 8, p. 18, 4 (1824); Zetterstedt, Dipt. Scand. Vol. 1, p. 384 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 495 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3025 (1849); Walker, Ins. Brit. Vol. 1, p. 93 (1851); Zetterstedt, Dipt. Scand. Vol. 11, p. 4271 (1852); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. Ins. Vol. 1, p. 167 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 107 (1862); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 78 (1893); Becker, Act. Soc. Fenn. Helsingfors. Vol. 26 (9), p. 28 (1900); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 27 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 114 [1910] (*Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 64 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 42 [1913] (*Pterempis*); Kuntze, Deutsche Ent. Zeitschr. p. 548 (1913). Europe, Siberia.
- hyalinata*, Meigen, Syst. Bes. Vol. 6, p. 338 (1830).
? leucoptera, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 221 (1804); Syst. Beschr. Vol. 3, p. 27 (1822); Curtis, Brit. Ent. Vol. 8, p. 18 (1824);

- Meigen, Syst. Beschr. Vol. 6, p. 337 (1830); Walker, List Dipt. Brit. Mus. Vol. 3, p. 495 (1849); Ins. Brit. Vol. 1, p. 94 (1851).
- pennaria*, Fallen, part, Empid. Suec. p. 20, part (1815); Zetterstedt, Fauna Ins. Lappon. p. 562, part (1838); Lundbeck, Dipt. Danica, Vol. 3, p. 118 [1910] (*Pterempis*); Wahlgren, Ent. Tidskr. Vol. 31, p. 63 (1910).
459. *E. vicina*, Arribálzaga, Natural. Argent. Buenos-Aires, Vol. 1, p. 294 (1878); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909). Argentina.
460. *E. villosa*, Macquart, Mém. Soc. Sc. Lille, p. 161 (1823). France.
461. *E. villosula*, Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 131 (1889); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 343 (1909). Chile.
462. *E. virgata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 408 (1895); Proc. Wash. Acad. Sc. Vol. 2, p. 408 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 312, f. 108 (1902). W. North America. Alaska.
463. *E. volucris*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 23 (1822); S. & C. Europe. Loew, Berl. Ent. Zeitschr. Vol. 11, p. 56 (1867); Strobl, Wien. Ent. Zeit. Vol. 18, p. 17 (1899); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 26 (1907). C. Europe.
- var. *vitripennis*, Meigen, Syst. Beschr. Vol. 3, p. 25 (1822); Curtis, Brit. Ent. Vol. 8, p. 18, 4 (1824); Macquart, Dipt. N. France, Vol. 3, p. 127 (1827); Hist. Nat. Vol. 1, p. 332 (1834); Boitard, Man. Ent. Vol. 3, p. 319 (1843); Walker, Ins. Brit. Vol. 1, p. 94 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 106 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 57, 163 (1867); ibidem, Vol. 13, p. 88 (1869); Raddatz, Arch. Ver. Freund Nat. Mecklenb Rostock, Vol. 27, p. 40 (1873); Elliot, Trans. Ent. Soc. Lond. 1896, p. 117-128 (1896); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 101 [1909] (*Coptophlebia*); Lundbeck, Dipt. Danica, Vol. 3, p. 106, f. 37 [1910] (*Coptophlebia*).
- nana*, Macquart, Mém. Soc. Sc. Lille, p. 161, 165 (1823); Bezzi, Deutsche Ent. Zeitschr. Beiheft, p. 91 (1909).
- turbida*, Meigen, Syst. Beschr. Vol. 7, p. 86 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 108 (1862); Loew, Berl. Ent. Zeitschr. Vol. 11, p. 61, 163 (1867); Neuhaus, Dipt. March. p. 72 (1886); Kuntze, Zeitschr. Hym. Dipt. Vol. 7, p. 26 [1907] (*turbida*).
464. *E. Walkeri*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 343 (1909). Magellan.
- fulva* Walker (not Macquart), Trans. Linn. Soc. Lond. (Zool.) Vol. 17, p. 341 (1837).
465. *E. xanthopyga*, Schiner, Novara Reise, Dipt. p. 204, 6 (1869); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 343, note (1904). Australia.
466. *E. xochitl*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, p. 370 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 292, f. 118 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 345 (1909). Mexico.

27. GENUS EMPIMORPHA, COQUILLETT

Empimorpha, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389, 396 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 329 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 249, 263 (1903); Melander, Williston Man. N. Amer. Dipt. p. 226 (1908); Kertész, Cat. Dipt. Vol. 6, p. 79 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 537 (1910).

Characters. — Robust species, differing from *Empis*, s. str., only in the greater development of the hairs which occur in conspicuous amount even on the face. Style of the antennæ much shorter than the third joint, basal joints of the antennæ hirsute; eyes of the male contiguous along the front, the upper facets larger than the lower but still small; proboscis twice as long as the head, vertical, palpi rather cylindrical, porrect, hairy. Disk of the thorax very hairy, not setose, about six notopleurals no

humeral or posthumeral; metapleuræ and pectus abundantly hairy. Abdomen with more or less hairs, stout, sometimes asymmetrical toward the end in the male, in which case the fifth ventral is large and posteriorly tuberculate. Legs strong but not deformed, hairy and bristly. Wings narrow, costa stopping at the tip of the wing, basal bristle present, auxiliary vein straight, ending before the costa, third vein with a short fork, discal cell complete, shorter than the basals, anal cell small, anal vein attaining the margin, axillar excision deep, alula small, with a short close fringe of fine hairs; calyptères nearly bare, their fringe almost microscopic.

Type species: *E. comantis*, Coquillett. This group was supposed by Coquillett to bear the same relation to *Empis* that *Neocota* did to *Rhamphomyia*. In both cases the distinguishing characters are confined to the hairy face and nearly bare calyptères. Species of *Empis* of the *obesa* group are very similar to *Empimorpha*, even to the reduction of the fringe of the calyptères. It seems more likely that *Neocota* has developed from *Empimorpha* by a loss of the fork of the third vein than that both of these genera are parallel outgrowths of *Rhamphomyia* and *Empis*. Such an evolution opens the interesting question as to whether *Rhamphomyia* is not a polyphyletic group, developing from several of the subgenera of *Empis* through a simplification of the third vein.

Geographical distribution.

1. *E. barbata*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 195 : Cent. 2, No. 19 California, Washington.
[1862] (*Empis*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 396
(1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 329, f. 106,
(1902).
2. *E. comantis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 396 (1895); California.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 330, 353 (1902).
3. *E. geneatis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 329, f. 105 (1902). California.
4. *E. plumipes*, Matsumura, Addit. Vol. 2, p. 356, pl. 21, f. 18 (1916). Japan.
5. *E. rufithorax*, Brunetti, Rec. Indian Mus. Vol. 9, p. 30 (1913); Fauna India.
Brit. India, Dipt. Vol. 1, p. 359 [1920] (*Hilara*).

28. GENUS NEOCOTA, COQUILLETT

Neocota, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 434 (1895); Proc. Ent. Soc. Wash. Vol. 5, p. 254, 260 (1903); Melander, Williston, N. Amer. Dipt. Man. p. 225 (1908); Kertész, Cat. Dipt. Vol. 6, p. 38 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 574 (1910).

Characters. — Large stout pilose species resembling *Rhamphomyia* but with hairy face and bare calyptères. Wholly opaque black. Eyes of the male entirely contiguous above the antennæ, bare, the facets minute and uniform, face broad and densely hairy; antennæ elongate, equaling the length of the head, the first joint cylindrical, long and hairy, the second joint globular and not setose, the third joint lanceolate, tapering to near the end, then narrowly cylindrical, its style one-third as long as the third joint, thick, consisting of two joints, with the basal segment quadrate and the outer segment conical and twice as long as the basal; proboscis slender, nearly vertical, slightly shorter than the head, palpi narrowly spatulate, black and densely hairy. Prothorax and mesothorax densely hairy, three notopleural bristles but no others, scutellum with a dense fringe, metapleuræ with a dense cluster of hairs. Basal segments of the abdomen with abundant hair, fourth to the seventh tergites velvety, second tergite with a double transverse row of pittings; lateral valves of the pygidium large and ovoid, dorsal valves small, penis rather slender, only its base visible. Legs robust, including the coxae densely hairy, no bristles but many of the hairs long and setiform. Fringe of the calyptères vestigial, the hairs of

microscopic size. Wings without a humeral bristle, the alula, anal incision and anal lobe well developed, costa stopping beyond the third vein but before the tip of the wing, auxiliary vein straight and vanishing at the tip, first vein long, third vein not forked, anal crossvein abruptly reflexed, continuous with the under side of the anal cell, anal vein faint, marginal cilia and setulæ microscopic.

Type species : *N. Weedi*, Coquillett, the only known species.

Geographical distribution.

1. *N. Weedi*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 434 (1895); S. E. United States. Howard, Ins. Book, pl. 18, f. 30 (1901).

29. GENUS RHAMPHOMYIA, MEIGEN

Rhamphomyia, Meigen, Syst. Beschr. Vol. 3, p. 42 (1822); Macquart, Dipt. N. France, Vol. 3, p. 129 (1827); Curtis, Brit. Ent. Vol. 8, p. 517 (1834); Macquart, Hist. Nat. Dipt. Vol. 1, p. 334 (1834); Zetterstedt, Fauna Ins. Lappon. p. 562 [1838] (*Rhamphomyza*); Westwood, Gen. Syn. p. 131 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 388 (1842); Boitard, Man. Ent. Vol. 3, p. 319 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 108 (1851); Berendt, Bernstein. Organ. Reste Vorw. [1856] (*Rumphomyia*); Rondani, Dipt. Ital. Vol. 1, p. 151 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 563 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862); Lioy, Atti Inst. Ven. 1864, p. 601 (1864); Beling, Arch. Naturg. Berlin, Vol. 48, p. 240 (1882); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 122 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390, 409 (1895); Proc. Ent. Soc. Wash. Vol. 5, p. 256 (1903); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 428 (1905); Melander, Williston Man. N. Amer. Dipt. p. 225 (1908); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 302, 320 (1909); Kertész, Cat. Dipt. Vol. 6, p. 15 (1909); Frey, Acta Soc. Sc. Fenn. Helsingfors. Vol. 31 (9), p. 17-21 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 599 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 30 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 48 (1910); White, Proc. Roy. Soc. Tasmania, 1916, p. 237 (1917); Malloch, Rept. Can. Arct. Exped. 1913-18, Dipt. p. 44 (1919); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 345 (1920); Frey, Notulæ Ent. Vol. 2, p. 1-10, 33-45, 65-77 (1922).

Choreodromia, Frey, Notulæ Ent. Vol. 2, p. 3, 6 (1922).

Dasyrhamphomyia, Frey, Notulæ Ent. Vol. 2, p. 4, 65 (1922).

Dionnæa, Meigen, Nouv. Classif. Mouches, p. 24, part. (1800); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 534 (1910).

Enicopteryx, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 264 (1829); Westwood, Gen. Syn. p. 131 (1840); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 249 (1903); Kertész, Cat. Dipt. Vol. 6, p. 15 [1908] (*Henicopteryx*); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 538 (1910).

Holoclera, Schiner, Wien. Ent. Monatschi. Vol. 4, p. 53 (1860); Fauna Dipt. Austr. Vol. 1, p. 80 (1862); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 122 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 251, 260 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 552 (1910); Frey, Notulæ Ent. Vol. 2, p. 3, 5 (1922).

Lundstroemiella, Frey, Notulæ Ent. Vol. 2, p. 3, 4 (1922).

Macrostromus, Wiedemann, Zool. Mag. Berlin, Vol. 1, Pt. 1, p. 60 (1817); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252, 260 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 564 (1910).

Megacyttarus, Bigot, Ann. Soc. Ent. France (5), Vol. 10, Bull. p. 47 (1880); Williston, Synopsis N. Amer. Dipt. p. 82 (1888); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 117 [1899] (*Megacillarus*);

Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 350 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 566 (1910).

Pararhamphomyia, Frey, Notulæ Ent. Vol. 2, p. 3, 33 (1922).

Platyptera, Meigen-Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 256 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 592 (1910).

Characters. — Usually rather slender insects measuring from three to nine millimeters, of dark often shining coloration, though sometimes yellow. Head globular, occiput hairy; eyes of the male usually contiguous or subcontiguous, if separated the front of the male is narrower than that of the female, when dichoptic the facets are uniform, when holoptic the upper facets are somewhat larger, eyes of the female always separated and with uniform facets; face quadrate or narrow, cheeks linear; antennæ inserted near the middle of the head, projecting horizontally about as much as the head-height, distinctly three-jointed, the first joint usually cylindrical and a little longer than the globose second joint, both more or less hairy, the third joint elongate conical, tipped by a two-segmented style, whose distal part is longer than the basal and tapers; proboscis ranging from as long as the head to twice as long, depending vertically or folded somewhat backward, palpi cylindrical upturned, hairy. Thorax often a little arched, metathorax small, notal bristles very variable, as in *Empis*, metapleuræ always more or less setose. Abdomen usually compressed, pygidium often large and elaborate, generally compressed and erect with slender central filament and broad lateral and dorsal valves, sometimes the pygidium is closed. Legs provided with hairs and bristles, sometimes the femora and tibiæ of the female pennate, sometimes the knees of the male armed. Wings of the female often darker and sometimes broader than of the male, costa stopping at the tip of the wing, auxiliary vein straight and interrupted at the end, third vein always simple, discal cell complete, except that rarely the posterior crossvein is lacking, in some species the discal cell varies with the sex, being extraordinarily large in the female, anal cell short, the crossvein strongly reflexed, anal vein discontinuous with the anal cell, anal angle of the wing large, but variable in size, axillar incision distinct, alula small or wanting, calypteres with a dense fringe.

Rhamphomyia is probably one of the most recent of the genera of Empididae, just as it is the most plastic and in alpine regions the dominant genus. With four hundred species described it has become one of the largest of genera and thus it is desirable that the well known and universally accepted name be maintained and that attempts to seek out and establish prior names be discouraged. In the interest of continuity I have accepted as the type species of *Rhamphomyia*, *sulcata* Meigen, the species selected by Curtis in 1834.

When Coquillett attempted to fix the nomenclature of the Empididæ in 1903 he resurrected Wiedemann's name *Macrostomus* for this genus. Seven years later, in his type species paper of 1910, he made *platyptera* the tautonymic type of *Platyptera* Meigen, 1803, and since this is supposed to be the same as *Dionnaea* Meigen, 1800, he would change the host of species known as *Rhamphomyia* from *Macrostomus* to *Dionnaea*, spp. I have here, following Curtis, adopted *borealis* as the type of *Platyptera*, thus placing it and the questionable *Dionnaea* in the synonymy under *Empis*. Until the Fabrician species *Hybos ferrugineus*, on which Wiedemann founded his genus *Macrostomus*, is rediscovered the name *Macrostomus* may well be kept in abeyance.

The species of *Rhamphomyia* are essentially boreal and seem to be unlimited in their number. Often a species is unusually common for a few days to disappear until the next year. Some species swarm in the open in an aerial dance; these are usually the species of summer or autumn. Others, especially the vernal forms, frequent the deep shade of woods, and may be found about herbage or lazily flying here and there. The genus is unusually plastic in characters, and hence a vast number

of species results. A day's collecting in the haunts of these insects has rarely failed to produce a new form. I have before me several hundred undescribed species of *Rhamphomyia*, and as the number of combinations of possible characters is far from being exhausted, it can be stated that we are just beginning our acquaintance with these interesting species.

Whether *Rhamphomyia* is a monophyletic or a polyphyletic group is unsettled. The trend of evolution in the genus is paralleled to a great degree by species of *Empis*. The pennate structure of the legs of the females, the development of ventral prongs on the abdomen of the male, the structure of his pygidium and of the knee armature, the tendency to holopticism and the reduction or amplification of the bristles may be the result of convergent evolution or may indicate that the various groups of species we call *Rhamphomyia* are offshoots of the subgenera of *Empis*, specialized by a simplification of the third vein. Richard Frey of Helsingfors has recently published a segregation of this complex group into subgenera. His study was based mainly on palaeartic species and accordingly his groups do not adapt themselves as readily to the tropical forms as they do to the nearctic species. The divisions proposed by Dr. Frey are based mainly on the arrangement of the notal bristles, character of axillar incision of the wing, form of pygidium and contiguity of the male eyes. The subgenera are diagnosed below and a list of species assignable to each is given. The list is obviously incomplete, being made up from Frey's paper and such additional species as are before me. Some species, like *R. cyanogaster*, Wh. & M., *pulchriventris*, Bez., *Sauteri*, Bez., and *tolteca*, Wh. & M., do not agree well with the subgenera as defined.

SUBGENUS DASYRHAMPHOMYIA, FREY

Characters. — Males rarely dichoptic, the eyes usually broadly contiguous; ocellar triangle developed; antennæ not excessively lengthened. Body rather stout, usually dull in color, dorsocentrals usually weak and pluriseriate, acrostichals present, scutellars many. Pygidium characteristically of two types, either with a long erect vesiculate structure, or small and closed, broadly deformed, with the seventh tergite convex and usually set in the wide sixth tergite, in the latter case armed with two prongs below and the last sternite usually fringed with two bunches of yellow setæ. Legs rather strong, rarely pennate in female, the hind femora rarely setose below. Wings commonly broad and brown in female, axillar incision usually acute.

Type species: *R. vesiculosa*, Fallen. Additional species include: *R. Brussneri*, Frey, *conservativa*, Mall., *coracina*, Zett., *Hambergi*, Frey, *Hovgaardii*, Holmgr., *mirifica*, Frey, *nigrita*, Zett., *Pokornyi*, Bez., *reflexa*, Zett., *vara*, Lw., *vespertilio*, Zett., *virgata*, Coq. and *Wuorentausi*, Frey.

SUBGENUS HOLOCLERA, SCHINER

Characters. — Males holoptic, ocellar triangle and bristles evident; antennæ not greatly lengthened. Small species with slender body, sometimes differing in color in the sexes, sometimes yellow in both male and female; dorsocentrals uniseriate, acrostichals present or absent. Pygidium small and closed. Legs slender. Axillar incision of wing ranging from obtuse to rather acute.

Type species: *R. nigripennis*, Fallen. Other species, all from the palaeartic region, include: *R. culicina*, Fall., *flava*, Fall., *flaviventris*, Macq., *heterochroma*, Fab., *pallidiventris*, Fall., *sciarina*, Fall., *tenuirostris*, Fall. and *umbripennis*, Meig.

SUBGENUS LUNDSTROEMIELLA, FREY

Characters. — Males dichoptic; the ocellar triangle not elevated, ocellar bristles very weak; antennæ greatly lengthened, nearly twice as long as the head, the third joint excessively long. Body

delicately built, without evident dorsocentrals and acrostichals. Legs slender. Wings narrow, axillar incision obtuse to rectangular, alulae nearly wanting.

Type species : *R. hybotina*, Zetterstedt. Additional species included in this subgenus are: *R. aterrima*, Frey, *limbipennis*, Bez., *longefilata*, Str., *magellensis*, Bez., *sphenoptera*, Lw. and *tumiditarsis* Old. All of these are palæarctic except *limbipennis*, which comes from South America.

SUBGENUS MEGACYTTARUS, BIGOT

Characters. — Males dichoptic, ocellar triangle and ocellar bristles developed; antennæ not abnormally lengthened. Body rather robust, dorsocentrals usually weak and pluriseriate, acrostichals present. Abdominal hairs usually pale or brown, pygidium usually open, with filiform penis. Legs normal, bristles few, not pennate in female. Wings with discal cell of female often greatly enlarged, axillar incision obtuse to rectangular, rarely acute.

Type species : *R. limbata*, Loew. Bigot erected the genus *Megacyttarus* on a female which he described as *argentea*. Coquillett has placed this as a synonym of *limbata*. Lw. It is to this group that Frey gave the subgeneric name *Choreodromia*. The species are mainly characterized by the separated eyes of the males and the abnormally enlarged discal cell of the females. The flies are usually met with in an aerial dance over pools and small streams. Males are usually more abundant than the females, the females often being silvery in color.

Additional species of *Megacyttarus* include: *R. anomala*, Old., *anomalina*, Zett., *anomalipennis*, Meig., *disparilis*, Coq., *fulvolanata*, Frey, *gutiflar*, Frey, *hirtipes*, Lw., *irregularis*, Lw., *kamtschatica*, Frey, *liturata*, Lw., *maculipennis*, Zett., *nigripes*, Fab., *nodipes*, Fab., *paradoxa*, Wahlb., *scaurissima*, Wh. and *tephraea*, Meig.

SUBGENUS PARARHAMPHOMYIA, FREY

Characters. — Males holoptic, ocellar triangle and bristles developed; antennæ not greatly lengthened. Body of regular build, dorsocentrals usually placed in one or two rows, if pluriseriate the thorax is shining black, acrostichals present. Pygidium regularly open, with penis usually filiform. Legs rather strong, often pennate in female. Wings with discal cell normal, axillary incision obtuse to rectangular, rarely acute.

Type species : *R. plumipes*, Fallen. This is the dominant group, abundantly represented in palæarctic and nearctic regions. The following species can be assigned to *Pararhamphomyia*: *R. albata*, Coq., *albibennis*, Fall., *albissima*, Frey, *amoena*, Lw., *anfractuosa*, Bz., *antennata*, Frey, *atra*, Meig., *barypoda*, Coq., *bifilata*, Coq., *bipila*, Str., *breviventris*, Frey, *brevis*, Lw., *candicans*, Lw., *caudata*, Zett., *chibinensis*, Frey, *cinefacta*, Coq., *cineracea*, Coq., *clavator*, Coq., *compta*, Coq., *costata*, Zett., *crassicauda*, Str., *curvula*, Frey, *debilis*, Lw., *dentata*, Old., *dentipes*, Zett., *dimidiata*, Lw., *dispar*, Zett., *diversipennis*, Beck., *effera*, Coq., *eupterota*, Lw., *fascipeunis*, Zett., *filata*, Zett., *flexuosa*, Coq., *fuscipennis*, Zett., *fuscula*, Zett., *galactoptera*, Str., *gibba*, Fall., *hilariformis*, Frey, *Helleni*, Frey, *hirtula*, Zett., *impedita*, Lw., *intermedia*, Frey, *kaninensis*, Frey, *leucoptera*, Lw., *limbata*, Lw., *lividiventris*, Zett., *longestylata*, Frey, *longicauda*, Lw., *longipes*, Meig., *lucidula*, Zett., *luteiventris*, Lw., *manca*, Coq., *nana*, Lw., *Nasoni*, Coq., *nitidicollis*, Frey, *niveipennis*, Zett., *novecarolina*, Beut., *nox*, Old., *obscura*, Zett., *omissinervis*, Beck., *ozernajensis*, Frey, *pennata*, Macq., *physoprocta*, Frey, *plumifera*, Zett., *plumipes*, Fall., *polita*, Lw., *poplitea*, Wahlb., *priapulus*, Lw., *pseudogibba*, Str., *pulchra*, Lw., *pusio*, Lw., *pusilla*, Zett., *rufipes*, Zett., *scolopacea*, Say, *serotina*, Old., *setosa*, Coq., *simplex*, Zett., *soccata*, Lw., *subglaucella*, Frey, *subsultans*, Frey, *tarsata*, Meig., *tenuiterfilata*, Beck., *tipularia*, Fall., *truncata*, Frey, *umbilicata*, Lw., *umbripes*, Beck., *unguiculata*, Frey, *ungulata*, Lw., *valga*, Coq., *vittata*, Lw. and *Woldstedti*, Frey.

SUBGENUS RHAMPHOMYIA, S. STR., FREY

Characters. — Males nearly always holoptic, ocellar triangle and bristles evident; antennæ not greatly lengthened. Body rather stout, acrostichals present, dorsocentrals in one or more rows. Pygidium usually small, half-open, penis usually thick and more or less hidden, rarely exposed and filiform. Legs rarely distinctly slender, usually rather robust, the hind femora of both sexes usually bearing bristles or spinules, sometimes in the females without bristles below and in the corresponding males with bristles only at base or middle, legs of females sometimes pannate. Axillar incision of wings acute, the alulæ usually well developed.

Type species: *R. sulcata*, Meigen. This group is well represented in palæarctic and nearctic regions. Dr. Frey notes (in litt.) « es ist auffallend, wie viele gelbbeinige *Rhamphomyia*-Arten in Nord-Amerika vorkommen; diese Arten sind wahrscheinlich älter als die schwarzbeinigen Arten ». The following species can be assigned to this subgenus: *R. albosegmentata*, Zett., *alpina*, Zett., *amplipedis*, Coq., *anthracina*, Meig., *argentata*, Roed., *armata*, Beck., *attenuata*, Frey, *avida*, Coq., *basalis*, Lw., *Buzzi*, Frey, *chionoptera*, Meig., *ciliata*, Coq., *cinerascens*, Meig., *clauda*, Coq., *colorata*, Coq., *conformis*, Kow., *corvina*, Lw., *crinita*, Beck., *curvinervis*, Old., *curvipes*, Coq., *discoidalis*, Beck., *fimbriata*, Coq., *givipes*, Lw., *glabra*, Lw., *gracilis*, Lw., *grammoptera*, Frey, *lævipes*, Frey, *latifrons*, Str., *longicornis*, Lw., *loripedis*, Coq., *luridipennis*, Now., *macilenta*, Lw., *malaria*, Beck., *minytus*, Walk., *modesta*, Wahlb., *montana*, Old., *morio*, Zett., *mutabilis*, Lw., *nitidolineata*, Frey, *nitidula*, Zett., *Oldenbergi*, Frey, *Palméni*, Frey, *pareicellulata*, Frey, *pectinata*, Lw., *phanerostigma*, Frey, *platycnemis*, Frey, *propinquua*, Meig., *pulla*, Lw., *quinquelineata*, Say, *rava*, Lw., *ravida*, Coq., *robustior*, Frey, *rustica*, Lw., *Sancti-Mauritii*, Beck., *scitula*, Frey, *scutellaris*, Coq., *serpentata*, Lw., *soccata*, Lw., *sociabilis*, Will., *spectabilis*, Frey, *spinipes*, Fall., *stigmosa*, Macq., *sudigeronis*, Coq., *testacea*, Lw., *tibialis*, Meig., *trilineata*, Zett., *tristriolata*, Now., *umbrosa*, Lw., *ursina*, Old. and *villosa*, Zett.

Geographical distribution.

1. *R. abdominalis*, Bezz, Ann. Mus. Hungar. Vol. 3, p. 433 (1905); Nova Bolivia.
Acta Akad. Naturf. Halle, Vol. 91, p. 323 (1909)
2. *R. *ablata*, Meunier, Ann. Sc. Nat. (Zool.) (9), Vol. 7, p. 92, 114, pl. 9, f. 5, 6 (1908).
3. *R. advera*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 418 (1900). Alaska.
4. *R. agasicles*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 499 (1849). Canada.
5. *R. albata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 25, p. 103 (1902). Arizona.
6. *R. albiventris*, Strobl, Glasnik Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 404 (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 565 (1900). Bosnia.
7. *R. albipennis*, Fallen, Empid. Suec. p. 30 [1816] *Empis*; Meigen, Syst. Beschr. Vol. 3, p. 59 (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 408 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 501 (1849); Bonsdorff, Finl. tvåv. Ins. Dipt. p. 171 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Becker, Act. Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 24, f. 24, 25 (1900); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Frey, Notulæ Ent. Vol. 2, p. 41 [1922] (*Pararhamphomyia*). N. & C. Europe, Siberia.
- niveipennis, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908).
- picipes, Zetterstedt, Dipt. Scand. Vol. 1, p. 410 (1842).
8. *R. albissima*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 17 (1913); Ent. Tidskr. 1914, p. 78 (1914); Notulæ Ent. Vol. 2, p. 37 [1922] (*Pararhamphomyia*). N. & C. Europe.
- niveipennis, Zetterstedt, part, Dipt. Scand. Vol. 1, p. 409 (1842).

9. *R. albogeniculata*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Wuertemburg. Stuttgart. Vol. 1, p. 53 (1840)
10. *R. albopilosa*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 418 (1900); Malloch, Report Canad. Arct. Exped. 1913-18, Dipt. p. 47 (1919). Alaska.
11. *R. albosegmentata*, Zetterstedt, Fauna Ins. Lappon. p. 563 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 393 (1842); ibidem, Vol. 8, p. 3033 (1849); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 153 (1864); Verrall, Ent. M. Mag. London, Vol. 19, p. 224 (1883); Meijere, Tijdschr. v. Ent. Vol. 50, p. 176 (1907); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 32 (1913); Notulæ Ent. Vol. 2, p. 74 (1922).
- arctica*, Zetterstedt, Fauna Ins. Lappon. p. 566 [1838] (*Rhamphomyza*)
- morio*, Wahlberg, in litt. (not Zetterstedt), Dipt. Scand. Vol. 8, p. 3034 (1849).
12. *R. alipes*, Meigen, Syst. Beschr. Vol. 3, p. 45 (1822); Macquart, Hist. Nat. Dip. Vol. 1, p. 339 (1834); Boitard, Man. Ent. Vol. 3, p. 320 (1843). C. Europe.
13. *R. alpina*, Zetterstedt, Fauna Ins. Lappon. p. 565 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 416 (1842); ibidem, Vol. 8, p. 3041 (1849); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 154 (1864); Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 22, f. 20, 21 (1900); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 34 (1913); Notulæ Ent. Vol. 2, p. 71 (1922). N. Europe, Siberia.
14. *R. americana*, Wiedemann, Aussereur. Zweifl. Ins. Vol. 2, p. 8 (1829); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 413 (1895). North America.
15. *R. amœna*, Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 21 (1840); Isis, Vol. 7, p. 548 (1840); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Frey, Notulæ Ent. Vol. 2, p. 38 [1922] (*Pararhamphomyia*). Germany.
- setigera*, Stein, Wien. Ent. Zeit. Vol. 9, p. 108 (1890)
16. *R. amplicella*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 431 (1895). California.
17. *R. amplipedis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 422 (1895); Frey, Notulæ Ent. Vol. 2, p. 68 (1922). E. United States.
18. *R. anaxo*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 500 (1849). Canada.
19. *R. andalusiaca*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 12 (1899). Spain.
20. *R. anfractuosa*, Beazzi, Ann. Mus. Hungar. Vol. 2, p. 199 (1904); Olden-berg, Arch. Naturg. (Berlin), p. 83, A, 6, 18 (1919); Frey, Notulæ Ent. Vol. 2, p. 35, 36 [1922] (*Pararhamphomyia*). Hungary.
21. *R. angulifera*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 15 (1913); Notulæ Ent. Vol. 2, p. 36 [1922] (*Pararhamphomyia*). Finland.
22. *R. *angusta*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 93, 119, pl. 9, f. 15, 16; ibidem, pl. 10, f. 12 (1908). Baltic Amber.
23. *R. angustipennis*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 336 : Cent. 1, No. 55 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895). E. United States.
24. *R. anomala*, Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 77 (1915); Frey, Notulæ Ent. Vol. 2, p. 8, 10 [1922] (*Choreodromia*). Germany.
25. *R. anomalina*, Zetterstedt, Fauna Ins. Lappon, p. 563 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 391 (1842); ibidem, Vol. 13, p. 5010 (1859); Becker, Act. Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 16, f. 6 (1900); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 18 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 55 (1910); Frey, Acta
- N. Europe, Siberia.

- Soc. Fenn. Helsingfors, Vol. 37 (3), p. 11 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 79 (1915); Frey, Notulæ Ent. Vol. 2, p. 8, 9 [1922] (*Choreodromia*).
26. *R. anomalipennis*, Meigen, Syst. Beschr. Vol. 3, p. 55, pl. 23, f. 4 (1822); C. & N. Europe.
Zetterstedt, Dipt. Scand. Vol. 1, p. 411 (1842); ibidem, Vol. 13, p. 5016 (1859); Coquillett, Proc. Wash. Ent. Soc. Vol. 5, p. 249 (1903); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 12 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 81 (1915); Frey, Notulæ Ent. Vol. 2, p. 8, 10 [1922] (*Choreodromia*).
hyalipennis, Stephens, Cat. Brit. Ins. Vol. 2, p. 264 [1829] (*Enicopteryx*), no description.
27. *R. antennata*, Frey, Mém. Acad. Sc. Russ. Vol. 29 (10), p. 9 (1915); N. Siberia.
Notulæ Ent. Vol. 2, p. 44 [1922] (*Pararhamphomyia*).
28. *R. anthracina*, Meigen, Syst. Beschr. Vol. 3, p. 54 (1822); Loew, Isis, C. Europe.
Vol. 7, p. 547, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 20 (1840); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1852); Schiner, Fauna Dipt. Austr. Vol. 1, p. 100 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 113 (1887); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 51 (1892); Czizek, Zeitschr. Mähr. Landesmus. Brünn, Vol. 7, p. 165 (1907); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A. 6, p. 24 (1919); Frey, Notulæ Ent. Vol. 2, p. 76 (1922).
alpestris, Schummel, Arb. Veränd. Schles. Ges. p. 189 (1843).
? *Loewi*, Nowicki, Verh. Naturf. Ver. Brünn, Vol. 6, p. 78 [1869] (*Löwi*); Mik, Fauna Hernst. Vol. 2, p. 2, 56 (1885); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 113, 119 (1887).
29. *R. anthracinella*, Strobl, Glasnik. Mus. Bosn. Herceg. Sarajevo, Vol. 10, Bosnia.
p. 405 (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 566 (1900).
30. *R. anthracodes*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 420 (1900). Alaska.
- . *R. *antipedalis*, Loew, Bernsteinfauna p. 41 (1850) no description; Baltic Amber, Lower Oligocene.
Giebel, Ins. Vorwelt, p. 208 (1866) no description.
31. *R. aperta*, Loew (not Zetterstedt), Berl. Ent. Zeitschr. Vol. 6, p. 199: C. United States.
Cent. 2, No. 27 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 418 (1895).
32. *R. apicalis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 330 (1909). Peru.
33. *R. aprilis*, White, Proc. Roy. Soc. Tasmania, 1916, p. 238, f. 46 (1917). Tasmania.
34. *R. arcuata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 421 (1895). Massachusetts.
35. *R. arcuincta*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 331 (1909). South America.
36. *R. argentata*, Roeder, Wien. Ent. Zeit. Vol. 6, p. 113 (1887); Frey, Notulæ Ent. Vol. 2, p. 73 (1922). C. Europe.
37. *R. argyrina*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 334 (1909). Bolivia.
38. *R. argyrotarsis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 324, f. 2 (1909). Bolivia.
39. *R. armata*, Becker, Mém. Acad. Sc. Petrograd, Vol. 28 (7), p. 56, f. 2 (1915); Frey, Notulæ Ent. Vol. 2, p. 69 (1922). W. Siberia Tundra.
40. *R. armimana*, Oldenberg, Ann. Mus. Hungar. Vol. 8, p. 346, f. 1 (1910). C. Europe.
41. *R. aterrima*, Frey, Notulæ Ent. Vol. 2, p. 5 [1922] (*Lundstroemiella*). Alps.
? *nigripes*, Strobl (not Fabricius), Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 34, p. 198 (1897); ibidem, Vol. 46, p. 58 (1909).
42. *R. atra*, Meigen, Syst. Beschr. Vol. 3, p. 45 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 339 (1834); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Jaroschewsky, Arb. Ges. Naturf. Univ. Kharkow, Vol. 11, p. 353 (1877); Frey, Notulæ Ent. Vol. 2, p. 43 [1922] (*Pararhamphomyia*). C. Europe.
43. *R. atrata*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 419 (1900). Alaska.

44. *R. attenuata*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 33 (1913); Ent. Tidskr. 1914, p. 79 (1914); Notulæ Ent. Vol. 2, p. 73 (1922). N. & C. Europe.
45. *R. aucta*, Oldenberg, Arch. Naturg. Berlin, Vol. 82, 1, p. 157 (1916). Tyrol.
46. *R. avida*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 425 (1895). Massachusetts.
47. *R. barypoda*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 417 (1900). Alaska.
48. *R. basalis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 77 : Cent. 5, No. 54 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 410 (1895). E. United States.
49. *R. Bezzii*, Frey, Notulæ Ent. Vol. 2, p. 74 (1922). Italy.
50. *R. bibioniformis*, Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 25 (1919). Austria.
51. *R. bicolor*, Macquart (not Wiedemann), Mém. Soc. Sc. Lille, p. 163 [1823] (*Empis*); Dipt. N. France, Vol. 3, p. 135 (1827); Hist. Nat. Dipt. Vol. 1, p. 338 (1834); Meigen, Syst. Beschr. Vol. 7, p. 92 (1838). France.
52. *R. bicolor*, Wiedemann (not Macquart), Aussereur. Zweifl. Ins. Vol. 2, p. 8 (1829); Macquart, Dipt. Exot. Vol. 1, 2, p. 164 (1838); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 26 (1919). Locality unknown.
53. *R. biflata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 424 (1895). California.
54. *R. bilineata*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 230 (1804); Syst. Beschr. Vol. 3, p. 56 (1822). C. Europe.
55. *R. bipila*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 171 (1909); Frey, Notulæ Ent. Vol. 2, p. 45 [1922] (*Pararhamphomyia*). Spain.
56. *R. Biroi*, Bezzii, Ann. Mus. Hungar. Vol. 6, p. 392 (1908). Crete.
57. *R. boliviiana*, Bezzii, Ann. Mus. Hungar. Vol. 3, p. 429 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 322 (1909). Bolivia.
58. *R. brevis*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 334 : Cent. 1, No. 52 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895). E. United States.
59. *R. breviventris*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3) p. 21 (1913); Ent. Tidskr. 1914, p. 78 (1914); Not. Ent. Vol. 2, p. 39, 40 [1922] (*Pararhamphomyia*). Sweden, Finland, Siberia.
60. *R. Brusewitsii*, Holmgren, Spec. Ins. N. Semlia, p. 20 (1880); Ent. Tidskr. Vol. 4, p. 163 (1883). Nova Zemla.
- var. *anomalinervis*, Frey, Mém. Acad. Sc. Russ. Vol. 29 (10), p. 11 (1915). Arctic Siberia.
61. *R. Brussnewii*, Frey, Mém. Acad. Sc. Russ. Vol. 29 (10), p. 10 (1915); Dipt. Sarekgebiet, p. 684 (1916); Notulæ Ent. Vol. 2, p. 67 [1922] (*Dasyrhaphomyia*). N. Siberia.
62. *R. caesia*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 56 (1822); ibidem, Vol. 6, p. 339 (1830); Macquart, Hist. Nat. Dipt. Vol. 1, p. 338 (1834); Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 53 (1840); Walker, List Dipt. Brit. Mus. Vol. 3, p. 501 (1849); Ins. Brit. Vol. 1, p. 109 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Oldenberg, Arch. Naturg. Berlin, Vol. 63, A, 6, p. 19 (1919). C. & N. Europe.
- schistacea*, Meigen, Syst. Beschr. Vol. 3, p. 57 (1822).
63. *R. californica*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 420 (1895). California.
64. *R. calvimonitis*, Cockerell, Canad. Ent. Vol. 48, p. 123 (1916). Colorado.
65. *R. cana*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3043 (1849); ibidem, Vol. 11, p. 4272 (1852); Lundbeck, Dipt. Danica, Vol. 3, p. 78 (1910). Denmark.
66. *R. canaliculata*, Macquart, Dipt. N. France, Vol. 3, p. 131 (1827); Hist. Nat. Dipt. Vol. 1, p. 335 (1834); Meigen, Syst. Beschr. Vol. 7, p. 90 (1838); Gimmerthal, Bull. Soc. Nat. Moscou, Vol. 15, p. 666 (1842); Boitard, Man. Ent. Vol. 3, p. 319 (1843). France.
67. *R. candicans*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 81 : Cent. 5, No. 61 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18 p. 412 (1895). E. United States.
68. *R. carbonaria*, Wiedemann-Meigen, Syst. Beschr. Vol. 3, p. 59 (1822); Walker, List Dipt. Brit. Mus. Vol. 3, p. 503 (1849); Zetterstedt, Dipt. Scand. Vol. 13, p. 5018 (1859). N. Europe.

69. *R. carenifera*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 338 Chile.
(1909).
70. *R. caudata*. Zetterstedt, Fauna Ins. Lappon. p. 565 (1838); Dipt. Vol. 1, N. & C. Europe.
p. 417 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 172
(1861); Boheman, Oefv. Akad. Foerh. Stockholm, p. 570 (1865);
Holmgren, Svensk. Akad. Handl. Stockholm, Vol. 8, p. 26 (1876);
Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908);
Wahlgren, Ent. Tidskr. Vol. 31, p. 59 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 27 (1913); Notulæ Ent. Vol. 2,
p. 42 [1922] (*Pararhamphomyia*).
arthiops, Zetterstedt, Fauna Ins. Lappon. p. 568 [1838] (*Rhamphomyza*);
Dipt. Scand. Vol. 1, p. 436 (1842); ? Walker, Ins. Brit. Vol. 1, p. 110
(1851); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114
(1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 175 (1861); Siebke,
Nyt Mag. Naturvid. Vol. 14, p. 400 (1866); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 59 (1892); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910).
71. *R. chibinensis*. Frey, Notulæ Ent. Vol. 2, p. 42 [1922] (*Pararhamphomyia*) N. Russia.
72. *R. chionoptera*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 201 (1904) ♂; Olden-
berg, Ann. Mus. Hungar. Vol. 8, p. 346 (1910) ♀; Frey, Notulæ Ent. Vol. 2, p. 75 (1922). Tyrol.
73. *R. ciliata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 428 (1895). New Hampshire.
74. *R. ciliipes*, Say, Jour. Acad. Nat. Sc. Philad. Vol. 3, p. 95 [1823] (*Empis*);
Compl. Writ. Vol. 2, p. 83 [1859] (*Empis*); Wiedemann, Aussereur.
Zweifl. Ins. Vol. 2, p. 7 (1830). Ohio.
75. *R. cinefacta*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 419 (1900). Alaska.
76. *R. cineracea*, Coquillett, ibidem, Vol. 2, p. 416 (1900). Alaska.
77. *R. cinerascens*, Meigen, Class. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 230
[1804] (*Empis*); Syst. Beschr. Vol. 3, p. 48 (1822); ibidem, Vol. 6,
p. 339 (1830); Curtis, Brit. Ent. Vol. 8, p. 517 (1834); Scholz,
Zeischr. Ent. Breslau, Vol. 5 (19), p. 55 (1851); Walker, Ins. Brit.
Vol. 1, p. 109 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98
(1862); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 49
(1892); Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 20
(1900); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19
(1909); Kleine, Soc. Ent. Steglitz, Vol. 24, p. 65 (1909); Strobl,
Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 46, p. 59 (1909); Lund-
beck, Dipt. Danica, Vol. 3, p. 323 (1910); Meijere, Tijdschr. v.
Ent. Vol. 60, p. 134, pl. 8, f. 3, 4 (1918); Oldenberg, Arch. Naturg.
Berl. Vol. 83, A, 6, p. 22 (1919); Frey, Notulæ Ent. Vol. 2,
p. 72 (1922). C. Europe, Siberia.
78. *R. cinerea*, Fabricius (not Meigen), Syst. Ent. p. 802 [1775] (*Empis*);
Spec. Ins. Vol. 2, p. 472 [1781] (*Empis*); Mant. Ent. Vol. 2, p. 365
[1787] (*Empis*); Ent. Syst. Vol. 4, p. 405 [1794] (*Empis*); Gmelin,
Syst. Nat. Vol. 5, p. 2890 [1790] (*Empis*); Olivier, Encycl. Méth.
Vol. 6, p. 389 [1791] (*Empis*); Meigen, Class. Beschr. Eur. Zweifl.
Ins. Vol. 1, p. 232, d. [1804] (*Empis*); Fabricius, Syst. Antl. p. 141
[1805] (*Empis*); Billberg, Enumer. Ins. p. 120 [1820] (*Empis*);
Macquart, Dipt. N. France, Vol. 3, p. 130, pl. 3, f. 8 (1827); Meigen,
Syst. Beschr. Vol. 6, p. 340 (1830). C. Europe.
79. *R. clauda*, Coquillett, Proc. U. S. Nat. Mus. Vol. 23, p. 610 (1901). E. United States.
80. *R. clavator*, Coquillett, ibidem, Vol. 23, p. 611 (1901). Alaska.
macrura, Coquillett (not Loew), Proc. Wash. Acad. Sc. Vol. 2, p. 420
(1900).

81. *R. clavigera*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 335 : Cent. 1, No. 53 New York.
 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 417 (1895).
82. *R. clypeata*, Macquart, Hist. Nat. Dipt. Vol. 1, p. 339 (1834); Meigen, Sicily.
 Syst. Beschr. Vol. 7, p. 92 (1838).
83. *R. colorata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 8, p. 420 (1895). Texas.
84. *R. compta*, Coquillett, ibidem, Vol. 8, p. 423 (1895). E. United States.
85. *R. confinis*, Zetterstedt, Dipt. Scand. Vol. 11, p. 4271 (1852); Wahlberg, Sweden.
 Ent. Tidskr. Vol. 31, p. 57 (1910).
86. *R. conjuncta* Loew, Berl. Ent. Zeitschr. Vol. 5, p. 336 : Cent. 1, No. 56 E. United States.
 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895).
87. *R. conservativa*, Malloch, Report Canad. Arct. Exped. 1913-18, Dipt. Arctic America.
 p. 48 (1919).
88. *R. cophas*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 499 (1849). New York.
89. *R. coracina*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3037 (1849); Strobl, N. & C. Europe.
 Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 51 (1892); Frey,
 Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Wahlberg, Ent. Tidskr. Vol. 31, p. 57 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 28 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 20 (1919); Frey, Notulæ Ent. Vol. 2, p. 67 [1922] (*Dasyrhamphomyia*).
fuscipennis, Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 170 (1861); Frey,
 Acta Soc. Sc. Fenn. Helsingfors, Vol. 30 (9), p. 20 (1908).
90. *R. * corrupta*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 114, pl. 8, Baltic Amber.
 f. 11, 12 (1908).
91. *R. corvina*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 334 : Cent. 1, No. 51 North America.
 [1861] (*cervina*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 418
 (1895); Proc. Wash. Acad. Sc. Vol. 2, p. 414 (1900).
92. *R. costata*, Zetterstedt, Dipt. Scand. Vol. 1, p. 431 (1842); Verrall, Ent. N. Europe.
 Mag. London, Vol. 30, p. 140 (1894); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910).
simulum, Nowicki, Verh. Naturf. Ver. Brünn, Vol. 6, p. 81, pl. 2, f. 3
 (1867).
tibiella, Zetterstedt, Dipt. Scand. Vol. 1, p. 432, obs (1842); ibidem, Vol. 8,
 p. 3044 (1849); Loew, Jahrb. Ges. Krakau, Vol. 41, p. 12 (1870);
 Lundbeck, Dipt. Danica, Vol. 3, p. 57 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 22 (1913); Notulæ Ent. Vol. 2, p. 40
 [1922] (*Pararhamphomyia*).
93. *R. crassicauda*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 57 C. & S. Europe.
 (1892); Frey, Notulæ Ent. Vol. 2, p. 45 [1922] (*Pararhamphomyia*).
94. *R. crassimana*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 34, p. 199 Styria.
 (1898); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 24
 (1919) ♂.
95. *R. crinita*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 118 (1887); Strobl, C. Europe.
 Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 53 (1892);
 Frey, Notulæ Ent. Vol. 2, p. 75 (1922).
96. *R. * crinitarsis*, Loew, Bernsteina fauna, p. 41 (1850); Giebel, Ins. Vorwelt, Baltic Amber, Lower
 p. 208 (1856); Meunier, Miscell. Ent. Vol. 7, p. 14 [1899] (*Empis*
 or *Rhamphomyia*); Ann. Sc. Nat. (Zool.), Vol. 7, p. 92, 116, pl. 9,
 f. 13, pl. 10, f. 1 (1908).
97. *R. culicina*, Fallen, Empid. Suec. p. 28 [1816] (*Empis*); Meigen, Syst. N. & C. Europe.
 Beschr. Vol. 3, p. 52 (1822); Macquart, Dipt. N. France, Vol. 3,
 p. 134 (1827); Hist. Nat. Dipt. Vol. 1, p. 337 (1834); Curtis, Brit. Ent. Vol. 8, p. 517 (1834); Zetterstedt, Fauna Ins. Lappon. p. 568
 (1838); Dipt. Scand. Vol. 1, p. 421 (1842); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1,
 p. 173 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99, 100 (1862);

- Siebke, Nyt. Mag. Naturvid. Vol. 12, p. 154 (1864); ibidem, Vol. 14, p. 386, 400 (1866); Becker, Ent. Zeitschr. Vol. 31, p. 111 (1887); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 46 (1893); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 21 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 67 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 59 (1910); Verrall, Ent. Mo. Mag. London, Vol. 48, p. 23 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 10 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A. 6, p. 15 (1919); Frey, Notulæ Ent. Vol. 2, p. 5 [1922] (*Holocera*).
?avida, Harris (not Coquillett), Engl. Ins. p. 151, pl. 44, f. 10 [1782] (*Empis*).
?rufipes, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 231 part [1804] (*Empis*).
98. *R. curvinervis*, Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 84 (1915); St. Moritz.
Frey, Notulæ Ent. Vol. 2, p. 75 (1922).
99. *R. curvipes*, Coquillett, Invert. Pacif. Vol. 1, p. 24 (1904). California, Nevada.
100. *R. curvula*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 22 (1913); Notulæ Ent. Vol. 2, p. 41 [1922] (*Pararhamphomyia*). N. & C. Europe.
101. *R. cyanogaster*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, Mexico.
p. 371 (1901); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91 p. 322 (1909).
102. *R. dana*, Walker, List Dip. Brit. Mus. Vol. 3, p. 502 (1849). Canada.
103. *R. daria*, Walker, ibidem, Vol. 3, p. 503 (1849). New York.
104. *R. debilis*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 330: Cent. 1, No. 45 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895). Saskatchewan.
105. *R. dentata*, Oldenberg, Ann. Mus. Hungar. Vol. 8, p. 344 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 14 (1913); Ent. Tidskr. 1914, p. 78 (1914); Not. Ent. Vol. 2, p. 35, 36 [1922] (*Pararhamphomyia*). C. & N. Europe.
106. *R. dentipes*, Zetterstedt, Dipt. Scand. Vol. 1, p. 397 (1842); ibidem, Vol. 8, p. 3035 (1849); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 169 (1861); Beling, Arch. Naturg. Berlin, Vol. 48, p. 217 (1882); Becker, Berl. Ent. Zeischr. Vol. 31, p. 112 (1887); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 47 (1892); ibidem, Vol. 34, p. 198 (1898); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 64, f. 20 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 13 (1913); Oldenberg, Arch. Naturg. Vol. 83, A. 6, p. 17 (1919); Frey, Notulæ Ent. Vol. 2, p. 35, 36 [1922] (*Pararhamphomyia*). C. & N. Europe.
107. *R. dimidiata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 325: Cent. 1, No. 36 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895); Malloch, Illinois Labor. Nat. Hist. Bull. Urbana, Vol. 12, Art. 3, p. 401 [1917] (larva, pupa). E. United States.
108. *R. discoidalis*, Becker, Wien. Ent. Zeit. Vol. 8, p. 81, pl. 1, f. 6 (1889); ibidem, Vol. 10, p. 293 (1891); Frey, Notulæ Ent. Vol. 2, p. 77 (1922). C. Europe, Alps.
- Sancti-Mauritii*, Strobl (not Becker), Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 52 (1892), Vol. 46, p. 61 (1909).
109. *R. dispar*, Zetterstedt, Fauna Ins. Lappon. p. 570 (1838); Dipt. Scand. Vol. 1, p. 419 (1842); ibidem, Vol. 13, p. 5020 (1859); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 173 (1861); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 34, p. 200 (1898); Frey, Acta Soc. Sc. C. & N. Europe.

- Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 59 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors Vol. 37 (3), p. 20 (1913); Notulæ Ent. Vol. 2, p. 37, 38 [1922] (*Pararhamphomyia*).
aperta, Zetterstedt, Dipt. Scand. Vol. 13, p. 5021 (1859); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 116 (1887).
fuliginella, Zetterstedt, Dipt. Scand. Vol. 1, p. 420 (1842), Vol. 8, p. 3041 (1849), Vol. 11, p. 4272 (1852), Vol. 13, p. 5021 (1859); Siebke, Nyt Mag. Naturvid, Vol. 12, p. 109 (1864); Mik, Fauna Hernstein, Vol. 2 (2), p. 56 (1885); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 116 (1887); Wahlgren, Tidskr. Vol. 31, p. 59 (1910).
fuscula, Zetterstedt, part, Fauna Ins. Lappon. p. 571, part [1838]; Dipt. Scand. Vol. 1, p. 404, part (1842).
griseola, Zetterstedt, part, Fauna Ins. Lappon. p. 571, part [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 418, part (1842).
110. *R. disparilis*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 415 (1900). Alaska.
111. *R. *distans*, Loew, Bernsteinfauna, p. 41 (1850); Giebel, Ins. Vorwelt, p. 208 (1856). Baltic Amber.
112. *R. diversa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 23, p. 611 (1901). E. United States.
113. *R. diversipennis*, Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 19, pl. 1, f. 14, 17 (1901); Frey, Notulæ Ent. Vol. 2, p. 43 [1922] (*Pararhamphomyia*). N. Siberia.
114. *R. dolichocera*, Bezzii, Ann. Mûs. Hungar. Vol. 3, p. 431 (1905); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 324, 326 (1909). Peru.
115. *R. dolichoptera*, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 328 (1909). Bolivia.
116. *R. dorsata*, Becker, Mém. Acad. Sc. Pétrograd, Vol. 28, No. 7, p. 56 (1915). Arctic Ural.
117. *R. duplicitis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 424 (1895). California.
118. *R. ecetra*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 500 (1849). Georgia.
119. *R. effera*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 427 (1895). United States.
120. *R. empidiformis*, Becker, Bull. Mus. Hist. Nat. Paris, 1909, p. 116 (1909); Ann. Soc. Ent. France, Vol. 79, p. 25 (1910). E. Africa.
121. *R. *enena*, Cockerell, Proc. U. S. Nat. Mus. Vol. 59, p. 30 (1921). Eocene, Colorado.
122. *R. Erberi*, Mik, Jahrb. Akad. Gym. Wien, 1878, p. 22, f. 10, 11 (1878); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 16 (1919). C. Europe.
123. *R. erinacioides*, Malloch, Report Canad. Arct. Exped. 1913-18, Dipt. Alaska; Arctic America p. 45, f. 6 (1919).
124. *R. *errabunda*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 116, pl. 9, f. 14, Vol. 10, p. 2-4 (1908). Baltic Amber.
125. *R. erythrophthalma*, Meigen, Syst. Beschr. Vol. 6, p. 340 (1830); Schiner, Fauna Dipt. Austr. Vol. 1, p. 100 (1862); Strobl, Mitteil. Bosn. Herceg. Sarajevo, Vol. 8, p. 564 (1900). C. Europe.
126. *R. eupterota*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 41 (1873); Frey, Hungary. Notulæ Ent. Vol. 2, p. 37, 38 [1922] (*Pararhamphomyia*).
127. *R. exigua*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 201 : Cent. 2, No. 32 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895). E. & C. United States.
128. *R. expulsa*, Walker, Trans. Ent. Soc. Lond. n. s. Vol. 4, p. 148 (1857); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 417 (1895). E. United States.
129. *R. fascipennis*, Zetterstedt, Fauna Ins. Lappon. p. 564 (1838); Dipt. Scand. Vol. 1, p. 431 (1842); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 19 (1913); Notulæ Ent. Vol. 2, p. 37, 38 [1922] (*Pararhamphomyia*). Scandinavia.

130. *R. ferruginea*, Fabricius, Syst. Antl. p. 146 [1805] (*Hybos*); Wiedemann, South America.
Zool. Mag. Berlin, p. 59, pl. 2, f. 2 [1817] (*Macrostomus*); Anal. Ent. 27 (1824); Auss. Zweifl. Ins. Vol. 2, p. 9 (1830); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 321 (1909).
131. *R. ferruginea*, Meigen, Syst. Beschr. Vol. 3, p. 60 (1822); Macquart, C. Europe
Hist. Nat. Dipt. p. 334 (1834).
132. *R. ficana*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 501 (1849). Canada.
133. *R. filata*, Zetterstedt, Dipt. Scand. Vol. 1, p. 392 (1842); Verrall, Ent. Mag. London, Vol. 30, p. 140 (1894); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 21 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 19 (1919); Frey, Notulæ Ent. Vol. 2, p. 39, 40 [1922] (*Pararhamphomyia*).
134. *R. filicanda*, Lundbeck, Conspectus Fauna Groenl. Vol. 2, p. 608, f. (1918). Greenland.
135. *R. fimbriata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 429 (1895). California.
136. *R. flava*, Fallen, Empid. Suec. p. 30 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 59 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 335 (1834); ?Zetterstedt, Fauna Ins. Lappon. p. 572 (1838) (*Rhamphomyza*); Loew, Isis, Vol. 7, p. 549 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 438 (1842); Boitard, Man. Ent. Vol. 3, p. 319 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 503 (1849); Ins. Brit. Vol. 1, p. 110 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 97 (1862); Siebke, Nyt. Mag. Naturv. Vol. 14, p. 400 (1866); Verrall, Ent. Mag. London, Vol. 19, p. 224 (1883); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 45 (1892); Lundbeck, Dipt. Danica, Vol. 3, p. 65 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 59, f. 6 (1910); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 15 (1919); Frey, Notulæ Ent. Vol. 2, p. 6 [1922] (*Holoclera*).
var. *bistriata*, Strobl, Mitteil. Steiermark, Graz, Vol. 46, p. 58 (1909). C. Europe.
137. *R. flavicoxa*, Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 56 (1851); Schiner, Germany.
Fauna Dipt. Austr. Vol. 1, p. 101 (1862).
138. *R. flavipes*, Matsumura, Journal Coll. Sc. Sapporo, Vol. 4, p. 67 (1911); Sachalin.
—. *R. flavipes*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 264 (1829), no description.
139. *R. flavirostris*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 501 (1849); Canada, Alaska.
Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 414 (1900).
140. *R. flaviventris*, Macquart, Dipt. N. France, Vol. 3, p. 134 (1827); Hist. Nat. Dipt. Vol. 1, p. 337 (1834); Meigen, Syst. Beschr. Vol. 7, p. 91 (1838); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 109 (1887); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 46 (1892); ibidem, Vol. 46, p. 58 (1909); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 15 (1919); Frey, Notulæ Ent. Vol. 2, p. 6 [1922] (*Holoclera*).
var. *bivittata*, Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 46 (1892). Alps.
var. *univittata*, Strobl, ibidem, Vol. 29, p. 46 (1892). Alps.
141. *R. flexicauda*, Zetterstedt, Dipt. Scand. Vol. 1, p. 403 (1842); ibidem, Vol. 13, p. 5014 (1859); Wahlgren, Ent. Tidskr. Vol. 31, p. 57 (1910). Norway.
142. *R. flexuosa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 433 (1895). Colorado.
143. *R. forcipata*, Linnaeus, Fauna Suec. p. 466 [1761] (*Empis*); Fabricius, Syst. Ent. p. 801 [1775] (*Empis*); Sulzer, Abgek. Gesch. Insekt., p. 221, pl. 28, f. 16 Europe.

- [1776] (*Empis*); Fabricius, Spec. Ins. Vol. 2; p. 471 [1781] (*Empis*); Mant. Ins. Vol. 2, p. 366 [1788] (*Empis*); Gmelin, Syst. Nat. Vol. 5, p. 2889 [1790] (*Empis*); Olivier, Encycl. Méthod. Vol. 6, p. 387 [1791] (*Empis*); Fabricius, Ent. Syst. Vol. 4, p. 404 [1794] (*Empis*); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 232, b. [1804] (*Empis*); Fabricius, Antl. p. 139 [1805] (*Empis*); Macquart, Lille, p. 163 [1823] (*Empis*); Zetterstedt, Dipt. Sc. Vol. 1, p. 414 [1842] (*Empis*).
appendiculata, Macquart, Dipt. N. France, Vol. 3, p. 132 (1827); Hist. Nat. Dipt. Vol. 1, p. 336 (1834); Meigen, Syst. Beschr. Vol. 7, p. 90 (1838); Gimmerthal, Bull. Soc. Nat. Mosc. Vol. 15, p. 666 (1842); Boitard, Man. Ent. Vol. 3, p. 319 (1843).
- . *R. formosa*, Loew, Bernsteinfauna, p. 41 (1850), no description; Giebel, Lower Oligocene. Ins. Vorwelt, p. 258 (1856), no description.
144. *R. frontalis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 199: Cent. 2, No. 28 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 416 (1895). Illinois.
145. *R. fulvolanata*, Frey, Notulæ Ent. Vol. 2, p. 7 [1922] (*Choreodromia*). Kamchatka.
146. *R. fumipennis*, Zetterstedt, Dipt. Scand. Vol. 13, p. 5019 (1859); Seibke, Nyt. Mag. Naturvid. Vol. 12, p. 154 (1864). Lapland.
147. *R. fumosa*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 327: Cent. 1, No. 39 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1895). E. United States.
148. *R. furcifer*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, p. 371 (1901); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 323 (1909). Mexico.
- . *R. fusca*, Stephens, Cat. Brit. Ins. Vol. 2, p. 264 [1829] (*Enicopteryx*) no description.
149. *R. fuscipennis*, Zetterstedt, Fauna Ins. Lappon. p. 567 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 402 (1842); ibidem, Vol. 8, p. 3036 (1849); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 50 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 25 (1913); Notulæ Ent. Vol. 2, p. 42 [1922] (*Pararhamphomyia*). C. & N. Europe.
- lugubrina*, Zetterstedt, Dipt. Scand. Vol. 1, p. 401 (1842), Vol. 8, p. 3036 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19) p. 54 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Strobl, Progr. Seitenst Vol. 14, p. 9 (1880); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910).
150. *R. fuscula*, Zetterstedt, Fauna Ins. Lappon. p. 571, part [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 404, part (1842); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 21 (1913); Notulæ Ent. Vol. 2, p. 39, 40 [1922] (*Pararhamphomyia*). N. W. Europe.
151. *R. galactodes*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 332 (1909). Bolivia.
152. *R. galactoptera*, Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 56 (1892); ibidem, Vol. 34, p. 200 (1897); Wien. Ent. Zeitschr. Vol. 18, p. 13 (1899); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 25 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 20 (1919); Frey, Notulæ Ent. Vol. 2, p. 41 [1922] (*Pararhamphomyia*). C. & S. Europe, Finland.
- lacteipennis*, Strobl (not Zetterstedt), Progr. Seitenst. Vol. 14, p. 58 (1880).
153. *R. geniculata*, Bigot (not Meigen), Bull. Soc. Ent. France (6), Vol. 7, p. 142 (1887); Ann. Soc. Ent. France (6), Vol. 9, p. 134 (1889). California.
154. *R. geniculata*, Meigen, Syst. Beschr. Vol. 6, p. 340 (1830); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Verrall, Ent. Mag. London, C. Europe.

- Vol. 19, p. 224 (1883); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 112 (1887); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 19 (1919).
155. *R. gentilis*, Loew, Beschr. Eur. Dipt. Vol. 2, p. 242 (1871). Russia.
156. *R. gibba*, Fallen, Empid. Suec. p. 32 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 58 (1822), Vol. 7, p. 89 (1838); Zetterstedt, Dipt. Scand. Vol. 1, p. 437 (1842); ibidem, Vol. 8, p. 304 (1849); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 56 (1892); Vol. 34, p. 201 (1897); Verrall, Ent. Mag. London, Vol. 30, p. 140 (1894); Lundbeck, Dipt. Danica, Vol. 3, p. 77, f. 78 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 27 (1913); Notulæ Ent. Vol. 2, p. 45 [1922] (*Pararhamphomyia*). N. & C. Europe.
- holosericea*, Meigen, Syst. Beschr. Vol. 6, p. 339 (1838).
157. *R. gibbifera*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 306 (1906). Spain.
158. *R. gilvipes*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 332: Cent. 1, No. 48 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895). E. & C. United States.
159. *R. gilvipilosa*, Coquillett, ibidem, Vol. 18, p. 434 (1895). Illinois.
160. *R. glabra*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 328 : Cent. 1, No. 41 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1895); Frey, Notulæ Ent. Vol. 2, p. 68 (1922). E. & C. United States.
161. *R. glauca*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 416 (1900). Alaska.
162. *R. glaucella*, Zetterstedt, Dipt. Scand. Vol. 1, p. 405 (1842); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 15, part (1913). Scandinavia.
163. *R. gracilis*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 329 : Cent. 1, No. 43 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895); Frey, Notulæ Ent. Vol. 2, p. 68 (1922). E. United States.
164. *R. grallatrix*, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 326 (1909). Bolivia.
165. *R. grammoptera*, Frey, Notulæ Ent. Vol. 2, p. 70 (1922). Kamchatka.
166. *R. griseola*, Zetterstedt, Fauna Ins. Lappon. p. 571, part [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 418, part (1842), Vol. 13, p. 5020 (1859). N. Europe.
167. *R. griseonigra*, Brunetti, Rec. Indian Mus. Vol. 9, p. 29 (1913); Fauna Brit. India, Dipt. Vol. 1, p. 346, f. 28 (1920). W. Himalayas.
168. *R. guifilar*, Frey, Notulæ Ent. Vol. 2, p. 8 [1922] (*Choreodromia*). N. Europe.
- anomalina*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3) p. 11, part (1913).
169. *R. Hambergi*, Frey, Naturw. Unters. Sarekgebirges, Vol. 4 (6), p. 684 (1917); Notulæ Ent. Vol. 2, p. 65 [1922] (*Dasyrhamphomyia*). N. Europe.
170. *R. Helléni*, Frey, Notulæ Ent. Vol. 2, p. 39, 40 [1922] (*Pararhamphomyia*). Russia.
171. *R. Hershelli*, Malloch, Report Canad. Arct. Exped. 1913-18, Dipt. 47 (1919). Yukon Territory.
172. *R. heterochroma*, Bezzii, Term. Fuezet. Vol. 21, 439 (1898); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 16 (1919); Frey, Notulæ Ent. Vol. 2, p. 6 [1922] (*Holoclera*). C. & S. Europe.
173. *R. hilariformis*, Frey, Notulæ Ent. Vol. 2, p. 35 [1922] (*Pararhamphomyia*). N. Siberia.
174. *R. himalayana*, Brunetti, Rec. Indian Mus. Vol. 9, p. 28 (1913); Fauna Brit. India, Dipt. Vol. 1, p. 346 (1920). W. Himalayas.
175. *R. hirtipes*, Loew, Ent. Zeitschr. Vol. 8, p. 80 : Cent. 5, No. 59 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 417 (1895). New Hampshire.
176. *R. hirtula*, Zetterstedt, Dipt. Scand. Vol. 1, p. 421, obs. (1842); Lundbeck, Dipt. Groenl. Vol. 1, p. 298 (1898); Collin, Ent. Mag. London (2), Vol. 24, p. 105 (1913); Frey, Notulæ Ent. Vol. 2, p. 42 [1922] (*Pararhamphomyia*). Greenland, England.

177. *R. Hovgaardii*, Holmgren, Nov. Sp. Ins. Nov. Seml. p. 21 (1881); Ent. Nova Zemla. Tidskr. Vol. 4, p. 162 (1883); Frey, Notulæ Ent. Vol. 2, p. 66 [1922] (*Dasyrhamphomyia*).
178. *R. hyalina*, Brullé, Expéd. Morée (Zool.) Vol. 3 (1), p. 299 (1832). Greece.
179. *R. hybotina*, Zetterstedt, Fauna Ins. Lappon. p. 571 (1838); Dipt. Europe. Scand. Vol. 1, p. 412 (1842); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1851); Loew, Zeitschr. Ges. Naturw. Vol. 10, p. 100 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 46 (1892); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 123 (1899); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 21 (1908); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 46, p. 58 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 57 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 9 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 14 (1919); Frey, Notulæ Ent. Vol. 2, p. 4 [1922] (*Lundstroemiella*). *tenuicornis*, Zetterstedt, Fauna Ins. Lappon. p. 571 [1838] (*Rhamphomyza*). var. *alpina*, Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 47 [1892]. Alps.
- var. *australis*, Frey, Notulæ Ent. Vol. 2, p. 5 [1922] (*Lundstroemiella*). C. Europe.
180. *R. *hypolitha*, Cockerell, Proc. U. S. Mus. Nat. Vol. 52, p. 378, pl. 31, f. 8 (1917). Florissant, Miocene.
181. *R. ignobilis*, Zetterstedt, Dipt. Scand. Vol. 13, p. 5015 (1859). Lapland.
182. *R. impedita*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 201 : Cent. 2, No. 31 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 416 (1895). C. & E. United States.
183. *R. incompleta*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 17 : Cent. 3, No. 31 (1863); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, 414 (1895). E. United States.
184. *R. infumata*, Wiedemann, Aussereur. Zweifl. Ins. Vol. 2, p. 9 (1830). Locality unknown.
185. *R. infuscata*, Meigen, Syst. Beschr. Vol. 3, p. 53, pl. 23, f. 4 (1822); Stephens, Cat. Brit. Ins. Vol. 2, p. 264 [1829] (*Enicopteryx*); Westwood, Introd. Classif. Ins. Vol. 2, Synops. p. 131 [1840] (*Enicopteryx*). C. Europe.
186. *R. insecta*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 426 (1895). Texas.
187. *R. insignis*, Loew, Beschr. Eur. Dipt. Vol. 2, p. 246 (1871). Siberia.
188. *R. *insolita*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 118, pl. 10, f. 11 (1908). Baltic Amber.
189. *R. intermedia*, Frey, Notulæ Ent. Vol. 2, p. 35, 36 [1922] (*Pararhamphomyia*). Europe.
190. *R. *involuta*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 115, pl. 10, f. 7-9 (1908). Baltic Amber.
191. *R. irregularis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 81 : Cent. 5, No. 60 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895); Proc. Wash. Acad. Sc. Vol. 2, p. 414 (1900). N. America.
192. *R. kamtschatica*, Frey, Notulæ Ent. Vol. 2, p. 7 [1922] (*Choreodromia*). Kamchatka.
193. *R. hainensis*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (10), p. 7 (1913); Notulæ Ent. Vol. 2, p. 42 [1922] (*Pararhamphomyia*). N. Russia.
194. *R. Kjellmanii*, Holmgren, Nov. Sp. Ins. Nov. Zeml. p. 22 (1880); Ent. Tidsks. Vol. 4, p. 163 (1883). Nova Zemla.
195. *R. klekovacensis*, Strobl, Glasnik. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 403 (1898); Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 565 (1900). Bosnia.
196. *R. lavigata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 325 : Cent. 1, No. 37 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895). E. United States, Canada.
197. *R. lavigipes*, Fallen, Empid. Suec. p. 27 [1816] (*Empis*); ? Meigen, Syst. Europe.

- Beschr. Vol. 3, p. 49 (1822);? Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 22 (1840); Isis, Vol. 7, p. 550 (1840);? Zetterstedt, Dipt. Scand. Vol. 1, p. 398 (1842); Gimmerthal, Bull. Soc. Nat. Moscou, Vol. 20, p. 163 (1847); Walker, List Dipt. Brit. Mus. Vol. 3, p. 499 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3035 (1849); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 29 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 22 (1919); Frey, Notulæ Ent. Vol. 2, p. 69 (1922).
- conformis*, Kowarz, Verh. Zool.-bot. Gez. Wien, Vol. 17, p. 321 (1867); Lundbeck, Dipt. Danica, Vol. 3, p. 51, f. 15, 16 (1910); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 21 (1919).
- pseudotrilineata*, Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 49 (1892); Glasn. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 402 (1898); Mitteil. Bosn. Herzeg. Sarajevo, Vol. 7, p. 564 (1900).
198. *R. latifrons*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 29 Finland. Lapland. (1913); Notulæ Ent. Vol. 2, p. 72 (1922).
199. *R. leptopus*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 41 (1872); Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 124 (1899).
200. *R. leucophenga*, Bezzii, Ann. Mus. Hungar. Vol. 3, p. 430 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 322 (1909).
201. *R. leucoptera*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 340 : Cent. 1, No. 62 E. United States. (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895).
202. *R. limata*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 417 (1900). Alaska.
203. *R. limbata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 338 : Cent. 1, No. 60 North America. (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 388 (1895); Wash. Acad. Sc. Vol. 2, p. 414 (1900).
- argentea*, Bigot, Bull. Soc. Ent. France (5), Vol. 10, p. 47 [1880] (*Megacyttarus*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 388 [1895] (*Megacyttarus*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 350 [1902] (*Megacyttarus*).
204. *R. limbipennis*, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 329 Peru, Bolivia. (1909).
205. *R. liturata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 339 : Cent. 1, No. 61 E. United States. (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 413 (1895).
206. *R. lividiventris*, Zetterstedt, Fauna Ins. Lappon. p. 528 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 406 (1842); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 21 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 13 (1913); Notulæ Ent. Vol. 2, p. 34 [1922] (*Pararhamphomyia*).
207. *R. longefilata*, Strobl, Mem. Soc. Esp. Nat. Hist. Vol. 3, p. 305 (1906); Spain. Frey, Notulæ Ent. Vol. 2, p. 5 [1922] (*Lundstroemiella*).
208. *R. longestylata*, Frey, Naturw. Unters. Sarekgebirges, Vol. 4 (6), p. 682 N. Europe. (1917); Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (10), p. 8 (1913); Notulæ Ent. Vol. 2, p. 44 [1922] (*Pararhamphomyia*).
209. *R. longicauda*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 326 : Cent. 1, No. 38 E. United States. (1861); Glover, Manusc. Notes, p. 44, pl. 3, f. 23, 24 (1874); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1895); Kellogg, Amer. Ins. p. 334, f. 470 (1905); Frey, Notulæ Ent. Vol. 2, p. 44 [1922] (*Pararhamphomyia*).
210. *R. longicornis*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 332 : Cent. 1, No. 47 E. United States. (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 417 (1895).
211. *R. longipennis*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 331 : Cent. 1, No. 46 E. United States. (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895).

212. *R. longipes*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 231 C. Europe.
 [1804] (*Empis*); Syst. Beschr. Vol. 3, p. 55, pl. 23, f. 3 (1822);
 Macquart, Mém. Soc. Sc. Lille, p. 164 [1823] (*Empis*); Dipt. N.
 France, Vol. 3, p. 133 (1827); Hist. Nat. Dipt. p. 339 (1834); Boitard,
 Man. Ent. Vol. 3, p. 320 (1843); Walker, List Dipt. Brit. Mus.
 Vol. 3, p. 501 (1849); Glover, Manuscr. Notes, p. 44, pl. 11, f. 3
 (1874); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29
 p. 57 (1892); Frey, Notulæ Ent. Vol. 2, p. 45 [1922] (*Pararhamphomyia*).
globifera, Strobl, Gym. Progr. Seitenst. Vol. 14, p. 58 (1880).
213. *R. loripedis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 419 (1895). California.
214. *R. lucidula*, Zetterstedt, Dipt. Scand. Vol. 1, p. 422 (1842); ? Scholz,
 Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1855); Schiner, Fauna
 Dipt. Austr. Vol. 1, p. 100 (1862); Wahlgren, Ent. Tidskr. Vol. 31
 p. 59 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3)
 p. 27 (1913); Notulæ Ent. Vol. 2, p. 45 [1922] (*Pararhamphomyia*).
plumifera, Bonsdorff, Finl. tvåv. ins. Dipt Vol. 1, p. 175 (1861)
215. *R. luctifera*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 333 : Cent. 1, p. 50 New York.
 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 417 (1895).
216. *R. luctuosa*, Loew, Berl. Ent. Zeitschr. Vol. 16, p. 114 : Cent. 10, California.
 No. 290 (1872); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414
 (1895).
lugens, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 200 : Cent. 2, No. 30 (1862).
217. *R. luridipennis*, Nowicki, Verh. Naturf. Ver. Brünn, Vol. 6, p. 80 (1869);
 Strobl, Mitteil. Naturf. Steiermark, Graz, Vol. 29, p. 52 (1892);
 Frey, Notulæ Ent. Vol. 2, p. 77 (1922).
218. *R. luteiventris*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 79 : Cent. 5, No. 57 E. United States.
 (1864); Cocquillet, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895);
 Frey, Notulæ Ent. Vol. 2, p. 34 [1922] (*Pararhamphomyia*).
219. *R. macerrima*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 327 Bolivia.
 (1909).
220. *R. macilenta*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 78 : Cent. 5, No. 55 E. United States.
 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895).
221. *R. macrura*, Loew, Beschr. Eur. Dipt. Vol. 2, p. 247 (1871). Siberia.
222. *R. maculipennis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 399 (1842); Wahlgren,
 Ent. Tidskr. Vol. 31, p. 55 (1910); Frey, Acta Soc. Sc. Fenn.
 Helsingfors, Vol. 37 (3), p. 11 (1913); Notulæ Ent. Vol. 2, p. 7, 9
 [1922] (*Choreodromia*).
dissimilis, Zetterstedt, Dipt. Scand. Vol. 8, p. 303 (1849); Lundbeck, Dipt.
 Danica, Vol. 3, p. 43, f. 9 (1910); Wahlgren, Ent. Tidskr. Vol. 31,
 p. 55 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912).
223. *R. magellensis*, Bezzi-Frey, Notulæ Ent. Vol. 2, p. 4 [1922] (*Lundstroemella*). Alps, Italy.
224. *R. mallos*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 502 (1849). Canada.
225. *R. manca*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 427 (1895);
 Howard, Proc. Wash. Acad. Sc. Vol. 2, p. 561 (1900). E. United States.
226. *R. marginata*, Fabricius, Mant. Ins. Vol. 2, p. 364 [1787] (*Empis*); Gmelin,
 Syst. Nat. Vol. 5, 2889 [1790] (*Empis*); Olivier, Encycl. Mitteil.
 Vol. 6, p. 387 [1791] (*Empis*); Fabricius, Syst. Ent. Vol. 4, p. 403
 [1794] (*Empis*); Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1,
 p. 228, pl. 11, f. 29 [1804] (*Empis*); Fabricius, Syst. Antl. p. 138
 [1805] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 43 (1822); Macquart,
 Hist. Nat. Vol. 1, p. 336 (1834); Walker, List Dipt. Brit. Mus.
 Vol. 3, p. 498 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19),
 p. 54 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 5017 (1859);

- Loew, Wien. Ent. Monatschr. Vol. 8, p. 122 (1864); Glover, Manusc. Notes, p. 44 (1874); Leunis, Synop. Zool. Vol. 2, p. 403 (1886); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 18 (1919).
- latipennis*, Meigen, Syst. Beschr. Vol. 3, p. 44 (1822); Schiner, Vol. 1, p. 98 (1862); Verh. Zool.-bot. Ges. Wien, Vol. 22, p. 74 (1872).
- platyptera*, Panzer, Fauna Germ. p. 24 [1794] (*Empis*); Lefebre, Ann. Soc. Ent. France (2), Vol. 9, p. 125, pl. 4, f. 1-4 [1851] (*Empis*); Lucas, Bull. Ent. Soc. France (3), Vol. 3, p. 243 [1859] (*Empis*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 97 (1862); Verh. Zool.-bot. Ges. Wien, Vol. 22, p. 74 (1872); Neuhaus, Dipt. March. p. 73 (1886); Kleine, Zeitschr. Naturwiss. Jena, Vol. 31, p. 188 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 38 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 49 (1910); Frey, Notulæ Ent. Vol. 2, p. 38 [1922] (*Pararhamphomyia*).
227. *R. *media*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 93, 119, pl. 9, f. 17, pl. 10, f. 13 (1908). Baltic Amber.
228. *R. melania*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 119 (1887); Oldenberg, Arch. Naturg. Berlin, Vol. 82, 1, p. 161 (1916); Frey, Notulæ Ent. Vol. 2, p. 76 (1922). Alps.
229. *R. metatarsata*, Zetterstedt, Fauna Ins. Lappon. p. 569 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 403 (1842); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 17 (1909); ibidem, Vol. 37 (3), p. 12 (1913). Scandinavia.
230. *R. micans*, Oldenberg, Arch. Naturg. Berlin, Vol. 80, A, 9, p. 69 (1915). Germany.
231. *R. micrargyra*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 333 [1909] (*micrargyra*). Peru.
232. *R. minutus*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 502 (1849); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895); Proc. Wash. Acad. Sc. Vol. 2, p. 414 (1900). Canada.
233. *R. mirifica*, Frey, Notulæ Ent. Vol. 2, p. 66 [1922] (*Dasyrhamphomyia*). Kamchatka.
234. *R. modesta*, Wahlberg, Oefv. Vet. Akad. Förh. Stockholm, p. 107 (1844); Zetterstedt, Dipt. Scand. Vol. 8, p. 3039 (1849); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 13 (1913); Notulæ Ent. Vol. 2, p. 35, 36 [1922] (*Pararhamphomyia*). Lapland.
235. *R. monstrosa*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 339, f. 3 (1909). Peru.
236. *R. montana*, Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 87 (1915); Frey, Notulæ Ent. Vol. 2, p. 76 (1922). Alps.
237. *R. morio*, Zetterstedt, Fauna Ins. Lappon. p. 565 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 416 (1842), Vol. 8, p. 3040 (1849); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 34 (1913); Notulæ Ent. Vol. 2, p. 77 (1922). Norway.
238. *R. mutabilis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 198: Cent. 2, No. 26 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1895); Webster, Canad. Ent. Vol. 30, p. 18 (1898). C. & E. United States.
239. *R. nana*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 341: Cent. 1, No. 64 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895). E. United States.
240. *R. Nasoni*, Coquillett, ibidem, Vol. 18, p. 423 (1895); Frey, Notulæ Ent. Vol. 2, p. 34 [1922] (*Pararhamphomyia*). C. United States.
241. *R. nigricans*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 80: Cent. 5, No. 58 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 416 (1895). E. United States.
242. *R. nigricauda*, Becker, Ann. Mus. Zool. St. Pétersb. Vol. 12, p. 314 (1907). Tibet.
243. *R. nigripennis*, Fabricius, Ent. Syst. Vol. 4, p. 407 [1794] (*Empis*); Meigen, Classif. Eur. Zweifl. Ins. Vol. 1, p. 234 [1804]. Europe.

(*Empis*); Fabricius, Syst. Antl. p. 144 [1805] (*Tachydromia*); Fallen, Empid. Suec. p. 30 [1816] (*Empis*); Zetterstedt, Fauna Ins. Lappon. p. 569 (1838); Dipt. Scand. Vol. 1, p. 424 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 502 (1849); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 173 (1861); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 154 (1864); ibidem, Vol. 14, p. 386 (1866); Lundbeck, Dipt. Danica, Vol. 3, p. 6, 71, f. 21 (1910); Collin, Ent. M. Mag. London (2). Vol. 24, p. 105 (1913); Frey, Acta Soc. Sc. Fenn. Helsingfors Vol. 37 (3), p. 10 (1913); Notulæ Ent. Vol. 2, p. 5 [1922] (*Holoclera*).
obscuripennis, Meigen, Syst. Beschr. Vol. 6, p. 340 (1830); Zetterstedt, Dipt. Scand. Vol. 1, p. 426 (1842); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 59 (1892).
umbripennis, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 21 (1908).

244. *R. nigripes*, Fabricius, Ent. Syst. Vol. 4, p. 405 [1794] (*Empis*); Meigen, C. & N. Europe.

Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 229, f. 31 (1804); Fabricius, Syst. Antl. p. 141 [1805] (*Empis*); Billberg, Enumer. Ins. p. 120 (1820); Meigen, Syst. Beschr. Vol. 3, p. 48, pl. 23, f. 2 (1822); Curtis, Brit. Ent. Vol. 8, p. 517 (1834); Macquart, Hist. Nat. Dipt. Vol. 1, p. 336 (1834); Zetterstedt, Fauna Ins. Lappon. p. 562 (1838); Dipt. Scand. Vol. 1, p. 389 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 499 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3030 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 56 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 108 (1851); Bonsdorff, Finl. tvåv. Ins. Vol. 1, p. 168 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98 (1862); Glover, Manusc. notes, p. 44, pl. 11, f. 2 (1874); Jaroschewsky, Arb. Ges. Naturf. Univ. Kharkow, Vol. 11, p. 353 (1877); Verrall, Ent. Mag. London, Vol. 19, p. 224 (1883); Neuhaus, Dipt. March. p. 73 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 113 (1887); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 49 (1893); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 18 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 6, 39, f. 6, 7 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 55 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 11 (1913); Oldenberg, Arch. Naturg. Vol. 83, A, 6, p. 16 (1919); Frey, Notulæ Ent. Vol. 2, p. 7, 9 [1922] (*Choreodromia*).

crassirostris, Fallen, Empid. Suec. p. 31 [1816] (*Empis*).

vicana, Harris, Engl. Ins. Vol. 151, pl. 44, f. 9 [1782] (*Empis*).

245. *R. nigrata*, Bigot, Bull. Soc. Ent. France (6), Vol. 7, p. 142 (1887); Ann. California.

Soc. Ent. France (6), Vol. 9, p. 133 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 410 (1895).

246. *R. nigrata*, Zetterstedt, Fauna Ins. Lappon. p. 567 (1838); Dipt. Sc. Vol. 1, p. 414 (1842); Staeger, Kröyer Nat. Tidsskr. Kjöbenhavn, (2), Vol. 1, p. 357 (1845); Zetterstedt, Dipt. Scand. Vol. 8, p. 3037 (1849); Schiödte, Till. Rin. Grönl. p. 68 (1870); Holmgren, Oefv. Vet. Akad. Förh. Stockholm, p. 29 (1872); Lundbeck, Vidensk. Meddel. Kjöbenhavn, p. 297 (1898); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 59 (1909); Frey, Notulæ Ent. Vol. 2, p. 66 [1922] (*Dasyrhamphomyia*).

? *borealis*, Otto Fabricius, Fauna Groenl. p. 211 [1780] (*Empis*).

247. *R. nigriventris*, Macquart, Dipt. Exot. Suppl. Vol. 1, p. 96 (1846); Bezzi, Brazil.

Nova Acta Akad. Naturf. Halle, Vol. 91, p. 322 (1909).

248. *R. nigromaculata*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stutt- C. Europe.
gart, Vol. 1, p. 53 (1840).

249. *R. nitida*, Macquart, Dipt. N. France, Vol. 3, p. 135 (1827); Hist. Nat. Dipt. Vol. 1, p. 337 (1834); Meigen, Syst. Beschr. Vol. 7, p. 92 (1838); Zetterstedt, Dipt. Scand. Vol. 1, p. 401, note (1842).
250. *R. nitidicollis*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 26 (1913); Notulæ Ent. Vol. 2, p. 44 [1922] (*Pararhamphomyia*).
251. *R. nitidivittata*, Macquart, Dipt. Exot. Suppl. Vol. 1, p. 97 (1846). Texas.
252. *R. nitidolineata*, Frey, Acta Soc. Sc. Fenn. Helsingfors, p. 37 (10), p. 8 (1913); Notulæ Ent. Vol. 2, p. 69 (1822). N. Russia, Siberia.
253. *R. nitidula*, Zetterstedt, Dipt. Scand. Vol. 1, p. 400 (1842); ibidem, Vol. 8, p. 3036 (1849); Bonsdorff, Finl. tvåv. Ins. Dipt. p. 170 (1861); Beling, Arch. Naturg. Berlin, Vol. 48, p. 216 (1882); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 57 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 30 (1913); Notulæ Ent. Vol. 2, p. 73 (1922).
- nitida*, Zetterstedt, Fauna Ins. Lappon. p. 566 [1838] (*Rhamphomyza*).
254. *R. niveipennis*, Zetterstedt, Fauna Ins. Lappon. p. 570 [1838] (*Rhamphomyza*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 171 (1861); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 34, p. 199 (1897); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 19 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 17 (1919); Frey, Notulæ Ent. Vol. 2, p. 37 [1922] (*Pararhamphomyia*).
- albichata*, Frey, Acta Soc. Sc. Fenn. Helsingfors, p. 31 (9), 20 (1908).
- lacteipennis*, Zetterstedt, Fauna Ins. Lappon. p. 570 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 410 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 171 (1861); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908).
255. *R. nodipes*, Fallen, Empid. Suec. p. 25 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 50 (1822); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 16 (1919); Frey, Notulæ Ent. Vol. 2, p. 7 [1922] (*Choreodromia*).
- spissirostris*, Fallen, Empid. Suec. p. 31 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 57 (1822); Zetterstedt, Fauna Ins. Lappon. p. 563 (1838); Dipt. Scand. Vol. 1, p. 390 (1842); ibidem, Vol. 8, p. 3031 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 54 (1861); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 168 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98 (1862); Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), 17, pl. 1, f. 9, 10 (1900); Frey, ibidem, Vol. 31 (9), p. 18 (1908); Lundbeck, Dipt. Dan. Vol. 3, p. 41, f. 8 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 55, f. 5 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 11 (1913).
256. *R. Nordquistii*, Holmgren, Nov. Sp. Ins. Nov. Zeml. p. 23 (1881); Ent. Nova Zemla. Tidskr. Vol. 4, p. 164 (1883).
257. *R. novecarolina*, Beutenmueller, Insecutor Insc. Mens. Vol. 1, p. 130 N. Carolina. (1913).
258. *R. nox*, Oldenberg, Arch. Naturg. Berlin, Vol. 82, 1, p. 155 (1916); Germany. Frey, Notulæ Ent. Vol. 2, p. 41 [1922] (*Pararhamphomyia*).
259. *R. nubigena*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 200 (1904); Frey, Tyrol. Notulæ Ent. Vol. 2, p. 76 (1922).
- . *R. obscura*, Eversmann, Bull. Soc. Nat. Moscou, Vol. 7, p. 424 (1834), no description.
260. *R. obscura*, Loew (not Zetterstedt), Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 21, f. 23, 27 (1840); Isis, p. 548, f. 23-27 (1840); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862).
- C. Europe.
- Russia, Germany.
- Texas.
- N. Russia, Siberia.
- C. & N. Europe.
- N. & C. Europe.
- C. & N. Europe.

261. *R. obscura*, Zetterstedt, Fauna Ins. Lappon. p. 564 [1838] (*Rhamphomyza*); N. Europe.
 Dipt. Scand. Vol. 1, p. 429 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 174 (1861); Lundbeck, Dipt. Dan. Vol. 3, p. 55 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Collin, Ent. Mag. London (2), Vol. 24, p. 105 (1913); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 25 (1913); Notulæ Ent. Vol. 2, p. 41 [1922] (*Pararhamphomyia*).
 var. *eunordquisti*, Frey, ibidem, Vol. 2, p. 41 [1922] (*Pararhamphomyia*). N. Siberia, Kamchatka
262. *R. obscurella*, Zetterstedt, Dipt. Scand. Vol. 1, p. 432 (1842). Norway.
263. *R. *obtusa*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 118, pl. 10, f. 9, 10 (1908). Baltic Amber.
264. *R. *oedalooides*, Meunier, ibidem, Vol. 7, p. 92, 117, pl. 10, f. 7, 8 (1908). Baltic Amber.
265. *R. Oldenbergi*, Frey, Notulæ Ent. Vol. 2, p. 72 (1922). S. Europe.
266. *R. omissinervis*, Becker, Act. Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 18, f. 11, 13 (1900); Frey, Notulæ Ent. Vol. 2, p. 36 [1922] (*Pararhamphomyia*). Siberia.
267. *R. orthoneura*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 431 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 324, 326 (1909). Peru.
268. *R. otiosa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 425 (1895). Colorado.
269. *R. ozernajensis*, Frey, Notulæ Ent. Vol. 2, p. 42 [1922] (*Pararhamphomyia*). Kamchatka.
270. *R. pachymera*, Bigot, Bull. Soc. Ent. France (6), Vol. 7, p. 142 (1887); Ann. Soc. Ent. France (6), Vol. 9, p. 133 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 410, note (1895). California.
271. *R. pallidiventris*, Fallen, Empid. Suec. p. 28 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 53 (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 405 (1842); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 16 (1919); Frey, Notulæ Ent. Vol. 2, p. 6 [1922] (*Holoclera*). N. Europe.
272. *R. pallistigma*, Roser, Correspondenzenbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 53 (1840). C. Europe.
273. *R. Palméni*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 30 (1913); Notulæ Ent. Vol. 2, p. 73 (1922). Finland.
274. *R. paradoxa*, Wahlberg, Oefv. Vet. Acad. Förh. Stockholm, p. 107 (1844); Zetterstedt, Dipt. Scand. Vol. 8, p. 3033 (1849); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Verrall, Ent. Mag. London, Vol. 48, p. 24 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 11 (1913); ibidem (10), p. 6 (1913); Notulæ Ent. Vol. 2, p. 6, 9 [1922] (*Choreodromia*). N. Europe, Siberia.
- poeciloptera*, Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 16, pl. 1, f. 7, 8 (1900).
- tripes*, Becker, ibidem, Vol. 26 (9), p. 23 (1900)
- var. *batylimensis*, Frey, Notulæ Ent. Vol. 2, p. 6 [1922] (*Choreodromia*). Finland.
275. *R. parva*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 433 (1895). Massachusetts.
276. *R. parvicellulata*, Frey, Notulæ Ent. Vol. 2, p. 74 (1922). Austria.
277. *R. pectinata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 333 : Cent. 2, No. 49 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895). E. United States.
278. *R. pectoris*, Coquillett, ibidem, Vol. 18, p. 420 (1895). Georgia.
279. *R. penicillata*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 336 (1909). Bolivia.
280. *R. pennata*, Macquart, Dipt. N. France, Vol. 3, p. 133 (1827); Curtis, Brit. Ent. Vol. 8, p. 517 (1834); Macquart, Hist. Nat. Dipt. Vol. 1, p. 338 (1834); Guerin, Icon. Regn. Anim. Ins. p. 537, pl. 94, f. 7 (1835); Meigen, Syst. Beschr. Vol. 7, p. 91 (1838); Zetterstedt, Dipt. Scand. Vol. 1, p. 428 (1842); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 503 (1849); Ins. C. & N. Europe.

- Brit. Dipt. Vol. 1, p. 110, pl. 4, f. 2bcd (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 100 (1862); Giard, Traité Ent. Vol. 3, p. 992, pl. 108, f. 7 (1885); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 62, f. 19 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Frey, Notulæ Ent. Vol. 2, p. 34 [1922] (*Pararhamphomyia*).
compta, Zetterstedt, Dipt. Scand. Vol. 1, p. 429 (1842).
281. *R. perpulchra*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 334 Bolivia.
(1909).
282. *R. phanerostigma*, Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 46, p. 2, 8 N. Europe.
(1918); Notulæ Ent. Vol. 2, p. 70 (1922).
283. *R. phemius*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 500 (1849); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 418 (1895). Canada.
284. *R. physoprocta*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 18 Finland.
(1913); Notulæ Ent. Vol. 2, p. 37 [1922] (*Pararhamphomyia*).
285. *R. pictipennis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 330 Bolivia.
(1909).
286. *R. pilifer*, Meigen, Syst. Beschr. Vol. 7, p. 89 (1838). C. Europe.
287. *R. piligeronis*, Coquillet, Proc. U. S. Nat. Mus. Vol. 18, p. 432 (1895). Illinois.
288. *R. platycnemis*, Frey, Notulæ Ent. Vol. 2, p. 69 (1922). Siberia.
289. *R. plumifera*, Zetterstedt, Fauna Ins. Lappon. p. 567 (1838); Dipt. Scand. Vol. 1, p. 436 (1842); ibidem, Vol. 8, p. 3045 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 5025 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, p. 201 (1897); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 27 (1913); Notulæ Ent. Vol. 2, p. 44 [1922] (*Pararhamphomyia*). C. & N. Europe.
290. *R. plumipes*, ?Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 230 C. & N. Europe.
[1804] (*Empis*); Fallen, Empid. Suec. p. 25, part [1816] (*Empis*); ?Meigen, Syst. Beschr. Vol. 3, p. 47 (1822); Macquart, Hist. Nat. Dipt. Vol. 1, p. 339 (1834); Zetterstedt, Fauna Ins. Lappon. p. 563 (1838); Dipt. Scand. Vol. 1, p. 426 (1842); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 174 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 100 (1862); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 21 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 59, f. 18 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 20 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 18 (1919); Frey, Notulæ Ent. Vol. 2, p. 39, 40 [1922] (*Pararhamphomyia*).
geniculata, Bonsdorff (not Meigen or Zetterstedt), Finl. tvåv. Ins. Dipt. Vol. 1, p. 174 (1861).
291. *R. Pokornyi*, Bezzi, Ann. Mus. Hungar., Vol. 2, p. 198 (1904); Frey, Austria.
Notulæ Ent. Vol. 2, p. 67 [1922] (*Dasyrhamphomyia*).
292. *R. polita*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 200 : Cent. 2, No. 29 C. & E. United States.
(1862); Coquillet, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1895).
293. *R. *polymorpha*, Loew, Bernsteinfauna, p. 41 (1850) not named; Meunier, Miscell. Ent. Vol. 7, p. 178 (1899). Baltic Amber, Lower Oligocene.
294. *R. poplitea*, Wahlberg, Oefv. Vet. Akad. Förh. Stockholm, Vol. 1, p. 107 (1844); Zetterstedt, Dipt. Scand. Vol. 8, p. 3042 (1849); Wahlgren, Ent. Tidskr. Vol. 31, p. 55 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 13 (1913); Notulæ Ent. Vol. 2, p. 34 [1922] (*Pararhamphomyia*). Lapland.

295. *R. *porrecta*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 117, pl. 10, f. 5, 6 (1908). Baltic Amber.
296. *R. præstans*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 12 (1913); Notulæ Ent. Vol. 2, p. 8 [1922] (*Choreodromia*). Finland.
297. *R. priapulus*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 335 : Cent. 1, No. 54 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895). E. United States.
298. *R. propinquua*, Meijere, Tijdschr. v. Ent. Vol. 61, p. 134, pl. 8, f. 7, 8 (1918); Frey, Notulæ Ent. Vol. 2, p. 72 (1922). Europe, Siberia.
sulcata, auctorum, p. p.
299. *R. pseudocrinita*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 170 (1909). Spain.
300. *R. pseudogibba*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 62 (1909); Frey, Notulæ Ent. Vol. 2, p. 45 [1922] (*Pararhamphomyia*). C. Europe.
- *R. *pteropa*, Loew, Bernsteinfauna, p. 41 (1850), no description; Giebel, Ins. Vorwelt, p. 208 (1856), no description. Baltic Amber, Lower Oligocene.
301. *R. *ptilopa*, Loew, Bernsteinfauna, p. 41 (1850); Giebel, Ins. Vorwelt, p. 208 (1856). Baltic Amber, Lower Oligocene.
302. *R. pulchra*, Loew (not Egger), Berl. Ent. Zeitschr. Vol. 5, p. 327 : Cent. 1, No. 40 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895). E. United States.
303. *R. pulchriventris*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 432 (1905); Nova Acta Akad. Naturf. Halle, Vol. 89, p. 323 (1909). Peru.
304. *R. pulla*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 330 : Cent. 1, No. 44 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1895); Slosson, Ent. News, Philad. Vol. 14, p. 266 (1903). E. North America.
305. *R. pusilla*, Zetterstedt, Fauna Ins. Lappon. p. 569 (1838); Dipt. Scand. Vol. 1, p. 418 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 172 (1861); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 58 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 59 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 26 (1913); Notulæ Ent. Vol. 2, p. 42 [1922] (*Pararhamphomyia*). C. & N. Europe.
- erythrophthalma, Strobl (not Meigen), Progr. Seitenstett. Vol. 14, p. 58 (1880).
306. *R. pusio*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 340 : Cent. 1, No. 63 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895). E. United States.
307. *R. quinquelineata*, Say, Journ. Acad. Nat. Sc. Philad. Vol. 3, p. 95 [1823] (*Empis*); Wiedemann, Aussereurop. Zweifl. Ins. Vol. 2, p. 7 (1830); Say, Compl. Writ. Vol. 2, p. 83 [1859] (*Empis*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 412 (1895); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 26 (1919). C. United States.
308. *R. rava*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 198 : Cent. 2, No. 25 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 410 (1895).
? *Morissoni*, Bigot, Bull. Soc. Ent. France (6), Vol. 6, p. 141 (1887); Ann. Soc. Ent. France (6), Vol. 9, p. 132 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 410 (1895). C. United States.
309. *R. ravida*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 418 (1895). C. United States.
310. *R. reflexa*, Zetterstedt, Fauna Ins. Lappon. p. 566 (1838); Dipt. Scand. Vol. 1, p. 402 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 170 (1861); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 17, note (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 57 (1910); Frey, Notulæ Ent. Vol. 2, p. 66 [1922] (*Dasyrhamphomyia*). Lapland.
311. *R. *remitarsis*, Loew, Bernsteinfauna. p. 41 (1850), no description; Giebel, Ins. Vorwelt, p. 208 (1856), no description; Meunier, Miscell. Ent. Vol. 7, p. 14 (1899). Baltic Amber, Lower Oligocene.
312. *R. robustior*, Frey, Notulæ Ent. Vol. 2, p. 70 (1922). N. Siberia.

313. *R. rostrifera*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 465, f. 4 (1912); Formosa. Suppl. Ent. Berlin, Vol. 3, p. 70 (1914).
314. *R. rotundipennis*, Bezzii, Ann. Mus. Hungar. Vol. 3, p. 433 (1905); Nova Peru. Acta Akad. Naturf. Halle, Vol. 91, p. 323 (1909).
315. *R. rufipes*, Zetterstedt (not Meigen), Fauna Ins. Lappon. p. 564 (1838); N. & C. Europe. Dipt. Scand. Vol. 1, p. 394 (1842); ibidem, Vol. 8, p. 3034 (1849); Bonsdorff, Finl. tvåv. Ins. Dipt. p. 169 (1861); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 112 (1887); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 21 (1913); Notulæ Ent. Vol. 2, p. 39 [1922] (*Pararhamphomyia*).
316. *R. rufirostris*, Say, Journ. Acad. Nat. Sc. Philad. Vol. 6, p. 159 [1829] Indiana. (*rufirostra*); Compl. Writ. Vol. 2, p. 355 [1859] (*rufirostra*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 417 (1895).
317. *R. rufithorax*, Brunetti, Rec. India Mus. Vol. 9, p. 30 (1913). India.
318. *R. rugicollis*, Meigen, Syst. Beschr. Vol. 3, p. 46 (1822); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862).
319. *R. rustica*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 79 : Cent. 5, No. 56 E. North America. (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1895); Howard, Ins. Book, p. 18, f. 31 (1901).
320. *R. Sancti-Mauritii*, Becker (not Strobl), Berl. Ent. Zeitschr. Vol. 31, Switzerland. p. 113 (1887); Oldenberg, Zool. Jahrb. Vol. 43, Syst. p. 223 (1920).
321. *R. saniculae*, Curtis, Brit. Ent. Vol. 8, p. 517 (1834). England.
322. *R. sapporensis*, Matsumura, Addit. Vol. 2, p. 355, pl. 21, f. 17 (1916). Japan.
323. *R. Sauteri*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 466 (1912). Formosa.
324. *R. scaurissima*, Wheeler, Ent. News, Philad. Vol. 7, p. 189, f. 1-3 (1896). California.
325. *R. sciarina*, Fallen, Empid. Suec. p. 28 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 54 (1822); ibidem, Vol. 6, p. 339 [1830] (*sciaria*); Zetterstedt, Fauna Ins. Lappon. p. 568 (1838); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 21 (1840); Isis. Vol. 7, p. 549 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 423 (1842); ibidem, Vol. 8, p. 3041 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 55 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 173 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 100 (1862); Neuhaus, Dipt. March. p. 73 (1886); Verrall, Ent. Mag. London, Vol. 30, p. 140 (1894); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31, p. 21 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 59 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 10 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 15 (1919); Frey, Notulæ Ent. Vol. 2, p. 6 [1922] (*Holoclera*). *flava*, Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 175 (1861); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908). *hybrida*, Zetterstedt, Fauna Ins. Lappon. p. 572 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 438 (1842); Lundbeck, Dipt. Danica, Vol. 3, p. 69 (1910); Collin, Ent. Mag. London (2), Vol. 24, p. 105 (1913). *tifulariata*, Zetterstedt, Fauna Ins. Lappon. p. 569 (1838).
326. *R. scitula*, Frey, Notulæ Ent. Vol. 2, p. 71, 72 (1922). S. Europe.
327. *R. scolopacea*, Say, Journ. Acad. Nat. Sc. Philad. Vol. 3, p. 96 [1823] E. United States. (*Empis*); Wiedemann, Aussereur. Zweifl. Ins. Vol. 2, p. 8 (1830); Say, Compl. Writ. Vol. 2, p. 83 [1859] (*Empis*).
328. *R. scutellaris*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 429 (1895); California. Frey, Notulæ Ent. Vol. 2, p. 71 (1922).
329. *R. sellata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 328 : Cent. 1, No. 42 E. United States. (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1896).
330. *R. septembris*, White, Proc. Roy. Soc. Tasmania, 1916, p. 239 (1917). Tasmania.
331. *R. *sepulta*, Cockerell, Canad. Ent. Vol. 48 p. 123 (1916). Florissant, Miocene.

332. *R. serotina*, Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 71 (1915); Germany, Holland.
Meijere, Tijdschr. v. Ent. Vol. 61, p. 137, pl. 8, f. 9, 10 (1918);
Frey, Notulæ Ent. Vol. 2, p. 45 [1922] (*Pararhamphomyia*).
333. *R. serpentata*, Loew, Neue Beitr. Dipt. Vol. 4, p. 35 (1856); Schiner, C. & S. Europe.
Fauna Dipt. Austr. Vol. 1, p. 100, and footnote (1862); Strobl,
Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 54 (1893); Frey,
Notulæ Ent. Vol. 2, p. 77 (1922).
334. *R. setosa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 426 (1895); North America.
Proc. Wash. Acad. Sc. Vol. 2, p. 419 (1900).
335. *R. Siebecki*, Strobl, Glasnik Mus. Bosn. Herzeg. Sarajevo, Vol. 10, p. 402 S & C. Europe.
(1898); Wiss. Mitteil. Bosn. Herzeg. Sarajevo, Vol. 7, p. 564 (1900);
Frey, Notulæ Ent. Vol. 2, p. 73 (1922).
336. *R. similata*, Malloch, Report. Canad. Arct. Exped. 1913-18, Dipt. p. 46 Arctic America.
(1919).
337. *R. simplex*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3035 (1849); Lundbeck, N. & C. Europe.
Dipt. Danica, Vol. 3, p. 56 (1910); Frey, Acta Soc. Sc. Fenn.
Helsingfors, Vol. 37 (3), p. 20 (1913); Notulæ Ent. Vol. 2, p. 39
[1922] (*Pararhamphomyia*).
littoralis, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 15, 19
(1908).
var. *major*, Frey, Notulæ Ent. Vol. 2, p. 39 [1922] (*Pararhamphomyia*). Russia.
338. *R. soccata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 342 : Cent. 1, No. 67 E. & C. United States.
(1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895).
339. *R. sociabilis*, Williston, Kansas Univ. Quart. Vol. 2, p. 76 [1893] (*Empis*); W. United States.
Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 400 [1895] (*Empis*);
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 284 (1902); Frey,
Notulæ Ent. Vol. 2, p. 70 (1922).
abdicta, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 430 (1895).
340. *R. sordida*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 337 : Cent. 1, No. 58 E. United States.
(1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 405 and 415
(1895).
crassinervis, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 338 : Cent. 1, No. 59
(1861).
341. *R. spectabilis*, Frey, Notulæ Ent. Vol. 2, p. 70 (1922). N. Siberia.
342. *R. sphenoptera*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 40 (1873); Beschr. S. E. Europe.
Eur. Dipt. Vol. 3, p. 232 (1875); Frey, Notulæ Ent. Vol. 2, p. 4
[1922] (*Lundstroemiella*).
343. *R. spinipes*, Fallen, Empid. Suec. p. 26 [1816] (*Empis*); Meigen, Syst. Europé.
Beschr. Vol. 3, p. 49 (1822); Bouché, Naturg. Ins. Vol. 1, p. 48,
pl. 4, f. 26-30 (1834); Macquart, Hist. Nat. Dipt. Vol. 1, p. 337
(1834); Zetterstedt, Fauna Ins. Lappon. p. 564 (1838); Westwood,
Introd. Classif. Ins. Vol. 2, p. 547, f. 129 (1840); Zetterstedt, Dipt.
Scand. Vol. 1, p. 395 (1842); Gimmerthal, Bull. Soc. Nat. Mosc.
Vol. 20 (2), p. 163 (1847); Zetterstedt, Dipt. Scand. Vol. 8, p. 3034
(1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 54 (1851);
Walker, Ins. Brit. Vol. 1, p. 109 (1851); Pipping, Not. Sällsk.
Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Nylander, ibidem,
Vol. 4, p. 247 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 169
(1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98 (1862); Siebke,
Nyt Mag. Naturvid. Vol. 14, p. 386 (1866); Glover, Manusc. Notes,
p. 44, pl. 7, f. 18 (1874); Neuhaus, Dipt. March. p. 73 (1886);
Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 47 (1892);
Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908);
Lundbeck, Dipt. Danica, Vol. 3, p. 45, f. 10, 11 (1910); Wahlgren,

- Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 28 (1913); Notulæ Ent. Vol. 2, p. 69 (1922).
344. *R. spinosipes*, Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 74 (1915); Germany. Frey, Notulæ Ent. Vol. 2, p. 69, note (1922).
345. *R. squamigera*, Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 20 Europe. (1840); Isis, Vol. 7, p. 547 (1840); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Loew, Jahrb. Gel. Ges. Krakau, Vol. 41, p. 12 (1870).
- fimbriatipes*, Nowicki, Verh. Naturf. Ver. Brünn, Vol. 6, p. 82 (1868).
- geniculata*, Zetterstedt (not Meigen), Fauna Ins. Lappon. p. 564 [1838] (*Rhamphomyza*); Dipt. Scand. Vol. 1, p. 427 (1842); ibidem, Vol. 13, p. 5023 (1859).
- gracilipes*, Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 22, f. 21 (1840); Isis, Vol. 7, p. 549, f. 21 (1840); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Loew, Jahrb. Gel. Ges. Krakau, Vol. 42, p. 174 (1871); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910).
- plumipes*, Fallen (not Meigen), Emp. Suec. p. 25, part (1816).
- var. *squamosissima*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 59 (1909). C. Europe.
346. *R. stigmosa*, Macquart, Dipt. N. France, Vol. 3, p. 131 (1827); Hist. Nat. Dipt. Vol. 1, p. 333 (1834); Meigen, Syst. Beschr. Vol. 7, p. 90 (1838); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 115 (1887); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 50 (1892); Mik, Wien. Ent. Zeit. Vol. 15, p. 106 (1896); Frey, Notulæ Ent. Vol. 2, p. 68 (1922).
- angustifennis*, Becker (not Loew), Berl. Ent. Zeitschr. Vol. 31, p. 117 (1887).
- conformis*, Frey (not Kowarz), Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 28 (1913); Ent. Tidskr. 1914, p. 79 (1914).
- melanica*, Strobl (not Becker), Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 50 (1892).
- salictorum*, Loew (collection name) Bezzia, Kat. Palæarct. Dipt. Vol. 2, p. 220 (1903); Kertész, Cat. Dipt. Vol. 6, p. 35 (1909).
347. *R. stylata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 432 (1895). California.
348. *R. subglauella*, Frey, Notulæ Ent. Vol. 2, p. 36 [1922] (*Pararhamphomyia*). Lapland.
- glauella*, Frey ♂ (not Zetterstedt), Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 15, ♂ (1913).
349. *R. subsultans*, Frey, Notulæ Ent. Vol. 2, p. 43 [1922] (*Pararhamphomyia*). Kamchatka.
350. *R. sudigeronis*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 431 (1895); Frey, Notulæ Ent. Vol. 2, p. 73 (1922). California.
351. *R. sulcata*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 229 [1804] (*Empis*); Fallen, Empid. Suec. p. 26 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 46 (1822); Macquart, Dipt. N. France, Vol. 3, p. 131 (1827); Curtis, Brit. Ent. Vol. 8, p. 517 (1834); Macquart, Hist. Nat. Dipt. Vol. 1, p. 335, pl. 8, f. 3 (1834); Zetterstedt, Fauna Ins. Lappon. p. 565 (1838); Dipt. Scand. Vol. 1, p. 396 (1842); Boitard, Man. Ent. Vol. 3, p. 319 (1843); Gimmerthal, Bull. Soc. Nat. Mosc. Vol. 20 (2), p. 164 (1847); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 54 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 109, pl. 4, f. 2a (1851); Nylander, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 247 (1858); Pipping, ibidem, Vol. 4, p. 114 (1858); Zetterstedt, Dipt. Scand. Vol. 13, p. 5012 (1859); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 169 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 153 (1864); Beling, Arch. Naturg. Berlin, Vol. 48, p. 214 (1882); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 47 (1892); Bezzia, Bull. Soc. Ent. Ital. Vol. 30, p. 123 (1899); Strobl, Wien. Ent. Zeit. Vol. 18, p. 12 (1899); Frey, Acta

Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908); Hamm, Ent. Mag. London, Vol. 45, p. 161 (1909); Kleine, Soc. Ent. Steglitz, Vol. 24, p. 65 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 47, f. 12 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 56 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 29 (1913); Meijere, Tijdschr. v. Ent. Vol. 60, p. 134, pl. 8, f. 5, 6 (1918); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 23 (1919); Frey, Notulæ Ent. Vol. 2, p. 72 (1922).

? *fixa*, Harris, Engl. Ins. p. 151, pl. 44, f. 8 [1782] (*Empis*).

? *pusilla*, Scopoli, Ent. Carn. p. 362 [1763] (*Erax*); Olivier, Encycl. Méth. Vol. 4, p. 272 [1789] (*Asilus*); Schiner, Verh. Zool.-bot. Ges. Wien, Vol. 6, p. 420 (1856), gen.?

352. *R. tarsata*, Meigen, Syst. Beschr. Vol. 3, p. 45 (1822); Zetterstedt, Dipt. Scand. Vol. 1, p. 430 (1842); ibidem, Vol. 8, p. 3041 (1849); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 384 (1866); Verrall, Ent. Mag. London, Vol. 19, p. 224 (1883); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 53 f. 17 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Frey, Notulæ Ent. Vol. 2, p. 43 [1922] (*Pararhamphomyia*).
? longipes, Walker (not Meigen), Ins. Brit. Dipt. Vol. 1, p. 110 (1851).

353. *R. tenuipes*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 121 (1907). Algeria.

354. *R. tenuirostris*, Fallen, Empid. Suec. p. 29 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 52 (1822); Macquart, Dipt. N. France, Vol. 3, p. 135 (1827); Hist. Nat. Dipt. Vol. 1, p. 337 (1834); Curtis, Brit. Ent. Vol. 8, p. 517 (1834); Zetterstedt, Fauna Ins. Lappon. p. 568 (1838); Dipt. Scand. Vol. 1, p. 406 (1842); Boitard, Man. Ent. Vol. 3, p. 320 (1843); Gimmerthal, Bull. Soc. Nat. Mosc. Vol. 20, p. 164 (1847); Zetterstedt, Dipt. Scand. Vol. 12, p. 4613 (1855); Pipping, Not. Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Leunis, Synops. Zool. Vol. 2, p. 403 (1886); Becker, Berl. Ent. Zeischr. Vol. 31, p. 112, note (1887); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 46 (1892); Adams, Ent. Mag. London p. 94 (1905); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 21 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 9 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 14 (1919).

pallidiventris, Bonsdorff (not Fallen), Finl. tvåv. Ins. Dipt. Vol. 1, p. 170 (1861).

rufipes, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 231, part [1804] (*Empis*).

variabilis, Fallen, Empid. Suec. p. 29 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 3, p. 51 (1822); Macquart, Dipt. N. France, Vol. 3, p. 134 (1827); Hist. Nat. Dipt. Vol. 1, p. 337 (1834); Zetterstedt, Fauna Ins. Lappon. p. 569 (1838); Dipt. Scand. Vol. 1, p. 407 (1842); Walker, List Dipt. Brit. Mus. Vol. 3, p. 499 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 55 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 109 (1851); Pipping, Not. Sällsk. Fenn. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 171 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 154 (1864); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 21 (1908); Lundbeck, Dipt. Dan. Vol. 3, p. 75 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 58 (1910); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 15 (1919); Frey, Notulæ Ent. Vol. 2, p. 5 [1922] (*Holoclera*).

355. *R. tenuiterfilata*, Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), Siberia, Russia.
 p. 20, pl. 1, f. 18, 19 (1900); Frey, Notulæ Ent. Vol. 2, p. 43 [1922] (*Pararhamphomyia*).

356. *R. tephraea*, Meigen, Syst. Beschr. Vol. 3, p. 47 (1822); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98 (1862); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 49 (1892); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 16 (1919); Frey, Notulæ Ent. Vol. 2, p. 7 [1922] (*Choreodromia*). C. Europe.
357. *R. tessa*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 422 (1895). E. United States.
358. *R. testacea*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 197 : Cent. 2, No. 24 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1895). E. & C. United States.
359. *R. tibialis*, Meigen, Syst. Beschr. Vol. 3, p. 44 (1822); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 112 (1887); Collin, Ent. Mag. London, Vol. 49, p. 105 (1913); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 29 (1913); Ent. Tidskr. 1914, p. 79 (1914); Notulæ Ent. Vol. 2, p. 71 (1922). Europe.
360. *R. tipularia*, Fallen, Empid. Suec. p. 27 [1816] (*Empis*); Meigen, Syst. Beschr. Vol. 6, p. 339 [1830] (*Empis*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 336 (1834); Gimmerthal, Bull. Soc. Nat. Mosc. Vol. 15, p. 666 (1842); Zetterstedt, Dipt. Scand. Vol. 1, p. 411 (1842); Schiner, Fauna Dipt. Austr. Vol. 1, p. 99 (1862); Jaroschewsky, Arb. Ges. Naturf. Univ. Kharkow, Vol. 11, p. 353 (1877); Becker, Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 25 (1900); Frey, ibidem, Vol. 31 (9), p. 20 (1908); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 58 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 55 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 13 (1913); Oldenberg, Arch. Naturg. Berlin, Vol. 83, A, 6, p. 17 (1919); Frey, Notulæ Ent. Vol. 2, p. 33 [1922] (*Pararhamphomyia*). N. & C. Europe, Siberia.
- cinerea*, Meigen (not Fabricius), Syst. Beschr. Vol. 3, p. 43 (1822).
361. *R. tolteca*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. p. 371 (1901); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 323 (1909). Mexico.
362. *R. tonsa*, Loew, Beschr. Eur. Dipt. Vol. 2, p. 244 (1871). Siberia.
363. *R. trilineata*, Zetterstedt, Schummel, Verh. Schles. Ges. (1832) no description; Zetterstedt, Dipt. Scand. Vol. 13 p. 5013 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 98, foot note (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 47 (1892); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Wahlgren, Ent. Tidskr. Vol. 31, p. 57 (1910). N. & C. Europe.
364. *R. tristis*, Walker, Trans. Ent. Soc. Lond. n. s. Vol. 4, p. 148 (1857). United States.
365. *R. tristriolata*, Nowicki, Verh. Naturf. Ver. Brünn, Vol. 6, p. 77, pl. 2, f. 2 (1868); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 54 (1892); Frey, Notulæ Ent. Vol. 2, p. 75 (1922). C. Europe.
366. *R. truncata*, Frey, ibidem, Vol. 2, p. 40 [1922] (*Pararhamphomyia*). Kamchatka.
367. *R. tumiditarsis*, Oldenberg, Arch. Naturg. Berlin, Vol. 82, 1, p. 153 (1916); Frey, Notulæ Ent. Vol. 2, p. 5 [1922] (*Lundstroemiella*). Tirol.
368. *R. tympanica*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 337 (1909). Bolivia.
369. *R. umbilicata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 342 : Cent. 1, No. 65 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 409, 415 (1895); Slosson, Ent. News, Philad. Vol. 14, p. 269 (1903); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 321, 323 (1909); Frey, Notulæ Ent. Vol. 2, p. 37 [1922] (*Pararhamphomyia*). E. United States.
- ungulata*, Loew, Berl. Ent. Zeitschr. Vol. 5, p. 342 : Cent. 1, No. 66 (1861).

370. *R. umbripennis*, Meigen, Syst. Beschr. Vol. 3, p. 54 (1822); ibidem, Europe, N. Africa.
 Vol. 6, p. 339 (1830); Macquart, Hist. Nat. Dipt. Vol. 1, p. 338 (1834); Explor. Algér. Zool. Vol. 3, p. 444 (1849); Schiner, Fauna Dipt. Austr. Vol. 1, p. 101 (1862); Verrall, Ent. Mag. London, Vol. 19, p. 224 (1883); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 58 (1892); Wien. Ent. Zeit. Vol. 18, p. 13 (1899); Lundbeck, Dipt. Danica, Vol. 3, p. 73 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 59 (1910); Oldenberg, Arch. Naturg. Vol. 83, A, 6, p. 15 (1919); Frey, Notulæ Ent. Vol. 2, p. 5 [1922] (*Holoclera*).
pulchra, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 346 [1860] (*Holoclera*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 80 [1862] (*Holoclera*).
 var. *morena*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 13 (1899). Spain.
371. *R. umbripes*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 117 (1887); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 54 (1893); Wien. Ent. Zeit. Vol. 18, p. 12 (1899); Oldenberg, Arch. Naturg. Vol. 83, A, 6, p. 18 (1919); Frey, Notulæ Ent. Vol. 2, p. 37 [1922] (*Pararhamphomyia*). C. & S. Europe.
372. *R. umbrosa*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 77 : Cent. 5, No. 53 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 414 (1896). E. No. America.
373. *R. unguiculata*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 24 (1913); Ent. Tidskr. 1914, p. 79 (1914); Notulæ Ent. Vol. 2, p. 40 [1922] (*Pararhamphomyia*). Finland, Sweden.
374. *R. *ungulina*, Loew, Bernsteina fauna, p. 41 (1850); Giebel, Ins. Vorwelt, p. 208 (1856); Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 92, 115, pl. 9, p. 10-12 (1908). Baltic Amber, Lower Oligocene.
375. *R. unicolor*, Zetterstedt, Fauna Ins. Lappon. p. 568 (1838); Dipt. Scand. Vol. 1, p. 422 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 173 (1861); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 112 (1887). N. & C. Europe.
376. *R. unifasciata*, Brunetti, Rec. Indian Mus. Vol. 6, p. 29 (1913); Fauna Brit. Ind. Dipt. Vol. 1, p. 345 (1920). W. Himalayas.
377. *R. unimaculata*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 201 : Cent. 2, No. 33 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895). E. & C. United States.
378. *R. uralensis*, Becker, Mém. Acad. Sc. Petrograd, Vol. 28 (7), p. 58, f. 3 (1915). Artic Ural.
379. *R. ursinella*, new name. Arctic America.
ursina, Malloch (not Oldenberg), Report Canad. Arct. Exped. 1913-18, Dipt. p. 46 (1919).
380. *R. ursina*, Oldenberg, Arch. Naturg. Berlin, Vol. 80, 9, p. 91 (1915); Alps.
 Frey, Notulæ Ent. Vol. 2, p. 76 (1922).
villosipes, Strobl (not Bezzii), Mitteil. Nat. Ver. Steiermark, Graz. Vol. 46, p. 60 (1909).
381. *R. valga*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 428 (1895) E. United States.
382. *R. vara*, Berl. Ent. Zeitschr. Vol. 5, p. 337 : Cent. 1, No. 57 (1861); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 415 (1895). E. & C. United States.
383. *R. vesiculosus*, Fallen, Empid. Suec. p. 27 [1816] (*Empis*); Zetterstedt, Fauna Ins. Lappon. p. 566 (1838); Dipt. Scand. Vol. 1, p. 433 (1842); Pipping, Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 174 (1861); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 20 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 49 f. 13, 14 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 55 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, N. Europe.

Vol. 37 (3), p. 28 (1913); Notulæ Ent. Vol. 2, p. 66 [1922] (*Dasyrhamphomyia*).

alata, Zetterstedt, Fauna Ins. Lappon. p. 567 [1838] (*Rhamphomyza*); Dipt. Scand. p. 434 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 175 (1861).

anthracina, Zetterstedt (not Meigen), Fauna Ins. Lappon. p. 567 [1838] (*Rhamphomyza*).

atripennis, Zetterstedt, Dipt. Scand. Vol. 1, p. 434 (1842); ibidem, Vol. 8, p. 4045 (1849); ibidem, Vol. 11, p. 4273 (1852); Wahlgren, Ent. Tidskr. Vol. 31, p. 55 (1910).

Fallenii, Meigen, Syst. Beschr. Vol. 3, p. 50 (1822).

lengens, Zetterstedt, Dipt. Scand. Vol. 13, p. 5024 (1859); Wahlgren, Ent. Tidskr. Vol. 31, p. 55 (1910).

384. *R. vespertilio*, Zetterstedt, Dipt. Scand. Vol. 1, p. 435 (1842); Strobl, N. & C. Europe. Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 61 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 28 (1913); Notulæ Ent. Vol. 2, p. 66 [1922] (*Dasyrhamphomyia*).

385. *R. villipes*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 414 (1900). Alaska.

386. *R. villosa*, Zetterstedt, Fauna Ins. Lappon. p. 563 (1838); Dipt. Scand. Vol. 1, p. 415 (1842); Siebke, Nyt Mag. Naturv. Vol. 12, p. 154 (1864); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 19 (1908).

Sancti-Mauritii, Strobl (not Becker), Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 51 (1893).

387. *R. villosipes*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 430 (1905); Nova Acta Peru. Akad. Naturf. Halle, Vol. 91, p. 322 (1909).

388. *R. virgata*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 430 (1895). E. United States.

389. *R. vittata*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 197 : Cent. 2, No. 23 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 411 (1896).

390. *R. Woldstedti*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 14 (1913); Notulæ Ent. Vol. 2, p. 34 [1922] (*Pararamphomyia*). Finland.

391. *R. Wuorentausi*, Frey, Notulæ Ent. Vol. 2, p. 67 [1922] (*Dasyrhamphomyia*). N. Siberia.

392. *R. Zaitzevi*, Becker, Mém. Acad. Sc. Petrograd, Vol. 28 (7), p. 54, f. 1 (1915). W. Siberia Tundra.

SUBFAMILY CLINOCERATINÆ

Charactera. — Head globose, the eyes separated, their facets small; antennæ short, located above the middle of the head, three-jointed, the third joint oval, conical or onion-shaped, rarely lengthened, the arista almost always much longer than the third joint; proboscis shorter than the head, extending down from the lowest part of the head, usually fleshy and thick, but sometimes sharp and chitinized, palpi small and incumbent on the proboscis, except in *Synaphotera*. Thorax narrow and elongate, the mesonotum flattened in front of scutellum, almost always with macrochætae on the disk, never pilose, metapleuræ often bearing a cluster of fine hairs. Genitalia of the male of the form of either a pygidium or an epipygium, bilateral; no ovipositor. Legs slender, nowhere thickened, unarmed, but rarely the front femora may bear flexor setulae, which however are never arranged as definite rows of denticles. In *Ceratomerus*, whose position is questionable, the middle femora of the male bear apophyses. Wings cuneiform, no anal angle developed, costa encompassing the entire wing, neuration complete, basal cells small, third vein forked or not, petiole of the second and third veins arising nearer the humeral crossvein than to the anterior, anal vein reduced or absent, anal crossvein variable, sometimes perpendicular, sometimes rounding into the underside of the anal cell, stigma absent or very weak, but in *Wiedemannia* distinct.

TABLE OF THE GENERA OF THE CLINOCERATINÆ

- A. Antennæ inserted at the middle of the head, the third joint conical with a rather short style which terminates in a bristle-like segment; face not constricted from the cheeks by a suture; eyes bare; no ocellar tubercle; proboscis short, sharp, chitinous and incurved; anal crossvein when present perpendicular to the anal vein; bristles never strong, no humeral bristle, metapleuræ bare; legs yellowish 2.
- B. Antennæ with the third joint remarkably lengthened, without evident style; face not constricted from the cheeks by a suture; eyes bare; no ocellar tubercle; proboscis short and fleshy; bristles well developed including the humeral, metapleuræ setulose; dorsal valves of pygidium erect, end valves abruptly bent downward; anal crossvein rectangularly curving, third vein furcate. 4.
- C. Antennæ inserted above the middle of the head, the third joint usually with a lengthened arista; oral margin of the cheeks with a more or less distinct incision or suture extending toward the eyes; eyes usually densely pubescent; proboscis usually short, vertical, soft and haustellate; anal cell usually rounded at the tip; bristles of the body always well developed, metapleuræ usually bearing hairs; body dark and legs largely or wholly black, except in Roederioides 5.
2. Anal cell wanting, discal cell apically open, the second posterior cell with a long petiole, first vein ending before the middle of the wing, third vein furcate; middle femora ♂ with an apophysis on the anterior surface; antennæ elongate, the first joint as long as the head. (**Pl. 7, Fig. 67**) Genus CERATOMERUS, Philippi.
- Anal cell present, discal cell complete, second posterior cell sessile, second basal cell long; legs slender, without armature or bristles; antennæ not longer than the head, the first joint small. 3.
3. Third vein not furcate, anal vein distinct from the under side of the anal cell and incomplete, auxiliary vein evanescent; about ten small dorsocentral bristles, one supraalar, four scutellar, ocellar and vertical bristles present; costa with a basal bristle present; pygidium terminal; present; third antennal joint lanceolate. (**Pl. 3, Fig. 20**) Genus BOREODROMIA, Coquillett
- Third vein furcate, anal vein continuous with the underside of the anal cell and complete to the margin, auxiliary vein attaining the costa; head, thorax and costa without bristles; epipygium tumid and reflexed; third antennal joint ovate. (**Pl. 3, Fig. 23**) Genus SYNAMPHOTERA, Loew.
4. Antennæ inserted above the middle of the head, the third segment tipped with minute peg-like joint; a row of acrostichals;

- pygidium with cruciate terminal sharp processes. (Pl. 3,
Fig. 21)* Genus **NIPHOCERUS**, nov. gen.
- Antennæ inserted below middle of head, no trace of apical style;
no acrostichals; pygidium without cruciate terminal processes.
(Pl. 3, Fig. 22)* Genus **CERATEMPIS**, nov. gen.
5. *Third vein simple; opaque black species with black legs and
halteres* 6.
- Third vein branched* 7.
6. *Anal crossvein greatly reflexed, nearly parallel with the axis of
the wing, first vein ending before the middle of the wing, costa
with setulae; proboscis very short, haustellate; face broad; eyes
pubescent, the lower facets enlarged; bristles strong, post-
humeral and intraalar bristles present, metapleura with a
bunch of hairs; epipygium small and reflexed; front femora
with flexor thorn-like spines, empodium and pulvilli long.
(Pl. 2, Fig. 17)* Genus **OREOTHALIA**, Melander.
- Anal crossvein perpendicular to the axis of the wing, the first
vein ending beyond the middle of the wing, costa without
setulae; proboscis nearly as long as the head, pointed; face
narrow; eyes bare, facets uniform; bristles weak, posthumeral
and intraalar wanting, metapleura bare; pygidium globose
and terminal; femora not spinose, empodium and pulvilli
minute. (Pl. 7, Fig. 68)* Genus **HELEODROMIA**, Haliday.
7. *Fourth vein arising near the base of the anal cell, the second
basal cell therefore as long as the anal cell, auxiliary vein
evanescent apically, anal crossvein strongly recurved* 8.
- Fourth vein arising near the end of the basal third of the anal
cell, the second basal cell therefore shortened, auxiliary vein
ending in the costa; femora never incrassate; metapleura with
hairs* 10.
8. *Eyes of both sexes contiguous on the face and pubescent; third
joint of the antennæ greatly lengthened and continuous with
the arista; proboscis slender, nearly as long as the head;
front coxae half as long as their femora; calypteres without
fringe; metapleura bare. (Pl. 7, Fig. 69)* Genus **TRICHOPEZA**, Rondani.
- Eyes distinctly separated; third joint of the antennæ conical
or oval, with a distinct thickened arista; proboscis thick* 9.
9. *Front femora thickened, spinose at the base and four times as
long as their coxae; proboscis as long as the head; metapleura
bare* Genus **DIPSOMYIA**, Bezzi.
- Femora slender, not spinose; proboscis short; eyes bare; meta-
pleura with a row of setæ. (Pl. 7, Fig. 65; Pl. 8,
Fig. 76)* Genus **PROCLINOPYGA**, nov. gen.
10. *Proboscis as long as the head, slender, pointed, rigid, somewhat
inflected, without labella; cheeks rostriform; brownish species
with yellow legs; epipygium reflexed and with a ventral keel.
(Pl. 2, Fig. 19)* Genus **ROEDERIOIDES**, Coquillett.

- Proboscis usually short, thick, fleshy, not inflexed; cheeks usually narrow; black or blackish species* II.
11. *Head articulated to the thorax high up on the occiput, extending obliquely forward; small species, body shining or subshining; prosternum as long as the mesosternum, legs yellowish, pulvilli vestigial; no trace of anal vein* 12.
- Head articulated nearer the center of the occiput, hanging vertically; prosternum shorter than the mesosternum; wings narrow, usually hyaline, with or without nebulous markings, veins rarely undulating; legs usually wholly or in large part black or blackish; large species, body opaque* 13.
12. *Wings maculate with lighter spots on a dark ground, veins undulating, three submarginals; empodium present; body more or less pollinose. (Pl. 3, Fig. 24)* Genus DOLICOCEPHALA, Macquart.
- Wings broad, hyaline, not maculate, veins not undulating, two submarginals; empodium vestigial; body entirely shining, without pollen: cavernicolous. (Pl. 7, Fig. 64)* Genus LAMPOSOMA, Becker.
13. *Face separated from the lower occiput by an incision on the cheeks that extends to the eye; face not descending beneath the eyes; no acrostichals* Genus CLINOCERA, Meigen 14.
- Cheeks broader, without such a deep incision, the face fusing with the lower occiput and extending more or less beneath the level of the eyes, its front margin deeply emarginate or impressed and carinate; stigma more or less developed, discal cell sharp apically, acrostichals usually abundant* Genus WIEDEMANNIA, Zetterstedt. 18.
14. *Pulvilli and empodium vestigial, much shorter than the claws; dorsocentrals sometimes numerous; stigma present; large species* Subgenus BERGENSTAMMIA, Mik.
- Pulvilli and empodium present, the latter at least as long as the claws; at most five or six dorsocentrals, two scutellars.* 15.
15. *Lower edge of the face straight across; stigma wanting or weak; costal setulae minute* 16.
- Oral margin of the face with a median excision or depression; discal cell blunt at the tip.* 17.
16. *Discal cell rather sharp apically; wings not or scarcely spotted; front femora of male with flexor bristles. (Pl. 2, Fig. 18). Subgenus CLINOCERA, s. str.*
- Discal cell rather blunt apically; wings spotted with nebulous marks on the crossveins, etc.; sometimes three submarginal cells; front femora of both sexes with flexor bristles or spines.* Subgenus HYDRODROMIA, Macquart.
17. *Face with a few scattered hairs beneath the antennæ; stigma poorly developed, wings not spotted; front femora with preapical comb of setulae on inner side* Subgenus KOWARZIA, Mik.
- Face entirely bare; stigma and nebulous spottings more or less developed; front femora with three or four preapical setulae on inner side* Subgenus PHÆOBALIA, Mik.

18. Face projecting more than half the eye-height below the eyes. 19.
Face projecting less than half the eye-height below the eyes 21.
19. Stigma large and more or less circular, located much beyond
 the end of the first vein, the second vein curving beneath the
 stigma, sometimes a clouding from the stigma extending below
 the second vein; five or six dorsocentrals with rather long
 alternate bristles; scutellum with discal setulae Subgenus WIEDEMANNIA, S. str.
- Stigma elongate, beginning at end of first vein and not touching
 second vein* 20.
20. Eyes round; face in profile quadrangular Subgenus CAMELOPIS, Engel.
Eyes oval, longer than wide; face tapering below Subgenus CLINOCERELLA, Engel.
21. Stigma rounded, beginning much beyond end of first vein;
 alternate weak dorsocentrals present Subgenus PSEUDOWIEDEMANNIA, Engel.
Stigma elongate, beginning at end of first vein 22.
22. All femora bearing strong anterior and posterior preapical
 bristles; costal spines strongly projecting; acrostichals if
 present proclinate Subgenus EUCELIDIA, Mik.
*Femora without distinct preapical bristles but sometimes with
 short preapical setæ; costal setulae not strongly projecting,
 usually minute and inclined* 23.
23. Front femora with an oblique comb of about five setulae just before
 the knee on the inner side; acrostichals short and weak, only
 the anterior ones present; only the strong dorsocentrals present;
 scutellum with marginal setulae in addition to the pair of
 bristles Subgenus ROEDERELLA, Engel.
*Front femora without comb of setulae near knee; alternating
 smaller dorsocentrals present; acrostichals forming a complete
 row, reclinate in front and proclinate in back* 24.
24. Front femora with about four distinct setæ at apical third of
 inner side; scutellum rarely bearing more than the apical pair
 of bristles; arista usually blunt Subgenus CHAMARDIPSIA, Mik.
*Front femora devoid of pronounced setulae within; scutellum
 with setulae* Subgenus PHILOLUTRA, Mik.

I. GENUS CERATOMERUS, PHILIPPI

Ceratomerus, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 766 (1865); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 124 (1889); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 300 (1909); Kertész, Cat. Dipt. Vol. 6, p. 111 (1909).

Characters. — Brownish species with infumated wings whose venation is similar to that of *Hemerodromia*, with lengthened antennæ and in the male with deformed legs. Head globular; antennæ inserted at the middle of the head, the first antennal joint as long as the head, the second joint short, the third joint elongate oval, its arista about one-half as long as this joint. Proboscis perpendicular, twice as long as the head, palpi retracted. Abdomen cylindrical, attenuated before the tip, the apex

inflated in the male. Legs slender, front femora with a hooklike tooth at the base, the front tibiæ tuberculate below the knee and hairy, middle femora armed with a large apophysis a little beyond the middle on the upper anterior face, directed forward and bispinose at its tip, beneath with a fascicle of hairs and with three bristles, middle tibiæ biseriately ciliate, hind tibiæ dilated apically and produced as a lobe beyond the insertion of the tarsus, pubescent but not ciliate. Female unknown.

Type species : the South American *C. paradoxus*, Philippi, which has not been found since its original discovery in 1865. This fly is well named for it is impossible to assign its systematic position with satisfaction. I take it that the slender legs and shortened front coxæ bespeak a relationship with the Clinoceratinæ, and that the *Hemerodromia*-like neuration is the result of parallel evolution.

Geographical distribution.

1. *C. paradoxus*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 766, pl. 28, Chile.
f. 46 (1865). — **Pl. 7, Fig. 67.**

2. GENUS BOREODROMIA, COQUILLETT

Boreodromia, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 247, 260 (1903); ibidem, p. 264 [1903] (*Boreomyia*); Aldrich, Cat. Dipt. p. 316 [1905] (*Boreomyia*); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Kertész, Cat. Dipt. Vol. 6, p. 119 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 515 (1910).

Synamphota, Melander (not of Loew, 1858), Trans. Amer. Ent. Soc. Vol. 28, p. 231 (1902).

Characters. — Brownish species, measuring three millimeters in length. Head globular, occiput shallowly convex, below with sparse hairs, above with setæ and two vertical bristles; ocelli placed well up on the vertex, one pair of ocellar and two pairs of lesser ocellar bristles; sides of the front broadly diverging, of the face parallel, cheeks linear, mouthopening very small; eyes bare, the incision at the antennæ broad and shallow, lower facets becoming larger; antennæ inserted below the middle of the head, three-jointed, the third joint conical, microscopically hairy, the arista half as long as the third joint, three-jointed, the basal joint small, the terminal joint thin and half as long as the intermediate; proboscis short, rigid, somewhat incurved, the fleshy labella half as long as the proboscis, palpi obliquely descending, two-jointed, slender, hairy. Thorax quadrate, not greatly convex but without depressed prescutellar area, pollinose, not pubescent nor setulose, no humeral bristle, one notopleural, one postalar, four scutellar, about ten dorsocentrals, none large, no acrostichals; pleuræ bare, no pectal bristle. Abdomen compressed, not tapering, comprising eight segments; pygidium small, terminal, upper valves spatulate and erect, middle valves corneous and directed forward, central filament hidden; no ovipositor. Legs slender, unarmed, hairy but without bristles, setæ or spurs, front coxæ one and two-thirds times as long as the posterior pairs, apex of hind tibiæ closely fimbriate, pulvilli small. Wings cuneiform, narrow at the base, no anal angle or alula, costa encompassing the entire wing, a basal costal bristle, the costa biseriately hairy, auxiliary vein straight, evanescent at the tip, closer to the first vein than to the costa, humeral crossvein before the base of the second basal cell, second vein long, third vein not forked, its pedicel arising near the middle of the second basal cell, discal cell complete, emitting three veins, second posterior cell sessile, anal cell short, anal crossvein recurved around the back of the anal cell and separate from the ciliate margin, the anal vein represented by a fold which starts behind the anal cell.

Type species : *B. bicolor*, Loew (Pl. 3, Fig. 20), Coquillett's designation. This species occurs along the Pacific coast from Alaska to the State of Washington. It may be that Bigot's *Clinocera maculipes*, a species with yellow antennæ, halteres and legs, may belong here.

Geographical distribution.

1. *B. bicolor*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 18 : Cent. 3, No. 34 W. North America.
[1863] (*Synamphotera*); Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 423 [1900] (*Synamphotera*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 231 [1902] (*Synamphotera*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 247 (1903). — Pl. 3, Fig. 20.
2. *B. ?maculipes*, Bigot, Bull. Soc. Zool. France, Vol. 12, p. 118 [1887] California. (*Clinocera*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 245 [1902] (*Clinocera*); Engel, Deutsche Ent. Zeitschr. 1918, p. 35 [1918] (*Clinocera*).

3. GENUS SYNAMPHOTERA, LOEW

Synamphotera, Loew, Zeitschr. Ges. Naturw. Vol. 11, p. 454 (1858); Syst. Beschr. Vol. 2, p. 255 (1871); Roeder, Wien. Ent. Zeit. Vol. 6, p. 169 (1887); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 120 (1889); Williston, Man. N. Amer. Dipt. p. 75 (1896); Melander, Williston Man. N. Amer. Dipt. p. 222 (1908); Lundbeck, Dipt. Dan. Vol. 3, p. 226 (1910).

Dryodromia, Rondani, Dipt. Ital. Vol. 1, p. 150 [1856] (*Dryodromya* and *Driodromya*); Mik, Ent. Nachr. Berlin, Vol. 12, p. 324 (1886); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248 (1903); Kertész, Cat. Dipt. Vol. 6, p. 119 (1909); Engel, Deutsche Ent. Zeitschr. 1918, p. 3 (1918).

Characters. — An entirely yellow, bristleless insect with short antennæ and full neuration. Head globular, eyes moderately large, round, bare, facets uniform, no excision at the antennæ, well separated on the front and face in both sexes; cheeks linear, epistome long, face extremely short; proboscis shorter than the head, vertical, chitinous, tubular, the palpi long, projecting straight forward, cylindrical and hairy in the female, compressed in the male, in which sex they are conspicuous; antennæ located well down on the head, the basal two joints shorter than broad, the third joint short ovate, rather broad, and with a stumpy two-jointed terminal style which is tipped with a short bristle. Thorax moderately large, pollinose, without bristles, the scutellum with a few marginal setulae, prothorax short. Abdomen nearly bare; epipygium reflexed and large, penis very thick, two pairs of erect side valves; ovipositor conical, erect. Legs rather slender, without bristles, spurs or setæ, front coxae one-half as long as the femora and slightly longer than the posterior coxae, pulvilli moderate, empodium bristle-like, minute. Wings with narrowed base, axillary angle very blunt, costa continuing around the wing, no basal bristle, no costal setulae, stigma very faint, auxiliary vein attaining the costa, first vein ending beyond the middle of the wing, third vein forked and normally the upper branch angulate and connected with the second vein by an extra crossvein, discal cell complete and long, posterior cells sessile, anal crossvein slightly reflexed and recurved, abruptly ending in the complete anal vein which meets the wing-margin, the anal cell a little shorter than the second basal, fourth vein arising near the base of the anal cell, a fold present beneath the humeral crossvein, marginal cilia and costal hairs very short. The crossveins at the end of the wing are quite unstable, additional ones may be present or the normal one absent.

Taxonomy: In the analytical key to the genera of Empidina of his Prodromus of the Italian Diptera, Rondani established the genus *Dryodromia*, citing but not describing the species *testacea*. Coquillett was of the opinion that Rondani's species was *Hilara tenella* Fallen, with a supernumerary

crossvein. Although such aberrant specimens may be rare in the genus *Hilara*, so as to make it doubtful if Coquillett's conclusion is correct, yet his supposition is plausible enough to warrant the retention of the better-known name *Synamphotera* for this genus. Anyway, Rondani's species *testacea* was tabulated *sine descriptione* and would give way to Loew's well described *pallida*. Mik referred *Dryodromia* to *Tachyptera* in the reference above given.

Type species : *S. pallida*, is the only known form. It occurs in Europe and is considered rare. Its earlier stages are not known.

Geographical distribution.

1. *S. pallida*, Loew, Zeitschr. Ges. Naturw. Vol. 11, p. 455 (1858); Syst. C. & S. Europe.

Beschr. Vol. 2, p. 253 (1871); Lundbeck, Dipt. Danica, Vol. 3, p. 227, f. 95-97 (1910); Collin, Ent. Mag. London, Vol. 24 (2), p. 106 (1913). — **Pl. 3, Fig. 23.**

testacea, Rondani, Dipt. Ital. Vol. 1, p. 150 [1856] (*Dryodromia*), table name; Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248 (1903), as syn. of *Hilara tenella*, Fall.

4. GENUS NIPHOCENIA, NOV. GEN.

Characters. — Slender insects with lengthened antennæ. Head globose, eyes round, bare, facets uniform, face wide and convex, particularly so in the female, continuous with the small cheeks, its lower edge straight; front broad, half as long as the face, setulose above antennæ, ocellar triangle scarcely elevated, ocellar bristles reaching two-thirds the way to the antennæ, no vertical bristles but the uppermost occipital setæ bristle-like; first antennal joint more than twice as wide as the globose second, both joints loosely setulose, third joint as long as the head, slightly tapering, loosely pubescent, tipped by a microscopic setula; proboscis short and thick, palpi pendent. Thorax with full rows of uniseriate uniform dorsocentrals and acrostichals, several humeral bristles, and one each of intrahumeral, posthumeral, notopleural, supra-alar, intra-alar and postalar bristles, two scutellars, metapleuræ bearing a cluster of setulæ; prescutellar area not depressed. Abdomen cylindrical, seven segments before the pygidium, loosely setose, showing small pittings arranged across base of second segment and in pairs on sides of the segments; pygidium globular, not widening the abdomen, dorsal valves erect, terminal valves abruptly bent vertically down and flat, terminal processes long, slender, pointed and decussate, penis erect. Legs slender, not bristly, coxae loosely setose, tarsi longer than their tibiæ, pulvilli small, empodium microscopic but fleshy. Wings normal, anal angle not projecting, costa encompassing entire wing, costal bristle present, auxiliary vein evanescent before the anterior crossvein, second vein long, third vein forked, discal cell complete, emitting three veins, anterior crossvein at two-fifths the length of the discal cell, the first basal cell therefore much longer than the second, anal crossvein arising rectangularly and uniformly curving into the lower side of the anal cell, no anal vein, fold in wing-membrane at root of second basal cell visible but not pronounced; calypteres with nearly straight edge and weak fringe.

Genotype : *N. eucera*, nov. sp.

Geographical distribution.

1. *N. eucera*, nov. sp. (1). — **Pl. 3, Fig. 21.**

Washington.

(1) ***Niphogenia eucera*, nov. sp. (Pl. 3, Fig. 21).** — Length 3 mm. Piceous black, covered with brown-gray but not heavy pollen. Antennæ entirely black, the pubescence of the third joint brown. About ten bristles in dorsocentral row and eight acrostichals. Eighth segment of female abdomen edged with numerous setæ, eighth sternite of male with long

5. GENUS CERATEMPIS, NOV. GEN.

Characters. — Male. Head spherical, eyes round, low-set, obliterating the cheeks, widely separated on both front and face, facets nearly uniform and rather large; a single orbital bristle opposite antennæ, ocelli large, ocellar bristles pronounced, upper occipital bristles well developed; first antennal joint cylindrical, twice as long as wide, with a few apical setulae above, second joint globose, with a whorl of setulae, third joint excessively long, being twice as long as the head, very slightly tapering to the blunt tip, heavily pubescent; proboscis minute, fleshy, palpi single-jointed, directed forward and bearing two inferior setulae.

Thorax normal, the notal bristles well developed, including one humeral, post-humeral, presutural, notopleural, two intraalar, one postalar, a complete row of dorsocentral and apical scutellar pair, metapleuræ with a vertical row of five setulae.

Abdomen loosely setose, not pitted, showing six compressed segments, the genitalia erect, with narrow central piece, the base of which is overlaid by a pair of downward extending flat valves, dorsal valves narrow erect, and penis slender deformed and projecting; seventh tergite of female fringed with close-set blackish setæ, the corresponding sternite long and narrow, ovipositor short and stubby, ending in two small erect hooks.

Legs slender, coxae loosely setose, remainder of legs with usual covering, no bristles other than a flexor grouping of stiff setulae toward tip of middle tibiae and a single small flexor bristle at apex of front tibiae; tarsi somewhat longer than their tibiae; pulvilli small, empodium microscopically setiform.

Wings broad, hyaline, veins yellowish, costa encompassing wing, fringed with marginal hairs, no stigma, auxiliary vein complete, ending in costa opposite fork of third vein, second vein long, ending near tip of wing, third vein forked, discal cell complete and rather large, emitting three posterior veins, anal crossvein perpendicularly recurved, rounding into the anal vein which continues only as a fold of the membrane beyond the anal cell.

Genotype: The following new species, which is of very delicate structure.

Geographical distribution.

1. *C. longicornis*, nov. sp. (1). — Pl. 3, Fig. 22.

Washington.

6. GENUS TRICHOPEZA, RONDANI

Trichopeza, Rondani, Dipt. Ital. Vol. 1, p. 150 [1856] (*Tricopeza*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 86 (1862); Röder, Wien. Ent. Zeit. Vol. 6, p. 169 (1887); Bezzi, Ann. Soc. Ent. France (6), Vol. 9, p. 121 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258, 261 (1903);

loose setæ along margin, male terminal processes testaceous, with black tip, penis testaceous, its tip deflected posteriorly, other pygidial parts piceous. Coxæ brownish yellow, legs piceous brown, the front knees a little lighter, middle tibiae of male with small thorn-like seriate flexor spines extending between middle and apex. Wings infumated, veins dark brown; calypteres and halteres yellow.

One male and twenty-five females, found by the author principally about the rills formed from melting snow on the Sluiskin slope above Paradise Park, Mount Rainier, Washington, August, 1917. Of the type lot single specimens were found along Van Trump Creek and on Eagle Peak, both localities also on Mount Rainier.

(1) ***Ceratempis longicornis*, nov. sp.** — Male. Length 2.8 mm. Entirely yellow, largely shining, the third antennal joint black, the ocellar spot and last tarsal joint blackish. Pubescence of third antennal joint gray, bristles of body black, hairs of legs dark. Upper valves of pygidium with black hairs, a pair of stiff black backward-directed setæ arising from near middle of pygidium, the flaring lower valves covered with downward-directed hairs.

Female. Third joint of antennæ a little shorter than in the male.

Eight specimens, Nasel River, and Ilwaco, in Southwestern Washington, July (Melander).

Kertész, Cat. Dipt. Vol. 6, p. 111 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 241 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45, 85 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 3 (1918); Brunetti, Fauna Brit. India. Dipt. Vol. 1, p. 371 (1920).

Characters. — Slender, long legged, dark colored species measuring about five millimeters.

Head rounded, eyes very large, occiput little convex; front broad in both sexes, the sides strongly converging toward the antennæ, no fronto-orbitals, ocelli large and not elevated, face obliterated by the contiguity of the eyes except for a short narrow space beneath the antennæ and at the oral edge, no cheeks; eyes of both sexes contiguous below, the middle anterior facets largest, finely pubescent, antennal excision strong; antennæ located high up on the head, strikingly long, three-jointed, nearly bare, the first joint lengthened and cylindrical, the second globose, the third greatly drawn out, tapering gradually and without articulation into the arista; proboscis slender, vertical, slightly shorter than the head, the labrum as long as the slender labium, palpi about one-fourth as long as the proboscis, rather cylindrical, one-jointed, decumbent and little hairy; occipital hairs sparse, uniserrate, enlarging above as bristles, a pair of long fine ocellar bristles present. Thorax arched, metathorax not declivous, no pubescence, bristles sparse but fine and long, one humeral, one posthumeral, no acrostichals, about four dorsocentrals, two notopleurals, two scutellars, one supra-alar and one postalar; pleuræ bare. Abdomen long and slender, not bristly, the segments marked with distinct lateral as well as transverse rows of pits; pygidium reflexed over the small eighth segment, consisting of a ventral piece which bears on the left front edge a prominent erect process and is tipped by a pair of small leaf-like valves, penis hidden or long, slender and curved; seventh segment of the female broadly emarginate above, the sides lobose, the seventh sternite acute, eighth segment retracted. Legs long and slender, front coxæ one-half as long as their femora and nearly twice as long as the posterior pairs, femora with some long fine flexor setæ, posterior tibiæ with long bristles, pulvilli small. Wings narrow, costa continuing around the entire margin, basal bristle present, no stigma, auxiliary vein distinct, straight, vanishing just beyond the end of the anal cell, third vein with a long fork, discal cell located before the middle of the wing, the sections of the fifth vein nearly two to one, second basal cell longer than the first, anal cell small, its crossvein strongly recurved and continuous with the under side of the anal cell, anal vein replaced by a fold, not attaining the margin, discontinuous with the anal cell, no anal lobe, no alula, marginal fringe long, calypters not at all fringed, their edge straight.

Type species : *T. longicornis*, Meigen (Pl. 7, Fig. 69). Westwood, Gen. Syn. p. 132 (1840), cited this species as the type of *Brachystoma*, but since Blanchard the same year assigned *vesiculosum* as the type of *Brachystoma* it is best to accept his decision.

Geographical distribution.

1. *T. albocincta*, Boheman, Oefv. Vet. Akad. Förh. Stockholm, Vol. 20, p. 80 N. Europe.
[1863] (*Brachystoma*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37
(3), p. 66 [1913] (*albicincta*); Ent. Tidskr. 1914, p. 80 (1914).
2. *T. fusca*, Brunetti, Rec. Indian Mus. Vol. 9, p. 31 (1913); Fauna Brit. India.
India Dipt. Vol. 1, p. 372, f. 35 (1920).
3. *T. longicornis*, Meigen, Syst. Beschr. Vol. 3, p. 12, pl. 22, f. 6, 7 [1822] Europe.
(*Brachystoma*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 344 (1834)
(*Brachystoma*); Zetterstedt, Dipt. Scand. Vol. 1, p. 363 [1842] (*Brachystoma*); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 52 [1851] (*Brachystoma*); Walker, Ins. Brit. Dipt. Vol. 1, p. 103, pl. 3, f. 4 [1851]
(*Brachystoma*); Zetterstedt, Dipt. Scand. Vol. 13, p. 5004 [1859]
(*Brachystoma*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 87 (1862);
Stein, Wien. Ent. Zeit. Vol. 22, p. 225 (1903); Lundbeck, Dipt.

Dan. Vol. 3, p. 243, f. 106, 107 (1910); Wahlgren, Ent. Tidskr. p. 31, p. 85 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 66 (1913).

4. *T. *sucina*, new name. Baltic Amber.
longicornis, Meunier, Ann. Soc. Sc. Nat. (Zool.) Vol. 7, p. 90, 108, pl. 7, f. 3-5 (1908).

7. GENUS DIPSOMYIA, BEZZI

Dipsomyia, Bezzl, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 389 (1909).

Characters. — Head rounded, ocellar prominence with long, well-separated bristles; face bare, cheeks narrow; antenna short, the third joint very short, bearing a thick arista which is twice as long as the antennæ; proboscis thick, blunt, vertical or slightly inclined, the palpi large, exserted perpendicularly, black. Thorax but little convex, opaque cinereous; dorsocentral bristles strong, extending to the front margin, alternating large and small bristles, acrostichals distinct, biserrate, scutellum with marginal hairs in addition to the two bristles; metapleuræ bare. Legs black, front coxae one-fourth as long as their femora, with projecting setæ in front, front femora incrassate, spinose at their base, front tibiae pectinate within with short bristles, pulvilli distinct but small. Wings elongate, narrow, rather pointed, the axillary angle very obtuse, no stigma, costa extending around the hind margin, humeral crossvein present, auxiliary vein evanescent apically, first vein ending beyond the middle of the wing, third vein furcate, its pedicel arising near the middle of the second basal cell, fourth vein arising at the base of the anal cell, discal cell rather large, complete, emitting three posterior veins, the second posterior cell sessile, anal cell shorter than the second basal, the crossvein rounded and distinct from the anal fold.

Type species: *D. spinifera*, Bezzl, from South America, the only known species. Bezzl states that this genus seems intermediate between *Philotutra* and *Hilarempis*, resembling the former in its general venation, shortened antennæ and lengthened bristles, but differing in its lengthened proboscis, thickened front femora and long second basal cell.

Geographical distribution.

1. *D. spinifera*, Bezzl, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 389, f. 14 Chili.
 (1909).

8. GENUS PROCLINOPYGA, NOV. GEN.

Characters. — Slender olivaceous species with black legs. Head globular, eyes of both sexes widely separated, facets uniform, bare, front wide the sides converging toward the antennæ, ocelli large, not elevated, a pair of strong diverging distant ocellar bristles behind which on the level of the posterior ocelli is a pair of smaller diverging approximated lesser ocellars; sides of the face parallel, the face of the female broader than in the male; cheeks small and narrow but separated from the face; antennæ located above the middle of the head, plainly three-jointed, the basal joints subequal, the third joint conical but with concave sides, arista terminal, as long as the third joint, thickened, geniculate at its attachment, three-jointed, the basal segment minute and quadrate, the apical part consisting of a microscopic hair; proboscis very short, thick, fleshy, and blunt, palpi short, horizontal, rather clavate and hairy; occipital hairs biserrate, the uppermost bristle-like. Thorax broader than the head, entirely pollinose, no pubescence, bristles prominent, metathorax longer than the scutellum and not declivous, one humeral, one posthumeral, an oblique row of three notopleurals, several supra-alars, one postalar,

usually four scutellars, six to eleven uniseriate dorsocentrals, the foremost large and placed toward the humeri, acrostichals irregularly biserrate; pectus with a few microscopic hairs, metapleuræ with a vertical row of five setulæ. Abdomen slender, twice as long as the thorax, setulose, the lateral pairs of pits oblique, the sixth, seventh and eighth segments of the male shallow to accomodate the large reflexed epipygium, from the anterior base of the epipygium two strong dorsal prongs extend forward, middle valves large, convex, ovate, terminal valves small and hastate, penis thick, curving forward; eighth segment of the female entirely retracted within the full-sized and compressed seventh segment. Legs long and slender, specialized in the male by bristles and a lengthening of the tarsus, front coxæ half as long as the femora and one-third longer than the posterior pairs. Wings almost cuneiform, the anal lobe slightly evident, costa continuing around the entire wing, with a long basal bristle and small setulæ, auxiliary vein straight, vanishing apically opposite the anterior crossvein, first vein extending two-thirds the wing-length, third vein with a long fork, the main branch ending at the tip of the wing, discal cell located before the middle of the wing, scarcely longer than the second basal, the anterior crossvein near the middle of the discal cell, the first basal cell therefore much longer than the second, pedicel of the second and third veins arising near the humeral crossvein, pedicel of the fourth vein arising directly under the humeral crossvein and close to the base of the second basal, anal cell narrow, its crossvein strongly recurved, extending back one-fourth the length of the anal cell before fusing with the anal vein, which is weak, gradually evanescent and does not reach the hind margin, no alula, hairs of the hind margin of the wing very small; calypteres moderate, their hairs fine and silky.

Type species : *P. amplexens*, nov. sp. (Pl. 7, Fig. 65). The genus is apparently a close relative of *Dipsomyia*, and with it shows many resemblances to the Empidinæ. The neuration is quite constant among the following species.

Geographical distribution.

- | | |
|---|---|
| 1. <i>P. amplexens</i> , nov. sp. — Pl. 7, Fig. 65. | California. |
| 2. <i>P. exorrecta</i> , nov. sp. | Montana. |
| 3. <i>P. fistulator</i> , nov. sp. — Pl. 8, Fig. 75. | Quebec. |
| 4. <i>P. monogramma</i> , nov. sp.
var. <i>flavicoxa</i> , nov. var. | W. United States.
Idaho, Washington. |
| 5. <i>P. solivaga</i> , nov. sp. | Washington. |

KEY TO THE MALES OF PROCLINOPYGA

- i. Under side of middle femora bearing three pale setæ at base, middle tibiae swollen near middle and there bearing many flexor setulæ, middle metatarsus as long as tibia; sixth abdominal segment with many long black bristles, sides of abdomen with short pale hairs. P. AMPLECTENS, nov. sp. (1).

(1) ***Proclinepyga amplexens*, nov. sp. (PI. 7, FIG. 65).** — Male. Length 4 mm. Olivaceous above, cinereous, almost white pruinose, on the sides and beneath, face whitish, mouthparts black; first joint of the antennæ short-cylindrical, with a few setulæ above, second joint with a crown of setulæ, third joint pubescent, twice as long as wide; hairs of lower occiput sparse and white. Ten dorsocentrals, about nine irregular pairs of acrostichals not extending on prescutellar area, no scutellar setulæ, pronotum with a pale hair on each side, metapleural hairs white. Lateral and ventral hairs of the abdomen short and white, dorsal setulæ black; sixth tergite with a cluster of black hairs, epipygium black. Legs entirely black, coxal hairs white, front femora equal to their tibiæ, the front tarsi one-half longer, middle tibiae somewhat longer than their femora, closely setulose from middle to apical fourth, broadly excised within toward tip, middle tarsi greatly lengthened, nearly twice as long as their femora, the metatarsus nearly as long as the remainder, hind legs simple the femora with a few black extensor setulæ, posterior femora with a long white bristle beneath near the base, all the pulvilli of microscopic size, empodium similarly small, compressed. Wings three times as long as broad, widest beyond

- Under side of middle femora more extensively setose, middle tibiæ not deformed and as long as basal two joints of their tarsi; sixth segment of abdomen not densely setose 2.
2. Setæ of middle femora particularly conspicuous along the middle, middle tibiæ densely setulose within; dorsal prong of pygidium with black velvety knob at middle and bluntly tong-shaped tip
- Setæ of middle femora evenly distributed, middle tibiæ scarcely setulose; dorsal prong of pygidium without velvety knob. 3.
3. Setæ of middle femora distant, about eight in number; dorsal prong of pygidium jet black, quadrate at base and then suddenly narrowed, distally ribbon-like, preapical valve split at apex, apical valve strap-shaped, side valve large and setose behind
- Setæ of middle femora closer, about twelve in number; side valve not setose, preapical valve pointed 4.
4. Six to eight dorsocentrals, four scutellars; side valve subshining, rounded quadrate P. *FISTULATOR*, nov. sp. (2).
- P. *EXPORRECTA*, nov. sp. (1).
- P. *MONOGRAMMA*, nov. sp. (3). 5.

the discal cell, slightly infumated, a very faint stigma, sections of the fourth vein proportioned 1.1 : 0.8 : 1 ; 2. of the fifth vein equal, posterior crossvein not very oblique; hairs of the calypteres white.

Female. Seven dorsocentrals. Tarsi not lengthened, only a little longer than their tibiæ, no basal bristles on the femora, tibiæ simple, unarmed except that the middle ones have two extensor and one apical small setulæ. Seventh segment of the abdomen with ciliate margin.

One specimen of each sex, received from Professor Aldrich, who collected them at Woodside, California, April 25, 1909.

(1) ***Proclinoxyga exorrecta*, nov. sp.** — Male. Length 3 mm. Heavily dusted with blackish brown above, dark gray on sides. Facets uniform. About eight dorsocentrals, four scutellars, metapleural setulæ black. Abdomen setose, the basal and ventral hairs pale, sixth segment with long setæ. Epipygium stout, the dorsocentral prongs deformed, loosely long-ciliate in front, swollen and black-velvety at middle and curved and bluntly tong-like at tip, apical valves large spoon-shaped and densely hairy beneath, side valves triangular, setose behind, penis shaped like a long stalked open? - mark. Lower side of middle femora with a group of six long stout bristles near the middle and with a row of pale flexor setæ which form a short fringe before the knee, hind femora with about eight extensor and seventeen flexor outstanding setæ, middle tibiæ finely setulose within more pronounced at tip, middle tarsi not abnormally lengthened, the metatarsus about two-thirds as long as the tibia. Stalk of halteres brown; calypteres and fringe yellow.

Type, Saltese, Montana, August 22, 1916 (Melander). A female from the Gallatin Ranger Station of Yellowstone National Park, July 28, 1923 (Melander) offers no distinctive characters. Its last abdominal segment is compressed and shiny.

(2) ***Proclinoxyga fistulator*, nov. sp. (Pl. 8, Fig. 75).** — Male. Length 3 mm. Heavily dusted with blackish brown above, dark gray on sides. Lower facets very slightly larger than upper. About eight dorsocentral, acrostichals long, four scutellars, metapleural setulæ black. Abdominal hairs not long except on sixth segment, those at base and on venter pale; epipygium stout, dorsal prongs wide at base then suddenly narrowed to the curled ribbon-like end which is ciliate within at tip, preapical valves narrow chitinized and excavated at tip to suggest an opposing thumb and forefinger, apical valves narrow chitinized and loosely hairy, side valves quadrangular and setose behind, penis tubular rectangularly bent at basal third, erect over middle third and slightly curved on apical third. Middle femora with a long pale flexor seta at base, the black flexor setæ sparse and inconspicuous, hind femora with about four flexor and eight extensor short setæ along basal half, tibiæ and tarsi normal. Stalk of halteres brown; calypteres and fringe yellow.

Holotype, Megantic, Quebec, 18 July, 1923 (Curran), deposited in Canadian National Museum.

(3) ***Proclinoxyga monogramma*, nov. sp.** — Male. Length 2.2 mm. Piceous black, the lower pleuræ and coxae somewhat cinereous. Lower facets a little enlarged. Eight dorsocentrals, four scutellars, acrostichals long, three to six black metapleural setulæ. Abdomen coarsely black-hairy, the bristles of the sixth segment not pronounced, epipygium relatively small, compressed, subshining, lateral valves not setose, triangular, the inner corner connected with its mate by an arched chitinous bridge, dorsal process straight, preapical process long and rod-shaped, apical valve small, narrowly

- Eleven dorsocentrals, six scutellars; abdomen coarsely setose,
side valve gray pruinose, twice as high as wide P. SOLIVAGA, nov. sp. (1).
5. Coxæ, halteres and calyptères blackish P. MONOGRAMMA, s. str.
- Apex of coxæ, base of femora, stalk of halteres, and calyptères
yellowish var. FLAVICOXA, nov. var.

9. GENUS OREOTHALIA, MELANDER

Oreothalia, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 232 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 254, 260 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 224 (1908); Kertész, Cat. Dipt. Vol. 6, p. 119 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 580 (1910).

Characters. — Slender, opaque black species of the habitus of *Clinocera*, but with the third vein simple and the anal crossvein abruptly turned back. Head oval, hanging vertically; eyes finely pubescent, widely separated on the face and front, facets uniform, emargination at the antennæ distinct; antennæ inserted high up, the front very short, ocellar tubercle prominent, cheeks narrow, incised and separated from the face; proboscis very short, thick and flat, palpi small and broad; antennæ short, contiguous, three-jointed, but the basal joint inset, third joint oval, its terminal arista apparently single-jointed, thickened, not hairy, nearly twice as long as the antennæ; ocellar and upper occipital bristles present, lower occiput with sparse hairs. Thoracic bristles long and strong, no acrostichals, five dorsocentrals, one humeral, two intraalars, one supraalar, one postalar, a few notopleural setulae near the root of the wing, metapleuræ with a cluster of hairs; prosternum shorter than the mesosternum. Abdomen as long as the thorax, the first six segments of the male cylindrical, then suddenly smaller to accommodate the reflexed epipygium, penis conical, upper valves directed forward and thin and flat toward the tip, abdomen of the female blunt. Legs slender, front coxæ two times as long as the posterior pairs, front femora with strong thorn-like setulae below on the apical two-thirds, hind tibiæ setose, no tibial spurs; claws long and slender, empodium long. Wings slender, costa continuing around the entire margin, a pair of strong basal bristles, costal setulae minute, auxiliary vein ending in the costa, first vein terminating before the middle of the wing, stigma very faint, third vein simple, discal cell complete, pentagonal, posterior cells sessile, fourth vein arising at one-third the length of the anal cell,

triangular and short-hairy; penis thickened, curving forward, the tip recurved. Legs ordinary, the middle femora below and the hind ones above with a row of setæ. Stalk of halteres piceous; calyptères blackish

Female. Last abdominal segment compressed and shining.

Type, Troy, Montana, August 10, 1916 (Melander). Nine Paratypes, all collected by author, at St. Regis Pass, Idaho-Montana; Priest Lake, Idaho; Lake Keechelus, Lake Cushman, Ramparts of Mount Rainier, Washington; and Shasta Springs, California.

Var. *flavicoxa*, vat. n. Stalk of halteres, calyptères, apex of coxæ, trochanters and more or less of base of femora yellow.

Type, Priest Lake, Idaho, August 1, 1916 (Melander). The penis is abruptly elbowed at the middle and its tip is widely recurved. Paratypes, male and female, Lake Keechelus, Washington. The penis of the paratype is gently curved.

(1) **Proclinopyga solivaga**, nov. sp. — Male. Length 2.2 mm. Notum olivaceous black, cinereous brown toward notopleural suture and on middle of abdomen, pleuræ and sides of abdomen cinereous. Facets uniform. About eleven dorsocentrals, acrostichals long, six scutellars, two pale metapleural setulae. Abdomen coarsely hairy, especially on sixth segment, the hairs of first segment and of sides of second pale, venter nearly bare, pittings very large; epipygium robust, erect, side valves distorted-oblong, heavily pollinose, hairy but not setose behind, dorsal prongs thin, curved preapical valves small and wiry, apical valves with small pointed tip, penis flat, bending forward near middle, its compressed base pollinose. Legs without distinctive characters. Stalk of halteres piceous; calyptères blackish.

Holotype, Van Trump Park, Mount Rainier, Washington, 21 July, 1922 (Melander).

the second basal cell therefore small, no fold in the wing membrane beneath the humeral crossvein, anal crossvein abruptly reflexed, the outer border of the second basal and anal cells nearly continuous, oblique, anal vein thin and short, anal angle very poorly developed, blunt.

But a single species is known, occurring along streams in wooded regions of the North-western United States.

Geographical distribution.

1. *O. pelops*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 233 (1902). — Idaho, Washington.

Pl. 2, Fig. 17.

10. GENUS HELEODROMIA, HALIDAY

Heleodromia, Haliday, Ent. Mag. London, Vol. 1, p. 159 (1833); Curtis, Brit. Ent. Vol. 8, p. 513 (1834); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 250, 260 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 224 (1908); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 550 (1910); Engel, Deutsche Ent. Zeitschrift, 1918, p. 3 (1918).

Macrocera, Zetterstedt, MS. (not Loew, 1845 [*amber Tipulidae*]; not Reitter, 1873 [*Coleoptera Nitidulidae*]), Fauna Ins. Lappon. p. 572 (1838); Scudder, Nomencl. p. 199, Index, p. 185 (1882).

Microcera, Zetterstedt (not Meigen, 1803 [= *Pipunculus Latreille*]); not Mannerheim, 1830 [*Coleoptera*]; not Lioy, 1864 [= *Hydrotæa Desvoidy*]); Zetterstedt, Fauna Ins. Lappon. p. 572 (1838); Dipt. Scand. Vol. 1, p. 440 (1842); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 569 (1910).

Sciadromia, Haliday, Westwood: Gen. Syn. p. 132 (1840); Walker, Ins. Brit. Dipt. Vol. 1, p. 107 (1851); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 563 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 85 (1862); Röder, Wien. Ent. Zeit. Vol. 6, p. 169 (1887); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 122 [1889] (*Sciodromyia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 17, p. 390 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 345 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257 (1903); Kertész, Cat. Dipt. Vol. 6, p. 129 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 604 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45, 86 (1910).

Characters. — Slender, black dusted species with narrow face, lengthened proboscis, globose pygidium, simple third vein and truncate anal cell. Head rotund, eyes bare, narrowly separated on the face, the vertex broad, lower anterior facets much enlarged in both sexes; basal joint of the antennæ minute, middle joint spherical, third joint oval or ovate, with a slender, terminal, apparently two-jointed arista, in reality the first joint of the arista is a terminal process of the third joint; proboscis cylindrical, obliquely projecting, shorter than the head, somewhat fleshy, palpi small, oval, hairy; cheeks obliterated; ocellar and vertical bristles present, lower occiput nearly bare. Thorax short, completely coated with fine dust, bristles small to moderate, one humeral, four dorsocentral, one supraalar, one postalar two scutellar, no acrostichals; metathorax not declivous, prosternum shorter than the mesosternum, metapleuræ bare, at most with one or two minute hairs. Abdomen long, segments marked with pits, the terminal segments of the male not reduced, pygidium large, globose, terminal, the valves small, uncinate, horny, penis hidden; last (seventh) tergite of the female densely pectinate apically with long setæ. Legs long, slender and blackish, no bristles or spurs, the claws, pulvilli and empodium minute. Wings narrow, cuneate, the anal angle scarcely developed and broadly rounded, costa encompassing the entire wing, no basal bristle or costal setulae, auxiliary vein attaining the costa, first vein ending beyond the middle of the wing, third vein simple, discal cell complete and long, posterior cells sessile,

a fold below the humeral crossvein, ourth vein arising near the base of the anal cell, its origin not abrupt, anal cell apically truncate by the rectangular anal crossvein, the anal vein continuing as a spur beyond the lower angle of the anal cell.

Taxonomy. — Haliday erected the genus *Heleodromia* for four species, *immaculata*, *bipunctata*, *stagnalis* and *fontinalis*. Curtis (Brit. Ent. 1834) designated *immaculata* as the type for *Heleodromia*, and Macquart (Hist. Nat. Dipt. Vol. 2, p. 658, 1835) repeated this. Macquart removed *stagnalis* and *bipunctata* to a new genus *Hydrodromia*, to which also belongs *fontinalis*. Mik (Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 325, 1881) placed *bipunctata* in his new genus *Kowarzia*. Thus by elimination, *stagnalis* is left as the type of the group *Hydrodromia*. Haliday, in a manuscript note, published in Westwood's Introduction, Generic Synopsis, p. 132 (1840), proposed to substitute *Sciodromia* as the generic name for *immaculata*, but did not give a valid reason for the change. *Microcera* was erected by Zetterstedt for *rostrata*, which is a synonym of *immaculata*: his manuscript name for this insect was *Macronura*.

The species of *Heleodromia* occur in Europe and in Western North America.

Geographical distribution.

1. *H. curtipes*, Becker, Deutsche Ent. Zeitschr. p. 645 [1910] (*Sciodromia*). Corsica.
2. *H. immaculata*, Haliday, Ent. Mag. London, Vol. 1, p. 159 (1833); Curtis, Europe. Brit. Ent. Vol. 8, p. 513 (1834); Macquart, Hist. Nat. Dipt. Vol. 2, p. 658 (1835); Meigen, Syst. Beschr. Vol. 7, p. 93 [1838] (*Hemero-dromia*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 505 (1849); Scholz, Zeit. Ent. Breslau, Vol. 5 (19), p. 56 [1851] (*Sciodromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 107, pl. 4, f. 1 [1851] (*Sciodromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 86 [1862] (*Sciodromia*); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 98 [1892] (*Sciodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 86 [1910] (*Sciodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 68 [1913] (*Sciodromia*).
fuscipennis, Roser, Correspondenzbl. Landw. Ver. Würtemb.-Stuttgart, Vol. 1, p. 53 [1840] (*Hemero-dromia*).
rostrata, Zetterstedt, Fauna Ins. Lappon. p. 572 [1838] (*Microcera*); Dipt. Scand. Vol. 1, p. 440 [1842] (*Microcera*); ibidem, Vol. 8, p. 3046 [1849] (*Microcera*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 176 [1861] (*Microcera*); Siebke, Nyt. Mag. Naturv. Vol. 12, p. 109 [1864] (*Microcera*).
3. *H. pectinulata*, Strobl, Glasnik, Zem. Mus. Bosn. Herceg. Sarajevo, S. Europe. Vol. 10, p. 399 [1898] (*Sciodromia*); Wissensch. Mitth. Bosn. Herceg. Sarajevo, Vol. 7, p. 561 [1900] (*Sciodromia*).
4. *H. pullata*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 345 [1902] W. United States. (*Sciodromia*); Kertész, Cat. Dipt. Vol. 6, p. 123 [1909] (*Hydrodromia*); Engel, Deutsche Ent. Zeitschr. 1918, p. 48 [1918] (*Hydrodromia*). —
PI. 7, FIG. 68.
- . **Sp. innom.* Meunier, Miscell. Ent. Vol. 7, p. 178 (1899). Baltic Amber.

II. GENUS ROEDERIOIDES, COQUILLETT

Roederioides, Coquillett, New York State Mus. Bull. Vol. 47, p. 585 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 239 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257, 263 (1903); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 363, note (1905); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 601 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 264 (1918).

Characters. — Slender brownish black pollinose species with long yellow legs. Head vertically elongate, the cheeks rostriform, two-thirds the eye-height, not sutured from the face but with a small impression just beneath the eye, truncate at the oral margin, narrowly notched at the epistome, face bare; proboscis nearly as long as the height of the head, rather slender and tapering, inflexed, chitinized, filling the oral cavity, opaque black except at the apex, without trace of the labella; palpi small, projecting downward, incumbent; antennæ touching, the third joint compressed ovate, but nearly straight above, pointed apically, bearing a moderately thick, two-jointed, downward-directed arista whose length is slightly more than the third joint; one pair of projecting ocellar bristles, one pair of verticals, setulæ of occiput sparse. Thorax not elongate, the prosternum half as long as the mesosternum; five dorsocentrals, no acrostichals, one humeral, one posthumeral, one small notopleural, one small supraalar, one postalar, four scutellar and also discal and marginal setulæ on the scutellum; metapleuræ with hairs. Abdomen comprising eight segments (♀), two small terminal styles; epipygium of the form of *Clinocera*, middle and upper valves hemispherical, nearly equal in size, keel not compressed, penis recurved at the tip. Legs slender, without setulæ or long hairs, tarsi peculiar, the first joint long, equalling the remainder, the second, third and fourth joints very short and similar, of the anterior pairs not longer than thick, fifth joint of all the tarsi continued dorsally at the apex as a conical process, so that the claws do not arise at the end of the tarsi, empodium fleshy and larger than the pulvilli, tactile hairs much reduced, claws long and curved. Costa encompassing the entire wing, a basal costal bristle present, no costal setulæ, no stigma, third vein forked, discal cell complete, pointed, second posterior cell acute and with a short petiole, no fold under the humeral crossvein, second basal cell short, as in *Clinocera*, anal cell almost rounded at the apex, coextensive with the second basal, no anal vein, cilia of the hind margin longer than the anterior crossvein.

Type species : *R. juncta*, Coquillett (manuscript name *Needhami* Coquillett) is the only representative. The species measures two and a half millimeters and was found in the mountainous woods of North-eastern New York. Mr. C. W. Johnson has taken the species on Mt. Washington, New Hampshire.

Geographical distribution.

1. *R. juncta*, Coquillett, New York State Mus. Bull. Vol. 47, p. 585 (1901); E. United States.
Needham, ibidem, p. 581, pl. 15, f. 5-8 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 239 (1902); Engel, Deutsche Ent. Zeitschr. 1918, p. 264 (1918). — Pl. 2, Fig. 19.

12. GENUS CLINOCERA, MEIGEN

Clinocera, Meigen, Illiger Mag. Ins. Vol. 2, p. 271 (1803); Rondani, Dipt. Ital. Vol. 1, p. 149 (1856); Loew, Wien. Ent. Monatsschr. Vol. 2, p. 238-253, 257-262 (1858); Schiner, Fauna Dipt. Aust. Vol. 1, p. 84 (1862); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 320 (1881); Bigot, Bull. Soc. Ent. France, Vol. 12, p. 102, 103, 118 (1887); Mik, Ent. Nachr. Berlin, Vol. 14, p. 42 (1888); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 120 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 240 (1902); Williston, Man. N. Amer. Dipt. p. 224 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 300 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 213 (1910); Wahlgren, Ent. Tidsk. Vol. 31, p. 45, 81 (1910); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 65, 68 (1911); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 364 (1920).

Atalanta of authors, Meigen, Nouv. Classif. Mouches, p. 31 (1800); Hendel, Verh. Zool.-bot. Ges. Wien, Vol. 58, p. 60 (1908); Kertész, Cat. Dipt. Vol. 6, p. 120 (1909); Oldenberg, Arch. Naturges. Berlin, Vol. 80, A, p. 9, 92 (1915); Engel, Deutsche Ent. Zeitschr. 1918, p. 1-80, 197-268 (1918).

Bergenstammia, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 (1881); Bigot, Bull. Soc. Ent. France, Vol. 12, p. 102 (1887); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 246, 262 (1903); Kertész, Rovert. Lapok, Vol. 18, p. 67 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 75 (1918).

Clinocera, s. str., Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 320 (1881); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248, 262 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 525 (1910); Kertész, Rovert. Lapok, Vol. 18, p. 67 (1911).

Heleodromia of authors, Haliday, Ent. Mag. London, Vol. 1, p. 159 (1833); Westwood, Gen. Syn. p. 132 (1840); Walker, List Dipt. Brit. Mus. Vol. 3, p. 503 [1849] (*Heliodromia*); Ins. Brit. Dipt. Vol. 50, p. 104 (1851); Bigot, Ann. Ent. Soc. France (3), Vol. 5, p. 561 (1857); Bull. Soc. Ent. France, Vol. 12, p. 102 (1887); Lundbeck, Dipt. Dan. Vol. 3, p. 217 (1910); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 67 (1911).

Hydrodromia, Macquart (not *Hydrodromia*, Scudder, Nomenclator 167 (1882); which is mistake for *Hydromyia*, Desvoidy), Hist. Nat. Dipt. Vol. 2, p. 658 (1835); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 251, 262 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 554 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 37 (1918).

Kowarzia, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 325 (1881); Bigot, Bull. Ent. Soc. France, Vol. 12, p. 103 (1887); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 251, 262 (1903); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 67 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 64 (1918).

Paramesia, Macquart (not Stephens, *Lepidoptera*, 1829), Hist. Nat. Dipt. Vol. 2, p. 656 (1835); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 584 (1910).

Phæobalia, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 [1881] (*Phæobalia*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255, 262 [1902] (*Phæobalia*); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 67 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 50 (1918).

Characters. — Medium sized, slender, olive-gray tomentose, long-legged flies that occur along water courses and in damp woods. Head globular, slightly wider than high in those species with small cheeks, sometimes the mouth is prolonged far downward and the head is then much higher than wide, eyes widely separated, the face relatively broad. Occiput rather flat, the neck located above the center but not far from the center, bristles and setulæ prominent, usually three pairs of proclinate vertical bristles; front broad, much shorter than the face, the ocelli on a prominent tubercle, the ocellar bristles divergent and proclinate, inserted closer together than the posterior ocelli; face broad, the lower edge transverse or peculiarly emarginate at the middle, cheeks variable, a more or less distinct suture between the genæ and the buccæ, which sometimes occurs as a deep notch. Eyes round or oval, densely pubescent, the facets uniform, the emargination at the antennæ small. Antennæ short, three-jointed, located high on the head and almost touching, but the sockets distinctly separated, the second joint globular, the third joint oval or onion-shaped, sometimes drawn out as a short process, with a thickened, two-jointed bare arista, whose basal joint is minute and whose apex is either blunt or pointed. Proboscis very short and thick, directed vertically down, the labrum formed into a broad chitinous visor overlapping the proboscis; palpi short, somewhat clavate, projecting, and usually but little hairy. Thorax long and narrow, with a flattened area in front of the scutellum, thickly coated with olivaceous

tomentum, prothorax visible from above, the prosternum long, but a little shorter than the mesosternum, flared out around the base of the coxae and sometimes medially sulcate; bristles strong, one or more humeral, one posthumeral, five or more dorsocentral, one or two notopleural, one or no supralar, one postalar, two scutellar bristles, acrostichals biserrate or wanting, sometimes the disk or the margin of the scutellum setulose and sometimes there are alternating setulae in the dorsocentral rows; metapleuræ with a group of upturned hairs, prosternum more or less hairy. Abdomen tomentose or pruinose, of the male the last two segments are small and support the forward-directed epipygium, which comprises a hood-shaped ventral piece terminated by the geniculate slender penis and bearing the circular convex lateral valves, to which are articulated the various-shaped and complicated upper valves. Legs slender and elongate, front coxae about one-half their femora in length, more or less setose or hairy along their front surface, the front femora sometimes a little thicker basally and variously provided with setulae, preapical femoral bristles or setulae sometimes developed, sometimes a characteristic oblique comb of setulae on inner side of front femora just before the knee, tibiae without apical spurs, the hind tibiae sometimes bearing setæ, pulvilli small, empodium narrow and about as long as the claws except in *Bergenstamia* where it is practically absent. Wings cuneiform, no anal angle, the costa strong around the entire margin, costal hairs long, costal setulae more or less developed, in *Euclidia* the setulae are prominent, humeral crossvein distinct, auxiliary vein entire, gently curving into the costa, first vein terminating near the middle of the wing, stigma prominent or poorly developed, marginal cell elongate, third vein forked, the upper branch sometimes connected with the second vein by an adventitious crossvein, in which case there are three submarginal cells, discal cell complete, pointed or blunt at its apex, emitting three posterior veins, the second posterior cell sessile or acute at its base, second basal cell small the fourth vein starting beyond the base of the anal cell with an abrupt bend, anal and second basal cells coextensive, the anal crossvein recurved or reflexed and continuing uninterrupted under the anal cell, anal vein distinct from the anal cell, or wanting, marginal cilia usually short.

This group of the *Empididae* was subdivided by Mik in 1881 into ten genera. In doing this Mik was ahead of his time, for even yet many of his characterizations are not considered to have generic value. Two of his groups, based on the structure of the face and cheeks, are here preserved as of generic rank, the remainder being considered as subgenera of the primary two. Mik's distinctions have not always been correlated in the North American species, nor in the subsequently discovered European ones, but in the main his selection of such characters as the structure of the cheeks and of the epistome, the formation of the empodium and the variations in chaetotaxy, showed a keen appreciation of characters hitherto neglected. In 1918 Engel published an extended critical review of the subgenera and species of *Clinocera* under the name *Atalanta*.

The species of *Clinocera* are very distinctive in appearance. The elongate thorax with olive-brown back and pruinose sides, the long slender black legs, short thick proboscis and cuneate wings, which are often spotted, are characteristic of the Clinocerans. Further distinctions exist in the furcate third vein, the small second basal cell and the rounded anal cell. The species are all found about damp woods, flying along water-courses or sometimes about ponds. They are boreal, occurring mainly in the North, in mountainous regions, some even appearing with the melting of the snow. They rarely occur in swarms, but while they are not rare insects, they are not easily captured, for they remain close to the water, fearlessly frequenting waterfalls, or running over the wet stones or even on the surface of the water, and thus they can be taken in the net only with difficulty. As adaptations to this life there have been developed the peculiar tomentose body covering, the dense pubescence of the eyes, the long legs and wings and the strange sensitive hairs and empodium at the apex of the tarsi.

GENUS CLINOCERA, MEIGEN

Characters. — To this division of the group *Clinocera* belong those species in which the lower part of the head is not produced downward beneath the eyes. The cheeks are thus narrow, and are deeply incised, even to the margin of the eye, and thus the sides of the face are entirely separated from the lower occiput or cheeks. The pulvilli and empodium are developed except in the subgenus *Bergenstammia*. None of the species has acrostichal or interstitial dorsocentral setulæ, and except in *Bergenstammia* there are no setulæ on the disk of the scutellum.

The following subgenera are recognized :

Clinocera, in the strict sense. Lower edge of the face transverse and not excised; discal cell rather sharp apically, wings not or scarcely spotted, costal setulæ usually minute, stigma wanting or faint.

Type species : *nigra*, Meigen (Pl. 2, Fig. 18). The only species mentioned in the original description. Some recent papers, unfortunately including Kertész' Catalogue of the Diptera of the world, have adopted Meigen's 1800 name *Atalanta* in place of the universally known *Clinocera*, and would call the subfamily the *Atalantinae*. As Meigen mentions no species in this early paper the type dates from Hendel's interpretation in 1908, reason enough for disregarding the name *Atalanta*.

SUBGENUS HYDRODROMIA, MACQUART

Characters. — Lower edge of the face entire and transverse, not notched; discal cell blunt apically, wings more or less obviously spotted on the crossveins and on the forks of the veins but without a distinct stigma, costal setulæ variable, sometimes an extra crossvein connecting the upper fork of the third vein with the second vein and thus producing three submarginal cells.

Type species : *H. stagnalis*, Haliday. Macquart included two species in *Hydrodromia*, *stagnalis* and *bipunctata*, the latter being removed by Mik to his genus *Kowarzia*. By elimination, as Coquillett noted in 1903, *stagnalis* becomes the type of *Hydrodromia*. *Heleodromia* of various authors, based on *stagnalis*, is not the same as Genus 10, *Heleodromia*, based on *immaculata*. *Paramesia Wesmælii*, Macquart, which was designated by Coquillett in 1903 as the type of *Paramesia*, is congeneric with *H. stagnalis*.

SUBGENUS PHÆOBALIA, MIK

Characters. — Lower edge of the face with a median excision, the membrane filling the notch depressed and more or less carinate; discal cell blunt at the tip, wings spotted as in *Hydrodromia*, but the stigma is more distinct.

Type species : *P. trinotata*, Mik, by Coquillett's designation, 1903. Engel has unnecessarily specified *P. dimidiata*, Loew as the genotype.

SUBGENUS KOWARZIA, MIK

Characters. — Face with several scattered hairs, its lower edge incised in the middle, the notch filled by a depressed membrane which is more or less carinate; discal cell blunt at the tip, wings not spotted, the stigma poorly developed.

Type species : *K. barbatula*, Mik, by Coquillett's designation, 1903.

SUBGENUS BERGENSTAMMIA, MIK

Characters. — Pulvilli and empodium vestigial or wanting; face not continued beneath the eyes, the lower edge notched in the middle, the cheeks divided by a distinct suture; discal cell blunt apically, wings not marked; no acrostichals, five or more dorsocentrals, scutellum with setulae in addition to the apical pair of bristles; large sized species.

Type species: *B. nudipes*, Loew, the only one mentioned by Mik.

A KEY TO THE NORTH AMERICAN SPECIES OF CLINOCERA, S. LAT.

1. Three submarginal cells present; no acrostichals 2.
- Two submarginal cells present 8.
2. Discal cell usually longer than the second posterior;
posterior femora without bristles; thorax bivittate
and with a white median vitta, veins undulating . HYDRODROMIA UNDULATA, nov. sp. (1).
2. Discal cell shorter than the second posterior 3.
3. Wings lightly infumated, no stigma, veins not
undulating CLINOCERA TRUNCA, nov. sp. (2).
3. Wings marked with nebulous spots 4.

(1) **Clinocera (Hydrodromia) undulata**, nov. sp. — Male. Length 3 mm. Olivaceous above, cinereous below, thorax narrowly biliniate with brown and centrally in front with a conspicuous cinereous vitta. Lower half of the face white, upper part blackish, front olivaceous brown, the vertical bristles strong, cheeks completely separated from the face, narrow, palpi elliptical with short sparse black hairs. Five strong dorsocentral bristles, no setulae, metapleural hairs fine, dense and yellow, pectus nearly bare. Coxal hairs sparse and yellow, legs black, front tibiae as long as their femora, front femora with two sparse rows of about eight regularly placed, short setae, hind tibiae with about eight extensor setae on the apical half, pulvilli and empodium normally strong and whitish. Halteres black, calypteres dark and with a pale yellow fringe. Wings almost hyaline but with broad darker spaces about the crossveins, stigma elliptical, costal setulae small, veins thin but strong, undulating, three submarginal cells, the third twice as long as its costal margin, discal cell longer than the broadly sessile second posterior cell, its outer third with parallel sides, sections of the fifth vein equal, anal crossvein recurved, anal vein a weak fold.

Type, Moscow Mountain, Idaho, July 6, 1912; paratype, same locality, Sept. 16, 1917 (Melander).

A female from Mount Rainier, Washington, August 3, 1905, submitted by Professor Aldrich, differs in that the front femora lack the flexor bristling, and the darkening of the wings is more extended, the wings being somewhat infumated but with subhyaline spaces near the centers of the short cells and near the base and apex of the longer cells.

(2) **Clinocera trunca**, nov. sp. — Female. Length 3.5 mm. Black, coated above with greenish and on the sides and below with slate-colored pollen; legs black; wings brownish, veins strong. Occiput and vertex greenish, upper part of the face concave, dusky, lower part convex, white, separated from the very short dusky cheeks; eyes large; occipital bristles not numerous; palpi small, black, proboscis very short; antennae black, the third joint short, arista longer than the antennae. Mesonotum completely greenish pollinose, with faint indications of two dark vittae, no acrostichals, six uniform dorso-centrals, scutellum with two apical bristles; metapleurae with a few fine white hairs; metanotum slate blue. Legs entirely black, front femora with a few inconspicuous setulae beneath; claws, pulvilli and empodium uniformly small. Wings unspotted but lightly infumated, the centers of the cells a little paler, veins strong, nearly black, costal setulae very minute, two submarginal cells, the second somewhat broader at the end than the first, the anterior branch of the third vein imperfect at its origin in the type specimen, gently and evenly sinuous, first and second posterior cells nearly alike at their base, the last section of the fourth vein but slightly longer than the front edge of the discal cell, anterior crossvein placed at two-fifths the length of the discal cell, the externo-anterior edge of the second basal cell two-thirds the length of the first section of the front border of the discal, posterior crossvein elbowed at its posterior third, making nearly a right angle with the fifth vein, last two sections of the fifth vein equal, anal crossvein round, separately closing the anal cell, anal vein replaced by a fold; halteres black.

Type specimen collected by Professor J. M. Aldrich at Dewatto, Washington, June 7, 1906. Four paratypes from Olga and Friday Harbor, Washington, May and June, differ in having three submarginal cells. However, in two of these specimens there are three submarginals in one wing and two in the other. A poorly preserved specimen in the National Museum, from Ungava Bay, Hudson Strait, having three submarginal cells, differs in that the legs are brownish instead of fully black.

This species agrees fairly well with Walker's brief description of *longipes*, except for the decidedly brown and not gray color of the wings and the lack of gray pollinosity of the legs. Osten Sacken suggested that Walker's species is the same as Loew's *simplex*, but the latter has the mesonotum evittate.

4. Halteres reddish; veins feebly undulating; knees black 5.
 Halteres with black knob; veins not undulating;
 thorax bivittate; femora ♂ with setulæ 6.
5. Thorax bivittate with brown; stigma almost obsolete,
 anterior crossvein at basal fourth of the discal cell. HYDRODROMIA CONJUNCTA, Loew.
 Thorax with the median white vitta only; stigma
 distinct, anterior crossvein at the basal third of the
 discal cell. HYDRODROMIA TAOS, Melander.
6. Discal cell blunt; hind tibiæ setose 7.
 Discal cell with narrow end; hind tibiæ not setose;
 third submarginal cell four times as long as its
 costal breadth. HYDRODROMIA LONGIFURCA, nov. sp. (1).
7. Stigma well developed and distinct from the cloudings,
 second submarginal cell two and a half times as
 long as its breadth along the costa; knees black. PHÆOBALIA LECTA, Melander.
 No stigma, second submarginal cell four times as long
 as its breadth along the costa; knees reddish HYDRODROMIA BINOTATA, Loew.
8. Acrostichals biseriate, scutellum with discal setulæ;
 second posterior cell acutely pointed; face deeply
 excised at oral margin; femora and tibiæ without
 setulæ PHILOLUTRA SIMPLEX, Loew.
 Acrostichals wanting; face less deeply excised 9.
9. Legs entirely black; halteres black; veins not undu-
 lating 10.
 Legs brown or reddish, or at least the knees brownish;
 discal cell equal to or shorter than the second
 posterior 22.
10. Discal cell longer than the second posterior; thorax
 not vittate; stigma faint; front femora with a sub-
 apical group of setulæ, middle femora setulose,
 pulvilli vestigial II.
 Discal cell equal to or shorter than the second posterior 12.
11. Empodium of posterior legs short but visible; wings
 clear hyaline, anal vein stronger. BERGENSTAMMIA DOLICHERETMA, Melander.

(1) **Clinocera (Hydrodromia) longifurca**, nov. sp. — Male. Length 3.5 mm. Dark olivaceous brown above, slaty gray below. Front uniformly dark brown, face white pruinose, a narrow crescent-shaped mark beneath the antennæ extending to the orbits, vertical bristles moderately coarse; cheeks very small, separated from the face; third joint of the antennæ oval, arista rather slender, less than twice as long as the antennæ; palpal hairs inconspicuously black. Thorax with two dark vittæ; pectal hairs very short and sparse; all the hairs of the abdomen pale; ventral piece of the epipygium conical, middle valves small, dorsal valves long and narrow, shining black. Legs entirely black, hairs of the front coxæ short and sparse, front tibiæ nearly as long as the femora, front femora with seven fine short black setæ and six white hairs, all uniformly distributed, both sides of the hind tibiæ with short setæ along the outer half, empodium prominent and yellow. Halteres black. Wings rather strongly brown, veins firm and blackish, three submarginal cells, the second narrow and four times as long as its width along the costa, discal cell three times as long as broad, a little shorter than the sessile second posterior cell, its posterior oblique crossvein strongly curved so that the apex of the discal cell has parallel sides, sections of the fifth vein equal, anal crossvein recurved, anal vein short and rather faint.

Type from Mount Washington, New Hampshire, Mrs. Annie T. Slosson, collector. Two females from same locality were collected by C. W. Johnson, 4 July, 1914. A female in the collection of C. W. Johnson, from Nain, Labrador, has no setæ on the femora and the discal cell is a trifle less drawn out at the apex.

Empodium of posterior legs wanting; wings infumated, anal vein weak	BERGENSTAMMIA BRUNNIPENNIS, NOV. SP. (1).
12. Front femora with conspicuous spinous setulae	13.
Front femora with no or but few setulae	18.
13. Face entirely white; stigma more or less evident	14.
Face darker in the middle or yellow above, vertex without a dark picture; thorax at most weakly bivittate; no stigma (if stigma is well developed and wings are spotted see <i>lecta</i> ♂)	15.
14. Vertex pictured; thorax not vittate; front tibiae ♂ with a long comb of hairs; arista longer than the antennae	CHAMÆDIPSIA COMATA, NOV. SP. (2).
Vertex uniformly brown; thorax bivittate; arista as long as the antennae.	CLINOCERA PRASINATA, NOV. SP. (3).

(1) **Clinocera (Bergenstammia) brunnpennis**, nov. sp. — Male. Length 5 mm. Front, vertex and upper occiput dark brown, ocellar tubercle flanked by a weak cinereous spot, face entirely white, rather narrow, not carinate or tuberculate, the lower edge nearly transverse, distinctly separated from the cheeks which measure about one-eighth the eye-height; upper occiput with bristles and setulae, sides with silky yellow hairs, lower part bare; palpi with short black hairs, proboscis rather large; third joint of the antennæ oval, shorter than deep, without an end-process, the arista more than twice as long as the antenna, slender and tapering. Thorax not vittate, dull brown above, including the metanotum, bristles rather weak, five or six dorsocentrals, no setulae, only one notopleural; pectus nearly bare. Sides of the abdominal segments marked with cinereous, centrally brown, epipygium small, ventral piece and base of the penis thickened, black, middle valve small, shining apically, dorsal valve narrow, corneous, shining black, incurved apically, the inner surface rugose. Legs black, hairs of the front coxae short and pale, front femora of the male with a row of flexor setulae on the basal half and distally with an irregular group of thorn-like setæ, front tibiae nearly as long as the femora, denticulate within, posterior femora with loose irregular short but strong setæ, middle tibiae with scattered setulae, hind tibiae with a preapical pair of bristles, pulvilli wanting, empodium of the front legs one-half as long as the claws and brown, of the posterior legs microscopic, claws black. Wings short and narrow, with a brown tinge, stigma weak, brown, but distinct, basal bristle short, costal setulae small, first vein ending at the middle, two submarginal cells, the second small, discal cell blunt, long and narrow, longer than the sessile second posterior cell, sections of the fifth vein 2 : 1, anal cell with curved apex, anal vein very faint but nearly reaching the margin.

Five specimens; Palo Alto, California, August, received from Professor Doane.

(2) **Wiedemannia (Chamædipsia) comata**, nov. sp. — Male. Length 4 mm. Of the usual olivaceous brown and slaty color, the front tibiae ♂ bearing a row of long hairs on the apical half, front pictured. Face entirely white pruinose, carinate below, the oral margin not notched at the middle, cheeks one-fourth the eyeheight, notched, but separated from the face only by a suture; front and vertex velvety black, but the front marked with a white spot almost enclosing the ocellar tubercle and the vertex with a post-ocellar stripe which meets the gray of the occiput; upper bristles short but strong and with a few intermixed setulae, lower hairs long fine sparse and yellow; third antennal joint rounded oval, without a terminal process, the arista thick, even enlarged apically, one-half longer than the antenna; hairs of the palpi black and conspicuous. Two or three minute acrostichal setulae in front, six dorsocentrals, four scutellars and a few discal and marginal fine setulae, a group of five notopleurals, metapleural hairs numerous and yellowish-brown, pectal hairs sparse. Discal setulae of the abdomen sparse and black, ventral and lateral hairs yellow; epipygium large, erect, ventral piece almost cylindrical, the penis blackish, middle valves convex, oval, bearing black hairs, surmounted by the irregular five-pronged jet black upper valve. Hairs of the front coxae short and yellow, legs black, of the male the front femora provided with a dense brush of setæ at the basal third of the under side beyond which are a few spine-like setæ, the front tibiae with an extensor row of very long closely placed black hairs and within with numerous stiff setæ and with preapical denticles, posterior femora almost bristleless, hind tibiae with a few short stiff preapical setæ, in the female the front legs lack the excessive ornamentation, the inner side of the femora with loose but stiff setæ and the under side of the tibiae with minute closely placed setulae, empodium yellow and long, claws entirely black. Wings very long, slender, with light brownish tinge, basal bristle small, costal setulae minute, stigma very faint, first vein ending at the basal two-fifths of the wing, two submarginal cells, the second long and rather narrow, discal cell narrow, not pointed, three-fourths as long as the second posterior, sections of the fifth vein nearly equal, anal cell apically with a uniform curve, anal vein thin but strong, vanishing half way to the margin.

Four specimens of this interesting species were taken by Professor J. M. Aldrich on Mount Rainier, Washington, August 3, 1905, at an altitude of 5000 feet.

(3) **Clinocera prasinata**, nov. sp. — Male. Length 4 mm. Olivaceous green, changeable in color from cinereous along the flanks to brownish dorsally. Face silvery white, front brownish, more gray when viewed from behind, no

15. Center of the face with a black spot; wings slightly infumated, auxiliary cell dark; setæ of the front femora and the hind tibiæ less pronounced basally. 16.

Face unicolorous; wings hyaline; front femora with about nine uniform short setæ, bristles of the hind tibiæ on the apical three-fifths

CLINOCERA OLIVACEA, nov. sp. (1).

16. Palpi with numerous black hairs; under side of the front femora ♂ with many minute thorn-like setulæ on the basal half, hind femora with three or four subapical flexor bristles 17.

Palpi with short sparse pale hairs; front femora without thorn-like setulæ, hind femora with two subapical flexor bristles

CHAMÆDIPSIA MINOR, nov. sp. (2).

17. Thorn-like setulæ of the front femora not in a regular row, with some coarse setæ intermixed; facial tubercle distant from the oral margin, the center of the face carinate beneath the tubercle, the upper part of the face brownish ♂ or cinereous ♀. CHAMÆDIPSIA HAMIFERA, nov. sp. (3).

vertical bristles, the occipital series sparse; cheeks narrow, with complete suture, palpi large, clavate, with sparse short black hairs; third joint of the antennæ ovate, shorter than broad, without an end-process, the arista thick and scarcely longer than the antenna. Viewed from in front the mesonotum is bilineate, the middle stripe not cinereous, five dorsocentrals, no setulæ, metanotum and pleuræ cinereous, metapleural hairs dense and yellow, pectus and front coxæ with sparse yellow hairs. Abdomen greenish cinereous; epipygium large, the lower piece large, conical and projecting, penis black, slender from its base, its middle third hidden, the apical deflected part thin, middle lamellæ broadly oval, erect, rather hairy, the upper pieces broad but short, shining black. Legs black, more or less cinereous basally, front femora with regular biseriate short flexor bristles, front tibiæ as long as their femora, with close uniform short flexor and extensor setulæ, posterior femora without bristles, hind tibiæ with about ten extensor setulæ on the outer half, pulvilli and empodium moderately large. Calypteres blackish, with a pale fringe; halteres black. Wings with a slight infumation, veins strong and blackish, costal setulæ small, stigma very weak, two submarginal cells, third and fourth sections of the costa subequal, discal cell shorter than the sessile second posterior cell, sections of the fifth vein equal, anal vein a weak fold nearly reaching the margin.

One specimen, received from Dr. J. M. Aldrich, who collected it at Mono Lake, California, July 23, 1911.

(1) **Clinocera olivacea**, nov. sp. — Female. Length 3.5 mm. Olivaceous brown, abdomen, pleurae and base of the legs cinereous-white, face and cheeks cinereous, separated by a suture, vertex brown, with strong ocellar vertical and occipital bristles, third joint of the antennae triangularly oval, as long as broad, without a process, the arista thick and slightly longer than the antenna; palpi with short brownish hair. Five dorsocentral bristles, no setulae, metapleural hairs pale yellow. Prosternum and coxae with conspicuous golden hair, under side of the front femora with two series of about nine short setae disposed along its length, front tibiae nearly as long as their femora, outwardly with short uniform setae, posterior femora bristleless, hind tibiae with about eleven extensor setae on the apical three-fifths, pulvilli and empodium moderate. Wings hyaline, veins weak, no stigma, costal setulae small, two submarginal cells, the second to the fifth costal sections proportioned 4 : 1 : 1.4 : 1.2, discal cell rather broad, a little shorter than the sessile second posterior cell, sections of the fifth vein proportioned 1 : 1.2, anal vein reflexed more than usual, the anal vein rather strong continuing as a fold almost to the margin.

One specimen, Muir Inlet, Alaska, Trevor Kincaid, collector.

(2) **Wiedemannia (Chamaedipsia) minor**, nov. sp. — Male. Length 3 mm. Related to *hamifera*, but differing as follows: face brownish cinereous, the lower sides when viewed from in front browner than the rest, when viewed from below concolorous, the dark central stripe and the carina less evident, margin excised; palpi with fewer and pale hairs; apical process of the third antennal joint less pronounced. No humeral setulae, metapleural and pectal hairs less developed. Front femora and tibiae without the flexor denticles, the front femora with nine setae uniformly distributed, no long yellow hairs. Discal cell three-fourths as long as the second posterior, anal crossvein less strongly recurved, the lower end of the anal cell obtusely angulate.

One specimen, received from Professor O. A. Johannsen, who collected it at Ithaca, New York.

(3) **Wiedemannia (Chamaedipsa) hamifera**, nov. sp. — Male. Length 4 mm. Brownish above, the thorax not vittate, cinereous below. Upper half of the face light brown in the male and cinereous in the female, sides below cinereous white, but slightly longer than broad between the antennae and the center of the tubercle, center tubercle black.

- Thorn-like setulae ♂ forming a regular, uniform, conspicuous row, of ♀ seriate but microscopic in size; facial tubercle not conspicuous, located nearer the emarginate oral margin, the lower part of the face scarcely carinate, the upper part white CHAMÆDIPSIA CTENISTES, nov. sp. (1).
18. Thorax bivittate 19.
Thorax not vittate; wings hyaline, no stigma 21.
19. Wings maculate, stigma very distinct; face entirely white HYDRODROMIA LECTA, Melander.
Wings not spotted, no stigma; face brownish above 20.
20. Legs black; wings lightly infumated CLINOCERA TRUNCA, nov. sp
Legs hoary; wings slightly gray ? HYDRODROMIA LONGIPES, Walker.
21. Second posterior cell very acute at the proximal end;
face not tuberculate. CLINOCERA LEPIDA, Melander.
Second posterior cell sessile; center of the face black
and strongly tuberculate CHAMÆDIPSIA GUBERNANS, nov. sp. (2).

and epistome strongly carinate, its margin not incised, cheeks one-fourth the eye-height, a faint suture present, palpi cylindrical, with abundant rather long black hair, upper part of the head brown, the upper bristles strong and interspersed with setulae; third joint of the antennæ onion-shaped, the apical process long, arista one-half longer than the antennæ, thick at the base only. Six dorsocentral bristles, no acrostichals, no scutellar or lateral setulae except a couple on the humeri; prosternum and metapleuræ with numerous long yellow hairs. Abdomen with yellow lateral hairs and black discal setulae, epipygium small, the terminal keel short, lateral valves with long loose yellow hairs, upper valves shining black, with slender pointed converging end-processes. Front coxae with abundant yellow hair, legs black, front tibiae nearly as long as their femora, in both sexes the under side of the front femora densely and irregularly set with short thorn-like setulae, interspersed with stronger setæ especially toward the apex, and basally with a few long yellow hairs also, inner side of the front tibiae uniformly and densely provided with minute setulae, posterior femora with a few strong flexor setæ, and hind tibiae with a preapical bristle on each side, pulvilli small but the empodium large and white. Halteres with black knob and reddish root; cilia of the calypteres very sparse and yellow. Wings long and narrow, with a brownish tinge, costal setulae moderate, no stigma, two submarginal cells, the second very narrow, discal cell blunt, two-thirds as long as the sessile second posterior cell, sections of the fifth vein 1 : 0.6, anal crossvein strongly recurved, anal vein faint, vanishing half way to the margin.

Thirty specimens; Beaverkill and Ithaca, New-York; Middletown, Connecticut; Moosehead Lake, Maine: from E. T. Cresson, Jr., C. W. Johnson, and O. A. Johannsen. Mr. Johnson states that the species swarmed over cut logs in a log-run on Moosehead Lake, so as to blacken the surface by their numbers.

(1) **Wiedemannia (Chamædipsia) ctenistes**, nov. sp. — Male. Length 3 mm. Closely related to *hamifera* but differs as follows: face narrower, from the antennæ to the center of the tubercle the face of the male is obviously longer than wide, the tubercle much less pronounced and located near the strongly notched oral margin, carina of the epistome not conspicuous, face of both sexes when viewed from in front uniformly white pruinose, except the tubercle, when viewed from below with a narrow dark median vitta. Palpi with finer hairs. Thorax with a trace of two vittæ. End process of the dorsal valves of the epipygium nearly straight. The flexor setulae of the front femora of the male are stronger and uniformly arranged in a conspicuous single straight row along the basal three-fifths, beyond which occur four or five strong thorn-like setæ; in the female the setulae are greatly reduced but still are seriatel arranged, without interspersed strong setæ; hind femora with about three preapical bristles.

Five specimens, collected by C. W. Johnson at Hanover, New Hampshire. Paratypes placed in the Boston Society of Natural History and in Mr. Johnson's collection.

(2) **Wiedemannia (Chamædipsia) gubernans**, nov. sp. — Male. Length 3.5 mm. Olivaceous brown above, slaty below. Face dark cinereous, strongly tuberculate, lower margin carinate and entire, a fairly deep incision separating the cheeks, which measure about one-seventh the eye-height; palpi large, villose and with a few short dark hairs, sides of the proboscis likewise villose; third joint of the antennæ spongy pubescent, short, onion-shaped, with a lengthened terminal process, the arista thick and blunt and slightly longer than the antenna; bristles of the upper half of the head strong, four pairs of postocellar setulae. Setulae present in the dorsocentral rows, no scutellar setulae, some posthumeral setulae present, metapleural hairs yellowish, pectus bare but the propleurae with four or five setulae. Epipygium with a long drawn-out keel, penis slender and brown, middle valves ovate, upper valves rather large, convex, opaque. Coxæ with short sparse blackish hairs, legs black, front tibiae seven-eighths as long as their femora, femora without setæ, hind tibiae scarcely setulose.

22. Thorax with two conspicuous shining vittæ; veins undulating, wings brown; legs without bristles; face broad and pure white CLINOCERA LINEATA, Loew.
- Thorax pollinose, no shining vittæ. 23.
23. Front femora with a basal group of setulæ, hind tibiæ with flexor setulæ; wings more or less spotted 24.
24. Thorax with a median cinereous line; upper part the face dark HYDRODROMIA MACULATA, Loew.
- Thorax bivittate; face entirely silvery. HYDRODROMIA STAGNALIS, Haliday.
25. Thorax not vittate; face dark; wings uniformly subfuscous, costal setulæ barely visible; three small flexor setulæ on the outer third of the hind tibiæ; arista thickened CLINOCERA FUSCIPENNIS, Loew.
- Thorax bivittate; face entirely white 26.
26. Wings with a spot at the base of the second submarginal cell and at the apex of the discal cell, the second submarginal cell as long as the second basal; legs black, the knees brown HYDRODROMIA GENUALIS, Coquillett.
- Wings uniformly brown; second submarginal cell longer than the second basal; legs brown. PHÆOBALIA BREVITIBIA, nov. sp. (1).

Geographical distribution.

SUBGENUS BERGENSTAMMIA

1. *B. brunniipennis*, nov. sp. California.
2. *B. dolicheretma*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 241 [1902] Idaho.
(*Clinocera*); Kertész, Cat. Dipt. Vol. 6, p. 125 [1909] (*Röderia*);
Engel, Deutsche Ent. Zeitschr. 1918, p. 198 [1918] (*Roederella*).

pulvilli and empodium moderate in size and brown. Halteres black, hairs of the calypteres sparse and brownish. Wings with a brownish tinge, a very faint darker cloud suffusing the crossveins, veins strongly brown, no costal setulæ, first vein ending beyond the middle of the wing, two submarginal cells, second to the seventh sections of the costa proportioned, $4:1:1.2:1:1.8:1.9$, discal cell much shorter than the broadly sessile second posterior cell, sections of the fifth vein, $1:0.6$, anal vein represented only by a very faint fold, anal crossvein strongly recurved.

One specimen, Nelson, British Columbia, July 17, 1910.

(1) ***Clinocera (Phæobalia) brevitibia*, nov. sp.** — Length 2.5 mm. Front, vertex and upper occiput a dark slaty brown, ocellar bristles long, a single row along the occipital and vertical orbits, one pair of central vertical bristles, setulæ sparse, lower occiput nearly bare; face cinereous, separated from the narrow cheeks, the lower margin transverse and not emarginate; hairs of the palpi minute, sparse, brown; third joint of the antennæ onion-shaped, with a distinct apical projection, arista thick, very slightly tapering, nearly twice as long as the antenna. Dorsum dark brown, marked with two velvety black vittæ which are abruptly interrupted at the prescutellar depression, intervittal space a little more grayish anteriorly, a white spot in front of the wings, metanotum cinereous; five strong dorsocentral bristles, no setulæ, pectus with a few hairs, upper pleurae brownish. Abdomen entirely blackish gray, its hairs very sparse, epipygium small, incumbent, ventral piece conical, penis brown, middle valve oval, subshining, dorsal valve clavate, villose, only subshining. Legs including the coxae brownish, hairs of the front coxae sparse, in the male the front femora with about six spinous flexor setæ, the front tibiæ about five-sixths as long as the femur and slightly shorter than the basal two joints of their tarsi, closely and minutely setulose within, apical half of the hind tibiæ setose on both sides, hind metatarsi with eight flexor setulæ; in the female the front legs are devoid of the flexor bristling and the setæ of the hind legs are much reduced. Wings oval, with a decided brown tinge, stigma elongate, weak, basal bristle long, costal setulæ minute, first vein ending at the middle of the wing, two submarginal cells present, discal cell blunt, as long as the broadly sessile second posterior cell, sections of the fifth vein nearly $3:1$, anal crossvein recurved, no anal vein.

Five males, one female; Olga and Tacoma, Washington, July and August.

3. *B. multiseta*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 104 C. Europe.
 (1892); Strobl, ibidem, Vol. 34, p. 209 (1898); Engel, Deutsche Ent. Zeitschr. 1918, p. 78, f. 25 (1918).
4. *B. nudipes*, Loew, Wien. Ent. Monatschr. Vol. 2, p. 386 [1858] (*Clinocera*); C. & S. Europe.
 Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 (1881); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 103 (1898); Engel, Deutsche Ent. Zeitschr. 1918, p. 76, f. 23, 24 (1918).

SUBGENUS CLINOCERA

1. *C. appendiculata*, Zetterstedt, Fauna Ins. Lappon. p. 559 [1838] (*Wiedemannia*); Dipt. Scand. Vol. 1, p. 367 [1842] (*Wiedemannia*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 247, note (1858); Zetterstedt, Dipt. Scand. Vol. 13, p. 5005 [1859] (*Wiedemannia*); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 153 [1864] (*Wiedemannia*); Mik, Verh. Zool-bot. Ges. Wien, Vol. 31, p. 325 (1881); Pokorny, ibidem, Vol. 37, p. 394 (1887); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, 1892, p. 100 (1893); Bezzu, Bull. Soc. Ent. Ital. Vol. 30, p. 148, 279 (1899); Wahlgren, Ent. Tidskr. Vol. 31, p. 82 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 61, f. 10, 11 (1913); Engel, Deutsche Ent. Zeitschr. 1918, p. 25, f. 7 [1918] (*Atalanta*).
 var. *aucta*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3019 [1849] (*Brachystoma*); Loew, N. Europe
 Wien. Ent. Monatschr. Vol. 2, p. 247, note 4 (1858); Wahlgren, Ent. Tidskr. Vol. 31, p. 28, f. 1, p. 82, f. 13 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 61, f. 12 (1913); Engel, Deutsche Ent. Zeitschr. 1918, p. 29 [1918] (*Atalanta*).
 var. *simplicinervis*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 62, f. 13 Lapland.
 (1913).
 var. *Storckii*, Mik, Verh. Zool-bot. Ges. Wien, Vol. 30, p. 348 (1880); Mik, ibidem, Vol. 31, p. 325 (1881); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 100 (1893); Engel, Deutsche Ent. Zeitschr. 1918, p. 28 [1918] (*Atalanta*).
2. *C. aquatica*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 122 [1914] (*Atalanta*); Oldenberg, Deutsche Ent. Zeitschr. 1919, p. 392 [1919] (Atalanta). E. Africa.
3. *C. bivittata*, Loew, Wien. Ent. Monatschr. Vol. 8, p. 259 (1865); Mik, Siberia, Europe.
 Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 325 (1881); Engel, Deutsche Ent. Zeitschr. 1918, p. 31 [1918] (*Atalanta*).
4. *C. fluviatilis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 34 (1913); Engel, W. Himalayas.
 Deutsche Ent. Zeitschr. 1918, p. 32 (1918); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 365, f. 33 (1920).
5. *C. fuscipennis*, Loew, Zeitschr. Ges. Naturw. Vol. 48, p. 324 (1876); E. United States.
 Melander, Trans. Am. Ent. Soc. Vol. 28, p. 245 (1902); Engel, Deutsche Ent. Zeitschr. 1918, p. 33 (1918).
6. *C. lineata*, Loew, Berl. Ent. Zeitschr. 1862, p. 207; Cent. 2, No. 50 (1862); N. America.
 Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 244, f. 61, 62 (1902); Engel, Deutsche Ent. Zeitschr. 1918, p. 34 (1918).
7. *C. nigra*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 292, 1, Europe.
 pl. 14, f. 20-25 (1804); Syst. Beschr. Vol. 2, p. 113, 1, pl. 16, f. 4 (1820); Macquart, Hist. Nat. Dipt. Vol. 1, p. 433, 1, pl. 10, f. 17 (1834); Blanchard, Hist. Nat. Ins. Vol. 3, p. 604 (1840); Loew, Wien. Ent. Monatschr. Vol. 2, p. 254, note 3 (1858); Schiner, Fauna Dipt. Austr. Vol. 1, p. 85 (1862); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 325, pl. 16, f. 4, 5 (1881); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 101 (1893); Lundbeck, Dipt. Dan.

- Vol. 3, p. 217, f. 89 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 21, f. 5 [1918] (*Atalanta*).
Robertii, Macquart, Hist. Nat. Dipt. Vol. 2, p. 657 [1835] (*Paramesia*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 254 (1858); Meigen, Syst. Beschr. Vol. 7, p. 79 [1858] (*Brachystoma*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 84 (1862).
unicolor, Curtis, Brit. Ent. p. 513 [1834] (*Heliodromia*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 504 [1849] (*Heleodromia*); Ins. Brit. Vol. 1, p. 106 [1851] (*Heleodromia*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 245, note (1858); Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 [1862] (*Hemerodromia*).
var. *rufipes*, Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 147 (1899); Engel, Deutsche Ent. Zeitschr. 1918, p. 24 [1918] (*Atalanta*). S. Europe.
8. *C. ?obscura*, Brunetti, Rec. Indian Mus. Vol. 9, p. 34 (1913); Engel, Deutsche Ent. Zeitschr. 1918, p. 36 (1918); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 366 (1920). W. Himalayas.
9. *C. olivacea*, nov. sp. Alaska.
10. *C. prasinata*, nov. sp. California.
11. *C. riparia*, Robert, Ann. Soc. Ent. France (1), Vol. 5, p. 537 [1836] (*Paramesia*). S. W. Europe.
12. *C. trunca*, nov. sp. Washington.

SUBGENUS HYDRODROMIA

- . *H. bicincta*, Tucker, Kansas Univ. Sc. Bull. Vol. 14, p. 97 [1907] (*Clinocera*, no description).
1. *H. binotata*, Loew, Zeitschr. Ges. Naturw. Vol. 48, p. 325 [1876] (*Clinocera*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 243 [1902] (*Clinocera*); Engel, Deutsche Ent. Zeitschr. 1918, p. 46 (1918). North America.
2. *H. conjuncta*, Loew, Wien. Ent. Monatschr. Vol. 4, p. 80 [1860] (*Clinocera*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 242, 346 [1902] (*Clinocera*); Engel, Deutsche Ent. Zeitschr. 1918, p. 62 [1918] (? *Phaeobalia*). E. North America.
3. *H. dilatata*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 387, f. 13 [1909] (*Heleodromia*); Engel, Deutsche Ent. Zeitschr. 1918, p. 46 (1918). Bolivia.
4. *H. fontinalis*, Haliday, Ent. Mag. London, Vol. 1, p. 160 [1833] (*Heleodromia*); Curtis, Brit. Ent. p. 513 [1834] (*Heleodromia*); Walker, Ins. Brit. Vol. 1, p. 105 [1851] (*Heleodromia*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 256 [1858] (*Clinocera*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 84 [1862] (*Clinocera*); Loew, Beschr. Eur. Dipt. Vol. 1, p. 271, note [1869] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 [1881] (*Heleodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 83 [1910] (*Heleodromia*); Engel, Deutsche Ent. Zeitschr. 1918, p. 41 (1918). Europe.
- Wesmælii, Zetterstedt, Dipt. Scand. Vol. 1, p. 364, part [1842] (*Brachystoma*).
5. *H. genualis*, Coquillett, Proc. Ent. Soc. Wash. Vol. 12, p. 124 [1910] (*Clinocera*); Engel, Deutsche Ent. Zeitschr. 1918, p. 37 [1918]. Alberta.
6. *H. longifurca*, nov. sp. New Hampshire,
Canada. [Labrador.]
7. *H. ?longipes*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 504 [1849] (*Heleodromia*); Osten-Sacken, Cat. Dipt. N. Amer. p. 106 [1878] (*Clinocera*).
8. *H. maculata*, Loew, Wien. Ent. Monatschr. Vol. 4, p. 79 [1860] (*Clinocera*); Glover, Manusc. Notes, p. 15, pl. 6, f. 8 [1874] (*Clinocera*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 244 [1902] (*Clinocera*); Engel, Deutsche Ent. Zeitschr. 1918, p. 48 (1918). United States.

9. *H. nivalis*, Zetterstedt, Fauna Ins. Lappon. p. 545 [1838] (*Hemerodromia*); Europe. Dipt. Scand. Vol. 1, p. 325 [1842] (*Ardoptera*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 8 [1858] (*Clinocera*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 155 [1861] (*Ardoptera*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 85 [1862] (*Clinocera*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 62, f. 15 (1913); Engel, Deutsche Ent. Zeitschr. 1918, p. 44 (1918).
10. *H. stagnalis*, Haliday, Ent. Mag. London, Vol. 1, p. 159 [1833] (*Heleodromia*); Curtis, Brit. Ent. p. 513 [1834] (*Heleodromia*); Macquart, Hist. Nat. Dipt. Vol. 2, p. 658 (1835); Meigen, Syst. Beschr. Vol. 7, p. 93 [1838] (*Hemerodromia*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 504 [1849] (*Heleodromia*); Ins. Brit. Vol. 1, p. 105, pl. 3, f. 6 [1851] (*Heleodromia*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 246, 4 [1858] (*Clinocera*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 84 [1862] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326, pl. 16, f. 6-7, [1881] (*Heleodromia*); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 101 [1892] (*Heleodromia*); Lundbeck, Vidensk. Meddel. p. 297 [1898] (*Clinocera*); Strobl, Wien. Ent. Zeit. Vol. 18, p. 77 [1899] (*Heleodromia*); Coquillett, Proc. Ent. Soc. Washington, Vol. 5, p. 264 (1903); Lundbeck, Dipt. Dan. Vol. 3, p. 219, f. 90 [1910] (*Heleodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 82 [1910] (*Heleodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 63 [1913] (*Heleodromia*); Engel, Deutsche Ent. Zeitschr. 1918, p. 39, f. 9 (1918).
appendiculata, Hansen (not Zetterstedt), Naturh. Tidskr. (3), Vol. 13, p. 259 [1880] (*Heleodromia*).
Westermannii, Zetterstedt, Fauna Ins. Lappon. Vol. 558 [1838] (*Brachystoma*); Dipt. Scand. Vol. 1, p. 360 [1842] (*Brachystoma*); ibidem, Vol. 8, p. 309 [1849] (*Brachystoma*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 162 [1861] (*Brachystoma*).
11. *H. taos*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 242 [1902] (*Clinocera*); Engel, Deutsche Ent. Zeitschr. 1918, p. 49 (1918). New Hampshire.
12. *H. undulata*, nov. sp. Idaho, Washington.
13. *H. Wesmælii*, Macquart, Hist. Nat. Dipt. Vol. 2, p. 656, pl. 24, f. 17 [1835] (*Paramesmia*); Meigen, Syst. Beschr. Vol. 7, p. 79 [1838] (*Brachystoma*); Zetterstedt, Dipt. Scand. Vol. 1, p. 364, part [1842] (*Brachystoma*); Loew, Zeitschr. Ges. Naturw. Vol. 8, p. 100 [1857] (*Clinocera*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 257 [1858] (*Clinocera*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 85 [1862] (*Clinocera*); Nowicki, Verh. Nat. Ver. Brünn, Vol. 6, pl. 2, f. 5 [1868] (*Clinocera*); Loew, Beschr. Dipt. Vol. 1, p. 271, note [1869] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 362 (1881); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 101 (1893); Lundbeck, Dipt. Dan. Vol. 3, p. 220, f. 91 [1910] (*Heleodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 82 [1910] (*Heleodromia*); Verrall, Ent. Mag. London, Vol. 48, p. 26 [1912] (*Clinocera*); Engel, Deutsche Ent. Zeitschr. 1918 p. 43 [1918] (*Heleodromia*).
nebulosa, Zetterstedt, Fauna Ins. Lappon. p. 559, note [1838] (*Brachystoma*).

SUBGENUS KOWARZIA

1. *K. amarantha*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 43, pl. 1, f. 10 (1908); Canary Islands. Engel, Deutsche Ent. Zeitschr. 1918, p. 73 (1918).
2. *K. barbatula*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 347 [1880] Europe. (*Clinocera*); Mik, ibidem, Vol. 31, p. 325, pl. 16, f. 1-3 (1881); Brocher,

- Bull. Soc. Zool. Genève, Vol. 1, p. 115-117 (1908); Strobl, Verh. Zool.-bot. Wien, Vol. 59, p. 178 (1909); Bezzi, Arch. Zool. Expér. Paris (5), Vol. 8, p. 50 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 68, f. 20 (1918).
 var. *securigera*, Engel, Deutsche Ent. Zeitschr. 1918, p. 70 (1918). C. Europe.
 Europe.
3. *K. bipunctata*, Haliday, Ent. Mag. London, Vol. 1, p. 159 [1833] (*Heleodromia*); Curtis, Brit. Ent. Vol. 8, p. 513 [1834] (*Heliodromia*); Macquart, Hist. Nat. Dipt. Vol. 2, p. 658 [1835] (*Hydrodromia*); Meigen, Syst. Beschr. Vol. 8, p. 93 [1838] (*Hemerodromia*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 251 [1858] (*Clinocera*); Mik, Verh. Zool-bot. Ges. Wien, Vol. 31, p. 325 (1881); Wahlgren, Ent. Tidskr. Vol. 31, p. 83 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 65, f. 16 (1918).
tenella, Wahlberg, Ofv. Vet. Akad. Förh. Stockholm, Vol. 1, p. 107 [1844] (*Paramesia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3020 [1849] (*Brachystoma*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 252, note 1 [1858] (*Clinocera*); Wahlgren, Ent. Tidskr. Vol. 31, p. 28 (1910).
Zetterstedti, Walker (not Fallen), Ins. Brit. Vol. 1, p. 105, 252 [1851] (*Heleodromia*). C. Europe.
4. *K. haemorrhoidalis*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 189, pl. 4, f. 60 (1908); Engel, Deutsche Ent. Zeitschr. 1918, p. 74 (1918). Madeira.
5. *K. plectrum*, Mik, Verh. Zool-bot. Ges. Wien, Vol. 30, p. 347 [1880] (*Clinocera*); Mik, ibidem, Vol. 31, p. 325 (1881); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 101 (1893); Engel, Deutsche Ent. Zeitschr. 1918, p. 71, f. 22 (1918). C. Europe.
6. *K. Schnabli*, Becker, Deutsche Ent. Zeitschr. p. 646 (1910); Engel, ibidem, 1918, p. 74 (1918). Corsica.
7. *K. tibiella*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 348 [1880] (*Clinocera*); ibidem, Vol. 31, p. 325 (1881); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 101 (1892); Engel, Deutsche Ent. Zeitschr. 1918, p. 67, f. 17, 18, 19 (1918). C. & S. Europe.

SUBGENUS PHÆOBALIA

1. *P. brevitibia*, nov. sp. Washington.
2. *P. dimidiata*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 271 [1869] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 (1881); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 103 (1893); Engel, Deutsche Ent. Zeitschr. 1918, p. 51, f. 10 (1918); Oldenberg, ibidem, 1919, p. 391 (1919). C. & S. Europe.
- picta*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 102 (1893); Oldenberg, Arch. Naturg. Berlin, Vol. 80 (9), p. 92 (1915).
3. *P. inermis*, Loew, Wien. Ent. Monatschr. Vol. 5, p. 349 [1861] (*Clinocera*); Loew, Beschr. Eur. Dipt. Vol. 1, p. 271, note [1869] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 19, pl. 4, f. 15 [1869] (*Clinocera*); ibidem, Vol. 31, p. 326 (1881); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 103 (1893); Engel, Deutsche Ent. Zeitschr. 1918, p. 53 (1918). C. & S. Europe.
4. *P. lecta*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 243 (1902); Engel, Deutsche Ent. Zeitschr. 1918, p. 63 (1918). Idaho.
5. *P. penicissa*, Becker, Berl. Ent. Zeitschr. Vol. 33, p. 335, f. 1, 2 [1889] (*penicissa*); Bezzii, Arch. Zool. Expér. Paris (5), Vol. 8, p. 50 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 60, f. 15 (1918). Caves in Dalmatia.
6. *P. Pokornyi*, Mik, Wien. Ent. Zeit. Vol. 5, p. 22 (1886); Engel, Deutsche Ent. Zeitschr. 1918, p. 54, f. 11 (1918). Alps.

7. *P. trinotata*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 19, p. 24, pl. 4, f. 12, Alps. .
 14 [1869] (*Clinocera*); ibidem, Vol. 31, p. 326 (1881); Strobl, Mitteil.
 Naturw. Ver. Steiermark, Graz, Vol. 29, p. 101 (1893); Engel,
 Deutsche Ent. Zeitschr. 1918, p. 58, f. 13, 14 (1918).
8. *P. varipennis*, Nowicki, Verh. Nat. Ver. Brünn, Vol. 6, p. 85, pl. 2, f. 4 C. Europe.
 [1868] (*Clinocera*); Loew, Beschr. Eur. Dipt. Vol. 1, p. 269 [1869]
 (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 19, p. 326, pl. 4,
 f. 13 [1869] (*Clinocera*); Mik, ibidem, Vol. 31, p. 326 (1881); Strobl,
 Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 101 (1893); Engel,
 Deutsche Ent. Zeitschr. 1918, p. 56, f. 12 (1918).

13. GENUS WIEDEMANNIA, ZETTERSTEDT

Wiedemannia, Zetterstedt, Fauna Ins. Lappon. p. 559 (1838); Dipt. Scand. Vol. 1, p. 365 (1842); Walker, Ins. Brit. Dipt. Vol. 1, p. 106 (1851); Rondani, Dipt. Ital. Vol. 1, p. 150 (1856); Bigot, Bull. Soc. Ent. France, Vol. 12, p. 103 (1887); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258, 262 (1903); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 202 (1904); ibidem, Vol. 3, p. 363, note (1905); Lundbeck, Dipt. Dan. Vol. 3, p. 217 (1910); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 68 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 249 (1918).

Camelopsis, Engel, Deutsche Ent. Zeitschr. 1918, p. 14, 232 (1918).

Chamaedipsia, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 (1881); Bigot, Bull. Soc. Ent. France, Vol. 12, p. 103 (1887); Mik, Wien. Ent. Zeit. Vol. 7, p. 72, note (1889); Mik, ibidem, Vol. 11, p. 55 (1892); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 247, 262 (1903); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 67 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 207 (1918).

Clinocerella, Engel, ibidem, p. 14, 238 (1918).

Eucellidia, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 (1882); Bigot, Bull. Ent. Soc. France, Vol. 12, p. 103 (1887); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 250, 262 (1903); Lundbeck, Dipt. Dan. Vol. 3, p. 217 (1910); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 68 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 199 (1918).

Orthorhynchium, Mik, in litt. Engel, ibidem, p. 238 (1918).

Philolutra, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327 (1881); Bigot, Bull. Soc. Ent. France, Vol. 12, p. 103 (1887); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255 (1903); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 68 (1911); Engel, Deutsche Ent. Zeitschr. 1918, p. 222 (1918).

Pseudowiedemannia, Engel, ibidem, p. 14, 243 (1918).

Röderella, Engel, ibidem, p. 13, 79 (1918).

Röderia, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 326 (1881); Bigot, Bull. Soc. Ent. France, Vol. 12, p. 103 (1887); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257, 262 [1903] (*Röderia*); Kertész, Rovert. Lapok, Budapest, Vol. 18, p. 68 [1911] (*Röderia*).

Characters. — Lower part of the head more or less produced beneath the eyes, cheeks not deeply divided by a suture, so that the face is never wholly constricted from the lower occiput; lower edge of the face deeply and broadly excised, the notch filled by a depressed membrane which may be more or less carinate; pulvilli and empodium developed; acrostichals usually present; discal cell usually quite sharp apically, the stigma more or less developed.

The following subgenera are closely related and are not sharply differentiated. *Wiedemannia* of Zetterstedt is not the same genus as *Wiedemannia* of Meigen, 1838 : the latter is *Minho* Desvoidy.

Subgenus *Wiedemannia* in the strict sense. Stigma large and circular, commencing much beyond the end of the first vein and projecting into the wing so that it touches or passes the second vein, this vein being bent down around the stigma, five or six dorsocentrals alternating with rather long bristles, the anterior acrostichals reclinate unlike those of the prescutellar area, which are proclinate, scutellum with discal setulæ; head prolonged below eyes about the eye-height; terminal lamellæ of epipygium usually tipped with tuft of hairs.

Type species : *W. bistigma*, Curtis. Zetterstedt described two species, *borealis* and *appendiculata*. The former is synonymous with *bistigma* and becomes the type since the latter belongs to the earlier genus *Clinocera*.

SUBGENUS PSEUDOWIEDEMANNIA, ENGEL

Stigma more or less rounded, located much beyond end of first vein, smaller than in *Wiedemannia* and not passing below second vein; at most five dorsocentrals with very weak alternating setulæ; acrostichals biseriate onto the scutellum but either weak or strong; cheeks about half the eye-height.

Type species : *P. lamellata*, Loew, by Engel's designation.

SUBGENUS EUCELIDIA, MIK

Stigma elongate, beginning at the tip of the first vein and not touching the straightened second vein, costal spines strongly projecting; all the femora bearing preapical bristles on the front and hind sides; acrostichals if present all proclinate; scutellum with setulæ in addition to the apical pair of bristles; alternate dorsocentrals small.

Type species : *E. Zetterstedi*, Fallen, by Coquillett's designation, 1903. Mik founded his genus *Roederia* upon *C. longipennis*, Mik. Engel finds that this species in the same as *Eucelidia Escheri*, Zetterstedt. Therefore *Roederia* falls a synonym of *Eucelidia*.

SUBGENUS CLINOCERELLA, ENGEL

Stigma elongate as in *Eucelidia*; legs without distinctive bristles; five or six dorsocentrals with or without alternating setæ; acrostichals more or less developed, scutellum with or without setulæ; head projecting downward half the eye-height and tapering to the proboscis.

Type species : *C. sorex*, Engel, by Engel's designation. Mik used an unpublished name, *Orthorhynchium*, for this group.

SUBGENUS CAMELOPIS, ENGEL

Stigma elongate as in *Eucelidia*; five dorsocentrals with intermediate setæ, acrostichals reaching over scutellum, the anterior ones reclinate, the posterior ones proclinate, rarely the intermediate dorsocentrals weak or wanting; scutellum setulose; humeral bristle long and weak; head strongly projecting downward below the eyes nearly or quite the eye height, in profile rectangular.

Type species : *C. phantasma*, Mik, by Engel's designation.

SUBGENUS PHIOLUTRA, MIK

Stigma elongate, as in the preceding, costal spines smaller; preapical femoral bristles not strong; anterior acrostichals reclinate, the posterior ones proclinate, scutellum with setulæ, five dorsocentrals often with intermixed setulæ, head sometimes greatly drawn out beneath the eyes.

Type species : *P. phantasma*, Mik, Coquillett's designation in 1903. Coquillett made this purely a synonym of *Roederia*, since its main characteristic, the presence of the acrostichal setulae is inconstant. I retain the name here, like the two following groups, as a matter of convenience. The species are not phylogenetically distinct, and since the characters chosen are artificial combinations, species occur which do not conform in all respects to the definitions given.

SUBGENUS CHAMÆDIPSIA, MIK

Stigma elongate, as in *Eucelidia*, costal spines not projecting; preapical bristles of the femora with setæ; rows of acrostichals incomplete, at most some reclinate setulae present on the anterior part of the notum, scutellum typically without setulae, alternate smaller setulae usually present in the dorsocentral rows; arista usually blunt at the end.

Type species : *C. hastata*, Mik, the only species originally mentioned.

SUBGENUS RŒDERELLA, ENGEL

Stigma elongate, oval, costal spines rather prominent; preapical bristles of the femora absent, front femora with a comb of three or four bristles near tip; only short anterior acrostichals present, scutellum with marginal setulae, five dorsocentral bristles present and no intermediate ones.

Type species : *R. Csernyi*, Bezzi, by original designation.

Geographical distribution.

SUBGENUS CAMELOPIS

- | | |
|--|--|
| 1. <i>C. erminea</i> , Mik, Wien. Ent. Zeit. Vol. 6, p. 161 [1887] (<i>Philolutra</i>).
2. <i>C. phantasma</i> , Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 350 [1880]
(<i>Clinocera</i>); ibidem, Vol. 31, p. 327, pl. 16, f. 12, 13 [1881] (<i>Philolutra</i>); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 107 [1893] (<i>Philolutra</i>); Engel, Deutsche Ent. Zeitschr. 1918, p. 233, f. 54a, 54b, 55 (1918).
3. <i>C. quercifolia</i> , Engel, Deutsche Ent. Zeitschr. 1918, p. 235, f. 56 (1918). | C. Europe.
C. Europe.
C. Europe. |
|--|--|

SUBGENUS CHAMÆDIPSIA

- | | |
|---|--|
| 1. <i>Ch. alpina</i> , Engel, Deutsche Ent. Zeitschr. 1918, p. 220, f. 44-46 (1918).
2. <i>Ch. Beckeri</i> , Mik, Wien. Ent. Zeit. Vol. 8, p. 71 (1889); Becker, ibidem, Vol. 8, p. 83, pl. 1, f. 7, 8 (1889); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 107 (1893); ibidem, Vol. 34, p. 209 (1898); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 149 (1899); Engel, Deutsche Ent. Zeitschr. 1918, p. 215, f. 36, 37 (1918).
<i>jngorum</i> , Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 105 (1893); Pokorny, Verh. Zool.-bot. Ges. Wien, Vol. 43, p. 2 [1893] (<i>Phaeobalia</i>); Mik, Wien. Ent. Zeit. Vol. 13, p. 49 (1894); Strobl, Mitteil. Naturw. Nat. Ver. Steiermark, Graz, Vol. 34, p. 209 (1898)
var. <i>crinata</i> , Engel, Deutsche Ent. Zeitschr. 1918, p. 217 (1918).
3. <i>Ch. bicuspidata</i> , Engel, ibidem, 1918, p. 217, f. 38, 39 (1918).
var. <i>ornata</i> , Engel, ibidem, 1918, p. 219, f. 40-43 (1918).
4. <i>Ch. ctenistes</i> , nov. sp.
5. <i>Ch. comata</i> , nov. sp.
6. <i>Ch. gubernans</i> , nov. sp.
7. <i>Ch. hamifera</i> , nov. sp. | Tirol.
C. & S. Europe.
Hungary.
Hungary.
New Hampshire.
Washington.
British Columbia.
E. United States. |
|---|--|

8. *Ch. hastata*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 349 [1880] Austria.
(*Clinocera*); ibidem, Vol. 31, p. 326 (1881); Engel, Deutsche Ent.
Zeitschr. 1918, p. 214, f. 35 (1918).
9. *Ch. lepida*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 241 (1902); Idaho.
Engel, Deutsche Ent. Zeitschr. 1918, p. 33 [1918] (*Clinocera*).
10. *Ch. longicornis*, Mik, Wien. Ent. Zeit. Vol. 6, p. 162 (1887); Strobl, C. Europe.
Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 107 (1893); Engel,
Deutsche Ent. Zeitschr. 1918, p. 211, f. 33 (1918).
11. *Ch. lota*, Haliday, Ins. Brit. Vol. 1, p. 107, pl. 3, f. 7 [1851] C. Europe, Asia Minor.
(*lotæ*); ibidem, Vol. 3, p. 11 [1856] (*Wiedemannia*); Loew, Wien.
Ent. Monatschr. Vol. 2, p. 244 [1858] (*Clinocera*); Mik, Verh. Zool.-
bot. Ges. Wien, Vol. 31, p. 327, pl. 16, f. 8, 10 [1881] (*Philolutra*);
Wien. Ent. Zeitschr. Vol. 8, p. 72, note 2 (1888); Engel, Deutsche
Ent. Zeitschr. 1918, p. 209, f. 32 (1918).
pusilla, Loew, Wien. Ent. Monatschr. Vol. 2, p. 243 [1858] (*Clinocera*).
12. *Ch. Mikiana*, Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 149 (1899); Strobl, C. & S. Europe.
Tief Nachl. p. 33 (1900); Engel, Deutsche Ent. Zeitschr. 1918,
p. 213, f. 34 (1918).
13. *Ch. minor*, nov. sp. New York.

SUBGENUS CLINOCERELLA

1. *C. Oldenbergi*, Engel, Deutsche Ent. Zeitschr. 1918, p. 241, f. 60 (1918). C. Europe.
Tiepii, Mik, in litt. Engel, ibidem, 1918, p. 241 [1918] (*Orthorhynchium*).
2. *C. sorex*, Engel, ibidem, 1918, p. 239, f. 58, 59 (1918). C. Europe.
brevimana, Mik, in litt. Engel, ibidem, 1918, p. 239 [1918] (*Orthorhynchium*).

SUBGENUS EUCELIDIA

1. *E. Escheri*, Zetterstedt, Fauna Ins. Lappon. p. 558 [1838] (*Brachystoma*); C. & N. Europe.
Dipt Scand. Vol. 1, p. 362 [1842] (*Brachystoma*); Bonsdorff, Finl.
tvåv. Ins. Dipt. Vol. 1, p. 163 [1861] (*Brachystoma*); Mik, Verh.
Zool.-bot. Ges. Wien, Vol. 31, p. 327 (1881); Wahlgren, Ent.
Tidskr. Vol. 31, p. 83 (1910); Engel, Deutsche Ent. Zeitschr. 1918,
p. 201, f. 28, 30e (1918).
longipennis, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 349 [1880] (*Clinocera*); Mik, ibidem, Vol. 31, p. 326 [1881] (*Roederia*).
stigmaticalis, Zetterstedt, undescribed (*Brachystoma*).
2. *E. pirata*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 350 [1880] (*Clinocera*); C. Europe.
ibidem, Vol. 31, p. 327 (1881); Engel, Deutsche Ent.
Zeitschr. 1918, p. 205, f. 31 (1918).
3. *E. Zetterstedti*, Fallen, Dipt. Suec. Suppl. 1, p. 7 [1826] (*Empis*); Zetter-
stedt, Dipt. Scand. Vol. 1, p. 362 [1842] (*Brachystoma*); Loew, Wien.
Ent. Monatschr. Vol. 2, p. 249 [1858] (*Clinocera*); Schiner, Fauna
Dipt. Austr. Vol. 1, p. 84 [1862] (*Clinocera*); Mik, Verh. Zool.-bot.
Ges. Wien, Vol. 31, p. 327, pl. 16, f. 11 (1881); Strobl, Jarb. Mus.
Kärnten, Klagenfurt, Vol. 47, p. 203 (1901); Lundbeck, Dipt. Dan.
Vol. 3, p. 222, f. 92 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 83
(1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 63
(1913); Engel, Deutsche Ent. Zeitschr. 1918, p. 203, f. 29, 30
(1918).

SUBGENUS PHILOLUTRA

1. *Ph. aquilex*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 272 [1869] (*Clinocera*); C. Europe.
Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327, pl. 16, f. 16

DIPTERA

- (1881); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 107 (1893); Engel, Deutsche Ent. Zeitschr. 1918, p. 229, f. 51 (1918).
2. *Ph. Bohemani*, Zetterstedt, Fauna Ins. Lappon. p. 558 [1838] (*Brachystoma*); Europe. Dipt. Scand. Vol. 1, p. 361 [1842] (*Brachystoma*); Vol. 8, p. 3019 (1849); Loew, Wien. Ent. Monatschr. Vol. 2, p. 252 [1858] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327 (1881); in Beck, Fauna Hernstein, Vol. 2, p. 2 (1885); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 107 (1893); Bezz, Bull. Soc. Ent. Ital. Vol. 30, p. 151 (1899); Wahlgren, Ent. Tidskr. Vol. 31, p. 83 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 63 (1913); Engel, Deutsche Ent. Zeitschr. 1918, p. 225, f. 48, 49 (1918).
3. *Ph. fallaciosa*, Loew, Berl. Ent. Zeitschr. Vol. 17, p. 44 [1873] (*Clinocera*); C. Europe. Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327, pl. 16, f. 14, 18 (1881); Mik, in Beck, Fauna Hernstein, Vol. 2, p. 2 (1885); Engel, Deutsche Ent. Zeitschr. 1918, p. 227, f. 50 (1918). *impudica*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 351 [1880] (*Clinocera*) ibidem, Vol. 31, p. 327 (1881).
4. *Ph. hygrobia*, Loew, Wien. Ent. Monatschr. Vol. 2, p. 248 [1858] (*Clinocera*); C. & S. Europe. Loew, Beschr. Eur. Dipt. Vol. 1, p. 273, note [1869] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327 (1881); Mik, in Beck, Fauna Hernstein, Vol. 2, p. 2 (1885); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 107 (1893); Bezz, Bull. Soc. Ent. Ital. Vol. 30, p. 151 (1899); Engel, Deutsche Ent. Zeitschr. 1918, p. 223, f. 47 (1918).
5. *Ph. lagunæ*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 43, pl. 1, f. 8, 9 Canary Islands. (1908); Engel, Deutsche Ent. Zeitschr. 1918, p. 231, f. 53 (1918).
6. *Ph. simplex*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 207 : Cent. 2, No. 49 United States. (1862); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 240 (1902); Engel, Deutsche Ent. Zeitschr. 1918, p. 36 [1918] (*Clinocera*).
7. *Ph. Wachtlii*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 351 [1880] C. Europe. (*Clinocera*); ibidem, Vol. 31, p. 327 (1881); Engel, Deutsche Ent. Zeitschr. 1918, p. 230, f. 52 (1918).

SUBGENUS PSEUDOWIEDEMANNIA

1. *P. armata*, Engel, Deutsche Ent. Zeitschr. 1918, p. 244, f. 61 (1918). C. Europe.
2. *P. lamellata*, Loew, Beschr. Eur. Dipt. Vol. 1, p. 267 [1869] (*Clinocera*); C. Europe. Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327 [1881] (*Wiedemannia*); Becher, Denks. Akad. Wien. Vol. 45, p. 147, pl. 3, f. 14 [1882] (*Clinocera*); Bezz, Ann. Mus. Hungar. Vol. 2, p. 202 [1904] (*Wiedemannia*); Kertész, Rovart. Lappon. Vol. 18, p. 67 [1911] (*Wiedemannia*); Engel, Deutsche Ent. Zeitschr. 1918, p. 246, f. 63 (1918).
3. *P. microstigma*, Bezz, Ann. Mus. Nat. Hungar. 1904, p. 201 [1904] Bosnia. (*Wiedemannia*); Engel, Deutsche Ent. Zeitschr. 1918, p. 248, f. 64 (1918).

SUBGENUS ROEDERELLA

1. *R. Csernyi*, Bezz, Ann. Mus. Hungar. Vol. 3, p. 362, f. 1 [1905] (*Roederia*); Italy. Engel, Deutsche Ent. Zeitschr. 1918, p. 197, f. 26, 27 (1918). *longipennis*, Bezz (not Mik), Bull. Soc. Ent. Ital. Vol. 30, p. 150 [1899] (*Roederia*). var. *rufipes*, Oldenbergs Archiv. Naturges. Vol. 80 (9), p. 92 [1915] (*Roederia*). Apennines.

SUBGENUS WIEDEMANNIA

1. *W. bilobata*, Oldenberg, Ann. Mus. Hungar. Vol. 8, p. 349, f. 2, 3 (1910); C. Europe.
Engel, Deutsche Ent. Zeitschr. 1918, p. 257, f. 69 (1918).
2. *W. bistigma*, Curtis, Brit. Ent. Vol. 8, p. 513 [1834] (*Heleodromia*); Walker, C. & S. Europe.
Ins. Brit. Vol. 1, p. 107 (1851); Loew, Wien. Ent. Monatschr. Vol. 2, p. 243 [1858] (*Clinocera*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 84 [1862] (*Clinocera*); Nowicki, Verh. Nat. Ver. Brünn, Vol. 6, pl. 2, f. 7 [1868] (*Clinocera*); Loew, Beschr. Eur. Dipt. Vol. 1, p. 269, note [1869] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327 (1881); Bezzı, Ann. Mus. Hungar. Vol. 2, p. 202 (1904); Bezzı, ibidem, Vol. 3, p. 364 (1905); Lundbeck, Dipt. Dan. Vol. 3, p. 223, f. 93 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 83 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 63 (1913); Engel, Deutsche Ent. Zeitschr. 1918, p. 251, f. 66 (1918).
borealis, Zetterstedt, Fauna Ins. Lappon. p. 207 (1832) no description; ibidem, p. 559 (1838); Dipt. Scand. Vol. 1, p. 365 (1842); ibidem, Vol. 8, p. 3021 (1849); ibidem, Vol. 11, p. 4270 (1852); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 163 (1861).
juvenilis, Zetterstedt, Dipt. Scand. Vol. 1, p. 366 (1842); ibidem, Vol. 8, p. 3022 (1849).
3. *W. Braueri*, Mik, Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 352 [1880] Austria.
(*Clinocera*); Mik, ibidem, Vol. 31, p. 327 (1881); Bezzı, Ann. Mus. Hungar. Vol. 2, p. 202 (1904); Bezzı, ibidem, Vol. 3, p. 363 (1905); Engel, Deutsche Ent. Zeitschr. 1918, p. 261, f. 71 (1918).
4. *W. oxystoma*, Bezzı, Ann. Mus. Hungar. 1905, p. 364, f. 2 (1905); Engel, Italy.
Deutsche Ent. Zeitschr. 1918, p. 255, f. 68 (1918).
5. *W. rhynchos*, Nowicki, Verh. Nat. Ver. Brünn, Vol. 6, p. 89, pl. 2, f. 6 C. Europe.
[1868] (*Clinocera*); Loew, Beschr. Eur. Dipt. Vol. 1, p. 268 [1869] (*Clinocera*); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 327, pl. 16, f. 15 (1881); Mik, Beck. Fauna Hernstein, Vol. 2, p. 2 (1885); Bezzı, Ann. Mus. Hungar. Vol. 2, p. 202 (1904); Bezzı, ibidem, Vol. 3, p. 364 (1905); Lundbeck, Dipt. Dan. Vol. 3, p. 224, f. 94 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 258, f. 70 (1918).
6. *W. stylifera*, Mik, Wien. Ent. Zeit. Vol. 8, p. 150 (1889); Bezzı, Bull. C. & S. Europe.
Soc. Ent. Ital. Vol. 30, p. 151 (1899); Ann. Mus. Hungar, Vol. 2, p. 202 (1904); ibidem, Vol. 3, p. 364 (1905); Engel, Deutsche Ent. Zeitschr. 1918, p. 253, f. 67 (1918).
7. *W. tricuspidata*, Bezzı, Ann. Mus. Hungar, Vol. 3, p. 365, f. 3 (1905); Hungary.
Engel, Deutsche Ent. Zeitschr. 1918, p. 262, f. 72 (1918).

14. GENUS DOLICHOCEPHALA, MACQUART

Dolichocephala, Macquart, Mém. Soc. Sc. Lille, 1823, p. 147 (1823); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248, 262 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 224 (1908); Kertész, Cat. Dipt. Vol. 6, p. 127 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 535 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 238 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 84 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 3 (1918); Brunetti, Fauna Brit. India, Dipt. Vol. 1, p. 370 (1920).

Ardoptera, Macquart, Dipt. N. France, Vol. 3, p. 105 (1827); Hist. Nat. Dipt. Vol. 1, p. 358 (1834); Westwood, Gen. Syn. p. 132 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 323 (1842); Boitard, Man. Ent. Vol. 3, p. 325 (1843); Walker, Ins. Brit. Vol. 1, p. 103 (1851); Rondani, Dipt. Ital. Vol. 1, p. 149 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 561 (1857); Schiner, Fauna Dipt.

Austr. Vol. 1, p. 85 (1862); Lioy, Atti. Instit. Ven. 1864, p. 722 (1864); Röder, Wien. Ent. Zeitschr. Vol. 6, p. 169 (1887); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 238 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 246 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 509 (1910).

Leptosceles, Haliday, Ent. Mag. London, Vol. 1, p. 160 (1838); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 560 (1910).

Character. — Small, slender, delicate, dark colored, rather shining species with brown wings marked with hyaline spots, with strong bristles and with simple legs. Head produced downward, the neck attachment located high up on the occiput so that the head is borne projecting obliquely forward, the mouth-opening at the end of the head and not on the lower side as in *Hemerodromia*. Occiput with radiating bristles, those above long, one pair of proclinate frontoorbitals, ocellar bristles strong, curving upward and spaced apart as much as the posterior ocelli, no lesser or postvertical bristles; front very short and broad, the ocelli on a large prominent tubercle, face tapering beneath, cheeks narrow; eyes large and ovate, the emargination at the antennæ small, facets uniform, densely short-hairy; antennæ not quite touching, short and three-jointed, the second and third joints globular, the thickened arista not hairy, bending downward, two-jointed but the basal segment minute; proboscis very short, thick, palpi minute, curving forward. Thorax elongate, the prothorax visible from above, the prosternum rather long but not flared out around the base of the coxae and not sulcate medially; bristles long, comprising one humeral, one posthumeral, no notopleural, one supraalar, one small postalar, five dorsocentral and two scutellar; metapleuræ with a group of upturned hairs, sparse in *irrorata*. Abdomen depressed, shining, more or less hairy, pygidium rather small but robust, turned back over the small last two segments, comprising a ventral hood-shaped piece, two lateral valves and two chitinized superior forcipate valves, the penis with thickened base and circularly curved end; no ovipositor, the end of the female abdomen blunt. Legs unarmed, the front coxae half as long as their femora, no bristles or tibial spurs but the front and hind tibiae tipped with a pecten of close fitting setulae, pulvilli small, empodium linear. Wings cuneate, no anal angle, costa encompassing the entire margin, a strong basal costal bristle, costal setulae minute, fringe of the hind margin longer than the anterior crossvein; auxiliary vein complete, ending by a gentle curve in the costa, first vein ending before the middle of the wing, no stigma, longitudinal veins more or less undulating, three submarginal cells, the fork of the third vein angulate and there connected with the second vein by an adventitious crossvein, second basal cell small, the fourth vein originating near the middle of the anal cell, discal cell long, complete, emitting three separate posterior veins, anal cell a little shorter than the second basal, apically round, no anal vein.

Type species: *D. irrorata*, Fallen (Pl. 3, Fig. 24). Macquart gave a very clear description of *Dolichocephala* with its single species, *maculata*, which in the same paper he determined as a synonym of *irrorata*. Later he redescribed the genus as *Ardoptera*. The species of this genus are delicate little flies that frequent damp shady woods near water. They are not common, never occurring in swarms.

Geographical distribution.

- | | |
|---|-------------|
| 1. <i>D. argus</i> , nov. sp. (1). | Washington. |
| 2. <i>D. combinata</i> , Becker, Ann. Soc. Ent. France, Vol. 83, p. 122 (1914). | E. Africa. |

(1) **Dolichocephala argus**, nov. sp. — Male and female. Length 2 mm. Shining, bronzed black, wings with numerous round white spots, discal cell long, halteres black. Vertex lightly dusted with brown pollen and marked with a cinereous Y-shaped spot which starts at the neck and includes the ocellar prominence; frontoorbital bristles proclinate, the

3. *D. guttata*, Haliday, Ent. Mag. London, Vol. 1, p. 161 [1833] (*Leptosceles*); Europe.
 Walker, List. Dipt. Brit. Mus. Vol. 3, p. 504 [1849] (*Ardoptera*); Ins. Brit. Vol. 1, p. 104 [1851] (*Ardoptera*); Loew, Wien. Ent. Monatschr. Vol. 2, p. 8 [1858] (*Ardoptera*); Strobl, Mitteil. Nat. Ver. Steiermark. Graz, Vol. 29, p. 99 [1893] (*Ardoptera*); Wien. Ent. Zeit. Vol. 18, p. 77 [1899] (*Ardoptera*); Lundbeck, Dipt. Dan. Vol. 3, p. 241 (1910).
oblongo-guttata, Dale, Hist. of Glenville's Wootton, p. 264 [1878] (*Ardoptera*).
 var. *exoleta*, Haliday, Ent. Mag. London, Vol. 1, p. 161 [1833] (*Leptosceles*).
 var. *nigrohalterata*, Strobl, Glasnik Zem. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 399 [1898] (*Ardoptera*); Wiss. Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 561 [1900] (*Ardoptera*).
 England
 C. Europe.
4. *D. irrorata*, Fallen, Empid. Suec. p. 13 [1815] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 66, pl. 23, f. 11 [1822] (*Hemerodromia*); Macquart, Dipt. N. France, Vol. 3, p. 106, pl. 3, f. 2 [1827] (*Ardoptera*); Haliday, Ent. Mag. London, Vol. 1, p. 161 [1833] (*Leptosceles*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 358, pl. 8, f. 3 [1834] (*Ardoptera*); Zetterstedt, Fauna Ins. Lappon. p. 544 [1838] (*Hemerodromia*); Blanchard, Hist. Nat. Ins. Vol. 3, p. 583 [1840] (*Ardoptera*); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. 1840, p. 22, pl. 1, f. 31; Isis, Vol. 7, p. 551, f. 31 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 324, 1 [1842] (*Ardoptera*); Boitard, Man. Ent. Vol. 3, p. 325 [1843] (*Ardoptera*); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 504 [1849] (*Ardoptera*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3012 [1849] (*Ardoptera*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 59 [1851] (*Ardoptera*); Loew, Wien. Ent. Monatschr. Vol. 2 [1858] (*Ardoptera*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 155 [1861] (*Ardoptera*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 85 [1862] (*Ardoptera*); Strobl, Mitteil. Nat. Ver. Steiermark. Graz, Vol. 29, p. 98 [1892] (*Ardoptera*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 239 [1902] (*Ardoptera*); Czizek, Jahrb. Mähr. Landesmus. Klagenfurt, Vol. 7, p. 166 [1907] (*Ardoptera*); Lundbeck, Dipt. Dan. Vol. 3, p. 239, f. 104, 105 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 85 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 65 (1913). — PI. 3, Fig. 24.
anomala, Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 59 [1851] (*Ardoptera*).
maculata, Macquart, Mém. Soc. Sc. Lille, 1823, p. 147, 165 (1823).
 Europe, North America.
5. *D. ocellata*, Costa, Ann. Sc. Napoli, Vol. 1, p. 76 [1854] (*Ardoptera*); Wien. Ent. Monatschr. Vol. 3, p. 64 [1859] (*Ardoptera*); Verrall, Ent. Mag. London, Vol. 48, p. 26 [1912] (*Ardoptera*). C. & S. Europe.
novemguttata, Strobl, Mitteil. Nat. Ver. Steiermark. Graz, Vol. 29, p. 98 [1892] (*Ardoptera*); Wien. Ent. Zeit. Vol. 18, p. 77 [1899] (*Ardoptera*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31, p. 6 (1908); ibidem, Vol. 37 (3), p. 65 (1913).
oculata, Loew, Wien. Ent. Monatschr. Vol. 2, p. 7 [1858] (*Ardoptera*).
 var. *albohalterata*, Strobl, Glasnik Zem. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 399 [1898] (*Ardoptera*); Wiss. Mitteil. Bosn. Herceg. Sarajevo, Vol. 7, p. 561 [1900] (*Ardoptera*). C. Europe
6. *D. septemnotata*, Brunetti, Rec. Indian Mus. Vol. 9, p. 35 (1913); Fauna Brit. India Dipt. Vol. 1, p. 371, pl. 4, f. 19-21 (1920). W. Himalayas.
7. *D. sparsa*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 122 (1914). E. Africa.

occipitals erect; face cinereous, cheeks small, mouth-parts scarcely protruding. Dorsum very lightly dusted, a faint gray median stripe visible on the front portion, pleuræ largely shining but with a more or less evident cinereous horizontal stripe; metapleural hairs abundant; abdomen shining. Legs including the coxae pale fuscous, the last tarsal joints darker, the coxae apically with longer hairs. Wings infumated, with round white spots as follows: one in third submarginal, the fourth and the fifth posterior cells, two in the first submarginal, the discal and the second posterior cells, three in the second submarginal cell, five in the marginal and first posterior cells, the spots of the first submarginal cell largest, second vein sinuous, submarginal cells of equal extent on the costa, last two sections of the fourth vein equal, of the fifth vein, 3 : 1, no anal vein, the anal cell not extending as far as the second basal, marginal cilia twice as long as the anterior crossvein.

Seven specimens; Bellingham and Mount Constitution, Washington, May to August.

15. GENUS LAMPOSOMA, BECKER

Lamposoma, Becker, Berl. Ent. Zeitschr. Vol. 33, p. 338 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 251, 263 (1903); Kertész, Cat. Dipt. Vol. 6, p. 120 (1909); Engel, Deutsche Ent. Zeitschr. 1918, p. 3 (1918).

Characters. — A minute insect, measuring one millimeter, shining black in color and resembling *Clinocera* in form and venation, but differing in lacking a pollinose coating and in having broader wings. Head rounded, eyes large, finely pubescent, cheeks narrow, divided, face narrowed beneath, white pollinose and bare, the occiput shining black; third joint of the antennæ oval, with a long apical arista; proboscis very short. Thorax polished, a faint pollinose median vitta present, five strong dorsocentral bristles, two scutellars, no acrostichals; pleuræ and abdomen shining, bare except for a few hairs on the terminal segments; epipygium small, reflexed, with the lower piece hood-shaped. Legs slender, without bristles, the femora somewhat pale-ciliate beneath, the tibiæ slightly enlarged apically, no pulvilli, empodium very small. Wings clear hyaline, broad and bluntly rounded, the veins slender, no stigma, costal bristle large, first vein ending before the middle of the wing, fork of the third vein greatly bowed, the second submarginal cell broad, discal cell rather blunt, emitting three posterior veins, anal cell extending as far as the second basal, no trace of the anal vein.

But a single species is known, *L. cavaticum*, which was found in a cave in Dalmatia.

Geographical distribution.

1. *L. cavaticum*, Becker, Berl. Ent. Zeitschr. Vol. 33, p. 339, f. 3 (1889); S. Europe.
Bezzi, Arch. Zool. Exper. Paris (5), Vol. 8, p. 50 (1911). — (Pl. 7,
Fig. 64).

SUBFAMILY HEMERODROMIINÆ

Characters. — Very slender delicate species with raptorial front legs; usually of light coloration and with pale yellow appendages. Head globular, mouth-opening placed far back, proboscis inflexed, always shorter than the head and stout at the base, palpi small; eyes large, nearly or quite contiguous on the face in both sexes, but separated above the antennæ, broadly emarginate on the lower occiput, lower facets largest when the eyes encroach on the face; antennæ short, three-jointed, with a short style (*Hemerodromia*) or hairy two-jointed arista (*Chelipoda*), inserted below the middle of the head. Thorax narrow and diagonally elongate, scarcely arched, mesosternum much longer than the prosternum, front legs greatly distant from the posterior pairs and located close to the head, mesonotum nearly bare (*Hemerodromia*) or with bristles (*Chelipoda*), metapleuræ hairy in the latter group, otherwise bare, prescutellar depression less evident than in the *Clinoceratinae*. Abdomen slender, sometimes the last segments of the female drawn out to form an ovipositor; genitalia of the male either a terminal pygidium or reflexed epipygium. Legs long, the posterior pairs very slender, the front pair strongly raptorial, front coxæ at least twice as long as the others, front femora thickened and armed below. Wings cuneiform, the anal angle not developed, costa continuing around the wing, except in *Hemerodromia*, s. str., where it is attenuated beyond the fourth vein, third vein forked (*Hemerodromia*) or not (*Chelipoda*), discal and anal cells present or absent, petiole of the second and third veins short, arising nearer the anterior than the humeral crossvein, anal crossvein if present perpendicular to the axis of the wing, stigma absent or present. Calypteres with straight edge and small fringe, and closely united to the base of the wing.

SYNOPSIS OF THE GENERA OF THE HEMERODROMIINÆ

1. Style shorter than the third antennal joint; proboscis usually inflexed and pointed; no discal macrochætae, metapleuræ bare, mesosternum but little concave; pygidium terminal and more or less erect; ovipositor chitinized and sometimes produced; third vein forked, anal cell, when present, broader towards apex, the crossvein straight, basal cells more than one-third the length of the wing, costa with very short hairs HEMERODROMIA, s. latiss. 2.
- Arista more than twice as long as the third antennal joint; proboscis nearly vertical; some discal macrochætae present, metapleuræ with some setulæ, mesosternum more or less concave; pygidium reflexed over the abdomen; ovipositor not produced and chitinized; no spur at the end of the front tibiae; third vein not forked, anal cell with parallel sides, except in Cephalodromia, and apically rounded, basal cells small, less than one-third the length of the wing, the anal and second basal cells subequal in extent, costa hairy and usually with a basal bristle. CHELIPODA, s. latiss. 9.
2. Eyes more or less separated on face; anal cell, when present, much shorter than second basal, the anal crossvein oblique; proboscis inflexed; no prothoracic bristle 3.
- Eyes large, contiguous beneath antennæ; anal cell subequal to second basal, anal crossvein perpendicular; prothoracic and supra-alar bristles distinct; discal cell emitting three veins Genus DRYMODROMIA, Becker.
3. Humeral crossvein wanting, auxiliary vein fusing with the costa close to the base of the wing, first vein terminating before the middle of the wing, pedicel of the second and third veins atrophied, anal cell imperfect or wanting; thorax quadrangular, without a depressed prescutellar space; front short, face very narrow, the eyes subcontiguous 4.
- Humeral crossvein present, auxiliary vein well separated from the costa and parallel with it, first vein terminating at or beyond the middle of the wing, pedicel of the second and third veins distinct though short, anal cell complete, the anal vein separated from the thickened hind margin; thorax with a distinct depressed prescutellar space and with evident humeri; eyes plainly separated on the face; style very short; scutellum smaller than the metanotum Genus CHELIFERA, Macquart. 5.
4. Discal cell open outwardly, anal cell wanting, second posterior cell petiolate; auxiliary vein a little bent, fused along its middle course and slightly separating from the costa at its tip, pedicel of the second and third veins entirely atrophied; hind margin of the wing scarcely thickened, the costa abruptly thinner beyond the fourth vein; anal vein running into the hind margin at its very root; scutellum as large as and usually larger than the metanotum; humeri not constricted; ovipositor usually short and conical;

- style of antennæ at least half as long as the third joint; lower facets of the eyes large. (Pl. 3, Fig. 26)* Genus **HEMERODROMIA**, Meigen.
- Discal cell closed outwardly, anal crossvein present, second posterior cell sessile; auxiliary vein straight, completely fusing with the costa, base of the pedicel of the second and third veins distinct; hind margin of the wings thickened, the anal vein separate but weak; humeri strongly constricted; ovipositor long, ensiform; third antennal joint without evident style; lower facets small; front coxae of male as thick as their femora. (Pl. 3, Fig. 27)* Genus **COLABRIS**, nov. gen.
5. *Discal cell complete; front legs strongly raptorial; style of the antennæ very short.* 6.
- Discal cell fused with either the second basal or the third posterior cell; thorax not narrowed in front* 7.
6. *Second posterior cell petiolate; thorax narrowing in front; no ovipositor. (Pl. 3, Fig. 28)* Subgenus **CHELIFERA**, Macquart.
- Second posterior cell sessile or nearly so; thorax broad in front; ovipositor rather long and thick* Subgenus **CLADODROMIA**, Bezzi.
7. *Second posterior cell petiolate; front legs strongly raptorial, the front femora spinose* 8.
- Second posterior cell sessile, first vein ending at the middle of the wing, second vein very short, humeral crossvein vestigial; front femora but little thickened and not strongly spinose; ovipositor long, ensiform; face very narrow; thorax shining black and not pollinose; palpi spatulate and silky; style of the antennæ microscopic* Subgenus **NEOPLASTA**, Coquillett.
8. *Discal cell fused with the third posterior cell; style of the antennæ microscopic; no ovipositor, the abdomen of the female with a blunt termination* Subgenus **THANATEGIA**, nov. subgen.
- Discal cell fused with the second basal cell; style one-third as long as the third antennal joint; ovipositor ensiform* Subgenus **METACHELA**, Coquillett.
9. *Anal vein separated from the hind margin of the wing, the anal cell not broader than the second basal cell; posthumeral and vertical macrochætae present; upper occiput short and declivous; wings rarely plicated* 10.
- Anal cell broad, reaching the hind margin with which the anal vein is fused; vertical and all thoracic bristles except the supraalar absent; occiput horizontal, strongly elevated above; discal cell open apically, the fourth vein furcate* Genus **CEPHALODROMIA**, Becker.
10. *Arista rather thick, but not plumose; front coxae not tuberculate and armed only with a basal spine, middle legs not armed* Genus **CHELIPODA**, Macquart. 11.
- Antennæ elongate, the arista bearing a plume of hairs beneath; front coxae, middle tibiae and middle metatarsi ♂ armed with tubercles or spines; discal cell complete* Genus **Ptilophyllostromia**, Bezzi.
11. *Discal cell complete, emitting three veins apically; anal cell subequal to the second basal cell. (Pl. 3, Fig. 25)* Subgenus **CHELIPODA**, Macquart.
- Discal cell open, the posterior crossvein absent and the fourth vein acutely forked; anal cell a little shorter than the second basal cell* Subgenus **PHYLLODROMIA**, Zetterstedt.

HEMERODROMIA, MEIGEN, SENSU LATIORE

Hemerodromia, Meigen, Syst. Beschr. Vol. 3, p. 61 (1822); Macquart, Dipt. N. France, Vol. 3, p. 101 (1827); Hist. Nat. Dipt. Vol. 1, p. 347 (1834); Zetterstedt, Fauna Ins. Lappon. p. 542 (1838); Dipt. Scand. Vol. 1, p. 260 (1842); Boitard, Man. Ent. Vol. 3, p. 322 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 142 (1851); Rondani, Dipt. Ital. Vol. 1, p. 148 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 82 (1862); Loew, Wien. Ent. Monatschr. Vol. 8, p. 237-255 (1864); Lioy, Atti Instit. Ven. 1864, p. 719 (1864); Röder, Wien. Ent. Zeit. Vol. 6, p. 169 (1887); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 123 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389, 391 (1895); Williston, Trans. Ent. Soc. London, Vol. 3, p. 440 (1896); Man. N. Amer. Dipt. p. 75 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 233 (1902); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 452 (1905); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 123 (1889); Kertész, Cat. Dipt. Vol. 6, p. 112 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 228 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 44, 83 (1910).

Characters. — Slender, elongate, frail, nearly bare, little flies with elongate raptorial front legs and forked third vein, usually of reddish coloration, but sometimes the body is blackened. Head produced obliquely downward, somewhat flattened, nearly as broad as the thorax, ocelli placed rather far forward, eyes obliquely oval, bare, with a rather deep notch at the antennæ, facets more or less enlarged below, lower orbits usually with a conspicuous white pruinose spot, which is broader below, vertex large and broad, front varying from short to long, but broader than the short face, which is sometimes quite narrowed by the subcontiguous eyes, cheeks linear, oral opening and epistome with dense silky pubescence, vertex rarely with setulæ and with no or one bristle. Antennæ inserted low on the head, contiguous, three-jointed, the basal joints not setose, the third joint compressed ovate, with apical unjointed style which varies from one-twentieth to two-thirds the length of the third joint and is either bare or hairy. Proboscis about as long as the head, slender, incurved, the densely hairy, small, slender, one-jointed palpi more or less retracted, vertical or directed somewhat backward. Thorax conspicuously longer than wide, quadrate or tapering forward when viewed from above when the prothorax and metathorax are quite visible, humeral grooves more or less distinct, pollinose but not pubescent, with very sparse biseriate acrostichal and uniserrate dorsocentral setulæ, no prescutellar bristles, no humeral, one supraalar, one or no postalar bristles, several notopleural setulæ, sometimes two fine scutellar bristles, sometimes no, two or four scutellar hairs, setulæ in front of the scutellum proclinate; mesopleuræ very long between the front and middle coxal attachments, metapleuræ bare. Abdomen slender and cylindrical, comprising eight segments, the first very small and in the male the last also, ovipositor more or less conical, variable in length, pygidium globose or with a hood-shaped lower piece surmounted by a pair each of broad, usually forcipate lateral and dorsal valves, penis clavate or capitate, sometimes hidden. Legs slender, the posterior pairs simple but the front pair remarkably modified, front coxæ very long, usually extremely slender, the prosternum flared at the coxal cavities, front femora and front tibiæ raptorial, the former swollen and with biseriate flexor setulæ and usually with prominent lateral stiff setæ, the front tibiæ shorter than the femora, produced within as a sharp spur and with flexor setulæ which engage between the femoral setulæ when the leg is folded, otherwise no bristles or apical spines, tarsi simple and slender, the posterior metatarsi long, pulvilli minute. Wings slender, oval, the anal angle wanting, costa continuing around the entire wing, but in *Hemerodromia* s. str. very much attenuated beyond the fourth vein, no costal bristle or setulæ, the costal hairs short, but the hairs of the hind margin rather long, especially toward the base, auxiliary vein present but its

course variable, third vein forked, the upper branch rather erect, stigma present or absent, pedicel of the second and third veins very short; otherwise the neuration varies greatly as outlined below.

Ethology: These odd little flies with their semi-translucent legs and bodies, are the most delicate of the Empididae. They occur, never in large numbers, in shady sheltered spots in the woods, usually near water, running over herbage and low shrubbery. Sometimes the species occur in houses, where, as Mrs. Slosson has said: «as they run upon a window-pane with the light behind them they seem mere shadows, little ghosts, frail, elf-like things». Some species are nocturnal, and are attracted to lights.

Brocher, who has recently studied the life-history of *C. precatoria*, found the larva in the mud of streams. The larva is slender, but tapers in front, its posterior end is furnished with five or six diverging hairs, and ventrally it has seven pairs of retractile propodia. The pupa has paired tracheal filaments, each two-thirds as long as the body, attached at the spiracles, and including one prothoracic and seven abdominal pairs; it has no dorsal spines but merely girdles of setulae on the abdomen. The pupal period lasted thirteen days.

Taxonomy: The old genus *Hemerodromia* has had a host of synonyms. While many of the names were bestowed as subgenera on certain permutations of three neurational characters, — namely, the presence or absence of the discal cell, the presence or absence of the anal cell and the shape of the second posterior cell, whether sessile or petiolate, — others are pure homonyms. The usual custom has been to consider these groups as subgenera, since they were based on but slight variations of the venation. These so-called subgenera have been characterized as follows:

	Discal cell	Anal cell	Second posterior cell
<i>Hemerodromia</i> , Meigen, s. str.	imperfect	absent	petiolate
<i>Metachela</i> , Coquillett	imperfect	present	petiolate
<i>Neoplasta</i> , Coquillett	imperfect	present	sessile
<i>Chelifera</i> , Macquart	present	present	petiolate
<i>Cladodromia</i> , Bezzi	present	present	sessile

If such variations in wing venation were not correlated with other characters the groups would be artificial and could have at most but subgeneric rank, for the second posterior cell, e. g. in *Cladodromia*, varies from sessile to V-shaped, and even to Y-shaped with a short petiole, and *Metachela* as defined included two species, one with the discal cell fused with the second basal and the other with it fused with the third posterior cell. However, a closer inspection shows that the trivial differences in neuration are associated with more fundamental characters in the structure of the wing and correlated at the same time with differences in the form of the body. Accordingly, the old genus *Hemerodromia* has been segregated into two distinct groups, *Hemerodromia*, s. str. (i. e. *Microdromia*) and *Chelifera*. The remaining groups are less important subdivisions of the latter genus, although *Chelifera*, s. str., with its thorax tapering anteriorly, may ultimately be removed from the broad-shouldered others, and *Neoplasta* with its narrow face and slender front legs may likewise be segregated from the rest. In addition, another genus has been erected for two Central American species whose neuration superficially resembles that of *Neoplasta*.

Concerning which of the restricted groups should bear the name *Hemerodromia* there has been much discussion. As this can be settled only by ascertaining the type species, various writers have argued as variously for this or that genotype. The following statement shows the nomenclatural status of *Hemerodromia* and its allies.

Hemerodromia, Meigen. Meigen originally described nine species under this name; the fifth, *mantispa*, Panzer, was selected as the type by Westwood, in 1840. However, in 1823 Macquart had

designated *mantispa*, Panz. as the type of his genus *Chelipoda*, thus removing this species from consideration as the type of *Hemerodromia*. In 1856 Rondani then assigned *oratoria* Fallen, Meigen's third species, as the type of *Hemerodromia*, but subsequent writers have generally accepted Meigen's first species, *monostigma* Meigen, as the type, following the « first species » principle.

However, this is not the whole story and the status of the names is far from being as clear as might be. *Mantispa*, Panzer proves to be *melanocephala* Fabricius : Macquart's *mantispa* is *vocatoria*, Fallen, and *monostigma*, Meigen is *precatoria* Fallen. The following arguments can be made :

1. Macquart's genus *Chelipoda*, with *mantispa* Panzer (= *melanocephala*, Fabricius) specified as its type, may be considered as a complete synonym of *Hemerodromia*, thus using the name in a different sense than agreed on by general usage. Recent codes sanction this procedure, as they make no allowance for mistakes in identification.

2. Or, if we allow for Macquart's misidentification, since he states that his *mantispa* has a discal cell whereas Panzer's species had the discal cell open, and make *mantispa* Macquart (= *vocatoria*, Fallen) the type of *Chelipoda*, we reserve the two oldest names, *Hemerodromia* and *Chelipoda* for two valid genera. This seems certainly the logical proceeding, even though it is forbidden by certain codes of nomenclature.

3. Since Macquart's *mantispa* was erroneously identified, Coquillett in 1903, selected Macquart's second species, *minor* as the type for *Chelipoda*, and as this proved later to be the true *mantispa* of Panzer (= *melanocephala*, Fabricius), in 1910 he placed this genus as a synonym of *Hemerodromia*, both having the same type by late designation. However, in doing this he overlooked Macquart's definite words : « celle nommée *mantispa*, que je considère sous le nom de *Chéliopode*, comme le type d'un genre distinct ». While the codes of nomenclature permit the abandonment of a genus if its type should prove to be the same as the previously selected type of an older genus, yet when an author selects the genotype when publishing his genus, that species is no longer available as the type for a polytypical genus. Applying this principle in the present case, Westwood should not have designated *mantispa*, Panzer as type for *Hemerodromia*, as it had already been specified as the type of *Chelipoda*.

4. Many writers (e. g. Bigot, Loew, Williston, Lundbeck, Bezzi, Kertész) have accepted Meigen's first species, *monostigma*, which is either *precatoria*, Fallen or *stigmatica* Schiner, as the genotype; although their reason for doing so is apparently that it was Meigen's first species. This principle is not tenable in any of the present day codes of nomenclature.

5. Therefore Rondani's selection of *oratoria*, Fallen is apparently the earliest indication of a type species for *Hemerodromia*.

* * *

SYNONYMY OF *HEMERODROMIA*, SENSU LATIORE

1. *Hemerodromia*, Meigen, Coquillett (1895, 1903). Type : *oratoria*, Fallen. — **PI. 3, FIG. 26.**
Microdromia, Bigot. No species mentioned. Type : *oratoria*, Fallen, Coquillett's designation (1902).
2. *Colabris*, nov. gen. Type : *rufescens*, nov. sp.
3. *Chelifera*, Macquart. Type : *raptor*, Macquart, the only species, which is *precatoria*, Fallen. — **PI. 3, FIG. 28.**
Hemerodromia, Bigot, Loew, Williston, Bezzi, Kertész, Lundbeck. Type : *monostigma*, Meigen, the first species, which is *precatoria*, Fallen.

Mantipeza, Rondani, Coquillett, Melander. Type : *monostigma*, Meigen, the only species, which is *precatoria*, Fallen.

Polydromia, Bigot. No species mentioned. Type : *precatoria*, Fallen, Coquillett's selection in 1903.

Metachela, Coquillett. Type : *collusor*, Melander, Coquillett's designation.

Hemerodromia, Melander (1902).

Neoplasta, Coquillett. Type : *scapularis*, Loew, Coquillett's designation.

Cladodromia, Bezzi. Type : *inca*, Bezzi. Two species were described, *inca* was figured.

Thanategia, nov. subgen. Type : *defecta*, Loew.

4. *Drymodromia*, Becker. Type : *Jeanneli*, Becker, the only species.

I. GENUS HEMERODROMIA, MEIGEN, S. STR.

Hemerodromia, Meigen, Syst. Beschr. Vol. 3, p. 61 (1822); Rondani, Dipt. Ital. Vol. 1, p. 148 (1856); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 250, 261 (1903).

Microdromia, Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 557, 563 [1857] (*Microdromyia*); Williston, Trans. Ent. Soc. Lond. Vol. 3, p. 440 (1896); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253 (1903); Melander, Williston's Man. p. 223 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 299 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 569 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 4 (1918); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 366 (1920).

Characters. — Apart from the general characters possessed by all the Hemerodromias as given previously, the following structures especially pertain to this genus.

Eyes large, the lower facets conspicuously enlarged so as to bring the eyes very close together on the middle of the face, the front short; style one-half to two-thirds as long as the third joint of the antennæ and hairy. Thorax quadrate, when viewed from above, without distinct prescutellar depression, the humeral grooves distinct; scutellum usually rather flat, as large as or larger than the metanotum, with a pair of well-separated setæ; ovipositor usually short and conical. Front coxæ thin, front femora setose. Costa greatly thinned beyond the fourth vein but the hind margin thickened at the very base, humeral crossvein wanting, the auxiliary vein bending forward and fusing with the costa close to the base of the wing and later separating from it, the first vein ending before the middle of the wing, second vein long and straight, pedicel of the second and third veins very short and completely atrophied, discal cell open outwardly, second posterior cell with a long petiole, anal cell wanting, the anal vein passing into the hind margin at its very root.

Geographical distribution.

1. *H. albicornis*, Meigen, Syst. Beschr. Vol. 3, p. 64, pl. 23, f. 14 (1822); C. Europe.
Macquart, Hist. Nat. Dipt. Vol. 1, p. 348 (1834); Walker, Ins. Brit. Dipt. Vol. 1, p. 144 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 (1862); Getcke, Wien. Ent. Zeit. Vol. 5, p. 166, pl. 2, f. 12 (1886); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 95 [1892] (*Microdromia*).
- . *H. bipunctata*, Schummel, Arb. Schles. Ges. Vaterl. Kult. Breslau, p. 70 (1832), no description.
2. *H. bivittata*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 766 (1865); Chile.
Bezzi, Ann. Mus. Hungar. Vol. 3, p. 453 [1905] (*Microdromia*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 390 [1909] (*Microdromia*).

3. *H. brachialis*, nov. sp. (1). Costa Rica.
 4. *H. brunnea*, nov. sp. (2). Georgia.
 5. *H. captus*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 391 (1895); E. United States.
 Melander, Trans. Am. Ent. Soc. Vol. 28, p. 237 (1902); Coquillett,
 Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903).
 6. *H. defessa*, Williston, Trans. Ent. Soc. London, p. 439, pl. 14, f. 166 West Indies.
 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 237 (1902);
 Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903); Bezzi,
 Ann. Mus. Hungar. Vol. 3, p. 460 [1905] (*Microdromia*); Bezzi, Nova
 Acta Akad. Naturf. Halle, Vol. 91, p. 390 [1909] (*Microdromia*).
 7. *H. *detestata*, Meunier (not female), Ann. Sc. Nat. (Zool.) Vol. 7, p. 90, Baltic Amber.
 103, pl. 6, f. 6 (1908).
 8. *H. dorsalis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 33 [1913] (*Chelipoda*); W. Himalayas.
 Fauna Brit. India Dipt. Vol. 1, p. 367 [1920] (*Microdromia*).
 9. *H. dorsata*, nov. sp. (3). Java.
 10. *H. empiformis*, Say, Journ. Acad. Nat. Sc. Philad. Vol. 3, p. 99 [1823] North America.
 (*Ochthera*); Wiedemann, Aussereur. Zweifl. Ins. Vol. 3, p. 446
 [1830] (*Ochthera empiformis*); Macquart, Hist. Nat. Dipt. Vol. 2,
 p. 519 [1835] (*Ochthera empiformis*); Walker, List Dipt. Brit.
 Vol. 4, p. 1100 [1849] (*Ochthera empiformis*); Say, Compl. Writ.
 Vol. 2, p. 85 [1859] (*Ochthera*); Melander, Trans. Amer. Ent. Soc.
 Vol. 28, p. 236 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5,
 p. 264 (1903); Slosson, Ent. News, Philad. Vol. 14, p. 269 (1903);
 Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 391 (1905).

(1) ***Hemerodromia brachialis*, nov. sp.** — Length 3 mm. Head, thorax, and the greatly swollen front femora black, rather shining, remainder of legs, halteres and mouthparts whitish-yellow, antennæ brown. Face almost obliterated at the middle, white pruinose, lower occipital orbits white-pollinose. Thorax very lightly dusted, more noticeable in profile, scutellum and metanotum finely rugulose; abdominal hairs sparse and yellow, pygidium large, globose, the upper valves ham-shaped and apically pointed, the large middle valves curved, parallel-sided and apically truncate but terminating in a hook above, ovipositor small and black. Front femora slightly longer than their coxæ, with close small black teeth below which are fringed on each side with longer yellow spine-like hairs, the black color paler at knee, front tibiae two-thirds as long as femora. Wings hyaline, veins blackish except at yellowish base, pedicel of fourth vein as long as fork, sections of fifth vein 1 : 1.2.

Three males and one female; La Suiza de Turrialba, Costa Rica, April-July, Pablo Schild collector, in Melander Collection. Readily distinguished by the large black front femora.

(2) ***Hemerodromia brunnea*, nov. sp.** — Female. Length 3 mm. Near *captus* Coquillett, but the fork of the fourth vein is opposite the end of the second vein and thus the pedicel is longer than the second posterior cell. Head black cinereous, the lower orbits prominently white-pruinose; antennæ yellow, the third joint twice as long as broad and three times as long as its arista; eyes subcontiguous on the middle of the face, the lower facets large; proboscis yellow, its tip blackened. Mesonotum brown, opaque pollinose, with two faint indications of broad darker vittæ on the anterior portion; scutellum strongly convex, chocolate-brown like the metanotum; pleura brownish yellow; abdominal tergites brown, venter and incisures yellow, ovipositor conical and brown. Front femora subequal in length to their coxæ, with two rows of moderate yellow setæ beneath, front tibiae terminated by a long slender black spine. Wings normal, veins pale, the second to the fifth sections of the costa proportioned 4 : 1.2 : 3 : 1, sections of the fourth vein proportioned 1 : 0.7 : 2.3 : 1.7, of the fifth vein, 0.8 : 1, distance between the crossveins nearly three times as long as the posterior crossvein.

One specimen; Clayton, Georgia, May, collected by J. C. Bradley.

(3) ***Hemerodromia dorsata*, nov. sp.** — Female. Length 3.3 mm. Head, dorsum of the thorax and abdominal tergites black, pleurae, sternites and lateral membrane yellow. Mouthparts and antennæ yellow. Thorax almost shining, tergites matt. Legs yellow, front coxae slender, as long as their femora, the latter without basal tubercle, but with a pair of spines, front tibiae three-fifths as long as their femora, with a rather strong black terminal spine. Ovipositor not longer than the broad last abdominal segment. Halteres entirely pale yellow. Wings nearly hyaline, veins blackish except at the base, sections two to five of the costa proportioned 1 : 0.8 : 0.9 : 0.3, fork of the third vein much beyond the end of the second vein, fringe of the hind margin equal in length to the anterior crossvein.

One specimen; Tjibodas, Mount Gede, Java, from the Bryant and Palmer collection. Type in the U. S. National Museum.

- vittata*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 210; Cent. 2, No. 56 (1862).
species innom. Williston, Trans. Ent. Soc. London, Vol. 3, p. 440 (1896).
- Subsp. *coleophora*, nov. subsp. (1). Wyoming.
 11. *H. femorata*, nov. sp. (2). Panama.
 12. *H. gonatopus*, Speiser, Berl. Ent. Zeitschr. Vol. 52, p. 145 (1907). E. Africa.
 13. *H. jugulator*, nov. sp. (3). New York.
 —. *H. meigeniana*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 264 (1829), no description. —
 14. *H. mesomelana*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 390, Peru.
 392 [1909] (*Microdromia*).
 15. *H. ? nigrolineata*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 53 (1840). C. Europe.
 16. *H. nympha*, nov. sp. (4). Java.

(1) ***Hemerodromia empliformis***, subspecies *coleophora*, nov. subsp. — Male. Length 3 mm. Head black, lower orbits and face silvery, thorax reddish, the notum centrally black, tergites black except at sides, pygidium shining black: legs, antennæ, mouthparts, venter and halteres pale yellow. Six flexor spines on front femora in length equal to diameter of tibiae, basal tubercle of front femora vestigial. Pygidium large, erect, the most conspicuous portion being the obliquely ascending hamate lower valves, the upper valves small, cupuliform, pronged above and furnished with long hairs, base of penis visible, enclosed in a projecting horny sheath. Wings clear, veins yellowish, first posterior cell much constricted at tip, second posterior cell shorter than its petiole, the fork occurring much before the end of the second vein, sections of fourth vein 3 : 2 : 5.5 : 7, of fifth vein 3 : 4.

Female. Ovipositor longer than deep.

Thirteen males, ten females, Yellowstone Park, Wyoming, several places along streams, August 4 to 5, 1918, i. e. the Madison River near the Park West Entrance, and the Cascades of the Firehole River.

(2) ***Hemerodromia femorata***, nov. sp. — Female. Length 3.5 mm. Testaceous, including the antennæ and the mouthparts, head and metanotum black, thoracic dorsum with a well defined central blackish vitta which includes the base of the scutellum, abdominal tergites except the last, brown, legs and halteres yellow. Occiput not shining, the lower white-pruinoose spot not continuing far along the orbits, front rather narrow, more than twice as long as broad, its lower part reddish, eyes narrowly but entirely separated on the face; third joint of the antennæ ovate, nearly twice as long as wide and three times as long as the style. Thorax very sparsely coated with gray dust, its hairs very sparse and minute, the vitta tapering anteriorly and then suddenly expanding over the humeri: ovipositor shining black, compressed, pointed, as long as the preceding segment. Front coxae very thin and elongate, nearly as long as the greatly thickened femora, the latter with a prominent spine-bearing tubercle near the base beneath, with strong biseriate black flexor setulae and with a posterior row of six heavy yellow setæ, front tibiae three-fourths as long as their femora, tipped with a spur and with a sharp black spine, biseriately black-spinulose beneath, last two joints of the posterior tarsi infuscated. Bases of the third and fifth veins heavy and brown, the second to the fifth sections of the costa proportioned 2.1 : 1 : 1 : 0.3, sections of the third vein, 4 : 1, of the fourth vein, 0.5 : 0.4 : 1 : 1.2, its posterior fork weak, sections of the fifth vein, 0.9 : 1.

One specimen; Alhajuelo, Canal Zone, Panama, March 4, 1912, collected by August Busck for the U. S. National Museum.

(3) ***Hemerodromia jugulator***, nov. sp. — Length 3 mm. Head, dorsum and upper side of the abdomen black, pleurae, venter, legs, antennæ, mouthparts and halteres pale yellow. Occiput dusky with brown, not shining, the lower orbits and the underside of the head densely white pilose, front oblong, more than twice as long as the distance between the posterior ocelli, eyes briefly touching on the lower part of the face; third antennal joint ovate, straight above, two-thirds longer than broad, with the style two-thirds as long as the third joint. Mesonotum very finely and sparsely pollinose so as to appear shining black, its setulae very sparse; first and last abdominal segments testaceous; pygidium long, erect, subshining, loosely hairy, with two pairs of dorsal forcipate valves and a projecting penis; ovipositor very short, compressed, shining, glabrous. Legs elongated, slender, front coxae greatly lengthened and very thin, nearly equal to their femora in length, the front femora comparatively slender. Veins brown, thin, the second, third and fourth sections of the costa proportioned 1.3 : 1 : 1, last sections of the third vein proportioned 2.8 : 1, of the fourth vein, 0.5 : 1 : 1, of the fifth vein, 0.9 : 1.

Four specimens; Cold Spring Harbor, Long Island, New York, July, 1913 (Melander).

(4) ***Hemerodromia nympha***, nov. sp. — Female. Length, including the ovipositor, 3 mm. Very slender, the head, thorax and abdomen black, subshining; antennæ, mouthparts and legs yellow. Front coxae very slender, five-sixths as long as their femora; the front femora without a basal tubercle, but with a spine, front tibiae two-thirds as long as their femora, with a sharp apical spine. Base of the very slender and elongate ovipositor yellowish. Halteres black, with a reddish root. Wings somewhat infumated, the second vein longer than usual, fork of the third vein beyond the end of the second vein, fork of the fourth vein as long as its pedicel, second to the fifth sections of the costa proportioned 1 : 0.3 : 0.7 : 0.15, sections of the fifth vein, 1 : 1.6, fringe of hind margin twice as long as the anterior crossvein.

One specimen, in the U. S. National Museum; Tjibodas, Mt. Gede, Java; from the Bryant and Palmer collection.

The species is noteworthy among the restricted genus *Hemerodromia*, in having an elongate ovipositor.

17. *H. oratoria*, Fallen, Empid. p. 11 (1815); ibidem, p. 34 [1816] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 63, pl. 23, f. 12 (1822); Macquart, Dipt. N. France, Vol. 3, p. 103, pl. 2, f. 7 (1827); Hist. Nat. Dipt. Vol. 1, p. 348 (1834); Zetterstedt, Fauna Ins. Lappon. p. 542 (1838); Dipt. Scand. Vol. 1, p. 262 (1842); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 144, pl. 5, f. 8g (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 145 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 (1862); Loew, Wien. Ent. Monatschr. Vol. 8, p. 244 (1864); Lundbeck, Dipt. Danica, Vol. 3, p. 236, f. 103 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 84 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 65 [1913] (*Microdromia*). — (Pl. 3, Fig. 26). var. *cataluna*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 178 (1909). Spain.
18. *H. orientalis*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 330, pl. 19, f. 25 [1911] (*Microdromia*). Java.
19. *H. raptoria*, Meigen, Syst. Beschr. Vol. 6, p. 341 (1830); Macquart, Hist. Nat. Dipt. Vol. 1, p. 348 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 264 (1842); ibidem, Vol. 8, p. 300 (1849); Pipping, Not. Sællsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 (1858); Zetterstedt, Dipt. Scand. Vol. 13, p. 4984 (1859); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 146 (1861); Loew, Wien. Ent. Monatschr. Vol. 8, p. 243 (1864); Lundbeck, Dipt. Danica, Vol. 3, p. 235, f. 101-102 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 84 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 65 [1913] (*Microdromia*). C. & N. Europe.
20. *H. rogoratoris*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 392 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 236 f. 65 [1902] (*Microdromia*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903). North America.
21. *H. simplicinervis*, nov. sp. (1). Philippine Islands.
22. *H. superstitionis*, Say, Long's Exped. Vol. 2, Append. p. 376 (1824); Wiedemann, Aussereur. Zweifl. Ins. Vol. 2, p. 11 (1830); Say, Compl. Writ. Vol. 1, p. 256 (1859); Osten-Sacken, Cat. N. Amer. Dipt. p. 242 (1878); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 391 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 236 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 [1903] (*Microdromia*). United States.
23. *H. unilineata*, Zetterstedt, Dipt. Scand. Vol. 1, p. 263 (1842); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 144 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 (1862); Loew, Wien. Ent. Monatschr. Vol. 8, p. 247 (1864); Wahlgren, Ent. Tidskr. Vol. 31, p. 84 (1910). C. & N. Europe.
24. *H. xiphias*, Bezzi, Suppl. Ent. Berlin, Vol. 3, p. 74 (1914). Formosa.

(1) ***Hemerodromia simplicinervis*, nov. sp.** — Male. Length 2 mm. Head and body black, legs entirely pale yellow. Head lightly dusted, eyes contiguous below antennæ, lower orbits and the horizontal face white-coated, base of antennæ yellow, last joint broken. Thorax lightly dusted, the metathorax becoming grayish, bristles and sparse setulæ yellowish. Upper valves of genitalia long, slender, forcipate, middle valves triangular. Front coxæ equal to their femora, denticles of front femora and tibiae black, a few setæ at base of front femora beneath, front tibiae tipped with a black spur. Halteres pale yellow. Wings yellowish at base, slightly smoky at middle and hyaline at apex, cuneiform, only two posterior veins, no trace of anal cell, fork of third vein forming angle of sixty degrees and ending midway between ends of second and third veins, first posterior cell widest at middle, its veins slightly convergent apically, sections of fifth vein proportioned 1 : 1.5.

Holotype : Mt. Makling, Luzon, Philippine Islands, received from C. F. Baker. This species is remarkable in the fusion of the second and third posterior cells, the fourth vein being simple and not forked. The contiguity of the eyes is also noteworthy.

2. GENUS COLABRIS, NOV. GEN.

Characters. — Eyes large, subcontiguous beneath the antennæ; but the lower facets not greatly enlarged; front very short; antennal style wanting. Thorax not tapering toward the front, the humeral grooves deep, in front of the scutellum the notum is slightly flattened but is not at all concave, scutellum strongly convex, about as large as the metanotum and devoid of bristles; pygidium small, closed, not forcipate; ovipositor very long and slender, ensiform. Front legs raptorial, the front coxae as robust as their femora, the latter weakly setose. Auxiliary vein straight, completely fusing with the costa beyond its base, no humeral crossvein, basal half of the pedicel of the second and third veins abruptly rudimentary, discal cell fused with the second basal, emitting three simple posterior veins, anal crossvein practically wanting. — it is faintly indicated however and is distinctly separated from the thickened hind margin — costa not abruptly attenuated beyond the fourth vein but continuing around the hind margin.

Type species : *C. rufescens*, nov. sp. The two following species, *rufescens* and *coxalis*, are the only known representatives.

Geographical distribution.

- | | |
|--|---------|
| 1. <i>C. rufescens</i> , nov. sp. (1). | Panama. |
| 2. <i>C. coxalis</i> , nov. sp. (2). | Panama. |

3. GENUS CHELIFERA, MACQUART

A. SUBGENUS CHELIFERA, MACQUART

Chelifera, Macquart, Mém. Soc. Sc. Lille, 1823, p. 150 (1823); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 123 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 247, 263 (1903).

Hemerodromia of authors, Loew, Wien. Ent. Monatschr. Vol. 8, p. 237-255 (1864); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 562 (1857); Williston, Trans. Ent. Soc. Lond. Vol. 3, p. 440 (1896); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 452 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 299 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 228 (1910).

Mantipeza, Rondani, Dipt. Ital. Vol. 1, p. 148 (1856); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 124 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389, 392 (1895); Williston, Trans. Ent. Soc. Lond. Vol. 3, p. 440 (1896); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 565 (1910).

(1) **Colabris rufescens**, nov. sp. — Length 2 2 mm. Reddish yellow, the head, metanotum and genitalia black, a diffused prescutellar spot dark brown, the lower mesopleuræ, scutellum and halteres light brown. Lower occiput bearing the usual densely white pruinose orbital mark, front no longer than the distance between the posterior ocelli, eyes very narrowly separated on the face, mouthparts yellow, palpi short; antennæ yellow, the third joint ovoid, one-half longer than broad, with a microscopic style. Thoracic setulæ microscopic, the two rows of acrostichals almost contiguous, notum and pleuræ not pollinose, rather shining. Front coxae but slightly shorter than their femora, of the male inflated, nearly as thick as their femora, of the female slender. Abdomen more or less fuscous, ovipositor nearly as long as the abdomen. Veins strong, brown, the second third and fourth sections of the costa proportioned 2 : 4 : 3.5, sections of the fifth vein proportioned 1 : 2 : 1.

Three specimens; Alhajuelo, Canal Zone, Panama, collected by August Busck. Type in the U. S. National Museum.

(2) **Colabris coxalis**, nov. sp. (PI. 3, FIG. 27). — Length 2 2 mm. Same as *rufescens*, but the front coxae are blackish and their hairs are black. Second, third and fourth sections of the costa proportioned 2 : 3.5 : 4.

Two specimens; Alhajuelo, Panama, March. Type in the U. S. National Museum.

Polydromia, Bigot, Ann. Soc. Ent. France (3), Vol. 557, 562 [1857] (*Polydromya*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 256 (1903); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 593 [1910] (*Polydromya*).

B. SUBGENUS CLADODROMIA, BEZZI

Cladodromia, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 453 (1905); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 299 (1909).

C. SUBGENUS METACHELA, COQUILLETT

Metachela, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253, 263 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 568 (1910).

Hemerodromia, s. str. Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 233 (1902).

D. SUBGENUS NEOPLASTA, COQUILLETT

Neoplasta, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 389, 392 (1905); Williston, Trans. Ent. Soc. Lond. Vol. 3, p. 440 (1896); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 254, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 299 (1909); Coquillett, Proc. U. S. Nat. Mus. Naturf. Halle, Vol. 37, p. 575 (1910).

Characters. — Eyes rather large, the lower facets larger than the upper but not expanding the eyes so as to bring them almost together on the face, as in *Hemerodromia*, front relatively longer, the sides of the face parallel; style one-third to one-twentieth as long as the third antennal joint, bare. Thorax typically gradually narrowed anteriorly, but often the thorax is quadrate, the humeri marked off by distinct grooves, a flattened or depressed space before the scutellum, the scutellum narrower than the metanotum, with two apical setæ placed close together. Front coxae thin, front femora usually strongly raptorial and setose in front and back. Costa continuing around entire wing-margin, auxiliary vein straight, well-separated from the costa, the humeral crossvein distinct, first vein ending at or beyond the middle of the wing, pedicel of the second and third veins distinct though rather short, anal cell complete, the anal vein separated from the hind margin.

The subgenera of *Chelifera* are characterized as follows :

Chelifera, Macquart, s. str. Discal cell complete, emitting two veins, the anterior of which is forked, thus the second posterior cell has a long petiole; second vein curving around a more or less visible stigma; thorax distinctly tapering anteriorly in surface view; no ovipositor; front femora strongly setose; antennal style very short and bare, one-tenth to one-fourth as long as the third joint of the antennæ.

Type species : *C. precatoria*, Fallen (Pl. 4, Fig. 28).

Cladodromia, Bezzi. Discal cell complete, emitting three veins, the second posterior cell sessile or pointed at its base, rarely with a short petiole; thorax opaque, apparently quadrate; ovipositor rather thick and long; front legs strongly setose; antennal style very short.

Type species : *C. inca*, Bezzi.

Metachela, Coquillett. Discal cell fused with the second basal, the anterior crossvein far before the posterior, second posterior cell with a long petiole; thorax not tapering in front; ovipositor ensiform; front femora setose; style one-third as long as the third antennal joint and thick; pygidium erect.

Type species : *Hemerodromia collusor*, Melander.

Thanategia, nov. subgen. Discal cell open outwardly, the anterior crossvein, much beyond the apparent posterior crossvein, second posterior cell with a very long pedicel; thorax not tapering anteriorly; ovipositor blunt not protracted; front femora strongly setose; antennal style microscopic and bare, pygidium large, terminal, the upper valves long, recurved and forcipate.

Type species : *Hemerodromia defecta*, Loew.

SUBGENUS CHELIFERA, MACQUART

Geographical distribution.

1. *C. * detestata*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 90, 103 ♀, pl. 6, Baltic Amber. f. 3-5 [1908] (*Hemerodromia*).
2. *C. Frigelii*, Zetterstedt, Ins. Fauna Lappon. p. 542 [1838] (*Hemerodromia*); N. Europe, Siberia. Dipt. Scand. Vol. 1, p. 261 [1842] (*Hemerodromia*); ibidem, Vol. 8, p. 3001 [1849] (*Hemerodromia*); ibidem, Vol. 13, p. 4983 [1859] (*Hemerodromia*); Becker, Acta Soc. Sc. Fenn. Helsingfors Vol. 26(9), p. 33 [1900] (*Hemerodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 84 [1910] (*Hemerodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 64 [1913] (*Hemerodromia*).
3. *C. melanocephala*, Haliday, Ent. Mag. London, Vol. 1, p. 158 [1833] N. & C. Europe. (*Hemerodromia*); Loew, Wien. Ent. Monatschr. Vol. 8, p. 238 [1864] (*Hemerodromia*); Mik, Wien. Ent. Zeit. Vol. 1, p. 39, pl. 1, f. 16-18 [1882] (*Hemerodromia*); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 95 [1892] (*Hemerodromia*); Lundbeck, Dipt. Dan. Vol. 3, p. 234, f. 100 [1910] (*Hemerodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 84 [1910] (*Hemerodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 65 [1913] (*Hemerodromia*).
flavella, Zetterstedt, Fauna Ins. Lappon. p. 543 [1838] (*Hemerodromia*); Dipt. Scand. p. 265 [1842] (*Hemerodromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 56 [1851] (*Hemerodromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 144 [1851] (*Hemerodromia*); Bonsdorff, Finl. tväv. Ins. Dipt. Vol. 1, p. 146 [1861] (*Hemerodromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 [1862] (*Hemerodromia*); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 133 [1887] (*Hemerodromia*).
4. *C. notata*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 209 : Cent. 2, No. 53 [1862] United States. (*Hemerodromia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 392 [1895] (*Mantipesa*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 238 [1902] (*Mantipesa*); Coquillett, Proc. Wash. Ent. Soc. Vol. 5, p. 264 (1903).
5. *C. nubecula*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 42 [1908] Canary Islands. (*Hemerodromia*).
6. *C. obsoleta*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 208 : Cent. 2, No. 52 [1862] United States. (*Hemerodromia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 392 [1895] (*Mantipesa*); Melander, Trans Amer. Ent. Soc. Vol. 28, p. 238 [1902] (*Mantipesa*); Coquillett, Proc. Wash. Ent. Soc. Vol. 5, p. 264 (1903).
7. *C. palloris*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 392 [1895] (*Mantipesa*); North America. Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 238 [1902] (*Mantipesa*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903); Slosson, Ent. News Philad. Vol. 14, p. 269 [1903] (*Hemerodromia*).
8. *C. precatoria*, Fallen, Empid. Suec. p. 10 [1815] (*Tachydromia*); Meigen, Europe, North America. Syst. Beschr. Vol. 3, p. 63, pl. 23, f. 13 [1822] (*Hemerodromia*); Macquart, Dipt. N. France, Vol. 3, p. 104, pl. 2, f. 8 [1827] (*Heme-*

rodromia; Hist. Nat. Dipt. Vol. 1, p. 347 [1834] (*Hemerodromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 266 [1842] (*Hemerodromia*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 505 [1849] (*Hemerodromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3001 [1849] (*Hemerodromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 56 [1851] (*Hemerodromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 143, pl. 5, f. 8 [1851] (*Hemerodromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 [1862] (*Hemerodromia*); Loew, Wien. Ent. Monaschr. Vol. 8, p. 238 [1864] (*Hemerodromia*); Mik, Wien. Ent. Zeit. Vol. 1, p. 39, pl. 1, f. 13, 15 [1882] (*Hemerodromia*); Mik. Beck Fauna Hernstein, Vol. 2 (2), p. 59 [1885] (*Hemerodromia*); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 95 [1892] (*Hemerodromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 238 [1902] (*Mantipeza*); Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 311 [1906] (*Hemerodromia*); Bezzii, Wien. Ent. Zeit. Vol. 26, p. 296 [1907] (*Hemerodromia*); Brocher, Ann. Biol. Lacustr. Bruxelles, Vol. 4, p. 44, 45 [1909] (*Hemerodromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 231, f. 98, 99 [1910] (*Hemerodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 84, f. 14 [1910] (*Hemerodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 64 [1913] (*Hemerodromia*).

— PI. 3, Fig. 28).

?*lineata*, Gimmerthal, Bull. Soc. Nat. Moscou, Vol. 9, p. 430 [1836] (*Hemerodromia*).

?*mantiformis*, Cuvier, Millim. Mag. Encycl. Vol. 1, p. 205, pl. 2, f. 3 [1795] (*Asilus*).

monostigma, Meigen, Syst. Beschr. Vol. 3, p. 62, pl. 23, f. 6 [1822] (*Hemerodromia*); Macquart, Dipt. N. France, Vol. 3, p. 104, pl. 3, f. 1 [1827] (*Hemerodromia*); Hist. Nat. Dipt. Vol. 1, p. 347 [1834] (*Hemerodromia*); Zetterstedt, Fauna Ins. Lappon. p. 543 [1838] (*Hemerodromia*); Dipt. Scand. Vol. 1, p. 267 [1842] (*Hemerodromia*); Boitard, Man. Ent. Vol. 3, p. 322 [1843] (*Hemerodromia*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 504 [1849] (*Hemerodromia*); Scholz, Zeit. Ent. Breslau, Vol. 5 (19), p. 56 [1851] (*Hemerodromia*); Walker, Ins. Brit. Vol. 1, p. 143 [1851] (*Hemerodromia*); Bonsdorff, Finl. tvåv. Dipt. Vol. 1, p. 146 [1861] (*Hemerodromia*); Glover, Manuscr. Notes, p. 25, pl. 6, f. 7 [1874] (*Hemerodromia*); Strobl, Glasnik. Zem. Mus. Bosn. Herceg. Sarajevo, Vol. 10, p. 12 [1898] (*Hemerodromia*); Mitteil. Nat. Ver. Steiermark, Graz, Vol. 34, p. 207 [1898] (*Hemerodromia*); Wissenschafts. Mitth. Bosn. Herceg. Vol. 7, p. 560 [1900] (*Hemerodromia*).

raptor, Macquart, Mém. Soc. Sc. Lille, 1823, p. 151, 165 (1823).

9. *C. stigmatica*, Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 [1862] (*Hemerodromia*); C. & S. Europe.

Mik, Beck Faun. Hernstein, Vol. 2 (2), p. 59 [1885] (*Hemerodromia*); Strobl, Wien. Ent. Zeit. Vol. 18, p. 77 [1899] (*Hemerodromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 233 [1910] (*Hemerodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 64 [1913] (*Hemerodromia*).

10. *C. trapezina*, Zetterstedt, Fauna Ins. Lappon. p. 543 [1838] (*Hemerodromia*); C. & N. Europe.

Dipt. Scand. Vol. 1, p. 264 [1842] (*Hemerodromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 56 [1851] (*Hemerodromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 83 [1862] (*Hemerodromia*); Strobl, Jahrb. Mus. Kärnten Klagenfurt, Vol. 47, p. 202 [1901] (*Hemerodromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 84 [1910] (*Hemerodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 64, f. 15 [1913] (*Hemerodromia*).

11. *C. valida*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 208 : Cent. 2, No. 51 [1862] (*Hemerodromia*); North America.

Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 392 [1895] (*Mantipeza*); Proc. Wash. Acad. Sc. Vol. 2, p. 423 [1900] (*Mantipeza*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 238 [1902] (*Mantipeza*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903).

SUBGENUS CLADODROMIA, BEZZI**Geographical distribution.**

1. *C. analis*, Thomsen, Eugen. Resa. Dipt. p. 474 [1869] (*Hemerodromia*); Patagonia.
Bezzi, Ann. Mus. Hungar. Vol. 3, p. 454 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 392 (1909).
2. *C. bicolor*, Philippi, Verh. Zool.-bot. Ges. Wien. Vol. 15, p. 767 [1865] Chile.
(*Hemerodromia*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 454 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 392 (1909).
3. *C. boliviensis*, Bezzi, ibidem, Vol. 91, p. 391, 396 (1909). Bolivia.
- + *C. cana*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 454 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391 (1909). Peru.
5. *C. flavipes*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 776 [1865] Chile,
(*Hemerodromia*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 453 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391 (1909).
6. *C. fuscimana*, Bezzi, ibidem, Vol. 91, p. 394 (1909). Peru.
7. *C. inca*, Bezzi, Ann. Mus. Hungar. Vol. 3, p. 454, f. 3 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391 (1909). Peru.
8. *C. nigrimana*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 766 [1865] Chile.
(*Hemerodromia*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 453 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391, 394 (1909).
9. *C. pallida*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 766 (1865); Chile.
Bezzi, Ann. Mus. Hungar. Vol. 3, p. 454, 459 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 390, 391 (1909).
10. *C. plurivittata*, Bezzi, ibidem, Vol. 91, p. 391, 395, f. 16 (1909). Peru.
11. *C. pratincola*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 766 [1865] Chile.
(*Hemerodromia*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 454 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 392 (1909).

SUBGENUS NEOPLASTA, COQUILLETT**Geographical distribution.**

1. *N. femoralis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391, 393 Peru.
(1909).
2. *N. mexicana*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 235, f. 56, 60 Mexico.
(1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903);
Bezzi, Ann. Mus. Hungar. Vol. 3, p. 460 (1905); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391 (1909).
3. *N. scapularis*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 209 : Cent. 2, No. 54 United States.
[1862] (*Hemerodromia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 392 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 234, f. 59 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 (1903); Slosson, Ent. News Philad. Vol. 14, p. 269 [1903] (*Hemerodromia*); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391 (1909).
4. *N. semilugens*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 766 [1865] Chile.
(*Hemerodromia*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 453 [1905]
(*Hemerodromia*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 391, 393 (1909).

SUBGENUS METACHELA, COQUILLETT**Geographical distribution.**

1. *M. albipes*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 505 [1849] (*Hemerodromia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 391 [1895] Hudson's Bay.

- (*Hemerodromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 237 [1902] (*Mantipeza*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 [1903] (*Chelifera*).
 2. *M. collusor*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 235, f. 57, 58, W. United States.
 64 [1902] (*Hemerodromia*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 [1903].
 3. *M. nigriventris*, Loew, Wien. Ent. Monatschr. Vol. 8, p. 242 [1864] (*Hemero-* C. Europe.
dromia); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 95 [1892] (*Lepidomyia*).

SUBGENUS THANATEGIA, NOV. SUBGEN.

Geographical distribution.

1. *T. defecta*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 210 : Cent. 2, No. 55 United States.
 [1862] (*Hemerodromia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 391 [1895] (*Hemerodromia*); Melander, Trans. Amer. Soc. Vol. 28, p. 235 [1902] (*Hemerodromia*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 264 [1903] (*Metachela*).

4. GENUS DRYMODROMIA, BECKER

Drymodromia, Becker, Ann. Soc. Ent. France, Vol. 83, p. 121 (1914).

Characters. — Slender, yellow species with full wings and raptorial front legs. Head orbicular, eyes large contiguous below the antennæ; proboscis short and perpendicular; third antennal joint short-conical, longer than the apical style. Thorax short, the prothoracic and supra-alar bristles distinct, otherwise bare. Abdomen elongate, with seven segments. Legs slender, front coxae long. Wings broad, immaculate, first vein attaining middle of wing, third vein forked, discal cell emitting three posterior veins, anal and basal cells nearly equally long, the anal cell perpendicularly closed.

But one species is known.

Geographical distribution.

1. *D. Jeanneli*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 121 (1914). East Africa.

5. GENUS CHELIPODA, MACQUART

A. SUBGENUS CHELIPODA, MACQUART

- Chelipoda**, Macquart, Mém. Soc. Sc. Lille, 1823, p. 148 (1823); Kertész, Cat. Dipt. Vol. 6, p. 117 (1909).
Chiromantis, Rondani (not Peters, 1854, *Reptilia*), Dipt. Ital. Vol. 1, p. 148 [1856] (*Chyromantis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 344 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 524 [1910] (*Chyromantis*).
Litanomyia, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 231 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 562 (1910).
Phyllodromia, of authors, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255, 260 (1903); Bezzi, Cat. Palæarct. Dipt. Vol. 2, p. 271 (1903); Kertész, Cat. Dipt. Vol. 6, p. 118 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 247 (1910); Engel, Deutsche Ent. Zeitschr. 1918, p. 3 (1918).

B. SUBGENUS PHYLLODROMIA, ZETTERSTEDT

- Phyllodromia**, Zetterstedt (not Serville, 1839, *Blattidae*); Zetterstedt, Isis, Vol. 1, p. 31 (1837); Fauna Ins. Lappon. p. 544, note (1838); Westwood, Gen. Syn. p. 132 (1840); Zetterstedt, Dipt. Scand.

Vol. 1, p. 269 (1842); Walker, Ins. Brit. Dipt. Vol. 1, p. 142 (1851); Rondani, Dipt. Ital. Vol. 1, p. 148 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 563 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 86 (1862); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 590 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45 (1910); Brunetti, Fauna Brit. Ind. Dipt. Vol. 1, p. 367 (1920).

Chelipoda, of authors, Mik, Wien. Ent. Zeit. Vol. 7, p. 299, 327 (1888); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 247, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 299 (1909); Kertész, Cat. Dipt. Vol. 6, p. 117 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 522 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 245 (1910); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 476 (1912).

Hemerodromia, of authors, Westwood, Gen. Syn. p. 132 (1840); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 550 (1910).

Lepidomyia, Bigot (not Loew, 1864, *Syrphidae*). Ann. Ent. Soc. France (3), Vol. 5, p. 557, 563 [1857] (*Lepidomyia*); Röder, Wien. Ent. Zeitschr. Vol. 6, p. 169 (1887); Mik, Wien. Ent. Zeitschr. Vol. 7, p. 299, 327 (1888); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 124 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 559 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45, 85 (1910); Engel, Deutsche Ent. Zeitschr. p. 3 (1918).

Thamnodromia, Mik, Wien. Ent. Zeit. Vol. 5, p. 278 (1886); Röder, Wien. Ent. Zeit. Vol. 6, p. 169 (1887); Mik, Wien. Ent. Zeit. Vol. 7, p. 299, 327 (1888); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 123 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 614 (1910).

Characters. — Slender, reddish yellow species with strongly raptorial front legs. Head globular, but a little longer in an obliquely downward direction, the upper occiput small, the front broad; eyes bare, large, with their lower forward facets somewhat enlarged, thus making the face and cheeks narrow. Occiput hairy below but with a single row of setæ above, the uppermost three pairs of which form the vertical bristles; ocelli large, one pair of strong ocellar bristles and usually behind them a pair of lesser ocellars; mouth-opening small and circular, proboscis much shorter than the height of the head, slender and slightly incurved, palpi very small, cylindrical and tipped with several fine hairs: antennæ plainly three-jointed, inserted close together and below the middle of the head, the last joint compressed conical, with a rather thick, long, two-jointed arista, the basal joint of which is minute. Thorax elongate, quadrate, narrower in front than the head, scarcely convex and in front of the small scutellum even flattened, coated with more or less evident gray pollen, humeri small, prothorax visible from above; macrochaetæ well developed, one strong posthumeral (possibly to be considered as the foremost dorsocentral), no humeral, two widely separated pairs of dorsocentrals, the anterior of which is sometimes reduced, two scutellars, one supraalar, no postalar, no notopleural, but sometimes some notopleural setulæ, only a single prescutellar pair of the acrostichal setulæ present: metapleuræ with two or three upward-directed setulæ, the mesosternum in profile angulate in the middle, prosternum ridged between the coxae. Abdomen more or less cylindrical, consisting of seven segments and the genitalia, male with a large to very large epipygium which comprises a strongly compressed, carinate lower part bearing a pair of forcipate valves directed forward over the abdomen; ovipositor very short, fleshy, with two styles. Front coxae slender and long, over twice as long as the posterior pairs, usually (*Chelipoda*, s. str.) with a basal spine directed forward, front femora incrassate, biserately spinulose beneath, and in front and back seriatel setose, front tibiae fitted to close against their femora, no terminal spine or tibial spurs, posterior legs simple, pulvilli minute. Wings margined behind by the prolongation of the costa, in front with a basal costal bristle and with costal hairs, no stigma, auxiliary vein straight, parallel with the costa and vanishing apically, first vein short, second vein long, third

vein not forked, pedicel of the second and third veins arising at the middle of the first basal cell, discal cell complete and with three posterior veins (*Chelipoda*) or open apically and the fourth vein forked (*Phyllodromia*), anal cell subequal to the second basal cell or somewhat shorter, the anal crossvein curved outwardly, anal vein separate from the hind margin of the wing : calypteres with a few cilia.

Nomenclature. **Type species :** by Macquart's original designation, *mantispa*, Macquart (not Panzer's) = *vocatoria*, Fallen (Pl. 3, Fig. 25). Coquillett (1903), would make *minor*, Macquart = *melanocephala*, Fabricius, the type, and in 1910 considered *Chelipoda* a synonym of *Hemero-dromia*. Rondani in 1856 designated *melanocephala*, Fabricius, as the type of *Phyllodromia* and *vocatoria*, Fallen as the type of *Chiromantis*. The next year Bigot founded *Lepidomyia* (not *Lepidomyia*, Loew (1864), a Syrphid, changed by Mik (1886), to *Lepidostola*), on the same *melanocephala*, as *mantispa*, Meigen. *Thamnodromia*, Mik, was a change of name for *Phyllodromia*, Zetterstedt (1837, not 1842), which was thought preoccupied by *Phyllodromia*, Serville (1839). The type of *Phyllodromia* thus becomes the type of *Thamnodromia*. *Litanomyia*, Melander, with type *mexicana*, Wheeler and Melander, by Coquillett's designation, is not generically distinct from *Chelipoda*.

The synonymy of these genera may be thus shown :

Subgenus *Chelipoda*, Macquart, s. str. Type : *vocatoria*, Fallen.

Chiromantis, Rondani. Type : *vocatoria*, Fallen.

Litanomyia, Melander. Type : *mexicana*, Wheeler and Melander.

Phyllodromia of the catalogues.

Subgenus *Phyllodromia*, Zetterstedt. Type : *melanocephala*, Fabricius.

Chelipoda of the catalogues.

Hemerodromia, Westwood and Coquillett, 1910. Type : *melanocephala*, Fabricius.

Lepidomyia, Bigot. Type : *melanocephala*, Fabricius.

Thamnodromia, Mik. Type : *melanocephala*, Fabricius.

Vocatoria possesses a complete discal cell, while in *melanocephala* the discal cell is outwardly open ; otherwise these species are structurally very much alike, certainly belonging to the same genus. Bezzi has recently described a curious form from Formosa, which in its reduced chætotaxy and enlarged anal cell departs markedly from the other species of *Chelipoda*. This species, *pictipennis*, has here been removed to the recently erected genus, *Cephalodromia*, Becker.

Geographical distribution. — Both of the subgenera occur in Europe and North America and *Chelipoda* occurs also in South America and Oceania.

SUBGENUS CHELIPODA, MACQUART

1. *C. albiseta*, Zetterstedt, Fauna Ins. Lappon. p. 544 [1838] (*Hemerodromia*) ; N. & C. Europe, North Dipt. Scand. Vol. 1, p. 271 [1842] (*Phyllodromia*) ; Scholz, Zeitschr. America.
Ent. Breslau, Vol. 5 (19), p. 57 [1851] (*Phyllodromia*) ; Schiner, Fauna Dipt. Austr. Vol. 1, p. 86 [1862] (*Phyllodromia*) ; Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 96 [1892] (*Thamnodromia*) ; Lundbeck, Dipt. Dan. Vol. 3, p. 250 [1910] (*Phyllodromia*) ; Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 68 [1913] (*Phyllodromia*).
vocatoria, Meigen, Syst. Beschr. Vol. 3, p. 65, pl. 23, f. 15 [1822] (*Hemerodromia*) ; Macquart, Dipt. N. France, Vol. 3, p. 103, pl. 2, f. 6 [1827] (*Hemerodromia*) ; Hist. Nat. Dipt. Vol. 1, p. 347 [1834] (*Hemerodromia*) ; Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 [1851] (*Phyllodromia*) ; Wahlgren, Ent. Tidskr. Vol. 31, p. 86 [1910] (*Phyllodromia*).
2. *C. * delicata*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 90, 109, pl. 6, f. 15 Baltic Amber.
[1908] (*Phyllodromia*).

3. *C. * dolosa*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 90, 104, pl. 6, f. 7-11 Baltic Amber.
[1908] (*Phyllodromia*).
 4. *C. elongata*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 232 [1902] United States.
(*Litanomyia*).
 5. *C. fascipennis*, Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. 72 [1914] (*Phyllo-* Java.
dromia).
 6. *C. fuscicornis*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 478 [1912] (*Phyllodromia*). Formosa.
 7. *C. fusciseta*, Bezzi, ibidem, Vol. 10, p. 477 [1912] (*Phyllodromia*). Formosa.
 8. *C. indica*, Brunetti, Rec. Indian. Mus. Vol. 9, p. 36 [1913] (*Litanomyia*). India.
Fauna Brit. India, Dipt. Vol. 1, p. 369, pl. 4, f. 16-18 [1920] (*Phyllo-*
dromia).
 9. *C. mexicana*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, p. 376 Mexico.
[1901] (*Sciodromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28,
p. 232, f. 63 [1902] (*Litanomyia*); Bezzi, Ann. Mus. Hungar. Vol. 3,
p. 460 [1905] (*Phyllodromia*).
 10. *C. nigricans*, nov. sp. (1). Java, Luzon.
 11. *C. parva*, nov. sp. (2). Java.
 12. *C. rhabdoptera*, nov. sp. (3). Costa Rica.
 13. *C. * rustica*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 90, 109, pl. 6, Baltic Amber.
f. 12-14 [1908] (*Phyllodromia*).
 14. *C. ? supplicatoria*, Meigen, Syst. Beschr. Vol. 3, p. 65 [1822] (*Hemerodromia*); C Europe.
Loew, Beschr. Posen. Gegend Art. Zweifi. Gatt. p. 22 (1840);
Isis, Vol. 7, p. 550 [1840] (*Hemerodromia*).
 15. *C. villata*, Arribalzaga, Natur. Argent. Buenos Aires, Vol. 1, p. 294 [1878] Argentina.
(*Hemerodromia*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 454, 459 [1905]
(*Phyllodromia*); Brethes An. Mus. Hist. Nat. Buenos Aires, Vol. 16,
p. 290 (1907).

(1) ***Chelipoda nigrans*, nov. sp.** — Male. Length 2.3 mm. Dark brown, almost black, unicolorous, including the antennæ, mouthparts, genitalia, stiff setæ of the front femora and the general tinge of the wings, legs brownish. Thorax dusky, not shining, the foremost dorsocentral, the supraalar and the intraalar bristles alone strong, the pair of scutellar bristles small; pygidium large, the width of the abdomen not diminishing apically, in outline triangular, terminating above in two forward-projecting, narrow, acuminate valves. Front coxae tubular, as long as their femora, which are sturdy and strongly setose, front tibiae three-fifths as long as their femora, slightly paler in color than the rest of the legs. Wings uniformly tinged with brown, second basal and anal cells of equal extent, discal cell complete, first vein ending before the middle of the wing, sections two to six of the costa proportioned 1 : 0.4 : 0.3 : 0.3 : 0.3, sections of the fourth vein, 1 : 8 : 6, of the fifth vein, 1 : 0.3.

Type in the U. S. National Museum, Tjibodas, Mt. Gede, Java, from the Bryant and Palmer collection. A specimen received from Prof. C. F. Baker, from Mt. Makling, Luzon, agrees with this species.

(2) ***Chelipoda parva*, nov. sp.** — Length 1.8 mm. Head black, dorsum reddish, pleurae and legs testaceous, abdomen brown. Base of antennæ black, remainder wanting; mouthparts yellowish. Mesonotum subshining, dusky, anterior dorsocentral bristle, i. e. the posthumeral, strong, middle and prescutellar dorsocentrals minute, notopleural, supraalar, and intraalar bristles only moderate. Pygidium moderately large, globular, slightly increasing the diameter of the abdomen, its valves open and shining, medially with a pair of erect hooks, the upper valve bluntly oval. Front coxae five-sixths, front tibiae three-fourths, as long as their femora, the latter bearing only two setæ on the postero-flexor edge, the femoral setæ yellowish. Halteres yellow, but with brown knob, calypteres blackish. Wings hyaline, veins brown, the second, third and fourth sections of the costa proportioned 1 : 0.4 : 0.3, sections of the fourth vein 1 : 7 : 9, of the fifth vein, 1 : 0.6, discal cell complete, second basal and anal cells coextensive.

Three specimens; Tjibodas, Mt. Gede, Java. Type in the U. S. National Museum.

(3) ***Chelipoda rhabdoptera*, nov. sp.** — Male. Length 2 mm. Body black, legs yellow, wings hyaline, pictured with a blackish gray band at end of marginal cell widening to hind margin and another faint cloud at middle of discal cell. Face linearly triangular, basal joints of antennæ yellow, third joint ovate black, arista two times antennal length and black; mouthparts yellow; head bristles strong and yellow, sides of mesonotum broadly pruinose, bristles brown, pleurae becoming brown at insertion of the yellow coxae. Pygidium greatly enlarged, reflexed, valves terminating in a thin curved process, penis long, thread-like, reflexed over the genitalia. Inner flexor side of the swollen front femora armed with five yellow setæ, outer flexor side with about seven small yellow setæ, under side minutely black-denticulate, hind tibia with expanded tip. Veins pale yellow in the hyaline portions of the wing, dark in the cloudings; halteres yellow.

Holotype, La Suiza de Turrialba, Costa Rica, April, 1922 (P. Schild), in Melander Collection.

16. *C. vocatoria*, Fallen, Empid. Suec. p. 12 [1815] (*Tachydromia*); Zetterstedt, Europe, United States.

Fauna Ins. Lappon. p. 544 [1838] (*Hemerodromia*); Dipt. Scand. Vol. 1, p. 270 [1842] (*Phyllodromia*); Mik, Wien. Ent. Zeit. Vol. 7, p. 327 [1888] (*Thamnodromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 345 [1902] (*Chiromantis*); Lundbeck, Dipt. Danica, Vol. 3, p. 249, f. 110 [1910] (*Phyllodromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 68 [1913] (*Phyllodromia*). — PI. 3, Fig. 25.
mantispa, Macquart (not Panzer), Mém. Soc. Sc. Lille, 1823, p. 149 (1823); Dipt. France, Vol. 3, p. 102, pl. 2, f. 5 [1827] (*Hemerodromia*); Hist. Nat. Dipt. Vol. 1, p. 349, pl. 8, f. 8 [1834] (*Hemerodromia*).

SUBGENUS PHYLLODROMIA, ZETTERSTEDT

Geographical distribution.

1. *P. flava*, Brunetti, Rec. Indian Mus. Vol. 9, p. 33 [1913] (*Chelipoda*); India.
 Fauna Brit. India, Dipt. Vol. 1, p. 368, f. 34, pl. 4, f. 14, 15 (1920).
2. *P. fusca*, Bezzi, Suppl. Ent. Berlin, Vol. 3, p. 73 [1914] (*Chelipoda*). Formosa.
3. *P. melanocephala*, Fabricius, Ent. Syst. Vol. 4, p. 407 [1794] (*Empis*); Syst. Antl. p. 144 [1805] (*Tachydromia*); Fallen, Emp. Suec. p. 12 [1815] (*Tachydromia*); Haliday, Ent. Mag. London, Vol. 1, p. 158 [1833] (*Hemerodromia*); Zetterstedt, Fauna Ins. Lappon. p. 544 [1838] (*Hemerodromia*); Dipt. Scand. Vol. 1, p. 269 (1842); Walker, Ins. Brit. Dipt. Vol. 1, p. 142, pl. 5, f. 7 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 146 (1861); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 379, 380 (1866); Loew, Ber. Naturh. Ver. Augsburg, Vol. 20, p. 46 (1869); Siebke, Dipt. Norv. Cat. p. 20 (1877); Neuhaus, Dipt. March. p. 74 (1886); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 96 [1892] (*Chelipoda*); Bezzi, Kat. Palæarct. Dipt. Vol. 2, p. 271 [1903] (*Lepidomyia*); Lundbeck, Dipt. Danica, Vol. 3, p. 6, 247, f. 108, 109 [1910] (*Chelipoda*); Wahlgren, Ent. Tidskr. Vol. 31, p. 85 [1910] (*Lepidomyia*); Dahl, Fauna Chorin. p. 466 [1912] (*Lepidomyia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 68 [1913] (*Chelipoda*).
mantispa, Panzer, Fauna Germ. Vol. 103, p. 16 [1806] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 64, pl. 23, f. 9 [1822] (*Hemerodromia*); Boitard, Man. Ent. Vol. 3, p. 322 [1843] (*Hemerodromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 57 [1857] (*Hemerodromia*); Mik, Wien. Ent. Zeit. Vol. 7, p. 327 [1888] (*Chelipoda*); Bezzi, Kat. Palæarct. Dipt. Vol. 2, p. 271 [1903] (*Lepidomyia*).
minor, Macquart, Mém. Soc. Sc. Lille, 1823, p. 148, 165 [1823] (*Chelipoda*).
obscuratoria, Walker, Ent. Mag. London, Vol. 4, 228 [1837] (*Hemerodromia*); List Dipt. Brit. Mus. Vol. 3, p. 505 [1849] (*Hemerodromia*).
raptor, Latreille, Gen. Crust. Ins. Vol. 1, pl. 16, f. 11, 12 [1806] (*Sicus*); ibidem, Vol. 4, p. 304 [1809] (*Sicus*), no description; Loew, Wien. Ent. Monatschr. Vol. 8, p. 248, note [1864] (*Hemerodromia*).
4. *P. *vaga*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 90, 106, pl. 6, f. 16, Baltic Amber.
 pl. 7, f. 1 [1908] (*Lepidomyia*).

6. GENUS CEPHALODROMIA, BECKER

Cephalodromia, Becker, Ann. Soc. Ent. France, Vol. 83, p. 121 (1914).

Characters. — Occiput drawn out horizontally; eyes approximate beneath the antennæ, the ocelli midway between the antennæ and the vertex; antennæ three jointed, the third joint elongate, furnished with fine hairs and with a long pubescent terminal arista; mouthparts small, the proboscis

slightly porrect; no vertical bristles, a single pair of small ocellar bristles, lower occiput bearded with white hairs. Thorax bare except for a single pair of strong supraalar bristles. Pygidium large, reflexed over the abdomen, bare and closed. Front coxae much longer than the posterior pairs, legs slender, front femora somewhat thickened. Wings cuneiform, no axillar lobe, costa continuing around the hind margin, hairy, humeral crossvein present, first vein short, second vein long, third vein not forked, ending at the tip of the wing, pedicel of the second and third veins arising from the middle of the first basal cell, discal cell open outwardly the fourth vein forked, anal cell large, in length slightly shorter than the second basal, but in breadth three times as wide as the second basal and extending to the hind margin of the wing, the anal vein fused with the fimbriate hind margin.

Type species : *C. curvata*, Becker. The preceding generic description has been furnished mainly by the detailed description of *C. pictipennis*, Bezzi. Bezzi provisionally placed his species in *Chelipoda*, from which it differs in the enlarged basal cells, the long level occiput, the lack of vertical and discal bristles and the bifasciate wings. The genotype has the wings uniformly yellowish.

Geographical distribution.

1. *C. curvata*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 121 (1914). E. Africa.
2. *C. pictipennis*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 476, f. 5 [1912] (*Cheli-* Formosa. *poda*); Suppl. Ent. Beschr. Vol. 3, p. 73 [1914] (*Chelipoda*).

7. GENUS PTILOPHYLLODROMIA, BEZZI

Ptilophyllodromia, Bezzi, Ann. Mus. Nat. Hungar. Vol. 2, p. 344 (1904).

Characters. — Elongate, opaque black, cinereous pruinose species with yellow legs, the front coxae elongate, the front femora raptorial, the antennae lengthened and with plumose arista, the neuration quite as in *Chelipoda*, and the anterior legs of the male deformed. Head round, eyes broadly separated above, two ocellar and two vertical bristles which are robust but short; face narrower than the front; proboscis short, palpi small and yellow; antennae greatly lengthened, first joint long, obconic, three times the length of the spherical second joint, third joint pyriform, as long as the first joint, bearing a thick apical arista nearly as long as the antenna and which is heavily plumose below. Thorax a little longer than the abdomen, front legs placed up toward the head, distant from the posterior pairs which are close together, scutellum very small, metanotum well developed; one humeral, one prescutellar dorsocentral, one supraalar and two small scutellar bristles present; metapleuræ with three or four hairs. Abdomen compressed, with short hairs, pygidium strongly compressed, upper lamellæ rounded. Front coxae subequal to the femora in length, armed with black subapical tubercle and with a row of setæ, front femora incrassate, raptorial, denticulate and setose beneath and with cilia above, posterior femora without setæ, middle tibiae incurved, apically dilated and with long hairs and subapical pencil on the extensor side, middle metatarsi tuberculate above, tarsal claws small. Wings cuneate, no costal bristle, no stigma, third vein not furcate, discal cell complete, posterior cells sessile, anal cell rounded apically and as long as the second basal. According to Dr. Bezzi's figure the basal half of the costa bears outstanding hairs and the vein in front of the anal cell is setose.

The type species, *P. Biroi*, is the only known form of this remarkable genus. It was collected in New South Wales, Australia.

Geographical distribution.

1. *P. Biroi*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 345, f. 5 (1904). Australia.

SUBFAMILY TACHYDROMIINÆ

Characters. — Small species in which the neuration is greatly reduced. Head globular; eyes generally separated above the antennæ, but in *Symbalophthalmus* there contiguous, often more or less approached below the antennæ in both sexes, in which case the lower facets may be larger, often pubescent; antennæ short, two- or three-jointed, the arista one- or two-jointed, usually hairy, generally apical but sometimes subdorsal; proboscis short, incurved or perpendicular, in the latter case more slender, in the former case strong at the base and pointed, no labella, palpi one-jointed, incumbent, often broad; vertical bristles usually well developed.

Thorax stout and short as compared with the abdomen, often pubescent and with discal bristles, metapleuræ bare; pygidium asymmetrical, with an unpaired upper valve. Legs sometimes slender, as in *Tachydromia*, sometimes stout, as in *Platypalpus*, coxæ not long, middle legs sometimes raptorial and armed, femora often thickened. Wings with reduced anal angle, costa continuing to the fourth vein, auxiliary vein always rudimentary or imperfect, third vein unforked, no discal cell, that is, the discal cell is fused with the second basal, basal cells large, the first and second basals equal or the second longer, three posterior cells, anal cell weak or wanting, the anal vein incomplete even when best developed but most often wanting, anal crossvein if present perpendicular, when present the anal cell is shorter than the basal cells, usually no stigma, no alula. Calypteres closely united to the base of the wing and possessing a small fringe.

TABLE OF THE GENERA AND SUBGENERA OF THE TACHYDROMIINÆ

1. Thorax slender, much longer than broad, the humeri large, strongly constricted; proboscis vertical and slender, palpi usually narrow; legs slender, cursorial and not bristly, front femora usually thickened	2.
Thorax broader, the humeri rarely large; legs hairy and usually with bristles or setæ; proboscis usually inflexed and stronger at the base, palpi usually quite broad	6.
2. Wings and halteres present, marginal cell long	3.
Wings and halteres entirely lacking	Genus PIELTAINIA, Arias.
3. First basal cell evidently shorter than the second; black species; front femora thickened	4.
First basal cell slightly longer than the second; only the outer angle of the anal cell present; yellow species with black head; last tarsal joint broadened, femora slender	Genus DYSALETRIA, Loew.
4. Anal crossvein present; proboscis longer than the palpi; third antennal joint oval, with apical arista; front narrow, its sides parallel; thorax elongate, narrowed anteriorly, mesonotum more or less pruinose; wings more or less nebulous (Pl. 4, Fig. 31). Anal crossvein completely wanting; proboscis often shorter than the palpi	Genus TACHYPEZA, Meigen.
5. Front narrow, its sides straight and nearly parallel; palpi narrow and elongate, but usually shorter than the proboscis, subshining and usually with an apical seta; face linear; antennal emargination of the eyes small, lower facets but slightly larger than the	5.

- upper; wings pictured usually with crossbands, basal cells long, the posterior crossvein located near the middle of the wing, marginal cell long, the second section of the costa usually much longer than the third section; polished species with the thorax narrowed in front (Pl. 4, Fig. 29). Genus *TACHYDROMIA*, Meigen.
- Front relatively broad and V-shaped, its sides bowed outwardly; palpi oval or elongate oval, white, densely hairy, rarely with an apical seta, longer than the short proboscis; eyes contiguous below the antennae obliterating the face, the lower facets large, the antennal emargination relatively deep; wings nearly or quite hyaline, basal cells short, the posterior crossvein near the basal third of the wing, second and third sections of the costa usually subequal; species sometimes pollinose and with quadrate thorax (Pl. 4, Fig. 35) Genus *TACHYEMPIS*, nov. gen.
6. First basal cell equal to or longer than the second, anal cell not formed; opaque pollinose species; eyes widely separated on face and front; two pairs each of ocellar and vertical bristles 7.
- First basal cell shorter than the second, if the basal cells are subequal the eyes are close together either above or beneath the antennae or the middle tibiae are spurred; abdominal pits, especially the dorsal ones, rarely visible; one pair of diverging ocellar bristles and usually one pair of verticals II.
7. Arista apical; humeri constricted; legs not evidently bristly; cheeks linear, face narrower than the front; ocellar bristles diverging; notum with coarse hairs but no strong anterior dorsocentrals; abdominal segments without dorsal pits. Genus *CHARADRODROMIA*, nov. gen.
- Arista subapical or dorsal; humeri not constricted; legs bristly; cheeks at least one-fourth the eye-height, face at least as broad as the front; abdomen with dorsal as well as lateral pittings. 8.
8. Mesonotum with discal bristles; last joint of the antenna bare and with a subapical bare arista; eyes bare; lower ocellar bristles converging. 9.
- No discal bristles on the thorax; third antennal joint pubescent and bearing a dorsal pubescent arista; eyes pubescent; lower ocellar bristles diverging. Genus *HALSANALOTES*, Becker.
9. Wings as long as or surpassing the abdomen, their hind margin ciliate. 10.
- Wings aborted, about one-third as long as the abdomen and bare on their hind margin; arista dorsal (Pl. 7, Fig. 71). Genus *THINODROMIA*, Melander.
10. Cheeks one-third the eye-height; first vein long, ending at two-thirds the length of the wing, first section of the costa not ciliate; femora, especially the hind pair, with numerous strong bristles; two presutural dorsocentrals present; antennae plainly three-jointed; pulvilli large (Pl. 7, Fig. 70). Genus *COLOBONEURA*, Melander.
- Cheeks narrower, usually one-sixth the eye-height; first vein ending near the middle of the wing, the first section of the costa usually

- ciliate with long hairs; femora not remarkably bristly; one presutural dorsocentral; antennæ apparently two-jointed; pulvilli small
- Genus CHERSODROMIA, Walker.
11. The two basal cells of the same length; eyes nearly or quite contiguous in part; legs slender and simple, neither the middle nor the hind tibiae spurred; discal bristles of thorax rarely present
- First basal cell evidently shorter than the second; eyes separated at least on the front: if the basal cells are equal the middle femora and tibiae are seriatelose beneath
12. Anal cell completely wanting; sides of the front diverging above; basal cells short; third joint of the antennæ with a dorsal arista; one pair of converging preocellar bristles; arista geniculate (PI. 4, Fig. 32)
- Genus MICREMPIS, nov. gen.
- Anal vein and sometimes anal crossvein present; basal cells long; last segments of the female abdomen lengthened; third joint of the antennæ lanceolate and with a terminal arista
13. Anal crossvein present, wings slender, the anal angle much reduced; eyes bare; palpi minute; eyes contiguous on the front
- Anal crossvein absent, wings broad, the anal angle large; eyes pubescent; front and face linear; antennæ two-jointed, arista thickened; scutellum large and flat (PI. 4, Fig. 33)
- Genus SYMBALLOPHTHALMUS, Becker.
14. Anal cell more or less formed; sides of the front parallel or slightly diverging; arista terminal; middle femora strongest and with two rows of flexor denticles or setulae; middle tibiae with one row of flexor denticles and ending in a more or less sharp spur; pleurae usually largely pollinose; thorax longer than broad
- Genus PLATYPALPUS, Macquart. 15
- Anal cell completely wanting; front femora stronger than the middle femora, the middle femora rarely armed; thorax scarcely or not longer than broad
16. Last tarsal joint short and flattened (PI. 4, Fig. 30)
- Last joint of the front and middle tarsi greatly lengthened
- Subgenus PLATYPALPUS, Macquart.
- Subgenus CLEPTODROMIA, Corti.
16. Arista dorsal, the antennæ apparently two-jointed, the last joint minute; eyes contiguous below the antennæ, sides of the front usually parallel (PI. 4, Fig. 37)
- Genus STILPON, Loew.
- Arista terminal or subterminal, antennæ three-jointed; eyes more or less separated below the antennæ, sides of the front usually diverging above
- Genus DRAPESTIS, Meigen. 17.
17. More or less yellow species; first section of the costa with longer cilia than on the following sections; legs slender; antennæ horizontal, the second joint without a seta beneath; hind tibiae with an extensor bristle and apical spur
- Black species; costa uniformly short-ciliate; antennæ directed obliquely upward
18. Third antennal joint lanceolate, the arista relatively short; middle segments of abdomen laterally with flattened setulae; wings oval. Subgenus ELAPHROPEZA, Macquart.

- Third antennal joint not lengthened; middle segments of the abdomen with hair-like setulae; wings narrow, the anal angle greatly reduced; costa with long cilia (Pl. 4, Fig. 36) . . .* Subgenus *CTENODRAPETIS*, Bezzi.
19. *Legs without strong bristles, no tibial spur; second joint of antennae without a seta or with but a short one; notum densely covered with appressed pubescence, no humeral or discal bristles present.* Subgenus *DRAPETIS*, Meigen.
Legs with bristles and the hind tibiae with an evident terminal spur; second joint of antennae with a strong seta beneath; pubescence of thorax more erect, humeral and sometimes discal bristles present (Pl. 4, Fig. 34): Subgenus *EUDRAPETIS*, Melander.

I. GENUS TACHYPEZA, MEIGEN

Tachypeza, Meigen, Syst. Beschr. Vol. 6, p. 341 (1830); ibidem, Vol. 7, p. 94 (1838); Zetterstedt, Fauna Ins. Lappon. p. 345 (1838); Dipt. Scand. Vol. 1, p. 312 (1842); Rondani, Dipt. Ital. Vol. 1, p. 147 [1856] (*Tachypeza*); Bigot, Ann. Soc. France (3), Vol. 5, p. 563 (1857); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 15 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 124 (1889); Williston, Man. N. Amer. Dipt. p. 75 (1896); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 42 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 222 (1908); Kertész, Cat. Dipt. Vol. 6, p. 136 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 611 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 262 (1910); Melander, Psyche, Vol. 17, p. 40 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45, 87 (1910); Brunetti, Fauna Brit. India, Dipt. Vol. 1, p. 378 (1920).

Characters. — Slender, moderate-sized species of black coloration, with usually nebulous wings and nearly devoid of hairs and bustles. Head higher than broad, the eyes large, bare, in both sexes meeting beneath the antennae particularly along the middle of the face, the facets of nearly uniform size, the hinder margin of the eye with an anterior curve, but rather angulate at the bottom; front quite narrow, its sides nearly parallel, but bowed about the ocellar protuberance which is placed rather far forward and bears three large ocelli but no bristles; occiput conical, with several short stiff black bristles above and numerous white hairs below; cheeks linear. Antennae two-jointed, without bristles, the outer joint ovate, slightly pubescent, with a long, bare, one-jointed, apical arista. Proboscis slender, vertical, about one-half as long as the head-height, the labium trilobed at the tip; palpi narrow, one-half as long as the proboscis, tipped with several strong bristles. Thorax oblong, tapering in front of the wings, flattened, the humeri large and marked off by coarse grooves; no humeral bristle, one or two short stiff notopleural, two rows of microscopic acrostichal setulae, dorsocentrals uniserial, microscopic except the prescutellar one which is usually stiff and large, postalar callus strong but without bristle, two straight stiff scutellar bristles; pleurae shining or pollinose, without bristles; abdomen comprising eight segments, the first ventral not chitinized, in the female tipped with two slender styles, in the male the pygidium is terminal, globular below. Front legs with rather large coxae and swollen femora, hind legs long and slender, anterior femora and tibiae biseriately setulose beneath, middle femora with a basal flexor bristle sometimes borne on a tubercle, no tibial spurs, spines or bristles, pulvilli small. Wings slender, elongate, no alula, costa extending to the fourth vein, somewhat thickened at the termination of the first vein, its hairs very short, costal bristle minute, cilia of anal angle longer and looser; basal cells long, posterior crossvein perpendicular or oblique, located just before the middle of the wings and one or two times its length beyond the anterior crossvein, first vein ending beyond the middle of the wing,

marginal cell long, first posterior cell narrow, with parallel veins; anal crossvein strong but the anal vein wanting; calypteres fringed.

Type species: *T. nubila*, Meigen, by designation of Rondani in 1856. Meigen in 1822 separated his species of *Tachydromia* into two groups, calling them A and B. In 1827 Macquart gave the name *Platypalpus* to group B, and in 1830 Meigen called his group A *Tachypeza*. It is from group A that Loew in 1863 separated his genus *Tachista*, which, containing the genotype of *Tachydromia* has become a synonym of that genus.

The species of *Tachypeza* are often found running up and down the trunks of trees, especially such as have smooth bark. They carry their wings held against the abdomen and are loathe to take to flight. They can easily be collected by blowing them with a puff into the insect net. The earlier stages are not known with certainty.

KEY TO THE NEARCTIC SPECIES OF TACHYPEZA

1. Pleuræ opaque. 2.
- Mesopleuræ largely or wholly shining, devoid of pollen 6.
2. Crossveins separated by three times their length; front femora dark
 exteriorly and yellow interiorly, middle femora with basal tubercle *T. DISTANS*, nov. sp. (1).
- Crossveins separated by their own length 3.
3. Anterior femora ♂♀ yellow inside and dark outside 4.
- Front femora ♂ with black marks inside; crossveins not more smoky
 than longitudinal veins 5.
4. Crossveins more smoky than veins; hind femora blackish; middle
 femora brownish, in ♂ with basal tooth, tibiæ with blackened apex,
 middle metatarsi ♂ flat. *T. DISCIFERA*, nov. sp. (2).

(1) ***Tachypeza distans*, nov. sp.** — Male. Length 3.2 mm. Antennæ reddish yellow, tipped with brown; mouth parts yellowish, apical seta of palpi pale brown. Thorax entirely pollinose. Abdomen subshining. Legs mainly yellow, marked with brown as follows, outside of front femora and tibiæ, subbasal and apical bands of middle tibiæ, apical half of hind femora except above, base and apex of hind tibiæ and last joint of all tarsi; front femora spongy pubescent below, flexor side of front tibiæ spongy pubescent in front and seriatel black-setulose behind; middle femora with strong basal setigerous tubercle, near which are several yellow setæ. Halteres yellow. Wings hyaline, first vein thickened, first posterior cell coarctate, as wide at middle as the marginal and submarginal cells together, hind crossvein remote from the anterior and very oblique, the distance between the two more than three times the length of either, last section of fifth vein two-thirds the length of the preceding.

Holotype, Kamiac Butte, Washington (Melander).

(2) ***Tachypeza discifera*, nov. sp. (PI. 4, FIG. 31).** — Length 3 mm. Head and thorax uniformly opaque cinereous; sides of front parallel; basal antennal joint yellow, the outer joint brown, rounded ovate, with terminal arista nearly four times the length of the antennæ: palpi pale yellow, with sparse white hairs and with a single black bristle near the end; proboscis robust, vertical, nearly as long as the height of the head, black at base and apex. No humeral bristles, but the usual dorsocentral, scutellar and notopleural pairs present. Abdomen subshining, hypopygium small, not noticeably hairy, the last ventral segment fringed. Legs dusky yellow, the hind femora, the upper side of the middle femora, the tips of all the tibiae and the end of the tarsi blackish, posterior tibiæ with a broad darker ring toward the base; front femora ♂ white-ciliate beneath but not marked with black, front tibiæ scarcely thickened, the inside closely setulose, middle femora loosely setulose within, and in the ♂ near the base with a strong mamillate tubercle bearing a pair of yellow spines, middle tibiæ closely setulose within, and in the ♂ flatly excised toward the end, the middle metatarsi of the ♂ broadly flattened so as to be one-fifth as broad as long, and densely setulose on the edges. Halteres entirely yellowish. Outer two-thirds of the wings lightly infumated and with blackish veins, root of the fifth vein pale, third section of the costa about three times as long as the fourth section, the fourth vein bent forward near the tip, crossveins subequal, their separation less than their length, last section of the fifth vein three-fourths as long as the preceding section.

Numerous specimens from Powell Co, Montana, July, collected by Wm. M. Mann: one specimen from Moscow, Idaho, taken by Dr. J. M. Aldrich.

Crossveins smoky; outer side of anterior femora with central clearer space in the dark area (variation with front femora nearly black); hind femora and tibiae with middle pale ring	T. INUSTA, Melander.
5. Front femora ♂ with two round black marks inside; middle femora not tuberculate, hind femora dark; front tibiae closely setulose inside; second antennal joint black	T. CORTICALIS, Melander.
Front femora ♂ with one large and one or two small black spots inside; middle femora ♂ with tubercle at base beneath, hind femora with middle pale ring; front tibiae not setulose; antennae red	T. ANNULARIS, nov. sp. (1)
6. Palpi, halteres, legs and wings blackish	T. WINTHEMI, Zetterstedt.
Palpi, halteres, more or less of legs, and at least base of wings paler	7.
7. Second basal cell but little longer than the first, its crossvein perpendicular; proboscis reddish at base	8.
Second basal cell longer than the first by at least the length of the posterior crossvein; proboscis mostly or wholly black	12.
8. Front coxae and femora ♂ marked with black, front femora ciliate beneath with black hairs; sides of front ♂ parallel; veins strong ♂♀; hypopygium with black hairs above.	T. BRACHIALIS, Melander.
Front coxae and inside of front femora not or inconspicuously marked; veins not strong	9.
9. Front tibiae ♂ incrassate apically, ♂ legs largely blackish, ♀ legs more yellow, hind tibiae tipped with black.	T. CLAVIPES, Loew.
Front tibiae not especially incrassate	10.
10. Front femora ♂ not ciliated; wings dark beyond the base; hypopygium large; hind knees paler.	11.
Front femora ciliate beneath with pale hairs; wings uniformly gray; tibiae of uniform color	T. POSTICA, Walker.
11. Middle tibiae ♂ with a strong excision before the apex; palpi brown; hypopygium hairy; outer side of front femora with a black streak, two-fifths of the hind femora yellow; third section of the costa nearly four times the fourth.	T. EXCISA, nov. sp. (2).

(1) *Tachypeza annularis*, nov. sp. — Length 3 mm. Occiput opaque cinereous, with six black bristles above and numerous white ones below; antennae red, the outer joint rounded oval, the subterminal arista four times the length of the antennae; palpi white, with white hairs and two terminal whitish bristles; proboscis reddish, with dark tip. Notum and pleurae uniformly opaque cinereous, no humeral bristles. Abdomen opaque piceous, the shining black hypopygium relatively small, its upper lamella with a few short hairs. Legs moderately strong and mostly yellow, the front femora ♂ with a subapical black spot and a central round black dot on the flexor side, and with another black spot on the inside proximal to the middle, these three spots wanting in the ♀, the outside of the front femora with a preapical black spot, more or less prolonged below in the ♂, the hind femora and tibiae with the basal and apical thirds black, middle tibiae tipped with black and with a suffused dark band below the knee, front femora not ciliate or setulose, middle femora black-setulose beneath and in the ♂ with a strong basal tubercle bearing several pale bristles, the front tibiae scarcely thickened and not setulose, the middle tibiae closely setulose beneath and in the ♂ deeply excised near the tip. Halteres entirely pale yellow; calypteres with nine pale hairs. Wings hyaline, a little infumated at the middle, veins narrow, especially apically, the root of the fifth vein pale, the fourth vein gently curved forward, the third section of the costa scarcely three times the fourth section, posterior crossvein longer than the anterior, the distance between them equal to the anterior crossvein, the last section of the fifth vein three-fourths as long as the preceding section.

Fieldbrook and Santa Cruz Mountains, California (U. S. N. M.); Mount Constitution, Washington, 17 July, 1909.

(2) *Tachypeza excisa*, nov. sp. — Male. Length 4 mm. Occiput pollinose, cinereous below, above with four stiff proclinate black bristles, the lower occiput with numerous white hairs; front cinereous, moderately narrow, widest at the ocelli, its lower sides parallel, ocellar bristles minute; antennae yellow, the third joint lanceolate, the terminal brown

- Middle tibiæ ♂ slightly excised; palpi yellow; hypopygium without long hair; hind knees narrowly yellow; third section of costa two to three times the fourth section T. ROSTRATA, Loew.
12. Pruinosity of humeri encroaching on mesopleuræ; femora (♀) not striped with black, hind tibiæ blackish; antennæ red T. PRUINOSA, Coquillett.
- Pleuræ not pruinose beneath the notopleural suture 13.
13. Front femora ♂ marked with black, hind tibiæ yellow with black tip; antennæ yellow 14.
- Legs uniformly blackish, not marked with black spots; antennæ blackish T. DOLOROSA, nov. sp. (1).
14. Front trochanters and coxæ ♂ with black spot, front femora ♂ with a black line beneath; wings smoky except at base T. FENESTRATA, Say.
- Coxæ and trochanters not marked, front femora ♂ with two black spots; fifth vein strong even at base. 15.
15. Middle femora ♂ tuberculate at base; apex of posterior tibiæ and underside of hind femora black; veins smoky in middle of wing, the outer third of wing not dark; humeri pollinose T. BINOTATA, nov. sp. (2).

arista two and one-half times the length of the antennæ : palpi brown, with pale hairs and with a terminal black bristle ; proboscis yellow, its tip black. Mesonotum and humeri subshining through the scabrous coating which extends over the notopleural suture near the root of the wing and thinly covers the posterior half of the pleura ; the usual pair of stiff dorsocentral and scutellar bristles present. Abdomen opaque except the large globose hypopygium, whose upper appendages are furnished with dense curved hairs. Legs robust, the front femora especially so, yellowish, but with a blackish streak on the outer side of the anterior femora, also the middle part of the hind femora and both sides of all the tibiae blackish ; front femora not ciliate nor noticeably setulose, the flexor surface of the middle femora and tibiæ strongly setulose, of the hind femora less strongly so ; several long stiff black bristles bunched at the base of the middle femora, matching a pronounced excision near the tip of the tibiæ ; front tibiæ scarcely thickened. Halteres yellowish. Wings subhyaline, a little infumated on the costal half, veins blackish but not clouded, a little paler toward the base, the third and fourth veins converging, the fourth sections of the costa about one-fourth as long as the third section, the crossveins and the space between them all subequal, the outer sections of the fifth vein equal.

A single male collected by Nathan Banks at Ithaca, New York, July 7.

(1) *Tachypeza dolorosa*, nov. sp. — Length 3 mm. Front and occiput gray pruinose, the upper occipital bristles black, the lower ones white; no ocellar bristles : antennæ black, the outer joint ovoid, with subterminal arista, which is nearly four times as long as the antennæ ; palpi pure white and with white hairs ; proboscis black. Mesonotum and humeri except their lower edge opaque pollinose, the usual small humeral, strong notopleural, single pair of posterior dorsocentral and two scutellar bristles present; pleurae shining black except for a thin coating of pollen near the middle coxae. Abdomen subshining, the hypopygium moderate, its upper lamellæ sparsely hairy, the hairs of the last ventral rather long. Legs black, the inside of the front femora and the base of the tarsi brown, no spottings or peculiar armature in the ♂ except for a slight emargination near the tip of the middle tibia and a corresponding pair of setulæ at the base of the middle femora. Halteres white ; calypteres with a short fringe of about ten white hairs. Wings long and narrow, smoky except at the base, veins black, the fifth vein dark even at its root, third and fourth veins slightly converging, the third section of the costa three times as long as the fourth, the posterior crossvein about two-thirds the length of the oblique anterior crossvein, the distance between them nearly two times the length of the posterior crossvein, last section of the fifth vein three-fourths the length of the preceding section.

Cloudcroft, New Mexico. This is the species mentioned in the Transactions of the American Entomological Society, vol. 28, p. 343, as *rapax*, occurring in company with *corticalis*. This last species is common in the West. I have seen it from New Mexico, California, Washington, Idaho, Montana, Wyoming and British Columbia.

(2) *Tachypeza binotata*, nov. sp. — Male. Length 3.3 mm. Occiput cinereous, with four black bristles above and numerous white hairs below, the front cinereous, no ocellar bristles; antennæ yellowish, the outer joint oval, the terminal arista nearly four times the length of the antennæ : palpi white and white-hairy; proboscis yellow except the very tip. Mesonotum and humeri opaque cinereous, pleurae entirely polished black. Abdomen shining piceous black, the hypopygium large, globular, jet-black, its upper appendages nearly bare, the last ventral segment with a loose fringe of long brown hairs. Legs rather robust, including the coxae largely yellow, the front femora with two large subquadrate black spots below, middle tibiæ tipped with black, the basal two-thirds of the hind femora, the base and apex of the hind tibiæ and much of all the tarsi brown, femora not ciliate and scarcely setulose, the middle femora with a pronounced spine-tipped tubercle at the

Middle femora not tuberculate, legs yellow except for the two femoral spots; wings smoky except base: humeri shining *T. humeralis*, nov. sp. (1).

Geographical distribution.

- | | |
|--|-------------------|
| 1. <i>T. annularis</i> , nov. sp. | California. |
| 2. <i>T. binotata</i> , nov. sp. | Washington. |
| 3. <i>T. brachialis</i> , Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 343 [1902] (<i>Tachydromia</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903). | E. United States. |
| 4. <i>T. clavipes</i> , Loew, Berl. Ent. Zeitschr. Vol. 8, p. 86 : Cent. 5, No. 73 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 [1895] (<i>Tachydromia</i>); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 229 [1902] (<i>Tachydromia</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903). | North America. |
| 5. <i>T. corticalis</i> , Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 343 [1902] (<i>Tachydromia</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903).
? <i>portacula</i> , Walker, List Dipt. Brit. Mus. Vol. 3, p. 506 [1849] (<i>Tachydromia</i>); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 [1895] (<i>Tachydromia</i>); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 227 [1902] (<i>Tachydromia</i>). | W. United States. |
| 6. <i>T. costalis</i> , Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 53 [1840] (<i>Tachydromia</i>). | C. Europe. |
| 7. <i>T. discifera</i> , nov. sp. — PI. 4, Fig. 31. | W. United States. |
| 8. <i>T. distans</i> , nov. sp. | Washington. |
| 9. <i>T. dolorosa</i> , nov. sp. | New Mexico. |
| 10. <i>T. *egelata</i> , Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 89, 98, pl. 4, f. 8-10 [1908] (<i>Tachydromia</i>). | Baltic Amber. |
| 11. <i>T. excisa</i> , nov. sp. | New York. |
| 12. <i>T. fenestrata</i> , Say, Journ. Acad. Nat. Sc. Philad. Vol. 3, p. 95 [1823] (<i>Sicus</i>); Wiedemann, Aussereur. Zweiftl. Ins. Vol. 2, p. 12 [1830] (<i>Tachydromia</i>); Say, Compl. Writ. Vol. 2, p. 82 [1859] (<i>Sicus</i>); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 [1895] (<i>Tachydromia</i>); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 228 [1902] (<i>Tachydromia</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903).
<i>rapax</i> , Loew, Berl. Ent. Zeitschr. Vol. 8, p. 85 : Cent. 5, No. 71 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 [1905] (<i>Tachydromia</i>); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 229, f. 55 [1902] (<i>Tachydromia</i>); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903). | North America. |

base beneath, the end of the middle tibiae slightly but conspicuously excavated. Halteres white, the base of their stems yellowish; calypteres with five white hairs. Wings subhyaline, a little dusky in the middle where the veins are blackish, the fifth vein dark even at its base, the fourth vein parallel with the third except toward its end where it gently curves forward, the third section of the costa three times the length of the fourth section, posterior crossvein shorter than the anterior, the distance between them longer than either, outer sections of the fifth vein subequal.

Two males from Pullman, Washington, July, 1908; another received from M. C. Van Duzee from Sacramento, California, June, 1915.

(1) *Tachypeza humeralis*, nov. sp. — Male. Length 3 mm. Humeri glistening black, legs yellowish, the front femora marked beneath with two round black spots. Occiput lightly cinereous, except a denser orbital line, occipital hairs sparse, antennæ reddish, the terminal brown arista quite four times the length of the antennæ, palpi whitish, proboscis reddish brown. Mesonotum lightly dusted, subshining, pleuræ polished. Pygidium moderately large, shining black, its hairs sparse. Femora not ciliate, the femoral spots smaller than in *binotata*, front tibiae not incrassate, middle femora not tuberculate, but with three spine-like setulae near base, middle tibiae with a shallow preapical setulose excavation, apex of the hind tibiae indefinitely darker. Halteres pale, calypteres with five pale hairs. Base of the wings whitish, but the first and fifth veins dark to their root, fourth vein slightly convergent toward the third, distance between the crossveins equal to the anterior crossvein.

One specimen, Washington, D. C., August 17, 1913.

- similis*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 506 [1849] (*Tachydromia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 [1895] (*Tachydromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 228 [1902] (*Tachydromia*).
13. *T. fuscipennis*, Fallen, Empid. Suec. p. 14 [1815] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 71 [1822] (*Tachydromia*); Macquart, Dipt. N. France, Vol. 3, p. 91 [1827] (*Tachydromia*); Curtis, Brit. Ent. Vol. 8, p. 477 [1833] (*Tachydromia*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 350 [1834] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 7, p. 95 (1838); Zetterstedt, Fauna Ins. Lappon. p. 546 (1838); Dipt. Scand. Vol. 1, p. 315 (1842); Boitard, Man. Ent. Vol. 3, p. 322 [1843] (*Tachydromia*); Gimmerthal, Bul. Soc. Nat. Moscou, Vol. 20 (2), p. 164 (1847); Walker, List Dipt. Brit. Mus. Vol. 3, p. 506 [1849] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3011 (1849); Scholz, Zeitsch. Ent. Breslau, Vol. 5 (19), p. 59 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 141 [1851] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 153 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 93 [1862] (*Tachydromia fusipennis*); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 14 (1864); Lundbeck, Dipt. Dan. Vol. 3, p. 265 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 88 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 71 (1913). Europe.
14. *T. Heeri*, Zetterstedt, Fauna Ins. Lappon. p. 547 (1838); Dipt. Scand. Vol. 1, p. 318 (1842); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 10, note (1864); Siebke, Nyt Mag. Naturvid. Vol. 12, p. 108 (1864); ibidem, Vol. 14, p. 387 (1866); Wahlgren, Ent. Tidskr. Vol. 31, p. 88 (1910). N. Europe.
15. *T. humeralis*, nov. sp. E. United States.
16. *T. incisa*, Brunetti, Rec. Indian Mus. Vol. 9, p. 41 [1913] (*Platypalpus*); Fauna Brit. India Dipt. Vol. 1, p. 379 (1920). W. Himalayas.
17. *T. inusta*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 226, f. 50, 54 [1902] (*Tachydromia*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903); Tucker, Kansas. Univ. Sc. Bull. Vol. 4, p. 96 [1907] (*Tachydromia*). W. United States.
18. *T. nubila*, Meigen, Classif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 239 [1804] (*Tachydromia*); Syst. Beschr. Vol. 3, p. 71 [1822] (*Tachydromia*); Curtis, Brit. Ent. p. 477 [1833] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 7, p. 95 (1838); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 12 (1864); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 126 (1892); Lundbeck, Dipt. Dan. Vol. 3, p. 266, f. 120, 121 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 88, f. 16 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 71 (1913). Europe.
- ? *cursoria*, Scopoli, Ent. Carn. p. 338 [1763] (*Musca*); Schiner, Verh. Zool.-bot. Ver. Wien, Vol. 6, p. 413 (1856).
- tibialis*, Macquart, Mém. Soc. Sc. Lille, p. 154 [1823] (*Tachydromia*); Dipt. N. France, Vol. 3, p. 91 [1827] (*Tachydromia*); Hist. Nat. Dipt. Vol. 1, p. 350 (1834); Meigen, Syst. Beschr. Vol. 7, p. 95 (1838).
- truncorum*, Meigen (not Fallen), ibidem, Vol. 3, p. 71 [1822] (*Tachydromia*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 350 [1834] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 7, p. 95 (1838); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 59 (1851).
- var. *nervosa*, Meigen, Syst. Beschr. Vol. 3, p. 72 [1822] (*Tachydromia*); Curtis, Brit. Ent. Vol. 8, p. 477 [1833] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 7, p. 95 (1838); Zetterstedt, Fauna Ins. Lappon. p. 547 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 317 [1842] (*Tachydromia*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 506 [1849] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3011 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 59 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 141 [1851] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 154 (1861); Europe.

- Schiner, Fauna Dipt. Austr. Vol. 1, p. 94 [1862] (*Tachydromia*); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 387 (1866); Cat. Dipt. Norv. p. 25 (1877); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 126 (1892).
19. *T. palliditibia*, Brunetti, Rec. Indian Mus. Vol. 9, p. 41 [1913] (*Platy-*
palpus); Fauna Brit. India Dipt. Vol. 1, p. 379 (1920). W. Himalayas..
20. *T. postica*, Walker, Trans. Ent. Soc. Lond. n. s. Vol. 4, p. 149 [1857] North America.
(*Tachydromia*); Coquillett, Proc. U. S. Mus. Vol. 18, p. 440 [1895]
(*Tachydromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 227,
f. 53 [1902] (*Tachydromia*).
21. *T. ?praelusio*, Walker, Ent. Mag. London, Vol. 3, p. 180 (1835). Britain.
22. *T. pruinosa*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 267 (1903). Missouri.
23. *T. rostrata*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 86 : Cent. 5, No. 72 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 [1895]
(*Tachydromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 229
[1902] (*Tachydromia*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5,
p. 265 (1903). E. United States.
24. *T. sericeipalpis*, Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (10), p. 10 Finland.
(1913).
var. *dilutata*, Frey, Mém. Acad. Sc. Russ. Vol. 29 (10), p. 14 (1915).
25. *T. truncorum*, Fallen, Empid. Suec. p. 14 [1815] (*Tachydromia*); Zetterstedt, Fauna Ins. Lappon. p. 547 (1838); Dipt. Scand. Vol. 1, p. 316 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 154 (1861); Schiner, Fauna Dipt. Austr. Vol. 1, p. 94 [1862] (*Tachydromia*); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 7 (1864); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 126 (1892); Wahlgren, Ent. Tidskr. Vol. 31, p. 88 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors. Vol. 37 (3), p. 72 (1913).
Ravipes, Thunberg, part. Nova Acta Soc. Scient. Upsal, Vol. 4, p. 26, var. *a* [1784] (*Empis*). Arctic Siberia.
N. & C. Europe.
26. *T. umbripennis*, Meigen, Syst. Beschr. Vol. 3, p. 70 [1822] (*Tachydromia*); C. Europe.
ibidem, Vol. 7, p. 95, pl. 67, f. 18-20 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 94 [1862] (*Tachydromia*).
27. *T. vittipennis*, Walker, Trans. Ent. Soc. Lond. n. s. Vol. 4, p. 149 [1857] North America.
(*Tachydromia*); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 439 [1895] (*Tachydromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 226 (1902).
28. *T. *voracis*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 89, 99, pl. 4, f. 11, Baltic Amber.
12, pl. 5, f. 1, 2 [1908] (*Tachydromia*).
29. *T. Winthemi*, Zetterstedt, Fauna Ins. Lappon. p. 548 (1838); Dipt. Scand. Vol. 1, p. 321 (1842); ibidem, Vol. 8, p. 3011 (1842); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 155 (1861); Loew, Zeit. Ent. Breslau, Vol. 17, p. 14 (1864); Siebke, Nyt. Mag. Naturv. Vol. 12, p. 108 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 [1895] (*Tachydromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 227 [1902] (*Tachydromia*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903); Wahlgren, Ent. Tidskr. Vol. 31, p. 88 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3) p. 72 (1913). Europe, Siberia, North America.
obscura, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 53 (1840).

2. GENUS TACHYDROMIA, MEIGEN

Tachydromia, Meigen, Illiger, Mag. Ins. Vol. 2, 269 (1803); Syst. Beschr. Vol. 3, p. 67 (1822); Macquart, Mém. Soc. Sc. Lille, 1823, p. 151 (1823); Curtis, Brit. Ent. Vol. 8, p. 477 (1824);

Macquart, Dipt. N. France, Vol. 3, p. 89 (1827); Hist. Nat. Dipt. Vol. 1, p. 349 (1834); Zetterstedt, Fauna Ins. Lappon. p. 548 (1838); Westwood, Gen. Syn. p. 132 (1840); Boitard, Man. Ent. Vol. 3, p. 322 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 138 (1851); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 563 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 92 (1862); Lioy, Atti Ins. Veneto Sc. Venezia, 1864, p. 720 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 125 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 17, p. 390, 439 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 224 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 611 (1910); Melander, Psyche, Vol. 17, p. 42-62 (1910).

Coryneta, Meigen, Nouv. Class. Dipt. p. 27 (1800); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 528 (1910).

Phoneutisca, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 19 (1863); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 125 (1889); Becker, Wien. Ent. Zeitschr. Vol. 9, p. 35 (1890); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 204 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 2, p. 255, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Bezzi, Nova Acta Akad. Nat. Halle, Vol. 91, p. 300 (1909); Kertész, Cat. Dip. Vol. 6, p. 135 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 588 (1910); Melander, Psyche, Vol. 17, p. 49 (1910).

Sicus, Latreille, not of Scopoli, Conopidae (1763); Précis. Caract. Ins. p. 158 (1796); Consid. Génér. p. 443 (1810); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 605 (1910).

Tachista, Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 15 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 125 (1889); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 43 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 258 (1903); Bezzi, Nova Acta Akad. Nat. Halle, Vol. 91, p. 300 (1909); Kertész, Cat. Dip. Vol. 6, p. 139 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 611 (1910); Lundbeck, Dipt. Danica, Vol. 3, p. 267 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45, 88 (1910); Bezzi, Ann. Mus Hungar. Vol. 10, p. 453 (1912).

Characters. — Minute, slender flies of shining jet-black color and almost devoid of hairs and bristles. Head globular, the occiput hemispherical or conical, its hairs sparse; eyes large, with large facets, those on the upper part a little smaller, broadly contiguous on the face in both sexes; front narrow, its sides straight and nearly parallel, slightly broadening above; three large ocelli, ocellar and vertical bristles minute; cheeks wanting; proboscis shorter than the head, rigid, vertical, not broadened at the base, palpi slender, ribbon-like, in the male sometimes remarkably broadened apically, vertical and applied against the proboscis, whose length they equal or exceed, tipped with one or several bristles; antennæ short, two-jointed, the basal joint without setæ, the outer joint rounded oval, with long, slender, nearly bare, terminal or subterminal arista.

Thorax longer than broad, not greatly convex, considerably narrowed from the wings forward; humeri greatly enlarged and constricted by a furrow from the disk of the mesonotum; sometimes a minute humeral bristle present, one notopleural, one postalar on a prominent callus, two or four scutellars, the lateral pair normally small, acrostichal setulæ microscopic, practically absent, dorsocentral setulæ uniseriate, minute, the hindmost longest but not as strong as in *Tachypeza*; no pleural bristles. Pygidium small, more or less globular, or triangular in outline; abdominal segments not pitted laterally. Legs slender, cursorial, the front femora strongest, devoid of bristles but with microscopic hairs, those of underside of front tibiae serrately arranged, no spurs or spines present: the male of *Tacoma* with the middle legs deformed, but in the other species the males have at most small flexor spines on

the middle femora or tibiæ. Wings narrow, anal angle much reduced, costa a little thicker beyond the ending of the first vein, marginal cell usually long, the second vein ending at the outer fifth of the wing, but sometimes (*Phoneutisca*) recurved, the second basal cell surpassing the first usually by two or more times the length of the crossveins, both crossveins small, the posterior transverse, no trace of an anal cell, hind margin rather short-ciliate but sometimes fringed; the wings are usually marked with two dark crossbands leaving a central more or less complete hyaline fascia between them, the base and apex hyaline.

Type species : *T. connexa*, Meigen (Pl. 4, Fig. 29). The application of the name *Tachydromia* has been the subject of considerable dispute, the history of which I have given in Psyche (Vol. 17, p. 41-48 [1910]). When Meigen erected the genus *Tachydromia* in 1803 he assigned to it two species, *cursitans* and *cimicoides*, one or the other of which must be selected as its type. In 1827 Macquart formed the genus *Platypalpus* for those species grouped about *cursitans*, and thus, by elimination, *cimicoides* became the type of *Tachydromia*. But Meigen's identification of the Fabrician *cimicoides* was erroneous, and later he described this species as *connexa*. This species was definitely assigned as the type of *Tachydromia* by Coquillett in 1903. According to the established laws of priority there is no reason for rejecting the name *Platypalpus*, although this has been done by the majority of dipterists, apparently on the principle of « follow the leader ». The recently resuscitated name *Coryneta*, Meigen, 1800, has been used by Kertész to replace *Tachydromia* (= *Platypalpus*) in his Catalogus.

Loew's genus *Phoneutisca* was established on *bimaculata*, an Alaskan species with recurved second vein and subterminal arista. I have this species now from British Columbia and Alaska. The male has curiously formed palpi, which slightly surpass the proboscis, the basal half slender and white, the apical half wide and black, deeply and broadly emarginate on the outside. This sex also has the second section of the costa thickened and there is a brown spot somewhat smaller than the stigmal spot beneath where the first vein attains the costa. In as much as other species of *Tachydromia* have the arista subterminal and the second vein recurved (e. g. *enecator*, Melander, *incompleta*, Becker) and the male of *Tacoma*, which is a true *Tachydromia*, has similarly formed palpi, it is evident that the name *Phoneutisca* can not have even subgeneric standing. It may interest those entomologists who insist on using the name *Tachista* for this group that Loew erected the genus *Phoneutisca* a year before he published *Tachista*.

The species of *Tachydromia* are extremely agile, running about with quick zig-zag movements, on the ground, over stones, boards, grass, etc. They carry the wings flat over the back and rarely make use of them in flight. They are predatory and nimbly chase microdiptera for prey.

NORTH AMERICAN SPECIES OF TACHYDROMIA

1. A pruinose spot on propleuræ above front coxæ, rarely absent; wings with two dark bands which rarely are united in front; the distance between the two crossveins more than twice the length of the hind crossvein 2.
- Crossveins approximated or meeting 13.
2. Second vein abruptly curving or bending forward at apex, the marginal cell obliquely truncated 3.
- Second vein not shortened, the marginal cell rather pointed; palpi and halteres usually pale 5.
3. Male palpi expanded apically into two black lobes, yellow at base 4.
- Male palpi narrow and black; legs black except for tarsi, tibiae not deformed; second vein straight; halteres dark. (Alask., Wyo., Queb.) T. ENECATOR, Melander.

4. Second section of costa subequal to third, second vein not sinuous; anterior legs and hind tibiæ brown, the middle tibiæ of male not deformed; stalk of male palpi shorter than proboscis, palpi of female white. (Alask., B. C., Ida., Wyo.)	T. BIMACULATA, Loew.
Second section of costa longer than third, second vein sinuous; legs almost wholly black, the middle tibiæ of male with stout thumb-like process before apex; stalk of male palpi slender and as long as proboscis, palpi of female black. (Wash., Ida.)	T. TACOMA, nov. sp. (1).
5. Dark cross-bands of wings united in front. (N. H., W. Va., Col., Ore., Wash.)	T. VARIPENNIS, Coquillett.
Dark cross-bands separated, at least through submarginal cell.	6.
6. Propleuræ polished	7.
Propleuræ pruinose; wings slender, the marginal hairs short	8.
7. Wings slender, fringe short. (Cal.)	T. PSELIOPHORA, nov. sp. (2).
Wings blunt, fringe longer than crossveins. (Mexico)	T. CILIATA, Melander.
8. Hyaline middle band of wings with parallel sides; palpi blackish; occiput polished	9.
Hyaline band curved or angulate at fourth vein; occiput largely pruinose	11.

(1) **Tachydromia Tacoma**, nov. sp. — Length 3.3 mm. Occiput gray pruinose except below where it becomes shining jet black, its upper bristles black, its lower ones long and white; front glistening black, its sides parallel, ocelli minute : antennæ blackish, the outer joint short ovate, with terminal arista four times the length of the antennæ : palpi of ♂ longer than the proboscis, slender, yellow and tapering on the proximal two-thirds, then suddenly widened, jet black, and with a deep and broad emargination on the outside; palpi of ♀ as long as the proboscis, slender, black and black-hairy; proboscis one-half the height of the head, vertical. Thorax slender, jet black, the humeri prominent; propleuræ and posterior half of the metapleuræ white-pruinose, scutellum and supraalar spot cinereous; humeral bristles wanting, dorsocentral and a single row of acrostichal setulæ minute, two supraalar macrochæte, four scutellar bristles. Basal half of the abdominal segments lightly cinereous, apical half jet black; hypopygium small, with long loose black curved hairs. Halteres with white knob; calypteres blackish, fringed with six dusky hairs. Legs jet black, the front coxæ white-pruinose anteriorly and tipped with a bunch of white hairs; front femora robust, pollinose beneath, middle femora ♂ with a series of long bristles on the anterior flexor edge; the middle tibiæ inside with a prominent thumb-like setulose preapical projection and on the extensor side with a setigerous tubercle opposite. Wings with two brown crossbands jointed in the marginal cell, the second vein sinuous, arching forward at the central hyaline fascia and then bending backward so that the marginal cell before its apex is as broad as the submarginal cell, the second section of the costa about one and one-half times as long as the third, the third and fourth veins subparallel, the fourth vein curving forward at the very end of the wing, the distance between the crossveins three times as long as the posterior crossvein, fifth vein nearly straight, terminating at the end of the hyaline fascia, its sections proportioned nearly two to one.

Many specimens from Mount Rainier, Washington, collected by Messrs. Dyar and Caudell and by the author, and one specimen from Moscow Mountain, Idaho. The flies are to be found running over the large boulders in the rivulets and on the rocky cliffs close to the timber-line and have been taken on tents in camp at Paradise Park. The specific name is suggested by the Indian name for the large snow-capped mountain which is the home of these flies.

(2) **Tachydromia psephilophora**, nov. sp. — Male. Length 2.1 mm. Polished black, the upper occiput very thinly pollinose, the pectus above pruinose. Front shining, face obliterated by the contiguity of the eyes; basal joint of antennæ yellow, outer joint brown, short-ovate, palpi blackish, narrow, tipped with a black seta. Abdomen shining, pygidium large, the left valve prominent, curved and finger-like, last ventral segment fringed with long black hairs. Coxæ yellow, front femora yellow becoming brown on distal one-half, front tibiæ thicker than the others and marked with two black rings, front tarsi yellow, the last joint black, middle brownish, the femora paler at base and furnished with two flexor rows of black setulæ, those at the base of the posterior row longest, hind femora and tibiæ blackish, paler at base, the tarsi brownish. Halteres white. Wings elongate oval, marked with two broad smoky bands, veins uniformly brown, middle hyaline stripe angled, costal sections two to four proportioned 5 : 3 : 1, second vein straight to near the gently curving end, first posterior cell narrow, sections of fourth vein 2 : 1 : 4.5, last section of fifth vein curved and separating the dark and light stripes.

Female. Front tibiæ uniformly yellowish brown.

Four specimens, Los Cerritos, California, April 3, 1915, collected by M. C. Van Duzee.

9. Front tibiae of male marked with black tip. 10.
 Front tibiae of male with two spots. (Ill., Indiana) T. HARTI, Malloch.

10. Lighter portions of legs pale yellow, the black of male front tibiae
 abruptly filling the apical third. (Texas) T. DIVERSIPES, Melander.

Lighter portions of legs testaceous, the black of male front tibiae
 forming an oval spot on inner side. (Ont., Virginia) T. PHENGITES, nov. sp. (1).

11. Front legs of male heavily pubescent; palpi black but overlaid with
 long silvery hairs; propleurae polished; dark cross-bands of wings
 blackish; legs black; second vein strongly sinuous. (Ore.) T. HIRTIPES, nov. sp. (2).
 Legs not heavily pubescent; propleurae pollinose 12.

12. Second vein somewhat sinuous; palpi black. (Cal., Utah, Ore., Wash.,
 Ida.) T. SCHWARZII, Coquillett.
 Second vein straight up to its apical curve; palpi white. (Wash.) T. CHELANA, nov. sp. (3).

13. Wings with a single large cloud in anterior apical portion; crossveins
 approximated; notum shining. 14.

(1) **Tachydromia phengites**, nov. sp. — Male. Length 1.5 mm. Similar to *Schwarzii*, Coquillett, but the middle hyaline band of the wings forming a straight streak across the wings instead of being curved or angulate at the fourth vein. Mesonotum lightly scabrous, subshining, abdomen polished black. Antennæ brown, the outer joint short oval, with terminal arista. Palpi blackish, slender, in both sexes, with a terminal black bristle; proboscis short, blackish. A large white pruinose spot above the front coxae. Knobs of the halteres white. Legs brownish yellow, the hind femora and tibiae, except basally, black. front coxae and upper side of front femora dark, last tarsal joint black, front tibiae of male tipped within with a round black spot. Wings marked with two broad brown crossbands, leaving the base, middle and tip hyaline; the second section of the costa two and one-half times the third section which is three times as long as the fourth section, second vein very gently curved and not arched, posterior crossvein shorter than the anterior, the distance between them twice the length of the posterior crossvein. In *Schwarzii* it is more than twice. Last section of the fifth vein ending in the middle of the hyaline band.

Five specimens, Falls Church, Virginia, received from Nathan Banks, and Ridgeway, Ontario, received from M. C. Van Duzee. The species is a close relative of *T. Harti*, which has two black spots on the front side of the front tibiae of the male.

(2) **Tachydromia hirtipes**, nov. sp. — Male. Length 3 mm. Shining black, the abdomen dull. Occiput and front pollinose, only the chin polished; antennæ piceous, the outer joint oval with the arista almost terminal and six times length of the joint; palpi narrow, black, nearly as long as the proboscis, overlaid with long silvery hairs and tipped with a black seta. Thorax smooth, jet-black, only the pectus and a spot above hind coxae pruinose. Abdomen piceous, the genitalia black, asymmetrical, consisting of three narrow, curving valves, last ventral fringed with black hairs. Legs black, the front coxae, the knees and the hind metatarsi piceous, front femora very robust, marked with a translucent spot near middle of inner side, front legs covered with abundant brown fine hairs, middle femora with a small brush of setae at base beneath, middle tibiae with a shallow emargination just before tip on flexor side. Knob of halteres white. Wings with arched costa, the base, middle and apex narrowly hyaline, the intermediate areas forming two broad deeply fuliginous cross-bands, second vein sinuous, the marginal cell narrowest at the hyaline middle band, second section of costa more than two times third, anterior crossvein at two-thirds length of discal cell, sections of fifth vein 2 : 1, the last section forming a sharp boundary to the proximal dark band, the middle hyaline band therefore angulate at the fourth vein.

Female. first four segments of abdomen subshining; legs not conspicuously pubescent, the front femora lacking the translucent spot and the middle legs lacking the basal brush and apical emargination.

Eight males and five females, all collected on rocks in the mountain stream near Viento, Oregon, July 1, 1917, and Aug. 1, 1921.

(3) *Tachydromia chelana*, nov. sp. — Female. Length 2.5 mm. Polished black, the upper occiput lightly pruinose, pruinosity of pectus reaching back to forward portion of the propleura. Antennae piceous, arista subterminal and five times length of last antennal joint; palpi two-thirds length of proboscis, ribbon-like, white, the hairs microscopic, no apical seta. Mesonotum smoothly polished; two pairs of scutellar bristles. Coxæ and metatarsi yellowish, remainder of legs blackish, the hind pair darkest. Halteres with white knob and blackish root. Wings narrow, bifasciate with blackish, leaving the base, middle and apex hyaline, the outer dark band encroaching proximally in the first posterior cell, first and second veins ending at outer edges of respective dark bands, the second vein straight through its course up to its apical gentle curvature, distance between crossveins three and one-half times the length of posterior crossvein, last section of fifth vein two-thirds the preceding, ending at beginning of outer dark band.

Two specimens, Stehekin, at the head of Lake Chelan, Washington, July 30, 1919 (Melander).

- Wings with two crossbands; crossveins meeting; notum dull; palpi bearing a very long seta; humeri yellow. T. SACHEM, nov. sp. (1).
14. Wholly black; propleuræ pollinose. (Wash.). T. MONACHA, nov. sp. (2).
- Legs, halteres and veins at base of wings in part paler; propleuræ polished. (Canada, Mass., Vt., N. Y., N. J., Md., Fla., Ill., Wisc., Mo., S. D.). T. MACULIPENNIS, Walker.

Geographical distribution.

- | | |
|--|------------|
| 1. <i>T. œmula</i> , Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 22 (1864). | C. Europe. |
| 2. <i>T. aliterpicta</i> , Becker, Berl. Ent. Zeitschr., Vol. 33, p. 343 [1889] (<i>Tachista alteropicta</i>); Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 32 [1900] (<i>Tachista</i>). | S. Europe. |
| 3. <i>T. annulimana</i> , Meigen, Syst. Beschr. Vol. 3, p. 69 (1822); Curtis, Brit. Ent. Vol. 8, p. 477 (1833); Meigen, Syst. Beschr. Vol. 7, p. 95 (1838); Schiner, Fauna Dipt. Austr. Vol. 1, p. 93 (1862); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 28 [1864] (<i>Tachista</i>); Strobl, Progr. Seitenst. Vol. 14, p. 57 (1880); Lundbeck, Dipt. Dan. Vol. 3, p. 269 [1910] (<i>Tachista</i>); Melander, Psyche, Vol. 17, pl. 3, f. 12 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 89 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 75 [1913] (<i>Tachista</i>). | Europe. |
| <i>albitarsis</i> , Zetterstedt, Dipt. Scand. Vol. 3, p. 313 [1842] (<i>Tachypeza</i>); ibidem, Vol. 8, p. 3011 [1849] (<i>Tachypeza</i>); Pipping, Not Sällsk. Fenn. Förh. Helsingfors, Vol. 4, p. 114 [1858] (<i>Tachypeza</i>); Nylander, ibidem, p. 247 [1858] (<i>Tachypeza</i>); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 153 [1861] (<i>Tachypeza</i>); Siebke, Cat. Dipt. Norv. p. 24 [1877] (<i>Tachypeza</i>). | |
| <i>arrogans</i> , Zetterstedt (not Linnæus), Fauna Ins. Lappon. p. 546, var. d [1838] (<i>Tachypeza</i>). | |
| <i>cimicoides</i> , Walker (not Fabricius), List Dipt. Brit. Mus. Vol. 3, p. 506 (1849). | |
| <i>umbrarum</i> , Haliday, Ent. Mag. London, Vol. 1, p. 161 [1833] (<i>Tachypeza</i>). | |
| 4. <i>T. arrogans</i> , Linnæus, Fauna Suec. p. 457 [1761] (<i>Musca</i>); Syst. Nat. (ed. 12), Vol. 2, p. 995 [1767] (<i>Musca</i>); Fabricius, Syst. Ent. p. 783 [1775] (<i>Musca</i>); Schrank, Enum. Ins. Austr. p. 467 [1781] (<i>Musca</i>); Fabricius, Spec. Ins. Vol. 2, p. 448 [1781] (<i>Musca</i>); Mant. Ins. Vol. 2, p. 349 [1787] (<i>Musca</i>); Gmelin, Syst. Nat. Vol. 5, p. 2852 [1790] (<i>Musca</i>); Fabricius, Ent. Syst. Vol. 4, p. 341 [1874] (<i>Musca</i>); ?Syst. Ant. p. 263 [1805] (<i>Calobata</i>); Olivier, Encycl. Méth. Vol. 8, | Europe. |

(1) **Tachydromia sachem**, nov. sp. — Female. Length 1.3 mm. Head black, occiput cinereous dusted, front dull black, narrow, sides bulging around the ocelli, the eyes closer together on vertex than above antennæ, upper facets enlarged, face obliterated; occipital setæ flat, scale-like, white; basal antennal joint whitish yellow, outer joint short oval, dusky, arista nearly four times the antennal length; proboscis yellow, tipped with brown, palpi small, circular, brown, bearing an extraordinary black seta which is nearly twice as long as the proboscis. Thorax dull in color, humeri very large and yellowish, remainder of thorax piceous, pleuræ lightly cinereous, propleuræ shining, one strong pair each of notopleuræ, dorsocentral and scutellar bristles. Abdomen dull piceous, a little paler at base and apex, its sparse short hairs black. Coxæ and most of anterior legs yellow, last two joints of anterior tarsi black, hind femora yellow on basal third, remainder black, hind tibiae white, the distal third black, first two joints of hind tarsi white, the last three black, hind metatarsus as long as remaining joints together. Wings narrow, infumated, the base, apex and a central spot included between the third and fifth veins hyaline, crossveins meeting, third and fourth veins nearly parallel, sections of fifth vein subequal; halteres pale yellow.

A single specimen of this remarkable fly received from Pablo Schild, Turrialba, Costa Rica, November, 1922. The species departs from the others in *Tachydromia* in having the thorax dull and the crossveins attigent.

(2) **Tachydromia monacha**, nov. sp. — Female. Length 2 mm. Related to *maculipennis*, Walker, but entirely black, including the root of the wings, the pedicel of the halteres and the base of the legs. Wings broader, the blackish anterior cloud which stops along the middle of the first posterior cell coming exactly to the middle of the wing, — in *maculipennis* the middle of the wing is a little closer to the third vein; the white pruinose spot above the front coxæ extending on the propleuræ, — in *maculipennis* it fills the pectus only. The wings, aside from the blackish cloud, are decidedly smoky, and not subhyaline as in *maculipennis*.

One specimen, Pullman, Washington, May 30, 1913.

p. 29 [1811] (*Musca*); Fallen, Empid. Suec. p. 13 (1815); Billberg, Enumer. Ins. p. 119 [1820] (*Sicus*); Meigen, Syst. Beschr. Vol. 3, p. 68 (1822); Macquart, Dipt. N. France, Vol. 3, p. 90 (1827); Curtis, Brit. Ent. Vol. 8, p. 477, pl. (1833); Macquart, Hist. Nat. Dipt. Vol. 1, p. 349 (1834); Guérin, Icon. Règne Anim. Ins. p. 537, pl. 94, f. 8 (1835); Meigen, Syst. Beschr. Vol. 7, p. 94 [1838] (*Tachypesa*); Zetterstedt, Fauna Ins. Lappon. p. 546, part [1838] (*Tachypesa*); Dipt. Scand. Vol. 1, p. 312 [1842] (*Tachypesa*); ibidem, Vol. 8, p. 3010 [1849] (*Tachypesa*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 59 [1851] (*Tachypesa*); Walker, Ins. Brit. p. 1139, pl. 5, f. 6 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 153 [1861] (*Tachypesa*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 93 (1862); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 20 [1864] (*Tachista*); Giard, Traité Ent. Vol. 3, p. 992, pl. 108, f. 8 (1885); Neuhaus, Dipt. March. p. 74 (1886); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 124 [1892] (*Tachista*); Lundbeck, Dipt. Dan. Vol. 3, p. 267, f. 122 [1910] (*Tachista*); Melander, Psyche, Vol. 17, pl. 3, f. 10 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 89, f. 17 [1910] (*Tachista*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 73 [1913] (*Tachista*).

bifasciata, Rossi, Fauna Etrusc. Vol. 2, p. 77 [1794] (*Empis*); Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 152 [1899] (*Tachista*).

cimicoides, Fabricius, Spec. Ins. Vol. 2, p. 447 [1781] (*Musca*); Mant. Ins. Vol. 2, p. 349 [1787] (*Musea*); Gmelin, Syst. Nat. Vol. 5, p. 2852 [1790] (*Musca*); Fabricius, Ent. Syst. Vol. 4, p. 339 [1794] (*Musca*); Latreille, Hist. Nat. Crust. Ins. Vol. 14, p. 312 [1804] (*Sicus*); Fabricius, Syst. Antl. p. 144 (1805); Olivier, Encyclop. Méthod. Vol. 8, p. 30 [1811] (*Musca*); Lamarck, Hist. Nat. Anim. sans Vert. Vol. 3, p. 402 [1816] (*Empis*); Billberg, Enumer. Ins. p. 119 [1820] (*Sicus*); Macquart, Mém. Soc. Sc. Lille, 1823, p. 115 (1823); Boitard, Man. Ent. Vol. 2, p. 369 [1828] (*Sicus*); Meigen, Syst. Beschr. Vol. 6, p. 341 (1830); Macquart, Hist. Nat. Dipt. Vol. 1, p. 349 (1834); Meigen, Syst. Beschr. Vol. 7, p. 94 [1838] (*Tachypesa*); Boitard, Man. Ent. Vol. 3, p. 322 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 140 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 93 (1862).

var. *productipes*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 85 [1909] (*Tachysta*) C. Europe.

5. *T. barbata*, Oldenberg, Zool. Jahrb. Vol. 43, Syst. p. 230, f. c, d [1920] Alps.
(*Tachista*).

6. *T. bimaculata*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 19 : Cent. 3, No. 35 [1863] (*Phoneutrisca*); Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 423 [1900] (*Phoneutrisca*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 204, part [1902] (*Phoneutrisca*); Psyche, Vol. 17, p. 52 [1910] (*Phoneutrisca*).

maculipennis, Coquillett, 1903 (not Walker). Proc. Ent. Soc. Wash. Vol. 5, p. 266 (1903).

7. *T. bistigma*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 490 [1912] (*Tachista*); Formosa.
Suppl. Ent. Berlin, Vol. 3, p. 78 [1914] (*Tachista*).

8. *T. calcanea*, Meigen, Syst. Beschr., Vol. 7, p. 95 [1838] (*Tachypesa*); C. Europe.
Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 59 [1851] (*Tachypesa*);
Schiner, Fauna Dipt. Austr. Vol. 1, p. 93 (1862); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 126 [1893] (*Tachysta*);
Melander, Psyche, Vol. 17, pl. 3, f. 6 (1910); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, 85 [1910] (*Tachysta*).

longipennis, Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 29 [1864] (*Tachista*).

9. *T. calcarata*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 85 [1909] (*Tachysta*). C. Europe.

10. *T. catalonica*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 319 [1906] Spain.
(*Tachysta*); Kertész, Cat. Dipt. Vol. 6, p. 151 [1909] (*Corynetia*).
var. *striatipennis*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 320 [1906] Spain.
(*Tachysta*); Kertész, Cat. Dipt. Vol. 6, p. 131 [1909] (*Corynetia*).

11. *T. chelana*, nov. sp. Washington.
 12. *T. ciliata*, Melander, Psyche, Vol. 17, p. 55, f. 13 (1910). Mexico.
Schwarzii, Wheeler & Melander (not Coquillett), Biol. Cent. Amer. Dipt. Vol. 1, p. 375 (1901); Melander, Trans. Ent. Soc. Vol. 28, p. 225, part (1902).
13. *T. connexa*, Meigen, Syst. Beschr. Vol. 3, p. 70, pl. 23, f. 24 (1822); Europe, East Indies. Macquart, Dipt. N. France, Vol. 3, p. 91, pl. 2, f. 3 (1827); Curtis, Brit. Ent. Vol. 8, p. 477 (1833); Macquart, Hist. Nat. Dipt. Vol. 1, p. 350, pl. 8, f. 9 (1834); Meigen, Syst. Beschr. Vol. 7, p. 95 [1838] (*Tachypeza*); Boitard, Man. Ent. Vol. 3, p. 322 (1843); Scholz, Zeit. Ent. Breslau, Vol. 5 (19), p. 59 [1851] (*Tachypeza*); Walker, Ins. Brit. Dipt. Vol. 1, p. 140 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 93 (1862); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 23 [1864] (*Tachista*); Neuhaus, Dipt. March. p. 74 (1886); Leunis, Synops. Zool. Vol. 2, p. 401 (1886); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 123 [1892] (*Tachysta*); Melander, Psyche, Vol. 17, pl. 3, f. 7 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 88 [1910] (*Tachista*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 72 [1913] (*Tachista*); Beazzi, Suppl. Ent. Berlin, Vol. 3, p. 78 (1914). —
- Pl. 4, Fig. 29.**
- cimicoides*, Meigen (not Fabricius), Klassif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 239 (1804).
morio, Zetterstedt, Fauna Ins. Lappon. p. 546 [1838] (*Tachypeza*); Dipt. Scand. Vol. 1, p. 314 [1842] (*Tachypeza*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 153 [1851] (*Tachypeza*); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 125 [1892] (*Tachysta*).
14. *T. denticulata*, Oldenberg, Ent. Mitteil. Berlin, Vol. 1, p. 212, f. 1, 2 Alps. [1912] (*Tachista*)
15. *T. diversipes*, Melander, Psyche, Vol. 17, p. 55 [1910] (*Schwarzii*, var. *diversipes*). Texas.
16. *T. enecator*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 226 (1902); Alaska, Quebec. Melander, Psyche, Vol. 17, p. 54, f. 1, 4 (1910).
17. *T. excisa*, Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 27 [1864] (*Tachista*).
18. *T. fuscinervis*, Frey, Mém. Acad. Sc. Russ. Vol. 29 (10), p. 15 [1915] (*Tachista*). Arctic Siberia.
19. *T. Harti*, Malloch, Canad. Ent. Vol. 51, p. 248 (1919). Illinois, Indiana.
20. *T. hirtipes*, nov. sp. Oregon.
21. *T. incompleta*, Becker, Act. Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 33, pl. 2, f. 33 [1900] (*Tachista*). Siberia.
22. *T. interrupta*, Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 19 [1864] (*Tachysta*); C. & S. Europe. Mik. in Beck, Fauna Hernstein, Vol. 2, p. 2 [1885] (*Tachista*); Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 34, p. 213 [1898] (*Tachysta*); Beazzi, Bull. Soc. Ent. Ital. Vol. 30, p. 161 [1899] (*Tachista*); Melander, Psyche, Vol. 17, pl. 3, f. 9 (1910). var. *obsoleta*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 46, p. 85 [1909] (*Tachysta*). C. Europe.
styriaca, Strobl, ibidem, Vol. 29, p. 124, var. *d* [1892] (*Tachysta*).
23. *T. latifascipennis*, Brunetti, Rec. Indian Mus. Vol. 13, p. 81 (1917); India. Fauna Brit. India Dipt. Vol. 1, p. 375 (1920).
24. *T. Lundströmi*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 73, f. 20 [1913] (*Tachista*). Finland.
25. *T. maculipennis*, Walker, List. Dipt. Brit. Mus. Vol. 3, p. 507 (1849); North America. Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 440 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 228 (1902); Melander, Psyche, Vol. 17, p. 52, 57, f. 3, 5 (1910). *bimaculata*, Melander (not Loew), Trans. Amer. Ent. Soc. Vol. 28, p. 204, part, f. 1 [1902] (*Phoneutrisca*).

- pusilla*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 87 : Cent. 5, No. 74 [1864] (*Tachypesia*) ; Coquillet, Proc. U. S. Nat. Mus. Vol. 18, p. 439 (1895) ; Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 229, f. 51 (1902) ; Coquillet, Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903).
26. *T. microptera*, Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 26 [1864] C. Europe. (*Tachista*).
? brevipennis, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 53 [1840] (*Tachypesia*).
27. *T. minima*, Becker, Act. Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 32 Siberia. [1900] (*Tachista*).
28. *T. monacha*, nov. sp. Washington.
29. *T. monserratensis*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 318 [1906] Spain. (*Tachista*) ; Kertész, Cat. Dipt. Vol. 6, p. 162 [1909] (*Coryneta*).
30. *T. ? morio*, Walker (not Zetterstedt), Ins. Brit. Dipt. Vol. 1, p. 141 (1851). Britain.
31. *T. nigerrima*, Bezzi, Dittero Faun. nivale, p. 112, pl. 1, f. 2, 8 [1918] Alps. (*Tachista*).
32. *T. ornatipes*, Becker, Wien. Ent. Zeit. Vol. 9, p. 69, f. 1, 2 [1890] S. Europe. (*Tachista*) ; Oldenberg, Zool. Jahrb. Vol. 43, Syst. p. 227, f. a, b [1920] (*Tachysta*).
33. *T. phengites*, nov. sp. * Virginia.
34. *T. pseliophora*, nov. sp. California.
35. *T. punctifera*, Becker, Act. Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 32, pl. 2, f. 32 [1900] (*Tachista*) ; Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 72 [1913] (*Tachista*).
maculipennis, Frey (not Walker), Naturw. Untersuch. Sarekgebirg. Vol. 4 (6), p. 687 [1916] (*Tachista*).
36. *T. sabulosa*, Meigen, Syst. Beschr. Vol. 6, p. 342 (1830) ; ibidem, Vol. 7, p. 95 [1838] (*Tachypesia*) ; Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 59 [1851] (*Tachypesia*) ; Schiner, Fauna Dipt. Austr. Vol. 1, p. 93 (1862) ; Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 17 [1864] (*Tachista*) ; Lundbeck, Dipt. Dan. Vol. 3, p. 270 [1910] (*Tachista*) ; Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 72 [1913] (*Tachista*).
fenestrata, Zetterstedt, Dipt. Scand. Vol. 1, p. 318 [1842] (*Tachypesia*).
37. *T. sachem*, nov. sp. Costa Rica.
38. *T. Schwarzi*, Coquillet, Proc. U. S. Nat. Mus. Vol. 18, p. 440 (1895) ; Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 225, part, f. 52 (1902) ; Slosson, Ent. News, Philad. Vol. 14, p. 267 (1903) ; Bezzi, Ann. Mus. Hungar. Vol. 3, p. 460 [1905] (*Tachypesia*) ; Melander, Psyche, Vol. 17, p. 54, f. 14 (1910).
39. *T. *stilpon*, Meunier, Loew, Bernsteinfauna, p. 42, part (1850) ; Miscell. Ent. Vol. 7, p. 178 (1899). Baltic Amber.
40. *T. styriaca*, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 124 C. Europe. [1892] (*Tachista*) ; Melander, Psyche, Vol. 17, pl. 3, f. 8 (1910).
semifasciata, Strobl, Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 125 [1892] (*Tachista*).
41. *T. Tacoma*, nov. sp. Washington.
42. *T. terricola*, Zetterstedt, Vet. Akad. Handl. Stockholm, Vol. 1, p. 81 (1819) ; Meigen, Syst. Beschr. Vol. 3, p. 72 (1822) ; Fallen, Dipt. Suec. Suppl. p. 6 (1826) ; Meigen, Syst. Beschr. Vol. 6, p. 341 (1830) ; ibidem, Vol. 7, p. 95 [1838] (*Tachypesia*) ; Zetterstedt, Dipt. Scand. Vol. 1, p. 320 [1842] (*Tachypesia*) ; ibidem, Vol. 8, p. 3011 [1849] (*Tachypesia*) ; Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 17 [1864] (*Tachista*) ; Lundbeck, Dipt. Danica, Vol. 3, p. 271 [1910] (*Tachista*) ;

- Wahlgren, Ent. Tidskr. Vol. 31, p. 88 [1910] (*Tachista*); Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 72 [1913] (*Tachista*)
apicata, Stæger, in litt., Zetterstedt, Dipt. Scand. Vol. 8, p. 3011 [1849]
(*Tachypeza*).
43. *T. tuberculata*, Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 25 [1864] C. Europe.
(*Tachista*); Verrall, Ent. Mag. London, Vol. 48, p. 26 [1912]
(*Tachista*).
44. *T. undulata*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 317 [1906] Spain.
(*Tachysta*); Kertész, Cat. Dipt. Vol. 6, p. 168 [1909] (*Coryneta*);
Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 182 [1909] (*Tachysta*).
45. *T. varipennis*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 266 (1903); United States.
Slosson, Ent. News, Philad. Vol. 14, p. 267 (1903); Melander,
Psyche, Vol. 17, p. 56, f. 11 (1910).

3. GENUS PIELTAINIA, ARIAS

Pieltainia, Arias, Bol. Soc. Esp. Hist. Nat. Vol. 19, p. 479 (1919).

Characters. — Entirely dull black, measuring less than three millimeters, the tarsi and sometimes the knees alone paler. Proboscis vertical, palpi narrow; antennæ porrect, the apical joint acuminate and with terminal arista; eyes large, occupying the whole side of the head. Legs simple in both sexes, slender, cursorial, the front femora slightly the largest. Male abdomen with blunt termination, female abdomen pointed. Wings and halteres entirely wanting.

Genotype : *P. iberica*, Arias. This is the only Empid known that is entirely wingless. In several of the genera of this subfamily there is a tendency toward rudimentary wings, in some cases a concomitant or possibly the reaction of the specialized cursorial habit, in other cases, e. g. the maritime species, the result of persistent winds making flying dangerous. Even the coastal and cascadal species with functional wings are loathe to fly. The present species was found in early spring actively running about in pursuit of Sciara and other microscopic Diptera on which it preyed.

Geographical distribution.

1. *P. iberica*, Arias, Bol. Soc. Esp. Hist. Nat. Vol. 19, p. 479, 1. 1a & 2a Spain.
(1919).

4. GENUS DYSALETRIA, LOEW

Dysaletria, Loew, Zeitschr. Ent. Breslau, Vol. 14, p. 7 (1860); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 124 (1889); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 43 (1902); Coquillett, Proc. Wash. Ent. Soc. Vol. 5, p. 249 (1903); Kertész, Cat. Dipt. Vol. 6, p. 143 (1909); Melander, Psyche, Vol. 17, p. 49 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45 (1910).

Characters. — Small yellow species with black head. Head globose, proboscis short; antennæ small, the third joint oval, with a slender apical arista. Thorax subopaque, with short pubescence. Legs slender, the anterior femora but slightly thickened, last joint of the hind tarsi somewhat widened. Wings narrow, the anal angle much reduced, basal cells long and equal, the crossveins near the middle of the wing, third and fourth veins parallel and close together, ending at the wing-tip, anal cell wanting, or only the outer angle visible.

Type species : *D. atriceps*, Boheman.

Geographical distribution.

1. *D. atriceps*, Boheman, Vet. Akad. Förh. Stockholm, 1851, p. 190 [1852] C. & N. Europe.
(*Tachypeza*); Zetterstedt, Dipt. Scand. Vol. 12, p. 4603 [1855]
(*Tachypeza*); Roeder, Wien. Ent. Zeit. Vol. 3, p. 291 (1884);
Wahlgren, Ent. Tidskr. Vol. 31, p. 89, f. 18 (1910).
melanocephala, Loew, Zeit. Ent. Breslau, Vol. 17, p. 31 (1863).
2. *D. ?* diabolica*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 89, 100, pl. 5, Baltic Amber.
f. 3, 4 [1908] (*Elaphropeza*).

5. GENUS TACHYEMPIS, NOV. GEN.

Characters. — Very small species, less than two millimeters in size, usually cinereous pollinose on the pleuræ as well as on the notum. Occiput flattened, without bristles; front broadly V-shaped, its sides outwardly bowed; ocelli variable in size, one pair of diverging ocellar bristles and usually in front of them a pair of minute convergent setulæ; one pair of vertical bristles, all the bristles small. Eyes contiguous below the antennæ obliterating the face and cheeks, bare, the lower facets large, the emargination at the antennæ deep. Antennæ short, two-jointed, without a seta beneath, the last joint oval, with subterminal arista. Proboscis very short, vertical, the palpi variable in width, but typically oval, densely hairy and white. Thoracic bristles variable, a double row of acrostichals may be present or absent, one humeral, two notopleural, two or four scutellar bristles. Abdomen comprising seven segments and the rather large pygidium, the segments usually with evident lateral pits. Legs simple, not enlarged, short-pubescent, no spines, spurs or bristles on the tibiæ, last tarsal joint and the pulvilli small; the front femora are a little the stoutest. Wings hyaline or nearly so, the costa not fringed; basal cells short, the posterior crossvein near the basal third of the wing but beyond the anterior crossvein, second and third sections of the costa nearly equal.

Type species: *T. agens*, Melander.

The following North American species make a natural group with the exception of *calva* and *simplicior*, which are aberrant in any of the genera. These species all possess pittings at the lateral margins of the tergites of the abdomen and are more or less pollinose. Two African species with clear wings, described by Dr. Bezzi as *Tachista*, appear to belong here.

KEY TO THE AMERICAN SPECIES OF TACHYEMPIS

- | | |
|--|---------------------------------|
| 1. Disk of mesonotum polished, devoid of all dust; humeri prominent | 2. |
| Thorax and abdomen subglaucous, the disk more or less dusted; humeri generally smaller. | 6. |
| 2. Legs largely or wholly black; front of male fulvous-pollinose between antennæ and front ocellus; base of pygidial hook with two strong bristles on right side | <i>T. nigra</i> , nov. sp. (1). |

(1) ***Tachyempis nigra*, nov. sp.** — Male. Length 1.5 mm. Black, vertex and disk of thorax polished. Ocelli small, front fulvous-pollinose between antennæ and front ocellus, entire occiput with dark brown coating, one pair each of black vertical and ocellar bristles; basal joint of antennæ yellow to reddish, outer joint black, the subdorsal arista black, two and a half times the antennal length; palpi broad, glistening white, with a few yellow setæ. Mesonotum nearly bare, about three minute acrostichal and seven dorsocentral hairs, notopleural suture, scutellar area and pleuræ densely velvety dark brown pollinose, becoming silvery toward and on coxae, one notopleural, one supra-alar and one pair scutellar black

Legs in large part yellow; front polished; pygidium without specialized bristles	3.
3. Third and fourth veins subparallel	4.
Third vein curving forward, diverging from the fourth.	T. SIMPLICIOR, Wheeler & Melander.
4. Occiput coated with dust; wings nearly uniformly infumated; front tibiæ darker than femora	5.
A broad orbital stripe polished; base of wings pale, remainder infumated; front tibiæ and femora concolorous	T. GAGATINA, nov. sp. (1).
5. Front tibiæ black; fourth vein ending at wing-tip.	T. CAPNODES, nov. sp. (2).
Front tibiæ piceous; fourth vein ending beyond wing-tip . .	T. SCHILDI, nov. sp. (3).
6. Hind metatarsi with a long black spine; palpi broad; legs yellow	T. LONGISPINA, nov. sp. (4).
No metatarsal spine	7.

bristles. Abdomen overlaid with dark brown pollen, pygidium contorted and usually open, the curved apical process slender and bearing two strong black setæ at base on right side. Legs blackish, the tibiae and base of tarsi brown, coxal hairs yellow, other hairs brown. Halteres yellow, calypteres and fringe dusky. Wings smoky, especially dark in marginal and submarginal cells, veins black and strong, sections of costa proportioned 10 : 11 : 9 : 3, third and fourth veins parallel toward end, the fourth ending beyond wing-tip, first basal cell two-thirds as long as second, sections of fifth vein 7 : 10.

Female. Antennal joints concolorous, dark; front wholly polished.

Twenty-four specimens, procured from Pablo Schild, La Suiza de Turrialba, Costa Rica, collected in various months; two specimens, San Jose, Costa Rica (Sturtevant). The legs may sometimes be more or less brownish.

(1) **Tachyempis gagatina**, nov. sp. — Differs from *capnodes* as follows: occiput broadly polished along the orbits, the first antennal joint of male yellowish, the apical joint brown, of female both joints black, arista two and a quarter times antennal length. The gray tone to the pruinosity of lower pleuræ and coxæ scarcely evident; pygidium short, small, closed, hairy, the terminal curved process thin and devoid of specialized bristles; legs mostly testaceous yellow, the front tibiæ not darker than their femora, the hind femora only a little brownish on apical half. Base of wings subhyaline, remainder uniformly infumated, costal sections about 0.6 : 1 : 0.6 : 0.3, first posterior cell about three-fourths as wide as the submarginal opposite the end of the second vein, fourth vein ending beyond wing-tip, sections of fifth vein 2 : 3.

Six males, one female, La Suiza de Turrialba, Costa Rica (P. Schild).

(2) **Tachyempis capnodes**, nov. sp. — Female. Length 1.5 mm. Occiput opaque black, only a linear orbital stripe shining, front polished, at the vertex as wide as antennal length, below as wide as the outer antennal joint, the single ocellar and vertical bristles black; palpi glistening white, with a white seta, proboscis black; antennæ black, the subdorsal arista two and a half times the antennal length. Mesonotum polished, extreme margins of notum and all of pleuræ pollinose, black, becoming silvery gray toward and on posterior coxæ. Abdomen subopaque black. Legs very sturdy, entirely bristleless, brownish yellow, front coxae black at base merging to the yellow tip, touched with gray pruinosity on front face, front tibiae entirely black, hind femora blackish on apical third. Halteres with yellow stem and white knob. Wings strongly and uniformly infumated, sections of costa beyond humeral crossvein proportioned 1 : 1 : 0.6 : 0.3, first posterior cell relatively narrow, about two-thirds as wide as the submarginal opposite end of second vein, fourth vein gently curving forward, ending at wing-tip, sections of fifth vein about 4 : 5.

Holotype, La Suiza de Turrialba, Costa Rica, August, 1922 (P. Schild).

(3) **Tachyempis Schildi**, nov. sp. — Differs from *capnodes* as follows: Occiput dusted up to the eyes; arista nearly three times the antennal length, basal antennal joint of male yellow, the outer joint black, of female the antennæ wholly black; gray pruinosity of lower pleuræ and of coxæ scarcely evident; pygidium open, the curved terminal process without specialized bristles; front femora less robust, front tibiæ piceous, becoming paler at apex; costal sections proportioned 10 : 11 : 7 : 3, first posterior cell about three-fourths as wide as the submarginal opposite end of second vein, the fourth vein ending beyond apex of wing, sections of fifth vein 3 : 4.

One male and one female, La Suiza de Turrialba, Costa Rica, procured from Pablo Schild. The species is dedicated to Mr. Schild, whose diligent collecting has made possible the recording of seven of the new species of the genus.

(4) **Tachyempis longispina**, nov. sp. — Length 1 mm. Uniformly cinereous gray, the legs yellow, hind metatarsi with a long black spine, wings hyaline. Front narrowly V-shaped, ocelli moderate in size; no ocellar bristles; antennæ reddish, the outer joint short oval, the terminal arista nearly four times the length of the antennæ. Palpi broad, white, not fringed; proboscis short and vertical, black. Occipital hairs white, sparse. Thorax not elongate, the humeri evident but not markedly constricted; only a few microscopic bristles present on the notum before the scutellum; four minute scutellar bristles. Hypopygial claspers fringed with hairs beneath. Legs including the coxae yellow, the hind femora brownish toward tip, hairs sparse, minute and white, but the hind metatarsi of the male tipped with a slender black

7. Second section of costa longer than third, usually one and one-half to two times the third 8.
- Second and third sections of costa subequal, or the second shorter than the third 12.
8. Front almost linear above and below the ocellar triangle; wings more or less clouded; femora apically and front tibiae centrally dark T. PICTIPES, nov. sp. (1).
- Front more or less V-shaped 9.
9. Upper half of front polished, lower half gray pollinose; palpi longer than broad, white; sections of fifth vein equal T. CINEREA, nov. sp. (2).
- Upper part of front dusted quite as much as lower 10.
10. Sections of fifth vein equal, veins mostly strong and black; palpi longer than wide T. NERVOSA, nov. sp. (3).
- Last section of fifth vein longer than preceding, veins thin; palpi nearly as broad as long 11.

spine which is as long as the metatarsus and attains the end of the following joint. Halteres white; calyptères bare. Wings nearly hyaline, slightly clouded near the middle, the second and third sections of the costa subequal, third and fourth veins parallel, last section of the fifth vein slightly longer than the preceding section, posterior crossvein two-thirds as long as the distance between the crossveins.

Five specimens, Havana, Cuba (C. F. Baker), and Jamaica (R. Thaxter).

(1) **Tachyempis pictipes**, nov. sp. (PI. 4, FIG. 35). — Length 2 mm. Black, with brownish tinge. Head very narrow, higher than wide, occiput lightly but completely gray-dusted, its lower hairs pale; front cinereous, almost linear, bulged at the very narrow ocellar triangle, vertical bristles almost touching; antennæ yellowish, the arista three and a half times the antennal length; palpi narrow and long, white, tipped with a long yellow seta, proboscis black. Thorax narrow, uniformly and lightly dusted with grayish, one each of humeral, notopleural, dorsocentral and supra-alar bristles, apical scutellars distant. Abdomen dull brown, the recurved apical process narrow. Front femora robust, fringed with long yellow sete beneath, coxae and most of legs yellow, the middle of the anterior tibiae and the distal part of the posterior femora brown. Wings narrow, hyaline at base, otherwise lightly infumated, veins brown, sections of costa proportioned 9 : 10 : 7 : 3, of fifth vein 4 : 5, first posterior cell narrow, ending at apex of wing, but little narrower than the submarginal opposite end of second vein. Halteres pale yellow, calyptères brown.

Seven males, three females, La Suiza de Turrialba, Costa Rica (P. Schild).

(2) **Tachyempis cinerea**, nov. sp. — Female. Length 1.5 mm. Occiput, thorax and abdomen gray pruinose; legs yellowish; basal joint of the antennæ red; palpi elongate, white and with small white hairs; two scutellar bristles; no acrostichal or dorsocentral bristles; halteres white; wings clear hyaline.

Front V-shaped, the bottom as broad as an antennal joint, ocelli large, divergent black ocellar bristles present; outer antennal joint short, onion shaped, the arista subterminal, three times length of antennæ; eyes completely contiguous on the face, the lower facets large; proboscis short, black. Humeri evident and delimited by a distinct furrow. Pleuræ densely pruinose, the ground color a little reddish near the front coxae, no pleural hairs. Middle tibiae minutely and closely black-setulose within. Veins narrow and dark except at the base of the wing, marginal cell long, the second section of the costa one and three-fourths as long as the third section, first posterior cell ending at the wing-tip, its veins parallel, the outer sections of the fourth vein proportioned 1 : 6, of the fifth vein, subequal.

Two specimens; Alamogordo, New Mexico, type in the Academy of Natural Science, Philadelphia. This species differs from *apicis* Williston (*Drapetis* = *insularis* Melander) in that the palpi and halteres are not brownish, the wings show no trace of infuscation, the second section of the costa is not equal to the third, the outer section of the fifth vein is not longer than the preceding, and in that the front is somewhat broader.

(3) **Tachyempis nervosa**, nov. sp. — Female. Length 1.75 mm. Front narrowly V-shaped, cinereous, vertical bristles minute, occiput completely dusted; basal antennal joint brown, apical blackish, arista three times the antennal length; palpi elliptical, brownish, white at base and with glistening white coating, two small white setæ; notum covered with brownish gray dust, humeri pronounced, no humeral bristle, seriate setulae very small, posterior dorsocentral present, one notopleural, lateral scutellar seta small; pleuræ cinereous, not whiter below. Abdomen subshining. Coxæ and legs mostly yellowish, apical half of posterior femora and posterior tarsi brownish, anterior femora with rows of yellow setulae below. Halteres yellow, calyptères blackish. Wings hyaline, veins mostly black, the fourth vein thin, the pedicel of the second and third veins yellow, costal sections about 1 : 1 : 0.6 : 0.2, sections of fifth vein equal, first posterior cell ending at apex of wing, third and fourth veins straight, very slightly converging.

Holotype, Stanford University, California, July, 1915 (Melander).

11. Recurved process of ventral pygidial valve broad, the dorsal valve duckbill-shaped; notal setulae weak	T. UNIVERSALIS, MELANDER.
Recurved pygidial process narrow, dorsal valve with short narrow termination; notal setulae evident	T. AGENS, MELANDER.
12. Four pronounced scutellar bristles; abdomen bristly before pygidium; halteres black	T. TETRACHÆTA, nov. sp. (1). 13.
Lateral scutellar bristles reduced	
13. Halteres and palpi yellow; arista three times antennal length	T. TETRACHÆTA, nov. sp. (1). 14.
Halteres dull yellow to black; arista four times antennal length	
14. Third vein much stronger than fourth and clouded	T. RUFICORNIS nov. sp. (2).
Third and fourth veins equal, infumation of wings uniform except for the paler base	T. CALVA, MELANDER.
15. Wings hyaline	T. APICIS, Williston. 16.
Wings more or less infumated	
16. End process of pygidium leaf-shaped; halteres yellowish; hind femora yellow and brown; palpi white ♂, blackish ♀ . . .	T. EPIBOSCA, nov. sp. (3).

(1) **Tachyempis tetrachæta**, nov. sp. — Male. Length 1.75 mm. Front broadly V-shaped, pollinose, two pairs of ocellar bristles, vertical bristles small, occiput completely coated with brown; basal antennal joint yellowish, apical black, arista relatively strong, twice the antennal length; palpi white, elongate ovoid, apical seta long and whitish. Notum dusted with brownish, bristles stronger than usual, humeri large, four notopleural, the center two strong, acrostichal and dorsocentral rows of setulae evident though small, the last two dorsocentrals setiform, four equal and strong scutellar bristles. Abdomen subshining, its hairs stronger than usual, the last segment bristly, pygidium large, the recurved terminal process rather broad. Coxæ and front legs yellowish, posterior legs yellowish brown, front femora robust, anterior femora with usual flexor setulae. Halteres and calyptères black. Wings rather broad, with slight and uniform infumation, veins strong and blackish, costal sections proportioned about 1 : 0.6 : 0.6 : 0.3, first posterior cell ending at apex of wing, about five-sixths as wide as the submarginal at end of second vein, third and fourth veins apically parallel, distance between crossveins less than the length of the posterior crossvein, sections of fifth vein nearly 2 : 3.

Holotype, Concepcion, Chile (R. Thaxter), deposited in the Museum of Comparative Zoology of Harvard University.

(2) **Tachyempis ruficornis**, nov. sp. — Male. Length 1.9 mm. Body black, legs, antennæ, mouthparts, hairs, bristles and halteres yellowish, wings lightly infumated along the veins. Occiput shining, very lightly dusted, almost glabrous; front narrowly V-shaped, the ocelli rather large, ocellar bristles large. Antennæ reddish-yellow, short, the two joints of equal size, the outer globular, with subdorsal arista, which is three times the length of the antennæ. Proboscis brownish, short and vertical; palpi short, narrow, tapering, white and with a long white terminal bristle. Mesonotum polished black, a little dusted anteriorly; one humeral, some supraalar, a double row of about five acrostichal, and about seven dorsocentral bristles present, scutellum with four bristles, the inner pair long; pleuræ finely scabrous so as to appear subshining: abdomen polished. Legs including the coxæ testaceous, the outer half of the hind femora infuscated, the front femora strongest, setulae of middle femora long and fine. Halteres pale yellow, including the base. Wings narrow, slightly infumated except at the base, the darkening a little stronger along the third vein, the second section of the costa slightly longer than the third, curve of the second vein uniform, the marginal cell two-thirds as wide as the submarginal cell, third and fourth veins subparallel, the first posterior cell widest before its tip, anterior crossvein two-thirds as long as the posterior, the distance between the crossveins one-half greater than the length of the posterior crossvein, fifth vein not angulate at the posterior crossvein, its sections subequal.

One specimen collected by Dr. P. P. Calvert, March 4, 1910, at Cachi, Costa Rica, at 3450 ft altitude.

(3) **Tachyempis epibosca**, nov. sp. — Male. Length 1.75-2 mm. Occiput lightly dusted, lower orbits polished, front very broad above, vertical bristles smaller than the ocellar pair, sometimes minute, ocelli moderately large; antennæ brownish, arista thin and four times the antennal length or even longer; palpi narrow, white, with three small apical white setæ, proboscis black. Thorax completely dusted with brownish, pleuræ not becoming cinereous on lower part, the single notopleural and scutellar bristles black, the supra-alar setulae brown, seriate setulae of notum developed though small, posterior dorsocentral not developed, no humeral. Abdomen dull piceous, pygidium large, elongate, very complex when open, the recurved terminal process ending in a broad leaf-like expansion, no specialized setæ. Coxæ and legs pale yellow, the front tibiae and apical half of hind femora brownish, legs elongate, front femora setose below, middle femora with a row of short setæ below, the basal seta long and yellow, middle trochanters with two or three small black spinules. Halteres sordid yellow, the root yellowish, calyptères and fringe dark. Wings elongate, somewhat infumated over basal half of third vein, veins brown, sections of costa proportioned 1 : 1.3 : 1.6 : 0.5, fourth vein ending beyond wing-tip, sections of fifth vein 3 : 4.

Female, palpi blackish.

Three males and five females, La Suiza de Turrialba, Costa Rica (P. Schild).

End process slender; halteres black; hind femora mostly black; palpi ♂ black. T. HALTERATA, nov. sp. (1).

Geographical distribution.

1. *T. agens*, Melander, Psyche, Vol. 17, p. 59, f. 2, 19 [1910] (*Tachydromia*). Washington.
2. *T. apicis*, Williston, Trans. Ent. Soc. Lond. Vol. 3, p. 442, pl. 16, f. 167 [1896] (*Drapetis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 209 [1902] (*Drapetis*); Bezzi, Nova Acta Akad. Nat. Halle, Vol. 91, p. 397 [1909] (*Ctenodrapetis*). insularis, Melander, Psyche, Vol. 17, p. 58, f. 17 [1910] (*Tachydromia*).
3. *T. calva*, Melander, Psyche, Vol. 17, p. 58, f. 18 [1910] (*Tachydromia*). Georgia.
4. *T. capnodes*, nov. sp. Costa Rica.
5. *T. cinerea*, nov. sp. New Mexico.
6. *T. dichroa*, Bezzi, Denkschr. Med. Nat. Ges. Jena, Vol. 13, p. 183 [1908] (*Tachista*). S. W. Africa.
7. *T. epibosca*, nov. sp. Costa Rica.
8. *T. gagatina*, nov. sp. Costa Rica.
9. *T. halterata*, nov. sp. Costa Rica.
10. *T. hyalipennis*, Macquart, Dipt. N. France, Vol. 3, p. 92 [1827] (*Tachydromia*); Hist. Nat. Dipt. Vol. 1, p. 351 [1834] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 7, p. 96 [1838] (*Tachypeza*). W. Europe.
11. *T. longispina*, nov. sp. West Indies.
12. *T. nervosa*, nov. sp. California.
13. *T. nigra*, nov. sp. Costa Rica.
14. *T. pictipes*, nov. sp. — Pl. 4, Fig. 35. Costa Rica.
15. *T. ruficornis*, nov. sp. Costa Rica.
16. *T. Schildi*, nov. sp. Costa Rica.
17. *T. simplicior*, Wheeler & Melander, Biol. Centr. Amer. Dipt. Suppl. p. 375 [1901] (*Phoneutrisca*); Melander, Trans. Ann. Ent. Soc. Vol. 28 p. 205, f. 6 [1902] (*Phoneutrisca*); Psyche, Vol. 17, p. 57, f. 15 [1910] (*Tachydromia*). Mexico.
18. *T. tetracheta*, nov. sp. Chile.
19. *T. universalis*, Melander, Psyche, Vol. 17, p. 60 f. 16 [1910] (*Tachydromia*). United States.
20. *T. vitripennis*, Bezzi, Denkschr. Med. Nat. Ges. Jena, Vol. 13, p. 182 [1908] (*Tachista*). S. Africa.

6. GENUS CHARADRODROMIA, NOV. GEN.

Characters. — Rather robust, opaque pollinose species measuring one to two millimeters. Head narrower than the thorax, globular, occipital setulae sparse, two vertical bristles, the inner pair convergent, two divergent ocellar bristles; eyes well separated, the lower hind margin shallowly excised,

(1) *Tachyempis halterata*, nov. sp. — Male. Length 2 mm. Black, occiput lightly dusted over all, front relatively narrow, the sides parallel above the front ocellus where the front is no wider than the antennal length, vertical bristles a little smaller than the ocellar pair, both black, ocelli moderate in size; antennae brown, arista thin, four times the antennal length, brown at base, otherwise yellowish; palpi black, elongate oval, the outer face glistening white in certain lights. Thorax dusted with dark brown, the pleurae becoming grayish brown toward coxae, the two notopleural, single supra-alar and four scutellar bristles black, about six acrostichal and seven dorsocentral setulae. Abdomen black, with brown coating, pygidium rather long, the apical process with long, thin curved termination which bears four thin setae on the right side. Legs rather slender, coxae and anterior legs largely yellowish, the femora brown above knees, hind femora black except at base, hind tibiae black at base merging to yellow at apex, last joint of all tarsi brown, no flexor setae or thorns present. Halteres, calypteres and fringe black. Wings strongly infumated, paler between humeral crossvein and origin of second vein and along hind margin, veins black, costal sections proportioned about : 1 : 1 : 0.4, fourth vein ending beyond wing-tip, the first posterior cell about two-thirds as wide as the submarginal opposite end of second vein, sections of fifth vein 5 : 6.

Holotype, La Suiza de Turrialba, Costa Rica (P. Schild).

antennal emargination small, facets nearly uniform, bare; sides of front diverging, separated at the antennæ as much as the length of the outer antennal joint, face short, a little narrower than the front, epistome shining; antennæ two-jointed, the outer joint oval, with long coarse terminal arista; proboscis incurved, palpi small, oval, bisetose. Thorax quadrate, notum coarsely hairy, one humeral, one posterior dorsocentral, apical scutellars crossed, pleuræ completely pruinose; abdomen subshining, terminal styles of female small. Legs hairy but devoid of distinct bristles, the front femora enlarged. Costa continuing to fourth vein, first section hairy, basal seta present, first vein ending at middle of wing, second vein ending midway between first and third, basal cells rather small, first somewhat surpassing the second, posterior crossvein oblique, the lower-outer angle of the second basal cell about seventy degrees, a vague trace of the reflexed anal crossvein visible.

Genotype: *C. microphona*, nov. sp.

This genus in venation and general appearance suggests *Chersodromia*, but the legs lack the characteristic bristles of that group. The specialized humeri suggest *Tachyempis*, but the approximation of the crossveins preclude kinship to *Tachyempis* and its near relatives. It is regrettable that the male has not been discovered. Two species are here recorded.

Geographical distribution.

- | | |
|---|-------------|
| 1. <i>C. microphona</i> , nov. sp. (1). | Washington. |
| 2. <i>C. syletor</i> , nov. sp. (2). | Washington. |

7. GENUS COLOBONEURA, MELANDER

Coloboneura, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 299 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248, 261 (1903); Bezzi, Ann. Mus. Hungar. Vol. 2, 321 (1904); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Kertéssz, Cat. Dipt. Vol. 6, p. 143 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 526 (1910); Melander, Psyche, Vol. 17, p. 49 (1910).

Characters. — Thick-set, bristly, opaque pollinose species of three to five millimeters. Head globular, higher than broad, broader than deep, densely pollinose, the occiput large below and setose, two pairs of vertical, one pair of diverging ocellar and one of cruciate preocellar bristles. Eyes small, in both sexes widely separated on the face and less so on the front, face convex, cheeks one-third the eye-height with nearly straight lower margin, clypeus pollinose, semicircular; eyes with very short microscopic pubescence, the facets uniform, hind margin broadly excised below and lower angle round. Antennæ contiguous, plainly three-jointed, the second joint with a crown of setulae, the third joint short

(1) **Charadrodromia microphona**, nov. sp. — Female. Length 1.25 mm. Piceous black, head and thorax entirely overlaid with fine cinereous pollen, bristles and hairs brown; abdominal hairs sparse. Antennæ brown, arista twice the length of the outer antennal joint; palpi yellow, proboscis black. Legs yellowish, the posterior coxae and the femora more or less tinged with light brownish, apex of tarsi somewhat dusky, a small preapical seta on hind tibiæ. Wings hyaline, veins yellow, fourth vein ending at wing-tip, first posterior cell widest at margin, sections of fifth vein proportioned 2 : 3.

Seven specimens, Bickleton, Washington, 26 June, 1917 (Melander).

(2) **Charadrodromia syletor**, sp. nov. — Female. Length 2 mm. Jet black in ground-color, the head overlaid with dust, brownish above and cinereous below, mesonotum lightly brown-dusted, pleuræ completely gray-pruinose. Antennæ black, arista twice the length of the antenna, palpi blackish. Notal hairs shorter than in the preceding species; abdomen shining black, terminal styles slender and black. Legs black, hairs yellowish, front tibiæ thicker than posterior pairs. Wings brownish, veins brown, first posterior cell widest at middle, crossveins meeting, sections of fifth vein 5 : 6.

Holotype, Adna, Washington, 10 July, 1917 (Melander).

ovate with a subapical slender arista directed somewhat downward, its basal joint very short followed by a slightly thickened rudimentary middle joint which may not be evident. Proboscis conical, inflexed, two-thirds the height of the head; palpi vertical, oval, setulose and with apical bristle. Thorax quadrate, densely pollinose, not pubescent but setulose and bristly, one humeral, one posthumeral, three notopleural, two supraalar, one postalar, five dorsocentral, and four scutellar bristles; pleuræ pollinose, even the metapleuræ bare but the propleuræ with a bristle above and one below and the sternopleuræ with two strong intercoxal bristles. Abdomen pollinose, setulose, consisting of eight segments, with two dorsal and several lateral pittings to each segment; hypopygium relatively small. Legs robust, coxae setose, the front pair strong; femora stout, setulose and bristly, the hind pair bristly above and in front as well as below; all the tibiae with three extensor bristles but the hind ones with additional bristles in several rows, front tibiae with two apical, posterior tibiae with one apical bristle, that of the hind legs short but strong; tarsal segments becoming depressed distally, hind metatarsi setulose, pulvilli very large. Wings rather oblong and narrow, opalescent, the costa evanescent before the fourth vein, its first section long, extending beyond the middle of the wing, with strong basal bristle and microscopically setulose, the second basal cell a little shorter than the first, no anal cell but the outer portion of the sixth vein faintly visible as a fold in the wing.

Type species: *C. inusitata*, Melander (Pl. 7, Fig. 10). *Nubifera*, Coquillett, is a true *Coloboneura* as verified by an examination of the type. The species frequent the dry sands of the seashore, over which they run with great agility but do not take readily to the wing although they are able to fly well.

Geographical distribution.

1. *C. argyropalpa*, Bezzii, Ann. Mus. Hungar. Vol. 2, p. 358, f. 7 (1904). New Guinea.
2. *C. exquisita*, Malloch, Ent. News, Philad. Vol. 34, p. 5 (1923). Maryland.
3. *C. hirta*, Walker, Ent. Mag. London, Vol. 3, p. 180 [1836] (*Tachypesa*); Europe, N. Africa. Ins. Brit. Vol. 1, p. 137, pl. 5, f. 5 [1851] (*Chersodromia*); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 119 [1907] (*Chersodromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 227, f. 125, 126 [1910] (*Chersodromia*).
4. *C. ornatipes*, Bigot, Bull. Soc. Zool. France, Vol. 16, p. 277 [1891] (*Chersodromia*); Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 44 [1908] (*Chersodromia*). Canary Isl.
5. *C. inusitata*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 230, 344, f. 47-49 (1902); Slosson, Ent. News, Philad. Vol. 14, p. 268 (1902). — Atlantic Coast of United States.
Pl. 7, Fig. 10.
6. *C. nubifera*, Coquillett, Dipt. Commander Isl. p. 343 [1899] (*Tachydromia*); Bering Sea. Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 342 [1902] (*Tachydromia*); Coquillett, Proc. Ent. Soc. Wash., Vol. 5, p. 265, note (1903); Melander, Psyche, Vol. 17, p. 52 (1910).

8. GENUS CHERSODROMIA, WALKER

Chersodromia, Walker, Ins. Brit. Vol. 1, p. 137 (1851); Rondani, Dipt. Ital. Vol. 1, p. 147 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 564 (1857); Schiner, Fauna Dipt Austr. Vol. 1, p. 96 (1862); Bigot, Bull. Soc. Zool. France, Vol. 16, p. 277 [1891] (*Chersodromia*); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 43 (1902); Coquillett, Proc. Ent. Soc. Wash., Vol. 5, p. 247, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Kertész, Cat. Dipt. Vol. 6, p. 144 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 522 (1910); Lundbeck,

Dipt. Danica, Vol. 3, p. 274 (1910); Melander, Psyche, Vol. 17, p. 49 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45, 89 (1910).

Characters. — Rather robust, opaque pollinose species, measuring from one to three millimeters. Head narrower than the thorax, globular, the occiput flattened hemispherical and rather finely and sparsely setulose; eyes well separated on the front and face, the facets uniform, microscopically hairy, cheeks about one-sixth the eye-height, but somewhat variable in width, clypeus quadrate, pollinose; two pairs of verticals, one of diverging ocellar and one of converging preocellar bristles. Proboscis conical, inflexed, one-half the head-height; palpi broadly oval, sparsely pubescent and with an apical bristle. Antennæ contiguous, apparently two-jointed as the first joint is rudimentary and fused with the second, the third joint short oval, more or less pubescent, with a subdorsal process forming a pedicel to the geniculate pubescent arista. Thorax opaque pollinose, setulose rather than pubescent, usually with the following bristles prominent: one humeral, one posthumeral, three notopleural, four dorsocentral, three supraalar, one postalar, two or four scutellar, but any of these may be absent; pleuræ largely pollinose, usually with a shining area on the lower part of the sternopleuræ, neither bristly nor hairy. Abdomen pollinose, consisting of seven segments plus the hypopygium, which is large, deformed, twisted to the right and formed of various forcipate valves over a triangular under-piece; the individual segments with lateral pittings. Legs rather stout, setulose, the front femora a little thickened, all the femora with a preapical bristle, hind tibiæ with several series of bristles beyond the middle, pulvilli large. Wings translucent and never hyaline, costa extending to the fourth vein, first vein ending at the middle of the wing, the first section of the costa usually loosely ciliate and with a strong basal bristle, second basal cell a little shorter than the first, no anal cell.

Type species: *Ch. arenaria*, Haliday, has smaller wings than usual, the wings scarcely exceeding the abdomen. The species of *Chersodromia* frequent mainly the sandy seashore, but some may occur on the shores of fresh water. The darker species seem to prefer the wet sand near the water's edge; the gray species occur higher up on the dry sand and even run up and down the beach grass. The developmental stages are unknown.

TABLE OF THE KNOWN SPECIES OF *CHERSODROMIA*

1. Yellow, with black bristles; eyes meeting beneath the antennæ	CH. LUTESCENS, Bezzii.
Black species; eyes separated more or less on the face	2.
2. Wings not longer than the abdomen; legs black	CH. ARENARIA, Haliday.
Wings surpassing the abdomen	3.
3. Hind tibiæ with one pair of bristles at the middle of their extensor side; bristles of notum yellowish; costal hairs pale	CH. NANA, Coquillett.
Hind tibiæ with several bristles along extensor side	4.
4. Middle tibiæ with flattened scale-like flexor bristles; no dorsocentrals; legs brown, knees yellow	CH. GRATIOSA, Becker.
Middle tibiæ without flexor scales; dorsocentrals usually present.	5.
5. Legs and halteres black, wings dusky, body dark	6.
Legs and halteres more or less yellow, wings whitish, body lighter gray pollinose.	7.
6. Middle tibiæ with extensor bristles; costa with very short setulæ, its second and third sections equal	CH. DIFFICILIS, Lundbeck.
Middle tibiæ without extensor bristles; costa ciliate, its second section shorter	CH. CURSITANS, Zetterstedt.

7. Thorax without discal bristles, its hairs pale	8
Thorax with discal bristles, including one posthumeral	9.
8. Scutellum with two bristles; second antennal joint minute; legs yellow. <i>Ch. brevicornis</i> , Meijere.	
Scutellum without bristles; second antennal joint equal to the third; femora brown	<i>Ch. incana</i> , Walker.
9. Costal hairs short; middle tibiae with extensor bristle	<i>Ch. speculifera</i> , Walker.
Costal hairs long	10.
10. Middle tibiae with two, hind tibiae with seven extensor bristles; hairs black	<i>Ch. houghii</i> , Melander.
Middle tibiae without long extensor bristles, hind tibiae with five; hairs pale	<i>Ch. beckeri</i> , nov. sp. (1).

Geographical distribution.

1. <i>Ch. arenaria</i> , Haliday, Ent. Mag. London, Vol. 1, p. 161 [1833] (<i>Tachypesa</i>); Walker, ibidem, Vol. 3, p. 180 [1835] (<i>Tachypesa</i>); List Dipt. Brit. Mus. Vol. 3, p. 507 [1849] (<i>Tachydromia</i>); Ins. Brit. Vol. 1, p. 138 (1851); Lundbeck, Dipt. Danica, Vol. 3, p. 281, f. 129 (1910). <i>brevipennis</i> , Zetterstedt, Fauna Ins. Lappon. p. 548 [1838] (<i>Tachypesa</i>); Dipt. Scand. Vol. 1, p. 323 [1842] (<i>Tachypesa</i>); ibidem, Vol. 13, p. 499 [1859] (<i>Tachypesa</i>).	Europe.
2. <i>Ch. beckeri</i> , nov. sp. <i>incana</i> , Becker (not Walker), Zeitschr. Hym. Dipt. Vol. 7, p. 119 (1907).	Germany.
3. <i>Ch. brevicornis</i> , Meijere, Tijds. v. Ent. Vol. 1, p. 177, pl. 4, f. 6, 7 (1907).	C. Europe.
4. <i>Ch. cursistans</i> , Zetterstedt, Vet. Akad. Förh. Stockholm, 1819, p. 82 [1819] (<i>Empis</i>); Fallen, Dipt. Suec. Suppl. Vol. 1, p. 7 [1823] (<i>Tachydromia</i>); Zetterstedt, Dipt. Scand. Vol. 1, p. 322 [1842] (<i>Tachypesa</i>); Walker, Ins. Brit. Vol. 1, p. 137 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862); Lundbeck, Dipt. Danica, Vol. 3, p. 278, f. 127 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 89 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 75 (1913).	Europe.
5. <i>Ch. difficilis</i> , Lundbeck, Dipt. Danica, Vol. 3, p. 280, f. 128 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 75 (1913).	N. Europe.
6. <i>Ch. grataiosa</i> , Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 41 (1908).	Canary Islands.
7. <i>Ch. Houghii</i> , Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 206, pl. 5, f. 2, 3 [1902] (<i>Stilpon</i>); Aldrich, Cat. N. Amer. Dipt. p. 314 [1905] (<i>Coloboneura</i>); Melander, Ent. News, Philad. Vol. 17, p. 370 (1906).	E. United States.
8. <i>Ch. incana</i> , Walker, Ins. Brit. Vol. 1, p. 138 (1851); Strobl, Verh. Zool.-bot. Ges. Wien. Vol. 59, p. 179 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 282 (1910).	Ireland.

(1) ***Chersodromia beckeri*, nov. sp.** Syn.: *incana*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 119 (1907). — Female. Length 1.3 mm. Blackish, densely overlaid with fine gray pollen, the hairs and bristles golden. Face and front equally broad, cheeks one-sixth the eye-height: two pairs of vertical, one of ocellar and one of preocellar bristles; proboscis, palpi and antennae yellow. Thorax with one humeral, one posthumeral, one notopleural, two supraalar, three dorsocentral, one postalar and two scutellar bristles, all long, slender and yellowish, acrostichal setulae biserrate; sternopleurae largely shining. Legs yellowish, the tarsi a little dusky toward the end, middle tibiae with two short extensor bristles, hind tibiae with about five extensor bristles, in two series. Wings opalescent, veins whitish, costa with long basal bristle, its first segment with moderately long, closely placed pale cilia, the second, third and fourth sections of the costa proportioned 1 : 2 : 1.

Two specimens, seabeach at Swinemünde, Baltic Sea, received from Dr. Th. Becker as *incana*. The species is a close relative of *Houghii*, which has stronger tibial bristles; the male of *Houghii* lacks the extensor bristles on the middle tibiae and has the flexor surface of this joint closely black setulose.

9. *Ch. lutescens*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 358 (1904). New South Wales.
 10. *Ch. nana*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 166 [1903] (*Coloboneura*). Florida.
 11. *Ch. speculifera*, Walker, Ins. Brit. Vol. 1, p. 138 (1851); Bezzi, Bull. Soc. Ital. C. & S. Europe.
 Vol. 30, p. 151 (1899); Becker, Zeitschr. Hym. Dipt. p. 119 (1907).
 12. *Ch. species*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 491 (1912). Formosa.

9. GENUS THINODROMIA, MELANDER

Thinodromia, Melander, Ent. News Philad. Vol. 17, p. 370 (1906); Williston's Man. N. Amer. Dipt. p. 222 (1908); Kertész, Cat. Dipt. Vol. 6, p. 169 (1909); Melander, Psyche, Vol. 17, p. 49 (1910); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 614 (1910).

Characters. — Robust, opaque gray pollinose, black-bristly species with aborted wings. Head globular, occiput flattened hemispherical, not very bristly; two pairs of vertical bristles, one of diverging ocellar and one of cruciate preocellar; eyes widely separated, facets uniform, bare, the incision at the antennæ strong, cheeks one-fourth the eye-height. Antennæ three-jointed but the first joint very small, no setæ beneath, third joint round, apically pubescent, with dorsal projection bearing the two-jointed geniculate arista, the basal joint of the arista long, solid, the outer joint slender and pubescent. Proboscis strong, very robust, inflexed, the palpi elliptical, setulose on the anterior surface and with a strong apical bristle. Thoracic bristles strong, one humeral, one notopleural, two supraalar, three dorsocentral, one postalar, a double row of sparse acrostichal setulæ, scutellum with two cruciate apical and two smaller lateral bristles; pleuræ not hairy. Abdomen with eight segments, the individual segments with a single pit on each side and with several larger ones at the extreme lateral edge, the last segment of the female compressed wedge-shape, with very short oval styles, in the male the abdomen is very robust, the bristles becoming longer posteriorly, those of the seventh segment nearly as long as the last four segments; pygidium large, globose, closed, somewhat asymmetrical and twisted to the right, the small apical organs nearly dorsal. Legs short, robust, with black setulæ, those of the under side of the front femora and of the outer apical side of the hind femora more prominent, hind tibiæ with two extensor bristles, one before the middle and one preapical, outer half of the hind tibiæ with about six bristles irregularly placed, a pair of short apical bristles on the front tibiæ, tarsi flattened distally. Wings very small, triangular in outline, horizontally extended over the abdomen, reaching only to the third segment, the costal margin straight, stiff and minutely setulose, first vein ending near the apex of the wing, crossveins close together, the second basal cell a little shorter than the first, both extending to beyond the middle of the wing, no trace of an anal cell, hind margin entirely bare.

Genotype : *T. inchoata*, Melander (Pl. 7, Fig. 71). The two known species are inhabitants of sandy beaches on the Pacific coast of North America. They are utterly unable to fly, but run nimbly over the sand and driftwood.

Geographical distribution.

1. *Th. inchoata*, Melander, Ent. News Philad. Vol. 17, p. 370, fig. (1906). — Pacific Coast of
 North America.
Pl. 7, Fig. 71.
 2. *Th. parallela*, nov. sp. (1). Pacific Coast of
 North America.

(1) **Thinodromia parallela**, nov. sp. — Length 2.5 mm. A larger species than *inchoata* differing in the structure of the wings. The wings are narrower in length, measuring three times as long as wide, the first five veins parallel, equally strong and reaching the wing-margin, the third vein ending at the extreme apex, the costa curving around the apex of the wing and continuing to beyond the end of the third vein. The costal setulæ are longer and obliquely erect. The blackish color fills the apical third of the wing. In *inchoata* the wings are twice as long as broad; the third vein bends forward so as to end in the first vein; the second and fourth veins are obsolete; the costa becomes thin at the apex of the wing; the costal

10. GENUS HALSANALOTES, BECKER

Halsanalotes, Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 41 (1902); Bezzı, Ann. Mus. Hungar. Vol. 2, p. 321 (1904); Kertész, Cat. Dipt. Vol. 6, p. 143 (1909); Melander, Psyche, Vol. 17, p. 49 (1910); Bezzı, Ann. Mus. Hungar. Vol. 10, p. 453 (1912).

Characters. — Head globular, the occiput flattened-hemispherical, eyes pubescent, widely separated on the front and face, cheeks one-fourth the eye-height, face slightly convex; two pairs of vertical bristles, one of ocellar and one of diverging preocellar; antennæ rather widely separated, very short, the outer joint triangular, pubescent, with a long hairy dorsal arista; proboscis very short, vertical, palpi oval, with small apical bristle. Thorax not broader than the abdomen, pollinose, with short pubescence but no discal bristles, two scutellar, one humeral, one notopleural and one supraalar bristle. Abdomen opaque pollinose, comprising five segments and the hypopygium which is terminal, blunt and provided with several lamellæ and a short, erect, dorsal appendage. Legs short, not thickened, simple, front tibiæ with two apical bristles, middle tibiæ with one apical, hind tibiæ with a row of four or five extensor bristles on the apical half, last tarsal joint widened. Wings oval, basal cells equal, third and fourth veins diverging, anal cell weakly indicated, visible only when obliquely viewed.

Genotype: *H. amaurus*, Becker, is the only species known. It measures 0.5 to 0.75 mm. in length. Two species described by Bezzı as *Halsanalotes* are here placed in the new genus *Micrempis*.

Geographical distribution.

1. *H. amaurus*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 42, 64, pl. 4, N. Africa.
f. 1-3 (1902).

11. GENUS MICREMPIS, NOV. GEN.

Characters. — Very small, nearly shining, dark colored species with pale legs. Head spherical, the upper occiput a little flattened; front V-shaped, the eyes nearly meeting just above the antennæ, a pair each of vertical, diverging ocellar and cruciate preocellar bristles, excision of the eye at the antennæ angular and small, facets uniform, face more or less obliterated below the antennæ, broadening toward the linear cheeks. Antennæ short, two-jointed, the basal joint with a small or with no seta beneath, the outer joint round, compressed, with a dorsal prolongation forming the pedicel of the geniculate, pubescent arista. Proboscis short, vertical but a little inflexed, not thickened at the base, palpi broadly oval, with apical bristle, Thorax convex, broader than the head or abdomen, shining, no discal bristles, not pubescent, but with hairs in the dorsocentral and acrostichal rows, bare on the flattened area in front of the scutellum, humeral bristle present or absent, two notopleural, two scutellar, no postalar bristle; pleuræ shining or more or less pollinose. Abdomen opaque, nearly bare, comprising seven segments and the hypopygium. Legs normal, rather short, front femora strongest, no bristles spines or spurs, the hairs inconspicuous. Wings with strong bristle at the base of the costa, second vein short, ending near the middle of the wing, basal cells small, equal, third and fourth veins diverging and with an anterior swing, no anal cell or veins.

Type species: *M. nana*, nov. sp.

setulae are short and more appressed; and the dark color is concentrated about the third vein and the lower angle of the second basal cell. None of the veins of *inchoata* reach the hind margin of the wing.

This species was found by the author in numbers on the seashore of Fossil Island of the Sucia Islands off the coast of Washington, July 16, 1909, and again on the dry sands about driftwood on the ocean beach at Long Beach, Washington, May to August, various years.

Geographical distribution.

1. *M. *eocenica*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 89, 96, pl. 4, f. 3-5 Baltic Amber.
[1908] (*Phoneutisca*).
 2. *M. fuscipes*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 490 [1912] (*Halsanalotes*). Formosa.
 3. *M. minuta*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 339 [1902] New Mexico.
(. *Stilpon*).
 4. *M. nana*, nov. sp. (1). — **PI. 4, Fig. 32.** Texas.
 5. *M. obliqua*, nov. sp. (2). Iowa.
 6. *M. setifrons*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 357, 41 [1904] Australia.
(. *Halsanalotes*).
 7. *M. *suspiciosa*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 89, 98, pl. 4, f. 6, 7 [1908] (*Phoneutisca*). Baltic Amber.
 8. *M. testacea*, nov. sp. (3). Virginia.
 9. *M. varipes*, nov. sp. (4). Peru.

(1) ***Micrempis nana*, nov. sp. (PI. 4, Fig. 32).** — Male. Length 0.6 mm. Dark brownish, legs yellow, the tarsi darker apically. Vertex and lower occiput lightly cinereous pollinose; antennæ brown, the outer joint compressed, circular, with four apical hairs and with a dorsal process (the pedicel of the arista), the arista then bent down, rather thickly pubescent. Proboscis and palpi yellowish. Thorax very lightly dusted, nearly polished, not pubescent, one row of sparse dorsocentral hairs. Halteres black, with yellow stems. Legs simple, not bristly, their pubescence inconspicuous. Wings subhyaline, veins brown, the second, third and fourth sections of the costa proportioned 1 : 4 : 1, crossveins meeting, sections of the fifth vein proportioned 1 : 2.

Two specimens, Austin, Texas.

(2) ***Micrempis obliqua*, nov. sp.** — Female. Length 1.3 mm. Occiput black, lightly dusted, front V-shaped, at the bottom scarcely one-half the diameter of the first antennal joint; face triangular, the eyes barely separated just beneath the antennæ, cheeks narrow; palpi whitish, not hairy, proboscis short, brown; antennæ two jointed, without a seta beneath, the basal joint yellow, the outer joint blackish, sub-triangular, its upper edge straight, its lower edge rounded, one-fourth longer than broad, apically pubescent, the terminal arista one-fourth longer than the antenna, microscopically pubescent; vertical bristles twice as long as the widely separated ocellars, brown. Dorsum shining black, pleuræ except the polished sterno-pleuræ opaque gray pollinose, mesonotal hairs pale, regularly placed on the anterior two-thirds, posteriorly bare except for the three dorsocentrals, four rows of about seven acrostichals, the middle rows diverging behind, humeral bristle erect, two scutellars. Abdomen brown. Legs entirely yellow, only the middle femora with preapical bristle, no extensor bristles or apical spur on the hind tibiae. Halteres brown, calypteres with four cilia. Wings clear hyaline, veins narrow brownish, the first four veins curving forward and diverging from each other, the third section of the costa two and one-half times as long as either the second or fourth section, submarginal cell apically large, at the tip of the first vein one-third as wide as the marginal cell and opposite the tip of the second vein more than twice as wide as the first posterior, this cell gradually widens to its apex, posterior crossvein strongly oblique, the lower apical angle of the second basal cell about fifty degrees, on the fourth vein the two basal cells of equal length, sections of the fifth vein equal.

One specimen, submitted by Professor R. L. Webster, who found it at Ames, Iowa, July 16, 1908, attacking the apple leaf-hopper, *Empoasca mali*.

(3) ***Micrempis testacea*, nov. sp.** — Female. Length 1.3 mm. Testaceous, including the head, thorax, legs, antennæ and mouthparts; the abdomen and knob of the halteres brown. Front broadly V-shaped, face linear; one pair each of strong vertical, ocellar and preocellar bristles; palpi ovate, with a single black seta, proboscis one-fourth the head-height; outer antennal joint ovate, with subdorsal deflected arista a little over twice the length of the antenna. Thorax round, rather shining, the disc seriatily hairy, with the following moderate blackish bristles: one humeral, two notopleural, one supraalar, one postalar, one intraalar, and two scutellar; upper pleuræ lightly pruinose, lower pleuræ glistening; abdomen opaque, ovipositor short, compressed, its blackish base shining. Wings with a slight yellowish tinge, the first four costal sections proportioned 1 : 0.5 : 0.9 : 0.7, the first four veins uniformly diverging and curving forward, the sections of the fourth vein proportioned 1 : 3, of the fifth vein, 1 : 1.3, crossveins meeting, cilia of the hind margin shorter than the anterior crossvein.

Type, Falls Church, Virginia, Nathan Banks, collector; Paratype, Plummer's Island, Maryland, H. Barber, collector, in U. S. National Museum.

(4) ***Micrempis varipes*, nov. sp.** — Length 1 mm. Body black, legs largely yellow, halteres black, wings clear hyaline, veins narrow and brown. Head lightly pollinose, more shining about the ocelli, front V-shaped, the sides curving inward, eyes subcontiguous below the antennæ; palpi and proboscis black, the latter one-half the height of the head; antennæ black, the outer joint defective. Mesonotum polished, nearly bare, pleuræ lightly pollinose, leaving a circular central spot polished: abdomen subshining, pygidium small, ovipositor short and opaque. Anterior coxæ yellow, the hind coxæ black, front legs yellow, in the female the tarsi and a vague apical spot on the upper side of the femora infuscated, middle legs of the female blackish beyond the middle of the femora, of the male only the outer half of the femora darkened, hind legs

12. GENUS SYMBALLOPHTHALMUS, BECKER

Symballophtalmus, Becker, Wien. Ent. Zeit. Vol. 8, p. 285 (1889); ibidem, Vol. 9, p. 34 (1890); Mitteil. Zool. Mus. Berlin, Vol. 2, p. 42 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257, 261 (1903); Kertész, Cat. Dipt. Vol. 6, p. 145 (1909); Lundbeck, Dipt. Danica, Vol. 3, p. 283 (1910); Melander, Psyche, Vol. 17, p. 49 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 54 (1910).

Macroptera, Becker (not Lioy, 1864, which is *Ula*, Tipulidae), Wien. Ent. Zeit. Vol. 8, p. 80 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 252 (1903).

Characters. — Black, shining, slender, nearly bare species with slender, long legs and cuneate wings. Head spherical, eyes large, the forward facets coarse, broadly contiguous above the antennae thus obliterating the front, but widely separated below the antennae, the face very short, the epistome large, filling in the oral opening to the very narrow cheeks; proboscis slender, shorter than the head-height, inflexed, the palpi minute, slender, setulose. Antennae three-jointed, inserted below the middle of the head, the second joint without lengthened setae, the third joint elongate, straight above, rounded below, long ham-shaped, the two-jointed terminal arista closely and briefly pubescent. One supraalar bristle, six scutellars, sparse biserrate acrostichal setulae present, the hairs of the posterior part of the mesonotum proclinate, of the anterior part reclinate as usual; pleurae shining, the metapleuræ velvety. Last abdominal segments of the female protractile, pygidium globular and complexly forcipate above. Legs simple, the middle femora with biserrate weak setulae and the front tibiae with weak extensor setulae, middle tibiae as long as their femora, front metatarsi of the male with long flexor hairs, no true bristles present, tibiae without apical spur or spines. Wings with long slender base, costa not thickened and with a small basal bristle, the first vein ending beyond the middle of the wing, pedicel of the second and third veins long, crossveins meeting and nearly transverse, anal crossvein and outer part of anal vein present, anal angle very obtusely rounded and shallow.

Genotype : *S. dissimilis*, Fallen, is the unique species. It occurs from the Alps north to central Sweden.

Geographical distribution.

1. *S. dissimilis*, Fallen, Empid. Suec. p. 9 [1815] (*Tachydromia*); Meigen, Syst. Europe.

Beschr. Vol. 3, p. 85 [1822] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 308 [1842] (*Tachydromia*); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 509 [1849] (*Platypalpus*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3010 [1849] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 133 [1851] (*Platypalpus*); Lundbeck, Dipt. Dan. Vol. 3, p. 284, f. 130 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 90 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 75 (1913).

cyanophthalmus, Strobl, Progr. Seitenst. Vol. 14, p. 8 [1880] (*Platypalpus*); Wien. Ent. Zeit. Vol. 10, p. 267 (1891); Mitteil. Nat. Ver. Steiermark, Graz, Vol. 29, p. 123 (1892).

pictipes, Becker, Wien. Ent. Zeit. Vol. 8, p. 80, f. 3-5 (*Macroptera*); ibidem, p. 285 (1889).

of the female entirely black, of the male the basal half of the femora yellow, the remainder of the hind legs blackish brown. Second vein very strongly curving forward, the first four sections of the costa proportioned 1 : 0.5 : 1.2 : 0.6, third vein uniformly diverging from the fourth, basal cells small and of equal extent along the fourth vein, hinder crossvein oblique, sections of the fourth vein, 1 : 3.5, of the fifth vein, 1 : 1.3.

One specimen of each sex, taken by C. T. Brues in grass sweepings along a ditch at Matucana, Peru, July 1, 1913.

13. GENUS MEGAGRAPHÄ, NOV. GEN.

Characters. — Small, stout, pubescent species with broad wings and large basal cells. Head globular, the occiput hemispherical, densely hairy above; eyes nearly contiguous above and beneath the antennæ, separating at the ocelli which are placed up toward the vertex and have small ocellar bristles, on the face the eyes broadly round away about one-third the distance from the antennæ, cheeks linear, pubescence of the eyes dense, facets uniform, hind margin of the eye S-shaped, excision at the antennæ small. Proboscis vertical, one-half the head-height, not thickened at the base, the palpi broadly oval, shorter than the proboscis, pubescent but without apical bristles. Antennæ three-jointed, without bristles, the third joint lanceolate, but flat above, densely pubescent, the arista terminal, shorter than the antennæ, without basal segment, thick with close pubescence. Thorax broadly quadrate, humeri not swollen, densely and uniformly pubescent, the hairs obliquely reclinate, no discal bristles, no humeral, about six short notopleural bristles in a prealar series diminishing in size toward the front, one pair of widely separated prescutellar dorsocentrals, scutellum large, rounded triangular, flat above, with close short pubescence above and below and with six marginal bristles; pleurae bare except for a velvety patch of tomentum on the metapleuræ before the halteres. Abdomen short and broadly conical, comprising seven segments and the lengthened ovipositor which latter terminates in two slender styles, basal ventral segments not chitinized, abdomen hairy especially toward the end, the first segment not broadly swollen at the sides as is usually the case in the pale-colored species of *Drapetis*. Legs slender and short, without bristles but with abundant hairs, no tibial spurs, the hind tibiae seriatel spinulose on the extensor side, last tarsal joint small, pulvilli minute. Wings broad, the anal angle broadly rounded, costa not swollen, first vein ending at the middle of the wing, pedicel of the second and third veins long, basal cells large and of the same length, anal crossvein wanting, but the anal vein present close to the hind margin, no costal bristles but the marginal cilia longer toward the root of the wing.

Type species : *M. (Drapetis) pubescens*, Loew, the only species so far known. I know of but four specimens of this species, all females, from Georgia, New York and Massachusetts.

Geographical distribution.

1. *M. pubescens*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 201 : Cent. 2, No. 57 E. United States.
 [1862] (*Drapetis*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 210,
 340 [1902] (*Drapetis*). — PI. 4, Fig. 33.

14. GENUS STILPON, LOEW

Stilpon, Loew, Neue Beitr. Dipt. Vol. 6, p. 34 (1859); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 125 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Melander, Trans. Ent. Soc. Vol. 28, p. 205 (1902); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 42 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 257, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Kertész, Cat. Dipt. Vol. 6, p. 134 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 609 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 260 (1910); Melander, Psyche, Vol. 17, p. 49 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 46 (1910).

Agatachys, Meigen, Syst. Beschr. Vol. 6, p. 343 (1830); Scudder, Nomencl. p. 10 and Univ. Index, p. 9 [1882] (*Agatachys*); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 504 (1910).

Tetraneurella, Dahl, Sitzgber. Ges. Naturf. Freunde, Berlin, p. 362 (1909).

Characters. — Minute species, scarcely over one millimeter in length. Head large, globular, the eyes densely short-pubescent, enlarged below, obliterating the face, the lower facets large in both sexes, hind margin of the eye curving forward below, front narrow, its sides parallel, ocellar bristles small, two small vertical bristles present, cheeks narrow. Antennæ short, the first joint vestigial, the second joint swollen, with setæ beneath, the third joint very short and round, with a dorsal geniculate arista, or rather, there is a short dorsal process arising from the third joint which bears the single-jointed arista. Proboscis short, vertical, palpi slender, with a strong apical bristle. Thorax broadly round, the dorsum moderately convex, shining or slightly pollinose in front, no discal bristles, but the hairs seriatly arranged to form two acrostichal and one dorsocentral rows, one humeral bristle present, two notopleurals, no prescutellar, two scutellars; pleuræ bare and shining, the metapleuræ velvety. Abdomen without pits, comprising eight segments; the pygidium large and forcipate. Legs short, the front pair robust, in the males of some species the middle legs are ornamented and the hind legs are ciliate with long bristles, middle femora with a preapical bristle, no tibial apical spines. Wings narrow, first and second veins very short, third and fourth veins diverging, anal cell entirely wanting, costal bristle small.

Nomenclature. — Concluding his description of *Tachydromia celeripes*, Meigen stated that he had received « mehrere Exemplare von Herrn von Winthem, unter dem Namen *Agatachys flavipes*. » This induced Coquillett to adopt the name *Agatachys* for this genus. The catalogues have considered *Agatachys* as a manuscript name and as a synonym of *Stilpon*. While it is unfortunate that Loew overlooked Meigen's *celeripes* and Winthem's generic name when he formed the genus *Stilpon*, yet his name is so well known that it seems an unnecessary perversion of the rules to make Meigen the unwitting sponsor of *Agatachys* in order to have it established.

The genus *Tetraneurella* of Dahl represents a curious form with reduced venation. Dahl now believes that his specimens lacking the second vein are dimorphic individuals of *graminum*. The American species, *pleuritica*, here described, also lacks the second vein, but is certainly specifically different from any of the other known species of *Stilpon*. The short V-shaped front suggests *Drapetis*, but the antennæ are two-jointed and bear a dorsal arista, as in *Stilpon*.

Type species of *Stilpon*, *S. graminum*, Fallen (Pl. 4, Fig. 37), by Coquillett's designation in 1903.

SYNOPSIS OF THE AMERICAN SPECIES OF STILPON

1. Pleuræ opaque pollinose except a circular spot filling the sternopleura; second vein wanting; legs yellow, the hind femora brown on the apical third ST. PLEURITICA, nov. sp. (1).
- Pleuræ largely polished; second vein complete, ending in the costa about midway between the first and third veins 2.

(1) ***Stilpon pleuritica*, nov. sp.** — Female. Length 1 mm. Black, the abdomen testaceous, legs yellow, the outer third of the hind femora brown, mesonotum lightly pollinose, pleuræ brown-pollinose except a circular spot covering the sternopleura; second vein wanting. Head large, front relatively short, V-shaped, narrowed below, lightly pollinose like the vertex; antennæ uniformly testaceous; proboscis brown, palpi yellow with a small apical black bristle. Humeral bristle strong, notal hairs very short and sparse; lower part of the metapleuræ shining; one strong preapical bristle on the middle femora; hind metatarsi and last tarsal joint not darkened. Halteres entirely pale yellow. Wings with a slight infumation, the third and fifth veins more strongly bordered, veins brown becoming pale at their base, only a minute basal spur of the second vein remaining, proximal section of the fifth vein slightly shorter than the distal section.

A single specimen from Mrs. Annie T. Slosson, collected in the White Mountains, New Hampshire. This is a *T. transversella* form. The V-shaped front and pollinose pleuræ are different from the characters of the other species of *Stilpon*.

2. Outer half of the hind femora brown, front tibiæ brown except at the base, hind tibiæ wholly brown, last tarsal joint black; abdomen testaceous; veins largely bordered with brown. ST. VARIPES, Loew.
- Legs entirely yellow, except sometimes the hind metatarsi and last tarsal joint brownish 3.
3. Spine-like bristles of the under side of the middle femora ♂ black, spines of the middle tibiæ ♂ stronger near the middle of the tibiæ; upper side of hind femora ♂ setose near the middle; hind metatarsi brown 4.
- Spine-like bristles of the under side of middle femora ♂ yellow; spines of the middle tibiæ ♂ uniform to the end; extensor setæ of the hind femora ♂ not developed, or stronger apically 5.
4. Hind femora not distorted, their lower bristles in two rows, those of the outer half long and conspicuous, upper side of the hind femora with weak bristles only; hind tibiæ externally pectinate; abdomen black. ST. SPINIPES, nov. sp. (1).
- Hind femora ♂ with a strong bend at the middle, before which is a dorsal row of strong curved bristles, the under side with a single row of bristles, denser on the basal half; hind tibiæ not pectinate; abdomen testaceous ST. CURVIPES, nov. sp. (2).
5. Hind tibiæ ♂ with long dense hairs on the anterior face, flexor bristles of the hind femora closely set; two or three preapical

(1) **Stilpon spinipes**, nov. sp. — Male. Length 1.3 mm. Black, the legs yellow, but the last tarsal joint black. Occiput very lightly pollinose, ocellar bristles minute, front subshining, palpi and proboscis yellow, base of the antennæ yellow, the last joint black. Thorax shining, very lightly pollinose in front, acrostichal hairs wanting on the posterior half of the dorsum, one row of seven dorsocentral hairs, scutellum with two bristles. Abdomen black, the last segment with four long black hairs, the pygidium large, forcipate. Front femora with a row of pale flexor bristles, under side of the middle femora with four stiff black spines located near the beginning of the outer third but without pale basal spines, middle tibiæ with a cluster of six contiguous black flexor spines engaging those of the femora and with about five additional black spines occupying the distal third, a strong preapical bristle on the middle femora, hind femora doubly pectinate on the antero-flexor side, the upper row with fourteen pale bristles extending the whole length of the femur, the lower row of eight longer dark bristles occupying the distal half, a row of smaller erect extensor bristles on the basal half of the hind femora, hind tibiæ loosely pectinate on the extensor side, extensor side of the anterior tibiæ and the hind knees a little dusky. Halteres brown. Wings smoky, the second vein ending beyond the middle of the wing, the second, third and fourth sections of the costa subequal, sections of the fifth vein proportioned nearly 1 : 2.

One specimen, Waycross, Georgia.

The species is closely related to *pectiniger*, differing mainly in the structure of the legs. The male of *pectiniger* has a single row of shorter bristles on the antero-extensor edge of the hind femora, the bristles of the hind tibiæ shorter, the flexor spines of the middle femora silky yellow and those of the middle tibiæ forming a uniform row along the distal three-fourths.

(2) **Stilpon curvipes**, nov. sp. — Male. Length 1 mm. Head, thorax and pygidium shining black; abdomen testaceous; legs including the coxae light yellow, the hind metatarsi and last tarsal joint darkened. Basal joint of the antennæ yellow, outer joint broken, mouthparts yellow, front cinereous almost to the ocelli. Hairs of the thorax coarse, pleuræ mostly shining; the inflated sides of the first abdominal segment black and pollinose. Pygidium rather large, pollinose below, with a long slender apical process bearing four long hairs, the last abdominal segment fringed with long loose black hairs. Middle femora with an interrupted flexor row of black thorn-like setulae along the basal three-fifths and with a long preapical bristle, middle tibiæ with a similar but double row along the outer half, hind femora with a strong lateral bend just beyond the middle, before the bend with a row of extensor bristles, beyond it the extensor bristles are irregular, underneath with a row of fourteen coarse bristles, of which eight are before the bend, hind tibiæ ciliate with twelve hairs along the antero-extensor edge. Halteres black. Wings narrowly oval, veins brown, very slightly bordered with brown, the second, third and fourth sections of the costa proportioned 1 : 0.8 : 0.6, fourth vein with a very slight anterior curvature, almost straight, posterior crossvein oblique, forming an angle of fifty degrees with the fifth vein, sections of the fifth vein proportioned 0.8 : 1, marginal cilia two-thirds as long as the posterior crossvein.

One specimen, Wisconsin, July.

- bristles on the middle femora; hind metatarsi brown; veins light brown, wings not clouded; pollen of the front extending to the ocelli ST. PECTINIGER, Melander.
- Hind tibiæ not pectinate; about eight flexor bristles on the hind femora ♂; a single preapical bristle on the middle femora; metatarsi yellow; veins dark, wings lightly clouded in the middle; front pollinose only near the antennæ. ST. PAUCISETA, nov. sp. (1).

Geographical distribution.

1. *St. celeripes*, Meigen, Syst. Beschr. Vol. 6, p. 343 [1830] (*Tachydromia*); C. Europe.
ibidem, Vol. 7, p. 95 [1838] (*Tachypesa*); Neuhaus, Dipt. March.
p. 75 [1886] (*Drapetis*).
flavipes, Winthem, in litt., Meigen, Syst. Beschr. Vol. 6, p. 343 [1840]
(*Agatachys*).
2. *St. curvipes*, nov. sp. Wisconsin.
3. *St. graminum*, Fallen, Empid. p. 15 [1815] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 72 [1822] (*Tachydromia*); Haliday, Ent. Mag. London, Vol. 1, p. 161 [1833] (*Tachypesa*); Walker, ibidem, Vol. 3, p. 180 [1835] (*Tachypesa*); Meigen, Syst. Beschr. Vol. 7, p. 95 [1838] (*Tachypesa*); Zetterstedt, Dipt. Scand. Vol. 1, p. 320 [1842] (*Tachypesa*); Walker, List Dipt. Brit. Mus. Vol. 3, p. 507 [1849] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3011 [1849] (*Tachypesa*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 59 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 136 [1851] (*Drapetis*); Loew, Neue Beitr. Dipt. Vol. 6, p. 43 (1859); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 154 [1861] (*Tachypesa*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 95 [1862] (*Drapetis*); Raddatz, Arch. Ges. Freunde Naturg. Mecklemb.-Rostock, Vol. 27, p. 44 [1873] (*Drapetis*); Lundbeck, Dipt. Dan. Vol. 3, p. 262, f. 118, 119 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 87 (1910); Dahl, Fauna Chorin, 465 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 71 (1913). — **Pl. 4, Fig. 37.**
var. *Beckeri*, Dahl, Sitzber. Ges. Naturf. Freunde, Berlin, p. 362 [1909] (*Tetra-neurella*); Fauna Chorin, p. 465 (1912). Germany.
4. *St. lunata*, Walker, Ins. Brit. Dipt. Vol. 1, p. 136, pl. 5, f. 3 [1851] C. Europe.
(*Drapetis*); Loew, Neue Beitr. Dipt. Vol. 6, p. 43 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 95 [1862] (*Drapetis*); Mik, Wien. Ent. Zeit. Vol. 19, p. 145, note (1900).
femorata, Heeger, Sitzber. Akad. Wiss. Wien, Vol. 9, p. 779-781, pl. 55, f. 1-5 [1852] (*Hemistrodromia*).
5. *St. obscuripes*, Adams, Kansas Univ. Sc. Bull. Vol. 3, p. 156 (1905). Rhodesia.
6. *St. pauciseta*, nov. sp. E. United States.

(1) *Stilpon pauciseta*, nov. sp. — Male. Length 1 mm. Shining black, the abdomen more piceous, legs entirely yellow. Front pollinose only near the antennæ, mouthparts yellow, the palpi with a conspicuous black bristle, basal joints of the antennæ yellow, the apical joint small, dusky, the long arista microscopically but closely hairy. Notal hairs rather long; pleuræ pollinose in back. Last segment of the abdomen with a loose fringe of long black hairs, pygidium large, lightly pollinose below, apically with a pair of long thin clavate plate-like valves, devoid of long hairs. Front femora ciliate apically, middle femora beneath near the middle with a row of yellow thorn-like setulae and preapically with one or two black bristles, middle tibiæ with a close row of black thorns along the outer three-fifths, hind femora with a loose row of about eight strong flexor bristles, hind tibiæ not pectinate. Halteres black. Wings with a light infumation, veins dark, the second, third and fourth sections of the costa proportioned 1 : 0.7 : 0.8, fourth vein nearly straight, posterior crossvein nearly perpendicular, sections of the fifth vein proportioned 1 : 0.9, marginal cilia nearly as long as the posterior crossvein.

Female. Abdomen more or less testaceous. Halteres yellow to brown, hind femora with a few flexor hairs, legs otherwise devoid of the sexual ornamentation along the flexor side.

Seven specimens, Dorchester County, Maryland (H. S. Barber) and Woods Hole, Massachusetts, July.

7. *St. pectiniger*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 205, f. 4, 5 E. United States.
(1902).
8. *St. pleuritica*, nov. sp. New Hampshire.
9. *St. spinipes*, nov. sp. Georgia.
10. *St. varipes*, Loew, Berl. Ent. Zeitschr. Vol. 6, p. 211 : Cent. 2, No. 58 E. United States.
(1862); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 205 (1902).

15. GENUS DRAPETIS, MEIGEN

Drapetis, Meigen, Syst. Beschr. Vol. 3, p. 91 (1822); Curtis, Brit. Ent. Vol. 8, p. 397 (1824); Macquart, Dipt. N. France, Vol. 3, p. 87 (1827); Hist. Nat. Dipt. Vol. 1, p. 357 (1834); Zetterstedt, Fauna Ins. Lappon. p. 553 (1838); Westwood, Gen. Syn. p. 132 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 327 (1842); Boitard, Man. N. Amer. Dipt. Vol. 3, p. 325 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 135 (1851); Rondani, Dipt. Ital. Vol. 1, p. 147 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 564 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 95 (1862); Lioy, Atti Inst. Sc. Veneto, Venezia, 1864, p. 603 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 125 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390 (1895); Williston, Man. N. Amer. Dipt. p. 75 (1896); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 43 (1902); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 206 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248, 261 (1903); Bezzi, Wien. Ent. Zeit. Vol. 23, p. 143-146 (1904); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 320, 350 (1904); Melander, Williston's Man. N. Amer. Dipt. p. 223 (1908); Kertész, Cat. Dipt. Vol. 6, p. 129 (1909); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 300, 397 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 535 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 252 (1910); Melander, Psyche, Vol. 17, p. 49 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 46, 86 (1910); Melander, Ann. Amer. Ent. Soc. Vol. 11, p. 183-221 (1918); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 380 (1920).

Crossopalpus, Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 557, 563 (1857); Scudder, Nomencl. p. 149 [1882] (*Grossopalpus*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 248 (1903); Proc. U. S. Nat. Mus. Vol. 37, p. 528 (1910).

Ctenodrapetis, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 351, 357, note (1904); Melander, Psyche, Vol. 17, p. 49 (1910); Ann. Ent. Soc. Amer. Vol. 11, p. 187 (1918).

Elatphropeza, Macquart, Dipt. N. France, Vol. 3, p. 86 (1827); Hist. Nat. Dipt. Vol. 1, p. 359 (1834); Westwood, Gen. Syn. p. 132 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 326 (1842); Boitard, Man. Ent. Vol. 3, p. 325 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 134 (1851); Rondani, Dipt. Ital. Vol. 1, p. 147 (1856); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 564 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 94 (1862); Lioy, Atti Inst. Sc. Veneto, Venezia, 1864, p. 722 (1864); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 122 (1889); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 43 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 249, 261 (1903); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 321, 346 (1904); ibidem, Vol. 5, p. 567 (1907); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 537 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 272 (1910); Melander, Psyche, Vol. 17, p. 49 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 45 (1910); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 478 (1912); Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 187 (1918); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 384 (1920).

Eudrapetis, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 187 (1918).

Characters. — Head globular, sometimes slightly longer than high, the occiput more or less hemispherical; eyes large, bare, at most with scattered microscopic hairs, the facets of uniform size, beneath the antennæ the eyes of both sexes nearly touching, leaving the face linear or very narrow, above the eyes the front is V-shaped, always broader than the face, sometimes the front is quite narrow; three small ocelli present. Excision of the eyes at the level of the antennæ very large; cheeks sometimes entirely obliterated, usually about one-eighth the height of the eye, with the oral margin obliquely descending posteriorly. Proboscis thick, rather sharply pointed, perpendicular or directed backward, averaging one-half the height of the head, its labrum with prominent base and as long as the labium; palpi broadly oval, one-jointed, applied to the proboscis, the inside glabrous and shining, the outside pollinose and with more or less evident recumbent pubescence, generally tipped with a longer seta. Antennæ inserted close together a little above the middle of the head, three-jointed, very rarely the basal joint is wanting, the second joint with a circle of small setulae, the bottom hair of which is sometimes long, the third joint compressed, varying from short oval, not longer than deep and with subterminal arista, to lanceolate or conical, nearly twice as long as deep and with terminal arista; the arista usually slender, nearly bare, two-jointed, its basal joint small, its outer joint two or three times as long as the antennæ, rarely the hairy coating is dense and longer so as to give a thickened appearance to the arista. Occiput often with a pollinose band on the lower orbits including the cheeks excepting their lower margin, usually with sparse short hairs, typically a pair of ocellar and one of vertical bristles present, rarely two or three pairs of vertical and an additional pair of post-vertical bristles present, sometimes an oral bristle on the back part of the cheeks.

Thorax robust, polished, nearly bare to pubescent, dorsocentral bristles usually limited to a pair of prescutellars, humeral and posthumeral bristles present or absent, notopleural bristles present but variable, scutellum with two or four marginal bristles, the lateral ones always shorter; pleuræ bare, rarely with pollinose streaks following the sutures. Abdomen with eight segments, the first segment laterally inflated, the fourth segment usually large and more heavily chitinized than the others, its sides often pollinose and bearing scimitar-shaped, glistening setulae, the fifth segment short: pygidium varying from small to large, closed to widely open, asymmetrical, the right side usually with a broad, curved, furcate valve. Legs rather stout, the front femora a little the strongest, usually the anterior femora with a basal flexor hair and with a preapical bristle on the outer side, the hind femora with one to three preapical bristles on the anterior face, hind tibiæ with extensor bristles and more or less produced as an apical tooth-like spur, the anterior tibiæ with a pair of apical bristles; sometimes the legs are devoid of all bristles; pulvilli broad; in those species with broad tibial spur the inside of the hind metatarsi and of the end of the hind tibiæ spongy pubescent.

Wings with anal angle more or less developed, except very rarely the first basal cell shorter than the second which is always fused with the discal cell, five simple longitudinal veins, anal cell completely wanting, at most a short trace of the anal vein present toward the hind margin but the anal crossvein always lacking, usually the entire margin of the wing uniformly short-ciliate, sometimes the hairs of the first section of the costa are longer, no stigma and no definite color pattern.

The species of *Drapetis* are included in the following subgenera :

I. SUBGENUS DRAPETIS, MEIGEN, S. STR.

Type species : *Dr. exilis*, Meigen. Rather small, bristleless, black species, with short antennæ directed somewhat upward, the second joint without a conspicuous seta beneath, the third joint oval, with long, subterminal arista; legs without strong bristles, the hind tibiæ without the terminal spur; notum densely short-pubescent, the hairs appressed, no humeral or discal bristles; wings broad, costa short-pubescent, third vein sometimes curved forward; halteres usually black.

The genus *Crossopalpus*, Bigot, was founded on *Platypalpus ambiguus*, Macquart, which is regarded as the same species as *Drapetis flavipes*, Macquart, and therefore it is synonymous with this subgenus.

2. SUBGENUS EUDRAPETIS, MELANDER

Type species : *E. spectabilis*, Melander (Pl. 4, Fig. 34). Robust, larger species, more or less bristly and black; antennæ longer, directed obliquely upward, the second joint with a pronounced seta beneath, the third joint ovate to lanceolate, with a long terminal arista; femora with preapical bristles, anterior tibiae with apical bristles and the hind ones with a more or less evident terminal spur and usually with extensor or apical bristles; humeral, and sometimes discal bristles (e. g. posthumeral, supraalar and anterior dorsocentrals) present; wings broad, costa short-pubescent, third vein straight or recurved; halteres yellow.

3. SUBGENUS ELAPHROPEZA, MACQUART

Type species : *E. ephippiata*, Fallen. Usually yellow, sometimes black species; antennæ elongate, horizontal, the second joint without seta beneath, the third joint lanceolate, two or more times the length of the first two joints together, the terminal arista relatively short, sometimes no longer than the antennæ; abdomen broad, the middle segment with flattened setulæ; legs slender, the hind tibiae with extensor bristles and with a more or less evident terminal spur; wings narrower, the hairs of the base of the costa longer than elsewhere.

4. SUBGENUS CTENODRAPETIS, BEZZI

Type species : *Ct. ciliatocosta*, Bezzi. Body slender, elongate, usually yellow, the abdomen longer than the thorax; antennæ elongate, horizontal, the second joint without seta, the third joint lanceolate with relatively coarsely pubescent terminal arista; legs slender, with bristles and a long terminal spur on the hind tibiae; wings long, narrow, cuneate, the costal margin with long cilia.

* * *

The species of *Drapetis* are æstival and occur in meadowlands and among shrubbery. The developmental stages are unknown. In the tropics yellow-colored species predominate, but in the Temperate Zone the general species are black and more robust. Here the species are found during the spring and summer, sometimes swarming about flowers like wild cherry and plum, and sometimes running about singly over grass and low shrubbery. A synopsis of the American species of *Drapetis* was given by the writer in the June, 1918, issue of the « Annals of the Entomological Society of America ». The preceding generic characterization is taken from this account.

Geographical distribution.

SUBGENUS DRAPETIS, MEIGEN, S. STR.

1. *D. aliternigra*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 192 (1918). North America.
nigra, Melander (not Meigen), Trans. Amer. Ent. Soc. Vol. 28, p. 208,
f. 12, 17, 24 (1902); Lundbeck, Dipt. Dan. Vol. 3, p. 257, note (1910);
Malloch, Bull. Illinois Labor. Nat. Hist. Urbana, Vol. 12, Art. 3, p. 403
(1917) larva, pupa.
2. *D. assimilis*, Fallen, Dipt. Suec. Emp. p. 8, 34 [1815] (*Tachydromia*); Europe, ?North America.
Walker, Ins. Brit. p. 136 (1851); Schiner, Fauna Dipt. Austr.
Vol. 1, p. 95 (1862); Lundbeck, Dipt. Dan. Vol. 3, p. 256, f. 115
(1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 87 (1910); Frey, Acta

- Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 71, f. 19 (1913); Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 187 (1918).
- affinis*, Loew, Neue Beitr. Dipt. Vol. 6, p. 41 (1859).
- arcuata*, Loew, ibidem, Vol. 6, p. 40 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862); Wahlgren, Ent. Tidskr. Vol. 31, p. 87 (1910); Kuntze, Deutsche Ent. Zeitschr. p. 548 (1913).
- minima*, Meigen (not Zetterstedt), Syst. Beschr. Vol. 7, p. 100 (1838).
- nigra*, Meigen, ibidem, Vol. 6, p. 344 (1830); Curtis, Brit. Ent. Vol. 8, p. 397 (1832); Macquart, Hist. Nat. Dipt. Vol. 1, p. 358 (1834); Walker, List. Dipt. Brit. Mus. Vol. 3, p. 511 (1849), oc. in Canada; Ins. Brit. Vol. 1, p. 136 (1851).
- pygmaea*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 54 (1840).
3. *D. bispina*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 192 (1918). Indiana.
4. *D. brevior*, Brunetti, Rec. Indian Mus. Vol. 9, p. 37 (1913); Fauna Brit. India Dipt. Vol. 1, p. 382, pl. 4, f. 23, 24 (1920). India.
5. *D. *decolorata*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 95 (1908). Baltic Amber.
6. *D. distans*, Bezzi, Wien. Ent. Zeit. Vol. 23, p. 144 (1904); Ann. Mus. Hungar. Vol. 10, p. 480 (1912).
- divergens*, Bezzi (not Loew), Ann. Mus. Hungar. Vol. 2, p. 353 (1904).
7. *D. divergens*, Loew, Berl. Ent. Zeitschr. Vol. 16, p. 90 : Cent. 10, No. 62 (1872); Melander, Trans. Amer. Ent. Vol. 28, p. 210, f. 15 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 397 (1909).
- minuta*, Williston, Trans. Ent. Soc. London, Vol. 3, p. 442, f. 168 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 210 (1902).
8. *D. diversa*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 193 (1918). New Mexico.
9. *D. dividua*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 208, pl. 5, f. 16, 18, 23 (1902).
- nigripes*, Melander, ibidem, Vol. 28, p. 339 [1902] (*Stilpon*). W. North America.
10. *D. exilis*, Meigen, Syst. Beschr. Vol. 3, p. 91, pl. 23, f. 25-28 (1822); Serville & Lepeletier, Encycl. Méth. Vol. 10, p. 437 (1825); Macquart, Dipt. N. France, Vol. 3, p. 88, part, pl. 2, f. 2 (1827); Meigen, Syst. Beschr. Vol. 6, p. 344 (1830); Curtis, Brit. Ent. Vol. 8, p. 397, part (1832); Macquart, Hist. Nat. Dipt. Vol. 1, p. 357, part, pl. 8, f. 12a (1834); Zetterstedt, Fauna Ins. Lappon. p. 554 (1838); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 23 (1840); Isis, Vol. 7, p. 552 (1840); Zetterstedt, Dipt. Scand. Vol. 1, p. 328 (1842); Boitard, Man. Ent. Vol. 3, p. 325 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 511, part (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3012 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 60 (1851); Walker, Ins. Brit. Dipt. Vol. 1, p. 136 (1851); Loew, Neue Beitr. Dipt. Vol. 6, p. 42 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 95 (1862); Lundbeck, Dipt. Dan. Vol. 3, p. 258, f. 116 (1910). Europe.
11. *D. fascifemorata*, Brunetti, Rec. Indian Mus. Vol. 9, p. 38 (1913); Fauna Brit. India Dipt. Vol. 1, p. 383 (1920). Formosa.
12. *D. femorata*, Melander, Ann. Soc. Ent. Amer. Vol. 11, p. 215 (1918). Formosa.
- femoralis*, Bezzi (not Wheeler & Melander), Ann. Mus. Hungar. Vol. 10, p. 480, 483 (1912).
13. *D. flavigornis*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 193 (1918). Panama.
14. *D. flavipes*, Macquart, Hist. Nat. Dipt. Vol. 1, p. 357 (1834); Meigen, Syst. Beschr. Vol. 7, p. 100 (1838); Boitard, Man. Ent. Vol. 3, p. 325 (1843); Loew, Neue Beitr. Dipt. Vol. 6, p. 38 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 95 (1862); Strobl, Mitteil. Nat. Ver. Europe.

- Steiermark, Graz, Vol. 29, p. 108 (1892); Bezzi, Arch. Zool. Expér. Paris (5), Vol. 8, p. 50 (1911).
- ? *ambigua*, Macquart, Dipt. N. France, Vol. 3, p. 94 [1827] (*Platypalpus*); Hist. Nat. Dipt. Vol. 1, p. 351 (1834); Schiner, Fauna Dipt. Austr. Vol. 1, p. 92 (1862).
- ? *curvipes*, Meigen, Syst. Beschr. Vol. 3, p. 75 [1822] (*Tachydromia*).
- exilis*, Macquart, Dipt. N. France, Vol. 3, p. 88, part (1827).
- fascipes*, Roser, Correspondenzbl. Landw. Ver. Würtemb. Stuttgart, Vol. 1, p. 54 (1840).
- nigra*, Fallen, Dipt. Suec. Emp. p. 8 [1815] (*Tachydromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 90 [1862] (*Platypalpus*).
15. *D. inermis*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 194 (1918). Panama.
16. *D. infumata*, Melander, ibidem, Vol. 11, p. 194 (1918). British Columbia.
17. *D. levis*, Becker, Ann. Mus. Zool. Acad. Sc. St. Pétersb. Vol. 18, p. 84 (1913). N. Africa.
18. *D. latipennis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 209, pl. 5, f. 11, 19 (1902); Ann. Ent. Soc. Amer. Vol. 11, p. 189 (1918). C. United States.
19. *D. micropyga*, Melander, ibidem, Vol. 11, p. 195 (1918). W. United States.
20. *D. *mortua*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 97 (1908). Baltic Amber.
21. *D. naica*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 195 (1918). W. United States.
22. *D. nigripes*, Zetterstedt, Dipt. Scand. Vol. 13, p. 4997 (1859); Wahlgren, Ent. Tidskr. Vol. 31, p. 87 (1910). N. Europe.
23. *D. pilosa*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 196 (1918). C. United States.
24. *D. procurrens*, nov. sp. (1). Costa Rica.
25. *D. pusilla*, Loew, Neue Beitr. Dipt. Vol. 6, p. 36 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862); Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 42 (1908); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 179 (1909); Lundbeck, Dipt. Dan. Vol. 3, p. 259, f. 117 (1910); Dahl, Fauna Chorin, p. 465 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 71 (1913).
- exilis*, Zetterstedt (not Meigen), Fauna Ins. Lappon. p. 554, part (1838); Dipt. Scand. Vol. 1, p. 328 (1842); ibidem, Vol. 8, p. 3012 (1849).
- minima*, Zetterstedt (not Meigen), Dipt. Scand. Vol. 1, p. 327 (1842).
- var. *fumipennis* Strobl, Mem. Soc. Esp. N. Hist. Vol. 3, p. 312 (1906). Spain.
26. *D. rotundicornis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 38 (1913); Fauna Brit. India, Dipt. Vol. 1, p. 383, pl. 4, f. 25 (1920). India.
27. *D. rufipes*, Brunetti, Rec. Indian Mus. Vol. 9, p. 38 (1913); Fauna Brit. India, Dipt. Vol. 1, p. 383 (1920). Bengal.
28. *D. setulosa*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 196 (1918). W. North America.
29. *D. trichura*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 197 (1918). Texas
30. *D. xanthopyga*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 352 (1904). Vol. 10, p. 479, 485 (1912). New Guinea.

(1) **Drapetis (s. str.) procurrens**, nov. sp. — Male. Length 1.4 mm. Body including pleurae shining black, hairs and bristles of head and thorax black, legs entirely yellow, unarmed, third vein diverging from fourth, antennæ yellowish. Front rather narrowly V-shaped; face obliterated except at clypeus; third antennal joint slightly darkened, triangular, nearly two times long as broad, the blackish arista micropubescent but not as densely so as in *divergens*, two and one-half times length of antenna; palpi large, oval, whitish, proboscis dark. Thoracic pubescence closely and uniformly distributed, lateral scutellar bristles half length. Third abdominal segment chitinized laterally, with flattened setulae, pygidium large, contorted, the valves terminally fringed, a long curled brown penis exserted. No pronounced bristles on legs, hind tibiae with small brush-like termination. Halteres yellow; calypteres dark, with black fringe. Wings hyaline, veins brown, third vein nearly uniformly diverging from fourth, the costal sections proportioned 1 : 2 : 1.5, sections of fourth vein, 1 : 0.7 : 4, of fifth vein, 1 : 0.9.

Holotype, received from Pablo Schild, La Suiza de Turrialba, Costa Rica.

SUBGENUS EUDRAPETIS, MELANDER

1. *E. aenea*, Walker, Entom. London, Vol. 5, p. 273 (1871).
 2. *E. aenescens*, Wiedemann, Aussereur. Zweifl. Ins. Vol. 2, p. 649 (1830); Loew, Neue Beitr. Dipt. Vol. 6, p. 35 (1859); Dipterenf. Südafri. p. 270 (1860); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862); Giglio-Tos, Ann. Soc. Ent. France, Vol. 64, p. 359 (1895); Becker, Mitteil. Zool. Mus. Berl. Vol. 2, p. 40 (1902); Strobl, Glasnik Zem. Mus. Bosn. Herceg. Sarajevo, Vol. 14, p. 470 (1902); Mitteil. Bosn. Herceg. Sarajevo, Vol. 9, p. 528 (1904); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 351 (1904); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 119 (1907); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 479, 482 (1912); Suppl. Ent. Berlin, Vol. 3, p. 75 (1914); Meijere, Tijdschr. Ent. Vol. 56, Suppl. p. 75 (1914).
- brunnipes*, Macquart, Hist. Nat. Dipt. Vol. 1, p. 358 (1834); Meigen, Syst. Beschr. Vol. 7, p. 101 (1838); Loew, Bemerk. Posen. Gegend Art. Zweifl. Gatt. p. 23 (1840); Isis, Vol. 7, p. 552 (1840); Boitard, Man. Ent. Vol. 3, p. 325 (1843); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 60 (1851).
- crassa*, Loew, Oefv. Vet. Akad. Forhandl. Stockholm, Vol. 15, p. 341 (1858).
3. *E. angustata*, Collin, Ent. M. Mag. London, 699, p. 187 (1922). Seychelles.
 4. *E. basalis*, Collin, ibidem, 699, p. 187 (1922). Seychelles.
 5. *E. armata*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 197 (1918). United States.
 6. *E. armipes*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 398, f. 18 (1909). Chile.
7. *E. aterrima*, Curtis, Brit. Ent. Vol. 8, p. 397, tab. (1832); Walker, Ins. Brit. Dipt. Vol. 1, p. 136 (1851); Loew, Neue Beitr. Dipt. Vol. 6, p. 40, note (1859); Lundbeck, Dipt. Dan. Vol. 3, p. 254, f. 111-113 (1910); Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 68 (1913).
- atra*, Walker, Ins. Brit. Dipt. Vol. 1, pl. 5, f. 3g, h (1851).
nervosa, Loew, Neue Beitr. Dipt. Vol. 6, p. 37 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862).
nigritella, Zetterstedt, Dipt. Scand. Vol. 1, p. 298 [1842] (*Tachydromia*).
8. *E. brevicula*, new name. Formosa.
brevis, Bezzi (not Meunier), Ann. Mus. Hungar. Vol. 10, p. 480, 483 (1912).
 9. *E. * brevis*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 96 (1908). Baltic Amber.
 10. *E. * decorata*, Meunier, ibidem, Vol. 7, p. 96 (1908). Baltic Amber.
 11. *E. discalis*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 198 (1918). W. United States.
 12. *E. diversipes*, Melander, ibidem, Vol. 11, p. 198 (1918). California.
 13. *E. dorsiseta*, Melander, ibidem, Vol. 11, p. 199 (1918). Costa Rica.
 14. *E. facialis*, Melander, ibidem, Vol. 11, p. 200 (1918). Georgia, Alberta.
 15. *E. femoralis*, Wheeler & Melander, Biol. Centr.-Amer. Dipt. Vol. 1, p. 376 (1901); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 211, f. 13 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 398 (1909). Mexico.
16. *E. fortis*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 399 (1909). Peru.
 17. *E. gilvipes*, Loew, Berl. Ent. Zeitschr. Vol. 16, p. 89 : Cent. 10, No. 61 (1872); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 213, 340 (1902); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 397 (1909). Texas, West Indies.
18. *E. humilis*, Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 69, f. 16-18 (1913). Finland.
19. *E. inculta*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 439 [1896] (*Platypalpus*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 218 (1902); Ann. Ent. Soc. Amer. Vol. 11, p. 188 (1918). Texas, California.
20. *E. lata*, Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 266 [1903] (*Tachydromia*); Melander, Psyche, Vol. 17, p. 51 (1910); Ann. Ent. Soc. Amer. Vol. 11, p. 189 (1918). Florida.

21. *E. Loewi*, Dahl, Fauna Chorin, p. 465 (1912).
 22. *E. luteipes*, Lepeletier & Serville, Encycl. Méthod. Vol. 10, p. 437 (1825).
 23. *E. marginata*, Meigen, Syst. Beschr. Vol. 7, p. 100 (1838).
 24. *E. medetera*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 208, pl. 5, f. 22 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 [1903] (*Elaphropeza*); Melander, Ent. News, Philad. Vol. 17, p. 372 (1906); Bezzi, Ann. Mus. Hungar. Vol. 5, p. 567 [1907] (*Elaphropeza*); Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 189 (1918).
 25. *E. moriella*, Zetterstedt, Fauna Ins. Lappon. p. 552 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 299 [1842] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 86, f. 15 (1910).
curvinervis, Zetterstedt, Dipt. Scand. Vol. 1, p. 301 [1842] (*Tachydromia*).
flexuosa, Loew, Bemerk. Posen Gegend Art. Zweifl. Gatt. 1840, p. 23 : Isis, Vol. 7, p. 552 (1840); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 60 (1851); Loew, Neue Beitr. Dipt. Vol. 6, p. 39 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862).
geniculata, Fallén, Suec. Empid. 7, part [1815] (*Tachydromia*).
nigra, Zetterstedt, Dipt. Scand. Vol. 1, p. 297 [1842] (*Tachydromia*); Bonsdorff, Finl. tväv. Ins. Dipt. Vol. 1, p. 150 [1861] (*Tachydromia*).
picipes, Zetterstedt, Dipt. Scand. Vol. 1, p. 298 [1842] (*Tachydromia*).
 var. *dilutipes*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 312 (1906).
 var. *setigera*, Loew, Neue Beitr. Dipt. Vol. 6, p. 39 (1859); Schiner, Fauna Dipt. Austr. Vol. 1, p. 96 (1862); Lundbeck, Dipt. Dan. Vol. 3, p. 255, f. 114 (1910); Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 70 (1913).
 26. *E. nitens*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 200 (1918).
 27. *E. nuda*, Melander, ibidem, Vol. 11, p. 201 (1918).
 28. *E. obscuripennis*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 768 (1865); Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 397 (1909).
 29. *E. oedimera*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 201 (1918).
 30. *E. parvicornis*, Melander, ibidem, Vol. 11, p. 202 (1918).
 31. *E. pennescens*, Melander, ibidem, Vol. 11, p. 202 (1918).
 32. *E. phæoptera*, Bezzii, Wien. Ent. Zeit. Vol. 28, p. 145 (1904); Ann. Mus. Hungar. Vol. 10, p. 479 (1912).
obscurepennis, Bezzii (not Philippi), Ann. Mus. Hungar. Vol. 2, p. 351 (1904).
 33. *E. pilipes*, Loew, Neue Beitr. Dipt. Vol. 6, p. 36 (1859); Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 152 (1899); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 40 (1902).
 34. *E. plumipes*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 203 (1918).
 35. *E. pubicornis*, Bezzii, Ann. Mus. Hungar. Vol. 10, p. 479, 482 (1912).
 36. *E. quadrisetosa*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 203 (1918).
 37. *E. scissa*, Melander, ibidem, Vol. 11, p. 204 (1918).
medetera, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 208, part (1902).
 Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 [1903] (*Elaphropeza*); Melander, Ent. News, Philad. Vol. 17, p. 372, part (1906); Bezzii, Ann. Mus. Hungar. Vol. 5, p. 567 [1907] (*Elaphropeza*).
 38. *E. septentrionalis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 211 (1902); Ann. Ent. Soc. Amer. Vol. 11, p. 190 (1918).
 var. *mexicana*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 204 (1918).
 39. *E. spectabilis*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 212, f. 9, 20 (1902); Ann. Ent. Soc. Amer. Vol. 11, p. 189, 190 (1918). — PI. 4, Fig. 34.
 40. *E. spinipes*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 205 (1918).
 41. *E. unipila*, Loew, Berl. Ent. Zeitschr. Vol. 16, p. 88 : Cent. 10, No. 60 (1872); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 211 (1902); Ent. News, Philad. Vol. 17, p. 372 (1906); Ann. Ent. Soc. Amer. Vol. 11, p. 188 (1918).
 Germany.
 France.
 E. Europe.
 W. United States.
 N. and C. Europe.
 Spain.
 Europe.
 Mexico, Panama.
 Africa.
 Chile.
 Vera Cruz.
 Washington.
 Peru.
 New Guinea.
 S. Europe, N. Africa.
 Texas.
 Formosa.
 Mexico, Central America.
 W. United States.
 Michigan.
 Tampico.
 United States.
 Africa.
 W. United States.

- var. *sitida*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 207, 339, f. 10, 14, 21 (1902); Cole, Report Laguna Marine Labor. Vol. 1, p. 152 (1912); Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 188 (1918).
42. *E. * vitiosa*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 96, pl. 3, f. 10-13 (1908). Baltic Amber.
43. *E. xanthopoda*, Williston, part, Trans. Ent. Soc. London, Vol. 3, p. 308, pl. 4, f. 85 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 212, 340 (1902); Ann. Ent. Soc. Amer. Vol. 11, p. 190 (1918). *flavipes*, Williston (not Macquart), Trans. Ent. Soc. London, Vol. 3, Appendix, p. 441, tab. (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 212 (1902).

SUBGENUS CTENODRAPETIS, BEZZI

1. *C. aristalis*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 205 (1918). — Philippine Islands.
Pl. 4, Fig. 36.
2. *C. ciliatocosta*, Bezzl, Ann. Mus. Hungar. Vol. 2, p. 355, f. 6 (1904); Nova Act. Akad. Naturf. Halle, Vol. 91, p. 397, 400 (1909); Ann. Mus. Hungar. Vol. 10, p. 481 (1912).
3. *C. cuneipennis*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 206 (1918). Philippine Islands.
4. *C. discoidalis*, Bezzl, Ann. Mus. Hungar. Vol. 2, p. 355 (1904); ibidem, Vol. 10, p. 482 (1912); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 382 [1920] (*Drapetis*).
5. *C. gracilis*, Bezzl, Ann. Mus. Hungar. Vol. 2, p. 354 (1904); ibidem, Vol. 10, p. 480 (1912).
6. *C. hamifera*, Bezzl, ibidem, Vol. 10, p. 479, 485 (1912). Formosa.
7. *C. monocheta*, Bezzl, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 401 (1909). Peru.
8. *C. rubrithorax*, Bezzl, Ann. Mus. Hungar. Vol. 2, p. 356 (1904); ibidem, Vol. 10, p. 482 (1912). New Guinea.
9. *C. strigifera*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 333 (1911); ibidem, Vol. 56, Suppl. p. 75 (1914); Java.
10. *C. valdiviana*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 768 [1865] (*Drapetis*); Bezzl, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 397 [1909] (? *Ctenodrapetis*). Chile.

SUBGENUS ELAPHROPEZA, MACQUART

1. *E. abdominalis*, Wiedemann, Aussereur. Ins. Vol. 2, p. 12 [1830] (*Tachydromia*); Bezzl, Ann. Mus. Hungar. Vol. 2, p. 359, note [1904] (*Tachydromia*); Melander, Psyche, Vol. 17, p. 52 [1910] (*Ctenodrapetis*).
2. *E. acroductyla*, nov. sp. (1). Luzon.
3. *E. antennata*, Becker, Bul. Mus. Hist. Nat. Paris, p. 115 (1909); Ann. Soc. Ent. France, Vol. 79, p. 24 (1910); Bezzl, Ann. Mus. Hungar. Vol. 10, p. 481 (1912). E. Africa.

(1) *Drapetis (Elaphropeza) acroductyla*, nov. sp. — Female. Length 2 mm. Luteous, the occiput, linear front, third antennal joint, arista, broad third abdominal tergite, apex of hind tibiae within, and last joint of all tarsi black. Third antennal joint lanceolate, three times long as wide and five-ninths as long as the closely pubescent arista; palpi ovate. Posterior dorsocentral large, two scutellars, acrostichal and dorsocentral setulae totaling four distinct rows. Sides of black. Abdominal band with many strong flattened black setulae. Two extensor bristles on hind tibiae. Wings hyaline, veins yellow, sections of costa proportioned 1 : 0.3 : 0.8 : 0.3, of fourth vein 1 : 1 : 3, of fifth vein 1 : 0.5, first posterior cell gently widening to tip, fourth vein ending slightly beyond apex of wing.

Two specimens, Mt Makling, Luzon, C. F. Baker. Related to *formosa* Bezzl, but that species has the first posterior cell narrowing at the tip.

4. *E. bacis*, Walker, List Dipt. Brit. Mus. Vol. 3, p. 510 [1849] (*Platypalpus*); Jamaica. Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 439 [1896] (*Tachydromia*); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 226 [1902] (*Tachydromia*); Bezzi, Ann. Mus. Hungar. Vol. 3, p. 460 [1905] (*Tachypeza*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 402, note (1909); Melander, Psyche, Vol. 17, p. 52 [1910] (*Ctenodrapetis*).
 5. *E. basalis*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 349 (1904); ibidem, Ceylon. Vol. 5, p. 567 (1907); ibidem, Vol. 10, p. 479 (1912); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 384 (1920).
 6. *E. bicaudata*, nov. sp. (1). Luzon.
 7. *E. bicolor*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 349 (1904); ibidem, New South Wales. Vol. 5, p. 567 (1907); Vol. 10, p. 482 (1912).
 8. *E. bicoloripes*, Brunetti, Rec. Indian Mus. Vol. 9, p. 43 (1913); Fauna E. Himalayas. Brit. India Dipt. Vol. 1, p. 385 (1920).
 9. *E. bihamata*, Bezzi, Ann. Mus. Nat. Hungar. Vol. 2, p. 353 [1904] New Guinea. (*Drapetis*); Meijere, Tijdschr. v. Ent. Vol. 54, p. 332 [1911] (*Drapetis*); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 480 [1912] (*Drapetis*).
 10. *E. binotata*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 333 [1911] (*Drapetis*); Java. ibidem, Vol. 56, Suppl. p. 74 (1913).
 11. *E. biuncinata*, nov. sp. (2). Luzon.
 12. *E. calcarifera*, Bezzi, Ann. Mus. Hungar. Vol. 5, p. 267, 268 (1907); Formosa. ibidem, Vol. 10, p. 481, 488 (1912); Suppl. Ent. Berlin. Vol. 3, p. 77 (1914).
 13. *E. callositibia*, Bezzi, Ann. Mus. Hungar. Vol. 2, p. 354 [1904] (*Drapetis*); New Guinea. ibidem, Vol. 10, p. 480 [1912] (*Drapetis*).
 14. *E. calva*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 206 (1918). Panama.
 15. *E. comata*, Melander, ibidem, Vol. 11, p. 207 (1918). Trinidad.
 16. *E. dispar*, Adams, Kansas Univ. Sc. Bull. Vol. 3, p. 157 (1905); Bezzi, Ann. Mus. Hungar. Vol. 5, p. 567 (1907); ibidem, Vol. 10, p. 480 (1912). Rhodesia.
 17. *E. ephippiata*, Fallen, Empid. Suec. p. 11 [1815] (*Tachydromia*); Meigen, Europe. Syst. Beschr. Vol. 3, p. 65, pl. 23, f. 10 [1822] (*Hemerodromia*);

(1) **Drapetis (Elaphropeza) bicaudata**, nov. sp. — Male. Length 2 mm. Luteous, the head black, occiput and the V-shaped face cinereous, third antennal joint blackish, bluntly conical, less than twice as long as wide and one-fourth as long as the dark arista; palpi oval, twice as long as wide, no long apical seta. Posterior dorsocentral moderate, apical scutellars approximate. Third abdominal tergite and its flattened setulae brown, pygidium large and very long, tipped with two long strong black bristles. Legs simple, the hind metatarsi as long as the subsequent joints and two-thirds as long as their tibiae which have two strong extensor setae. Wings hyaline, veins yellowish, a strong seta at base of costa, costal sections proportioned 1 : 0.5 : 1 : 0.4, first posterior cell gradually widening, sections of fourth vein 1 : 1.2 : 3.5, of fifth vein 1 : 0.5.

Female. Caudal bristles absent.

Type and allotype, Mt. Makling, Luzon, Philippine Islands, C. F. Baker. A third specimen has somewhat shorter hind metatarsi.

(2) **Drapetis (Elaphropeza) biuncinata**, nov. sp. — Male. Length 1.5 mm. With *bihamata*, Bezzi, differing in its lengthened antennæ and darkened halteres. Head spherical, yellow, lightly pollinose, ocellar triangle brown, a pair each of ocellar and vertical bristles, strongly yellow; basal joint of antennæ yellow, crowned with black setulae, outer joint lanceolate, black, pubescent, three times as long as deep, arista black, pubescent, two and one-fourth times length of last joint; mouthparts small, yellow, palpi orbicular, with brown apical seta. Thorax shining testaceous yellow, mesonotum as wide as long, bristles and hairs yellow, metanotum castaneous below or wholly so, a postalar but no dorsocentral bristle, two scutellars. First five segments of abdomen castaneous, the remainder testaceous, seventh segment fringed with long pale setæ, pygidium very large, quadrate from above. Legs pale yellow, no outstanding setæ, hairs yellow, hind tibiae with two strong curved black bristles just before the apex on the outer side, which are close together and one before the other. Halteres with brown knob. Wings hyaline, veins yellowish, costal sections proportioned 1 : 0.9 : 0.9 : 0.5, third and fourth veins parallel beyond end of second vein, anterior crossvein at middle of discal cell, sections of fifth vein 1 : 0.9.

Two males, Mt. Makling, Luzon, C. F. Baker. The paratype has a faintly smoky cloud over the central part of the wings. A female from the same locality lacks the tibial spurs and has pale halteres.

DIPTERA

- Macquart, Dipt. N. France, Vol. 3, p. 87, pl. 2, f. 1 (1827); Hist. Nat. Dipt. Vol. 1, p. 359, pl. 8, f. 14a (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 326 (1842); Boitard, Man. Ent. Vol. 3, p. 325 (1843); Walker, List Dipt. Brit. Mus. Vol. 3, p. 505 (1849); Ins. Saunders, Dipt. Vol. 1, p. 134, pl. 5, f. 4 (1851); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 60 [1851] (*ephippium*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 95 (1862); Strobl, Mitteil. Naturf. Ver. Steiermark, Graz, Vol. 29, p. 127 (1892); Melander, Ent. News, Philad. Vol. 17, p. 372 [1906] (*Drapetis*); Bezzi, Ann. Mus. Hungar. Vol. 5, p. 567 (1907); Lundbeck, Dipt. Dan. Vol. 3, p. 273, f. 123, 124 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 89 (1910); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 482 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 75 [1913] (occ.).
? maculata, Macquart, Mem. Soc. Sc. Lille, p. 153 [1823] (*Tachydromia*).
18. *E. exul*, Osten-Sacken, Berl. Ent. Zeitschr. Vol. 26, p. 113 (1882); Bezzi, Ann. Mus. Hungar. Vol. 2, p. 347 (1904); ibidem, Vol. 5, p. 567 (1907); Vol. 10, p. 480 [1912] (*Drapetis*). Philippine Is.
19. *E. ferruginea*, Brunetti, Rec. Indian Mus. Vol. 4, p. 43 (1913); Fauna Brit. India, Dipt. Vol. 1, p. 386 (1920). India.
20. *E. flavicollis*, Becker, Bull. Mus. Hist. Nat. Paris, p. 116 [1909] (*Drapetis*); Ann. Soc. Ent. France, Vol. 79, p. 24 [1910] (*Drapetis*); Bezzi, Ann. Mus. Hungar. Vol. 10, p. 482 (1912). E. Africa.
21. *E. flavidia*, Williston, Trans. Ent. Soc. London, Vol. 3, p. 308, pl. 11, f. 86 [1896] (*Drapetis*); Coquillett, Proc. U. S. Nat. Mus. Vol. 22, p. 251 [1900] (*Tachydromia*); Wheeler & Melander, Biol. Centr. Amer. Dipt. Vol. 1, p. 376 [1901] (*Drapetis*); Melander, Mon. Empid. p. 213 [1902] (*Drapetis*); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 265 [1903] (*Tachydromia*); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 397, 398 [1909] (*Drapetis*); Melander, Psyche, Vol. 17, p. 52 [1910] (*Ctenodrapetis*); Ann. Ent. Soc. Amer. Vol. 11, p. 191 (1918). West Indies, C. & S. America.
fulvida, Bezzi, Wien. Ent. Zeit. Vol. 23, p. 144 [1904] (*Tachista*).
xanthopoda, Williston, Trans. Ent. Soc. London, Vol. 3, Appendix, 441, table [1896] (*Drapetis*).
22. *E. formosae*, Bezzi, Ann. Mus. Hungar. Vol. 5, p. 566 (1907); ibidem, Vol. 10, p. 480, 487 (1912). Formosa.
23. *E. fulvithorax*, Wulp, Termes. Fuzet. Vol. 20, p. 138 (1897); Tijdschr. Ent. Vol. 42, p. 49 (1899); Bezzii, Ann. Mus. Hungar. Vol. 2, p. 347 (1904); ibidem, Vol. 5, p. 567 (1907); ibidem, Vol. 10, p. 480 (1912); Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 73 (1914); Brunetti, Fauna Brit. India, Dipt. Vol. 1, p. 385 (1920). Ceylon, Java.
24. *E. hirsutitibia*, Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 73 (1914). Java.
bishamata, Meijere (not Bezzi), ibidem, Vol. 54, p. 332 [1911] (*Drapetis*).
25. *E. inflexa*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 207 (1918). Panama.
26. *E. Kerteszi*, Bezzi, Ann. Mus. Hungar. Vol. 10, p. 481, 486 (1912); Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 74 (1914). Formosa.
27. *E. lata*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 208 (1918). Panama.
var. *nigrocalcarata*, Melander, ibidem, Vol. 11, p. 208 (1918). Panama.
var. *ungulifera*, Melander, ibidem, Vol. 11, p. 208 (1918). Panama.
28. *E. lanuginosa*, Bezzi, Suppl. Ent. Berlin, Vol. 3, p. 75 (1914). Formosa.
29. *E. lineola*, Meijere, Tijdschr. v. Ent. Vol. 54, p. 331 (1911); Vol. 56, Suppl. p. 75 (1914). Java.
30. *E. lutea*, Meijere, ibidem, Vol. 54, p. 332 (1911); ibidem, Vol. 56, Suppl. p. 76 (1914). Java.
31. *E. luteicollis*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 209 (1918). Panama.

32. *E. marginalis*, Bezzia, Ann. Mus. Hungar. Vol. 10, p. 481, 489 (1912). Formosa.
33. *E. melanura*, Bezzia, ibidem, Vol. 10, p. 481, 489 (1912); Suppl. Ent. Formosa. Berlin, Vol. 3, p. 78 (1914).
34. *E. metatarsata*, Bezzia, Ann. Mus. Hungar. Vol. 2, p. 348 (1904); ibidem, Vol. 5, p. 567 (1907); ibidem, Vol. 10, p. 482 (1912); Brunetti, Fauna Brit. India Dipt. Vol. 1, p. 586 (1920).
35. *E. nigricans*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 209 (1918). Panama.
36. *E. obliquinervis*, Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 73, pl. 2, f. 9 (1914). Java.
37. *E. palpata*, Meijere, ibidem, Vol. 54, p. 330 (1911); ibidem, Vol. 56, Suppl. p. 76 (1914). Java.
38. *E. pictithorax*, Bezzia, Ann. Mus. Hungar. Vol. 10, p. 482, 486 (1912). Formosa.
39. *E. pleuralis*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 210 (1918). Vera Cruz, Panama.
40. *E. plumea*, Melander, ibidem, Vol. 11, p. 210 (1918). Mexico.
41. *E. rectineura*, Melander, ibidem, Vol. 11, p. 211 (1918). Panama.
42. *E. scutellaris*, Bezzia, Ann. Mus. Hungar. Vol. 10, p. 481, 482, 487 (1912). Formosa.
43. *E. seminigra*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 211 (1918). Costa Rica.
44. *E. simplicipes*, Melander, ibidem, Vol. 11, p. 212 (1918). West Indies.
45. *E. spuria*, Bezzia, Ann. Mus. Hungar. Vol. 2, p. 347 (1904); ibidem, Vol. 5, p. 567 (1907); ibidem, Vol. 10, p. 480 (1912). New Guinea.
46. *E. tenera*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 212 (1918). Trinidad, Panama.
47. *E. uniseta*, Melander, ibidem, Vol. 11, p. 213 (1918). Philippine Islands.
48. *E. epsilon*, Melander, ibidem, Vol. 11, p. 214 (1918). West Indies.
49. *E. variata*, Melander, ibidem, Vol. 11, p. 221 (1918). India.
- variegata* (bis), Brunetti, Rec. Indian Mus. Vol. 9, p. 42 (not 39) [1913]; Fauna Brit. India Dipt. Vol. 1, p. 387 [1920] (*Drapetis*).
50. *E. variegata*, Brunetti, Rec. Indian Mus. Vol. 9, p. 39 [1913] (*Drapetis*); Fauna Brit. India Dipt. Vol. 1, p. 381 [1920] (*Drapetis*). Burma.
51. *E. vittata*, Melander, Ann. Ent. Soc. Amer. Vol. 11, p. 214 (1918). Florida, West Indies.
52. *E. xanthocephala*, Bezzia, Ann. Mus. Hungar. Vol. 10, p. 481, 488 (1912). Formosa.

16. GENUS PLATYPALPUS, MACQUART

Platypalpus, Macquart, Dipt. N. France, Vol. 3; p. 92 (1827); Hist. Nat. Dipt. Vol. 1, p. 351 (1834); Westwood, Gen. Syn. p. 132 (1840); Boitard, Man. Ent. Vol. 3, p. 323 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 121 (1851); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 563 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 87 (1862); Lioy, Atti Inst. Sc. Veneto, Venezia, 1864, p. 721 (1864); Loew, Jahrb. Gel. Ges. Krakau, Vol. 41, p. 12 (1870); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 123 (1889); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 390, 438 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 214 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255, 261 (1903); Melander, Williston's Man. N. Amer. Dipt. p. 222 (1908); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 592 (1910); Melander, Psyche, Vol. 17, p. 49 (1910).

Brevios, Brunetti, Rec. Indian. Mus. Vol. 9, p. 22 (1913).

Cleptodromia, Corti, Wien. Ent. Zeit. Vol. 26, p. 101 (1907); Melander, Psyche, Vol. 17, p. 49 (1910).

Corynetea (of authors), Meigen, Nouv. Classif. Mouches, p. 27 (1800); Hendel, Verh. Zool.-bot. Ges. Wien, Vol. 58, p. 56 (1908); Kertész, Cat. Dipt. Vol. 6, p. 146 (1909); Bezzia, Ann. Mus. Hungar. Vol. 10, p. 453 (1912).

Howlettia, Brunetti, Rec. Indian Mus. Vol. 9, p. 23 (1913); Fauna Brit. India Dipt. Vol. 1, p. 360 (1920).

Phoroxypha, Rondani, Dipt. Ital. Vol. 1, p. 146 (1856); **Bezzi**, Bull. Soc. Ent. Ital. Vol. 30, p. 153 (1899); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 124 (1889); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255, 261 (1903).

Tachydromia (of authors), Meigen, Illiger's Mag. Ind. Vol. 2, p. 269 (1803); Syst. Beschr. Vol. 3, p. 67 (1822); Zetterstedt, Fauna Ins. Lappon. p. 548 (1838); Dipt. Scand. Vol. 1, p. 138 (1842); Rondani, Dipt. Ital. Vol. 1, p. 146 (1856); Loew, Zeitschr. Ent. Breslau, Vol. 17, p. 4 (1863); Williston, Man. N. Amer. Dipt. p. 75 (1896); Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 42 (1902); Bezzi, Kat. Palæarkt. Dipt. Vol. 2, p. 279 (1903); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 407 (1907); Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 300, 402 (1909); Wahlgren, Ent. Tidskr. Vol. 31, p. 45, 112 (1910); Lundbeck, Dipt. Dan. Vol. 3, p. 285 (1910); Brunetti, Fauna Brit. Indian Dipt. Vol. 1, p. 374 (1920).

Characters. — Usually robust, and with sturdy legs, the middle femora more or less incrassated and biseriately setulose or spinose beneath, the middle tibiae shorter than their femora, raptorial, uniseriately setulose beneath and terminating within in a more or less developed spur : in color usually black though some species are testaceous. Head globular, sometimes a little higher than broad, the front broader than the face, but of variable breadth, sometimes with parallel sides sometimes diverging above, cheeks linear; eyes reniform, facets uniform; palpi flat and broadly oval, appressed to the proboscis, usually with one or two apical setæ, the front side with short incumbent hairs, the inner side glabrous; proboscis vertical, shorter than the head. Antennæ close together, located near the middle of the head, two- or three-jointed, the second joint globose, with a whorl of minute hairs, the third joint compressed, varying from short ovate to subulate, slightly pubescent, and always with a terminal arista which is closely but briefly pubescent. Ocelli three in number, placed on a low tubercle, one pair of proclinate ocellar bristles, in back of which are sometimes a pair of minor setulæ; one or two pairs of vertical bristles differentiated, the upper occipital setulæ usually dark, the lower more abundant and white. Thorax quadrate, the dorsum slightly convex, humeri rather distinct, thorax usually dusted, typically with a shining glabrous space on the sterno-pleuræ, sometimes the disk of the mesonotum devoid of pollen and highly polished, rarely densely pubescent, usually with biseriate acrostichal setulæ and uniseriate dorsocentral and sparse lateral setulæ, the hindmost dorsocentral setiform, chætotaxy including one or two humeral bristles, one or several notopleural, one supraalar, one postalar, two or four scutellar bristles, the lateral scutellars small. Abdomen with eight segments, shining, rarely with pollinose fasciæ or lateral markings, the subdorsal and lateral pittings sometimes evident though usually not noticeable on the shining integument, hairs sparse but moderately long; ovipositor short, shining or pollinose; pygidium more or less globular, asymmetrical, closed, comprising a spherical lower piece, covered by a dorsal usually fringed valve. Legs without true bristles, but the middle femora sometimes seriatel setose along the postero-flexor face and less evidently so along the anterior surface, the front femora with more or less developed biseriate flexor cilia, tibiae without apical setæ, last tarsal joint a little flattened, tarsi unusually long in *Cleptodromia*. Calypteres fringed with a few cilia; halteres never black. Wings hyaline or nearly so, not pictured but sometimes clouded, tip of first vein and costa between the first and third veins sometimes thickened, first vein ends beyond the middle of the wing, third vein straight or recurved toward the fourth, crossveins approximate, usually when the crossveins meet on the fourth vein the posterior crossvein is nearly transverse but when the posterior crossvein is located beyond the anterior it is distinctly oblique, anal crossvein angularly projecting from the fifth vein, the anal vein always faint but visible for more or less of its course, axillary lobe somewhat developed but obtusely rounded.

Type species : *P. cursitans*, Fabricius (**Pl. 4, Fig. 30**), by designation of Westwood, Introduction, Vol. 2, Synopsis, p. 132 (1840).

The species of this dominant genus are able to run and to fly well. The more pruinose species are campestral and are found through the warmer summer months, the testaceous and the shining black species are more fond of shady woods. The species are predatory, using their curiously formed middle legs for holding their prey. The larva of *P. major*, has been found in the ground under moss (Beling).

The genus *Platypalpus*, includes so many species that it has become unwieldy from a practical viewpoint. Furthermore, in this assemblage there are species so diverse that if contrasting forms were selected they would present differences striking enough to warrant a division of the genus. The pale yellow species scarcely resemble the black ones; those with stout legs and strong tibial spur are unlike the delicate forms with slender legs and no tibial spur. Rondani attempted a segregation on the basis of antennal structure, forming his genus *Phoroxypha* for those few species which have a short dark arista, centering about *longicornis*. Bezzi, in 1899, formed two groups, one where the antennæ are longer than the head, the other where they are shorter. Frey, in 1907, based his primary divisions on the length of the tibial spur. All of these characters, however, present too many gradations to have even subgeneric value. Even Corti's recent subgenus, *Cleptodromia*, founded on an abnormal tarsal structure, is scarcely more secure, although so far its solitary species presents a unique characterization.

In 1913, Brunetti described two related Indian genera of Empididæ, *Brevios* and *Howlettia*, locating them near *Bicellaria*. Later he suppressed *Brevios* as a synonym of *Platypalpus*, but retained *Howlettia* as a genus of anomalous position, placing it provisionally in the Empidinæ. In its description *Howlettia* agrees with *Platypalpus* so completely that in view of the complexity of this dominant genus it is here included as a synonym. The only distinctive character specified by Brunetti is the complete anal cell and anal vein, and it may be that in the strength of the anal vein *Howlettia* presents a more archaic condition than found in the other forms of *Platypalpus*.

A discussion of the usage of the terms *Platypalpus*, *Tachydromia* and *Coryneta* has previously been given by the writer in *Psyche*, Vol. 17, p. 41-48 (1910) and need not be repeated here. Briefly the case is this: Meigen described *Tachydromia* in 1803, citing *cursitans* and *cimicoides*. In 1827 Macquart segregated the genus *Platypalpus* for *cursitans* and its allies. Thus *cimicoides* becomes the only available type of *Tachydromia*. *Coryneta* is one of the aggravating genera of Meigen's 1800 paper. It was erected, like the others in this paper, without designation of type and, in our opinion, its genotype and its validity date from Hendel's interpretation in 1908.

SYNOPSIS OF THE NEARCTIC SPECIES OF PLATYPALPUS

1. Coxæ and femora largely or entirely black; body and antennæ black; veins dark	2.
Coxæ, femora and tibiæ largely or entirely yellow	16.
2. Front tibiæ thickened, wider than the hind tibiæ; front femora thickened; spur of the middle tibiæ strong; third antennal joint oval; thorax pollinose, with yellow bristles	3.
Front tibiæ not incrassate	5.
3. Posterior crossvein short, located beyond the anterior a greater distance than its length, the under side of the second basal cell greatly bowed upward distally, the first posterior cell wide near the middle; legs variegated	4

Posterior crossvein located near the anterior, second basal and first posterior cells normal; legs black except the brownish trochanters, knees, hind tibiae and bases of the tarsal joints	P. LUPATUS, Melander.
4. Hind femora and all the tibiae yellow, tarsi sharply annulate. Hind femora middle tibiae and middle tarsi black, front tarsi dark at the base	P. COQUILLETTI, Melander, ♂.
5. Some of the femora in part at least yellowish; crossveins meeting	P. PACHYCNEMUS, Loew.
All the femora black, except sometimes at the knees	6. 9.
6. Thorax shining, its bristles yellow; tibiae yellow, tarsi piceous; third antennal joint oval	7.
Thorax pollinose and pubescent, its bristles black; front legs entirely yellow, the other tarsi more or less annulate, spur of the middle tibiae very strong; third antennal joint lanceolate, equal to arista	P. DISSIMILIPES, nov. sp. (1).
7. Front opaque pollinose; antennae reddish; front legs entirely yellow, the middle femora black and with strong black setae, the hind femora black on the apical half, spur of middle tibiae long	P. VERSIPES, nov. sp. (2).
Front shining; antennae black; front legs partly dark, the middle femora without black setae, tibial spur short	8.
8. Front femora above and the posterior femora apically black, the base of all the femora yellow, anterior femora yellow-ciliate beneath, front coxae yellow	P. DIVERSIPES, Coquillett.

(1) *Platypalpus dissimilipes*, nov. sp. — Length 2 mm. Black, cinereous, subpollinose, legs yellow, the posterior femora black, bristles black, crossveins contiguous. Front and face somewhat whitish when viewed from below, two pairs each of long ocellar and vertical bristles; palpi small, whitish, with a long subapical dark hair; third joint of antennae lanceolate, twice as long as wide, the rather strong arista equal to the antennal length. Dorsum of thorax subshining although evidently pollinose, its bristles strong, one humeral, one notopleural, one supraalar, one postalar, one dorsocentral, four scutellar bristles, four rows of acrostichals and numerous lateral setulae; glabrous spot of sternopleura large; abdomen shining, each segment with a single subdorsal pit, pygidium small, its hairs sparse and black. Middle femora stout, coarsely black-setulose and spinose beneath, front femora sometimes brownish along the middle, underneath subciliate with pale hairs, spur of middle tibiae long and strong and with a black tip, last tarsal joint blackened. Costa not swollen at the insertion of the first vein, its second, third and fourth sections proportioned 3.5 : 2.4 : 1, third and fourth veins parallel, crossveins meeting, the posterior crossvein nearly transverse, anal crossvein forming an angle of eighty degrees, anal vein represented by a pronounced fold.

Fifteen specimens, type from Moscow Mountain, Idaho, June 12, 1910 (Melander). Paratypes from Moscow, Troy and Waha, Idaho; Mount Constitution, Winlock, Pluvius and Ilwaco, Washington; Eagle Creek, near Portland, Oregon (Melander); Kaslo, British Columbia (R. P. Currie); and Colorado (C. F. Baker).

(2) *Platypalpus versipes*, nov. sp. — Male. Length 2.2 mm. Mesonotum shining, head pollinose, front legs yellow, middle femora and apical half of the hind femora black. Sides of the opaque front slightly diverging above, face short, as broad as the bottom of the front, anterior part of the epistome pollinose; palpi yellow, proboscis blackish; vertical and ocellar bristles black; antennae brown, the third joint short-ovate, the arista less than twice as long as the antenna. Sides of the thorax lightly pruinose, sternopleura and a central spot on the hypopleura glabrous, bristles blackish, notal hairs moderately long and blackish. Abdominal hairs short and sparse, pygidium small and nearly bare. Front coxae yellow, posterior coxae blackish, anterior femora robust, the front pair not ciliate, middle femora setulose with black bristles in back, posterior tibiae and tarsi yellow, the last joint of the tarsi dusky, the middle patella black, middle tibiae two-thirds as long as their femora, the spur longer than the diameter of the tibia and tipped with black. Veins yellowish, third and fourth veins parallel, the first posterior cell widest beyond its middle, second, third and fourth sections of the costa proportioned 1 : 0.6 : 0.2, crossveins meeting, sections of the fourth vein, 1 : 1.8; of the fifth vein 0.5 : 0.6 : 1, base of the anal vein wanting, marginal cilia shorter than the anterior crossvein.

One specimen, Falls Church, Virginia, May, received from Nathan Bauk.

Anterior femora brown, hind femora black, none of the femora yellow at the base, nor ciliate, the front coxae piceous . . .	P. DIREPTOR, nov. sp. (1).
9. Thorax pollinose, not or scarcely shining	10.
Mesonotum shining, except more or less at the humeri and along the lateral edge, its bristles yellow	14.
10. Bristles of the thorax black; third antennal joint lanceolate or elongate oval	11.
Bristles of the thorax yellow; third antennal joint oval	13.
11. Spur of the middle tibiae long; posterior crossvein beyond the anterior and oblique; anterior femora ciliate with yellow hairs beneath	P. PLUTO, Melander.
Spur of the middle tibiae moderate; crossveins meeting, the posterior transverse; femora not ciliate, but the middle femora black-spinose beneath	12.
12. Third antennal joint much shorter than arista and not hairy; halteres and calypteres blackish; front femora robust . . .	P. LACERTOSUS, nov. sp. (2).
Third antennal joint very long and pubescent, longer than arista; halteres and calypteres pale; front femora not very stout	P. SPINOSUS, nov. sp. (3).

(1) **Platypalpus direptor**, nov. sp. — Male. Length 1.9 mm. Black, dorsum shining, bristles yellowish, pleurae cinereous, antennae short, legs brown, the hind femora black, no tibial spur nor femoral cilia, crossveins nearly meeting and transverse. Occiput and face cinereous, front moderately broad and very lightly dusted, epistome shining; palpi short and blackish; antennae two-jointed, the outer joint short-ovate, half as long as the arista; two pairs of black vertical and one of ocellar bristles. Humeri, notopleural suture, scutellum, metathorax and pleurae densely cinereous pruinose, the sternopleurae largely glabrous; two humeral, one notopleural, one supraalar, one postalar, two pairs of scutellar and two of small dorsocentrals, setulae very sparse and fine, seven biseriate acrostichals; abdomen shining, pygidium rather large and spherical. Front coxae and anterior femora except the knees brown, posterior coxae and hind femora black, tibiae and bases of the tarsi yellowish brown; anterior femora stout, neither ciliate nor setose, front tibiae as thick as the others. Veins brown, costa not thickened, its second, third and fourth sections proportioned 2 : 1.6 : 1, third vein straight, ending much before the apex of the wing, fourth vein slightly undulating, posterior crossvein at two-fifths the length of the wing, slightly oblique and arising a little before the anterior, sections of the fifth vein proportioned 0.5 : 0.5 : 1, anal crossvein forming an angle of fifty degrees, anal vein indicated only as a faint fold, marginal cilia short.

One specimen, Nelson, British Columbia, July.

(2) **Platypalpus lacertosus**, nov. sp. — Female. Length 2 mm. Body, antennae, palpi, halteres, coxae and femora except knees, black; mesonotum densely brown-pollinose, bristles black, legs short and stout, tibial spur vestigial, middle femora setose beneath, crossveins contiguous. Occipital hairs setiform, black, the two vertical and the ocellar bristles strong and black, front and face each short and wide, pollinose, epistome shining, proboscis stout, black, palpi with subapical black setæ; antennæ very short the outer joint ovate, slightly longer than wide and three-fifths as long as the arista. Mesonotum with abundant black setulae, the bristles strong, including the humeral, scutellars widely separated, a strong lateral scutellar present; pollen of pleurae brownish gray, sternopleura almost wholly polished. Abdomen shining, the hairs short and inconspicuous, last two segments dull, styles long and slender. Knees, tibiae and tarsi brown, front femora robust, front tibiae strong though not incrassate, middle femora with about twenty-five minute setulae in each row and behind with a series of six long black bristles, middle tibiae scarcely two-thirds the length of the femora, tarsi not annulate and scarcely darker apically. Calypteres, fringe and halteres blackish. Wings hyaline, two and one-half times as long as wide, veins thin but firm and black, costal sections proportioned 5 : 3.5 : 2.5 : 1, first posterior cell ending at the wing-tip, a little widest at the middle, third vein straight, fourth vein lightly sinuous, sections of the fifth vein proportioned 1 : 1 : 2, anal crossvein forming an angle of fifty-five degrees, anal vein very weak, anal angle rather pronounced.

Two specimens, North end of Lake Coeur d'Alene, Idaho, August 24, 1916 (Melander). The species is related to *hians* and *luctator* but is readily distinct in its blackened legs.

(3) **Platypalpus spinosus**, nov. sp. — Male. Length 2.2 mm. Black, the legs piceous. The ocellar and the two pairs of vertical bristles strong and black; third antennal joint pubescent and slender, three and one-half times as long as broad, the arista equal in length to the third joint; face and front cinereous; palpi small, white,

13. Slender; legs including the coxae entirely black; pleurae and front coxae pruinose; discal dorsocentrals present *P. TACHISTIFORMIS*, nov. sp. (1).
- More robust; legs piceous, the front coxae, the middle trochanters and the tibiae yellowish brown; pleurae and front coxae silky-pollinose; bristles minute. *P. SERICATUS*, nov. sp. (2).
14. Anterior femora ciliate beneath; posterior crossvein beyond the anterior; middle tibiae with a sharp spur; four scutellar bristles 15.
- Femora not ciliate; crossveins meeting; no tibial spur; third antennal joint oval; two scutellar bristles *P. CONTIGUUS*, nov. sp. (3).

with a few fine black hairs. Thoracic dorsum with golden pollen, bristles black, one humeral, one notopleural, one supraalar, one small prescutellar and two scutellar bristles; pleurae more closely but finely pollinose, the sternopleurae largely shining; abdomen shining, white-hairy, pygidium large. Spur of the middle tibiae moderately large, black, the postero-flexor edge of the middle femora with thirteen black spine-like bristles, the anterior side of the middle tibiae also with smaller more scattered black bristles, under side of front femora with small pale separated cilia. Costa greatly thickened at the confluence with the first vein, the second, third and fourth sections proportioned 4 : 3 : 1, third and fourth veins subparallel, crossveins meeting, the posterior transverse, anal vein weak but uniformly perceptible, anal crossvein forming an angle of sixty degrees, abruptly attenuated beyond its basal third.

One specimen, Stanford, California, January 27, 1906, collected by Dr. J. M. Aldrich.

(1) *Platypalpus tachistiformis*, nov. sp. — Female. Length 2 mm. A slender black species that resembles certain species of *Tachyempis*, but has a distinct anal crossvein and a spur on the middle tibiae. Head and thorax cinereous pollinose becoming pruinose on the pectus and pleurae, legs black, the front knees narrowly brown. Palpi elliptical, one-half as long as the proboscis, black but thickly overlaid with white pruinosity and with a few white hairs; first antennal joint entirely wanting, the last joint globular, scarcely longer than wide, the rather thick arista one third longer than the antennae; the ocellar and two pairs of vertical bristles rather strong and blackish. Bristles of the thorax prominent and yellow, four dorsocentrals, six rows of acrostichals, six scutellar bristles; sternopleurae with a small rounded glabrous spot; sides of the first abdominal segment pollinose, the last three segments opaque brown pollinose. Front femora rather thick, middle femora loosely ciliate with short bristle-like yellow hairs on the postero-flexor edge, front coxae with abundant golden bristles. Costa not swollen, its second, third and fourth sections proportioned 2.5 : 2.5 : 1, third and fourth veins parallel, anterior crossvein slightly beyond the oblique posterior crossvein, sections of the fifth vein subequal, anal crossvein complete, forming an angle of seventy degrees, anal vein indicated by a fold.

Two specimens, Harlowton, Montana, June 14, 1913 (Melander).

(2) *Platypalpus sericatus*, nov. sp. — Female. Length 2 mm. Black, legs piceous, the front coxae, apex of posterior coxae, base of femora, apex of front femora, and the tibiae yellowish-brown. Third antennal joint oval, one-half longer than wide, the arista one and two-thirds times as long as the antennae; face, front and palpi white-pruinose, the latter one-third as long as the proboscis; ocellar bristles brown. Dorsum of the thorax with yellow pollen, the minute sparse hairs and the bristles yellow, one humeral, one notopleural, one prescutellar, four scutellar, the lateral pair small; pleurae and front coxae with silky whitish-yellow pollen, the sternopleurae largely shining; abdomen polished. Anterior femora ciliate with yellow hairs, spur of the middle tibiae long and strong. First vein swollen on entering the costa, the second, third and fourth sections of the costa proportioned 3.2 : 2.5 : 1, posterior crossvein oblique, located beyond the anterior a distance equal to the length of the anterior crossvein, fourth and fifth veins parallel, fifth vein interrupted before reaching the margin; base of anal vein completely wanting, the remainder indicated as a fold.

One specimen, Nelson, British Columbia, July, 1910.

(3) *Platypalpus contiguus*, nov. sp. — Length 2 mm. Black, largely shining, legs black, crossveins contiguous. Third antennal joint short-oval but broad, the arista not much longer than the antennae; face and front cinereous pollinose; palpi small, one-fourth as long as the proboscis, blackish; cephalic hairs blackish. Humeri, notopleural sutures, scutellum and metanotum cinereous pollinose, otherwise the notum is polished, the bristles and hairs yellowish, three humeral hairs, three notopleural, one supraalar, one scutellar, two pairs of dorsocentral bristles, acrostichals biserrate and small; pleurae white-pruinose, the sternopleurae polished; abdomen shining, the pygidium rather erect. Posterior coxae and femora and the tarsi apically, black, femora not ciliate, no tibial spur. Costa only slightly swollen at the confluence with the first vein, the second, third and fourth sections proportioned 2.5 : 1.6 : 1, third and fourth veins parallel, anal crossvein evanescent, forming an angle of fifty degrees, anal vein indicated only by a fold in its outer part.

Bear Lake, British Columbia, 7000 feet altitude, July 29, 1903, collected by R. P. Currie for the U. S. National Museum.

15. Legs entirely black; third antennal joint oval	P. MONTICOLA, Melander.
Tibiae and front knees brown, tarsi annulate; third antennal joint lanceolate	P. INFERIALIS, nov. sp. (1).
16. Thorax yellow in large part; veins yellowish.	17.
Thorax wholly black, although more or less overlaid with pale pollen	35.
17. Head entirely yellow; front femora with seriate flexor bristles	18.
Head globular, the occiput at least black	20.
18. Spur of middle tibiae strong; posterior crossvein oblique and distant from anterior; sides of front nearly or quite parallel	19.
Spur of middle tibiae undeveloped; crossveins meeting, the posterior transverse; head globular, sides of front diverging above.	
	P. DECOLOR, nov. sp. (2).
19. Head globular; antennæ tipped with black; thorax shining; third vein recurved	P. MELLEUS, nov. sp. (3).

(1) **Platypalpus inferialis**, nov. sp. — Length 2.2 mm. Occiput cinereous, one pair of brown vertical bristles, front and face cinereous when viewed from above, white when seen from below; palpi one-third as long as the proboscis, whitish and with a few white hairs; third antennal joint lanceolate, twice as long as broad, the arista slightly longer than the antennæ. Dorsum of the thorax shining black except that the lateral and posterior margins are cinereous pollinose, its hairs and bristles fine and yellow, one dorsocentral, one supraalar, one notopleural and two pairs of scutellar bristles; pleuræ whitish pruinose, the notopleural suture more olivaceous, sternopleuræ with large triangular shining space; abdomen subshining, its hairs not long, the pygidium rather small, the terminal valve with a fringe of silky hairs above and the left valve similarly fringed beneath. Front and middle femora stout, with long loose white seriate hairs beneath, spur of the middle tibiae large; femora black, knees and tibiae brown, the hind pair darkest, tarsi annulate, but usually the black predominating. Costa thickened at the end of the first vein, its second, third and fourth sections proportioned 3.5 : 2.5 : 1, fourth vein slightly sinuous but in the main subparallel with the third, posterior crossvein oblique, separated from the anterior by one-half the length of the anterior crossvein, sections of the fifth vein proportioned nearly 3 : 2, anal crossvein located at one-third the length of the second basal cell, nearly perpendicular, anal vein indicated merely by a fold.

Numerous specimens, type from Almota, Washington, April 17. Paratypes from Pullman and Oroville, Washington (Melander) and Juliaetta, Idaho (J. M. Aldrich).

(2) **Platypalpus decolor**, nov. sp. — Female. Length 2.5 mm. Entirely pale yellow, the arista, ocellar spot, apex of abdomen and last tarsal joint alone brownish. Head globular, sides of the pollinose front diverging, epistome shining, palpi small; last antennal joint slender, lanceolate, two and a half times as long as broad and slightly longer than the short-pubescent arista. Mesonotum polished except along the sides where it is lightly dusted, nearly bare, the few hairs seriate, humeral bristle strong, middle scutellars long; pleuræ dusted except most of the sternopleuræ; abdomen shining. Legs slender, front femora biseiately fringed beneath with about ten bristles, middle femora with about eighteen slender spinules in each flexor row, those behind longer, middle tibiae obliquely truncate without spur. Veins pale yellow, crossveins at two-fifths the wing-length, the posterior transverse and slightly proximal to the anterior, first posterior cell ending at the wing-tip, its veins parallel, base of the anal vein weakly indicated, anal crossvein slightly recurved, sections of the costa proportioned 6 : 4 : 2 : 1.

Two specimens, Foot of Lake McDonald, Glacier National Park, Montana, August 14, 1916 (Melander).

(3) **Platypalpus melleus**, nov. sp. — Female. Length 2 mm. Yellow, the ocellar spot, outer half of the third antennal joint with the arista, the eyes, flexor setulæ of middle legs, apex of tibial spur and last tarsal joint alone blackish. Back of occiput shining, front and face moderately broad, white, the face three-fourths as long as the front, epistome shining; third joint of the antennæ pyriform, one-half longer than broad and about one-third as long as the arista. Thorax shining above, setulæ very sparse and small, pleuræ white-pruinose except the moderate sternopleural spot; abdomen shining except the last two segments. Legs rather slender, the middle femora moderately inflated, front femora scarcely ciliate, middle femora with a posterior row of eight silky yellow setæ, middle tibiae two-thirds as long as their femora. In *tersus* the front femora are somewhat swollen and ciliate, and the middle tibiae are three-fourths as long as their femora. Wings narrow, the second, third and fourth sections of the costa proportioned 4.5 : 3.5 : 1, third vein recurved but ending just before the tip of the wing, posterior crossvein slightly oblique, located its own length beyond the anterior and at three-sevenths the length of the wing, anal crossvein perpendicular. sections of the fifth vein proportioned 0.4 : 0.6 : 1, base of the anal vein wanting, marginal cilia as long as the anterior crossvein.

One specimen, Gowanda, New-York, received from M. C. Van Duzee.

- Head narrow, the occiput conical; antennæ whitish; humeri pollinose; third and fourth veins parallel P. TERSUS, Coquillett.
20. Thorax with a median broad blackish vitta 21.
- Thorax without a median vitta 26.
21. Proboscis black; wings elongate, the posterior crossvein beyond the anterior and oblique; tibial spur large P. CALIGARIS, new name.
- Proboscis mostly yellow; posterior crossvein transverse 22.
22. At least sides of thorax lightly pollinose; front densely pollinose; anal crossvein nearly perpendicular 23.
- Thorax shining; front shining; middle femora with yellow setæ beneath in addition to the biseriate setulæ; anal crossvein strongly recurved 24.
23. Crossveins widely separated; middle femora setose as well as setulose P. RUBEFACTUS, nov. sp. (1).
- Crossveins meeting; middle femora not setose; northern species 25.
24. Third antennal joint short ovate, yellow, one-third the arista; flexor setæ of middle femora strong; crossveins at one-third length of wing; mesonotal hairs rather long and sparse; face silvery P. MESOGRAMMUS, Loew.
- Third antennal joint lanceolate, black, two-thirds the arista; flexor setæ weak; crossveins at two-fifths length of wing; hairs short and dense; face shining P. XANTHOCHITON, nov. sp. (2).

(1) *Platypalpus rubefactus*, nov. sp. — Male. Length 1.75 mm. Body largely yellowish. Occiput black, hoary, its hairs long, sparse and white, sides of the concolorous front somewhat diverging above, epistome bare, proboscis yellowish, palpi rather large, white-pubescent; last antennal joint slightly infuscated apically, short-conical, one-half longer than broad and one-third as long as the thickened dark arista. Mesonotum dusky, a dark vitta occupying the central third, dorsal setulæ very sparse and seriatly arranged in four rows, humeral bristle weak, two scutellars; pleuræ pruinose, about half of sternopleura bare. Abdominal tergites brownish yellow, hypopygium brown, terminal, larger than the diameter of the abdomen, its valves closely fringed with long curled yellow hairs. Legs entirely yellow, the apex of the tarsi infuscated, front femora with a half dozen outstanding hairs beneath, middle femora rather slender, with about twenty minute setulæ in each flexor row, in front and in back of which are outstanding yellow hairs, those in back fewer and much longer, spur of middle tibiae minute. Wings lutescent, three and a half times as long as wide, veins yellow, the third to fifth sections of the costa proportioned 6 : 3 : 1, first posterior cell narrow, occupying one-fourth the width of the wing, its veins parallel, crossveins before the middle of the wing, the posterior transverse, spaced its own length from the anterior, anal crossvein at the middle of the basal cell, perpendicular, anal vein evanescent basally.

Type: Chicago, Illinois, July, 1914 (Melander). Paratypes, Plummer's Island, Maryland (McAtee). The female has the telescopic tip of the abdomen opaque yellow.

(2) *Platypalpus xanthochiton*, nov. sp. — Female. Length 2.5 mm. Yellow, head black, mesonotum with a central brown vitta, tergites medially brown, legs entirely pale yellow, veins thin and blackish, crossveins meeting. Head globular, occiput dull, its upper hairs dense and pale, front moderately broad and shining, face and epistome shining; proboscis yellow, palpi large and pale yellow; base of the antennæ shining yellow, distal joint triangular, black, nearly twice as long as broad and about half the length of the finely pubescent arista. Disc of the mesonotum devoid of pollen but uniformly clothed with rather dense short fine whitish hairs, no humeral, two notopleural and the middle pair of scutellar bristles yellow. The dark vitta is narrow in front becoming broader and diffusing posteriorly; base of the scutellum brown. Abdominal hairs inconspicuous; terminal segments not ochraceous. Legs slender, front femora not ciliate, middle femora with twenty-five setulæ in the anterior row and twenty in the posterior and with about ten weak yellow posterior setæ, middle tibiae two-thirds as long as their femora, the spur small and yellow. Halteres, calypteres and root of wing pale yellow. Sections of the costa proportioned 10 : 7 : 5 : 2; of the fourth vein, 1 : 1.4, of the fifth vein, 2 : 3 : 4, third vein ending at extreme wing-tip, anal crossvein forming an angle of fifty degrees, anal vein faint, anal angle weak.

Two specimens, Almota, Washington, June 2, 1918 (Melander).

25. Mesonotum uniformly pollinose; third antennal joint tipped with black; anal crossvein and anal vein distinct; face as long as the front; middle tibiæ much shorter than the femora
- Sides of the notum pollinose; antennæ yellow; anal crossvein and anal vein very faint; face much shorter than the front; middle tibiæ nearly as long as their femora
26. Thorax densely yellow-pollinose; antennæ wholly or quite yellow; middle femora setose
- Thorax shining or subshining; third antennal joint usually darkened
27. Posterior crossvein beyond the anterior; tarsi yellow
- Crossveins meeting; tarsi narrowly annulate; proboscis yellow
28. Proboscis yellow; antennæ entirely pale yellow; abdomen ♀ sometimes with a narrow median vitta
- Proboscis black; third antennal joint tipped with black; middle segments of the abdomen black.
- P. FLAVIROSTRIS, Loew.
var. VITTIGER, nov. var. (1).
- P. MELANOGASTER, nov. sp. (2).
- 27.
- 29.
- 28.
- P. IMPEXUS, Melander.
- P. VERSUTUS, Melander.
- P. OCHRICOLLIS, nov. sp. (3).

(1) **Platypalpus flavirostris**, var. **vittiger**, nov. var. — Male. Length 3.6 mm. Head black, cinereous pollinose, mouth-parts and base of the antennæ yellow, the third antennal joint about one-third the arista, elongate-oval, mesothorax lightly pollinose over an ochraceous ground, a median blackish vitta about one-fourth the width of the notum; base of the pollinose scutellum narrowly blackish; most of the sternopleuræ smooth. Abdomen blackish, shining. Legs yellow, the last tarsal joint blackish, middle femora with five yellow bristles and twenty-six black setulæ on the antero-flexor edge and nineteen bristles on the postero-flexor edge, spur of middle tibiæ small. Wings yellowish hyaline, the veins pale, crossveins touching, the posterior nearly transverse, third vein ending at the wing-tip, reflexed and parallel with the fourth, under side of the second basal cell slightly shorter than the ultimate section of the fifth vein, anal vein rather weak and forming an angle of eighty degrees.

Type, Douglas, Alaska, August 7, 1901, collected by the late E. L. Jenne. A series of similar specimens was taken by the writer in Glacier National Park, Montana, August 14, 1916. A dozen additional specimens from the same locality lack the mesonotal vitta and approach the typical form of *flavirostris*.

Differs from *mesogrammus* Loew in having the thorax pollinose, not shining, the third antennal joint black and the middle femora more slender and with only the usual two rows of black setulæ beneath. In *mesogrammus* there is an additional row of about ten long pale bristles along the postero-flexor edge.

(2) **Platypalpus melanogaster**, nov. sp. — Length 1.8 mm. Head and abdomen black, thorax yellow with a median castaneous vitta, legs pale yellow. Occiput lightly cinereous, its hairs white; antennæ inserted low down on the head, the front about twice as long as the face; two pairs of ocellar bristles but apparently no vertical ones; front moderately broad, its sides slightly diverging above, minutely but densely whitish pubescent rather than pollinose, face one-half as broad as the front, whitish, epistome black; proboscis reddish; antennæ short, yellow, the third joint short-pyriform, apparently ovate but really pointed, the arista brown, rather coarse, two and one-half times the actual length of the third joint. Thorax short, quadrata, broader than the head, humeri large, squarish, sides of the notum thinly white pollinose, disk polished and nearly bare; pleuræ lightly white-pollinose, the sternopleuræ largely shining; abdomen shining, blackish, its hairs sparse, the pygidium moderate in size, the dorsal valve short and rugose apically; abdomen of female tapering, the last two segments opaque ochraceous, styles slender and yellow. Legs long and slender, simple, middle tibiæ five-sixths as long as their femora, the flexor setulæ of the middle legs rather long, thin and yellow, no extra setæ, no tibial spur. Veins pale but towards the apex becoming brownish, costa not thickened, its second, third and fourth sections proportioned 3 : 1.8 : 1, third and fourth veins rather widely separated, straight and parallel, posterior crossvein transverse, located a little before the anterior and at three-sevenths the wing length, fifth vein very long, its sections proportioned 0.5 : 0.5 : 1, anal crossvein perpendicular, very faint, anal vein very faint and close to the margin.

Three specimens; Ungava Bay, near northern Labrador, collected by L. M. Turner. Type in the U. S. National Museum.

(3) **Platypalpus ochricollis**, nov. sp. — Female. Length 2 mm. Head black, densely cinereous pollinose, the bristles pale yellow, proboscis black, palpi pale yellow, antennæ yellow, the lanceolate third joint tipped with black. Thorax ochraceous, overlaid with dense whitish pollen, sternopleuræ with a small glabrous spot;

29. Proboscis entirely black; spur of the middle tibiae large; anal crossvein perpendicular 30.
 Proboscis yellow; tibial spur rather small; first posterior cell not much narrowed 31.
30. Mesonotum shining, pleurae mostly shining; first posterior cell somewhat narrowed in the margin; tarsi weakly annulate. R. LETUS, Loew.
 Mesonotum dusted, subshining, pleurae pruinose except the sternopleural spot; fourth vein not noticeably curved forward; tarsi not at all annulate. P. LETABILIS, nov. sp. (1).
31. Front narrow, its sides nearly parallel; antennae entirely yellow; base of the anal vein wanting 32.
 Front rather broad, its sides diverging above; third antennal joint blackened; anal crossvein oblique, anal cell closed, crossveins nearly meeting (P. flavirostris, Loew) 33.
32. Anal crossvein perpendicular, crossveins separated. P. MIMUS, nov. sp. (2).
 Anal crossvein oblique, crossveins meeting; penis hook-like, right valve with thorn-tipped process P. UNCIATUS, nov. sp. (3).

abdomen glistening, yellow at base and apex, the fourth and fifth segments blackish, the third segment somewhat brown. Legs yellow, the last tarsal joint dusky, middle femora with sixteen minute black setulae on the antero-flexor edge, twenty on the postero-flexor edge, in back of which are about nine long yellow bristles, spur of middle tibiae long, acute and tipped with black. Wings yellowish hyaline, veins pale, the third and fourth veins parallel, the first posterior cell ending full at the tip of the wing, posterior crossvein beyond the anterior by the length of the latter, sections of fifth vein equal, anal vein weak.

One specimen, Pullman, Washington, June 16, 1912 (Melander).

(1) **Platypalpus letabilis**, nov. sp. — Female. Length 3.2 mm. Very close to *letus*, differing in having the mesonotum dusted so as to be subshining, the pleurae completely white-pruinose except for the sternopleural spot, and the third and fourth veins subparallel towards the end. Head cinereous, body testaceous, front and face narrow, densely white-pruinose; proboscis black, palpi white; third antennal joint blackish, ovate, one-half longer than wide and one-half as long as the arista. Thoracic setulae very sparse; abdomen shining except the terminal segments. Front femora with sixteen flexor cilia; middle femora yellow-setose in back, middle tibiae two-thirds as long as their femora, the terminal spur long and sharp and tipped with black, last tarsal joint blackish, otherwise the tarsi not annulated. Segments two, three and four of the costa proportioned 5:3:1, first vein a little thickened apically, crossveins almost meeting, the posterior slightly oblique, located at three-sevenths the length of the wing, sections of the fifth vein proportioned 0.6:0.8:1, anal crossvein perpendicular, base of the anal vein very weak.

One specimen, South Wales, New-York, presented by M. C. Van Duzee.

(2) **Platypalpus mimus**, nov. sp. — Female. Length 2 mm. Pale yellow, the arista and occiput gray, the minute flexor setulae of the middle legs black. Front and face narrow, white, their sides subparallel; antennae whitish, the third joint ovate, three-fifths as long as the arista; lower occiput and proboscis yellow, palpi small and white. Mesonotum very lightly dusted, a little more pronounced on the humeri, subshining, setulae very sparse, humeral bristle large; pleurae finely white-pruinose except the sternopleural spot; abdomen shining to the end. Anterior femora moderately thickened, the front ones with a dozen small flexor cilia, the middle ones with a posterior row of fine pale setae, middle tibiae two-thirds as long as their femora, the terminal spur short blunt and yellow. Wings narrow, third and fourth veins parallel, second, third and fourth sections of the costa proportioned 3.5:2.3:1, posterior crossvein located at three-sevenths the winglength, slightly oblique, placed beyond the anterior a distance equal to the latter, sections of the fifth vein proportioned 0.5:0.6:1, anal crossvein almost perpendicular, base of anal vein completely wanting, marginal cilia equal in length to the anterior crossvein.

Two specimens received from M. C. Van Duzee, who collected them at Gowanda and Niagara Falls, New York, June.

(3) **Platypalpus unciatus**, nov. sp. — Male. Length 2.2 mm. Light yellow, head black, occiput subshining, front narrow, shining, but slightly divergent above, mouthparts yellow, antennae pale yellow, outer joint triangular, arista brownish, twice as long as last joint. Thorax shining, scutellum sometimes brown, bristles brown, notal hairs pale and relatively long. Abdomen centrally dark, pygidium long, asymmetrical, open, the central filament consisting of a long strong curved hook, with black point, the right valve ending above in a long

33. Third antennal joint orbicular, one-fifth the arista; epistome more or less pollinose	var. <i>MICROCERUS</i> , nov. var. (1).
Third antennal joint tapering and about one-third the arista; epistome polished.	34.
34. Stout, abdomen brownish; third antennal joint ovoid; middle femora with 20 to 23 spines in the posterior flexor row	var. <i>FLAVIROSTRIS</i> , Loew, s. str.
Slender, abdomen yellow; third antennal joint lanceolate; middle femora with 18 to 20 flexor spines	var. <i>DILUTIOR</i> , nov. var. (2).
35. Mesonotum shining, pollinose only along the lateral and posterior edges, or with shining vittæ	36.
Mesonotum more or less densely but uniformly pollinose, not completely shining	67.
36. Crossveins nearly or quite contiguous, the posterior usually transverse; tibial spur small	37.
Posterior crossvein beyond the anterior and oblique; spur of the middle tibiæ large and strong; anterior femora ciliate.	62.
37. Arista white, contrasting with the antennæ; thorax nearly bare of hairs and bristles	38.
Arista black or brown, not contrasting	39.
38. Arista much longer than the lanceolate third antennal joint; front and face narrow; legs brown, middle femora ciliate with long black bristles; third and fourth veins converging, anal crossvein very oblique.	P. <i>CALLITHRIX</i> , nov. sp. (3).

finger-like process, which is tipped with a black thorn, the left valve large and broadly triangular. Middle femora closely mucronate below, tibial spur undeveloped. Wings hyaline, veins deep yellow, crossveins meeting, lower outer angle of second basal cell sixty degrees, anal crossvein oblique, base of anal vein obliterated.

Four males and four females, La Suiza, Costa Rica, April-August, Pablo Schild, collector.

(1) *Platypalpus flavirostris* Loew, var. *microcerus*, nov. var. — An interesting variation occurs on Moscow Mountain, Idaho. Five specimens caught during July and August, 1916 and 1918, have the front of the epistome pollinose like the face, the third antennal joint orbicular and shorter than wide so that it is one-fifth as long as the arista. Other specimens captured with these have the last antennal joint short-lanceolate as usual, making the arista about three times this joint, and the epistome is polished. About twenty-three spines occur in the posterior flexor row of the middle femora. The anal crossvein forms an angle of sixty degrees.

(2) *Platypalpus flavirostris* Loew, var. *dilutior*, nov. var. — A slender pale yellow form of the species from Vashon Island, and numerous other places in Western Washington, with lanceolate antennæ, eighteen to twenty flexor spines in the posterior row on the middle femora, the female with the underside of the second basal cell four-fifths as long as the continuation of that vein, the male with the two parts of the fifth vein nearly equal, the anal vein steeply inclined forming an angle of about seventy-five degrees, and the uppermost lamella of the pygidium broadly excised on the outer side, the proximal lobe with four golden bristles.

The typical form of *flavirostris* is stout, brownish, with ovoid third antennal joint, with twenty to twenty-three spines in the posterior flexor row of the middle femora, the two parts of the fifth vein equal and the anal crossvein forming an angle of sixty degrees.

(3) *Platypalpus callithrix*, nov. sp. — Female. Length 3 mm. Black, disk of the thorax shining and nearly bare, legs yellowish, slender, the middle femora with unusually long black flexor setæ, tibiæ without spur, third antennal joint elongate and about one-third as long as the white arista, front and face almost linear, third vein recurved, crossveins approximate, the posterior crossvein oblique, anal crossvein very steep. Occiput blackish, subshining, with a postocellar cordiform cinereous spot, its hairs sparse, facets of the eyes comparatively large, palpi brownish, with a subapical seta; antennæ three-jointed, black, the last joint pubescent, lanceolate, twice as long as wide, with a rather thick white and white-pubescent arista, the shaft of which in the type specimen is black on its middle third. Thoracic bristles short and black, sides of the thorax behind the humeri, and the scutellum rather thinly cinereous pruinose, pleuræ glabrous except their posterior margin, abdomen brownish black, the last two segments pollinose. Front femora with a few long black flexor setæ, middle femora with black setæ

- Arista much shorter than the elongate third antennal joint; front and face broad; hind femora blackish apically, middle femora with fine yellow cilia; third and fourth veins parallel P. FLAMMIFER, Melander.
39. Anal cell completely lacking; antennæ black, the third joint broadly oval and one-half the arista; palpi yellow; front shining; tarsi dusky distally, not annulate P. CUNEIPENNIS, Melander.
- Anal crossvein and part of anal vein evident, as usual 40.
40. The elongate third antennal joint and arista pubescent; wings marked with blackish spot over crossveins P. PICTIPENNIS, Bezzi.
- Arista bare; wings not spotted 41.
41. Base of the antennæ yellow; sides of front parallel or nearly so; coxae and legs yellow 42.
- Antennæ completely black: if the base of the antennæ is brownish the sides of the front diverge above and the face is broad 47.
42. Front opaque, pleurae mostly pollinose; tarsi weakly annulate. Front and greater part of the pleurae shining; only the last joint of the tarsi dark. 43.
43. Face much narrower than the front; proboscis yellowish; last two segments of the ♀ abdomen ochraceous; femora scarcely thickened, no tibial spur; hairs of the notum long; third antennal joint long, hairy and equal to the rather stout arista 46.
- Face much narrower than front; only last tarsal joint blackened; proboscis brown; abdominal segments shining except on incisures; legs rather slender, no tibial spur; notal hairs sparse and moderate; third antennal joint hairy and equal to the rather stout arista; pygidium robust, forming about one-third the abdomen P. TENUIS, nov. sp. (1).

on each side of the usual two rows of black setulae, those behind eight in number and longer than the diameter of the femur, middle tibiae two-thirds as long as their femora. Wings lightly infumated, veins brown, costa not at all thickened, its second, third and fourth sections proportioned 4.2 : 2.2 : 1, pedicel of the second and third vein arising near the base of the second basal cell, third vein ending beyond the apex of the wing, convergent with the fourth, the first posterior cell widest at three-fifths its length, crossveins near the basal third of the wing, the posterior arising slightly before the short anterior cross vein, sections of the fifth vein proportioned 0.4 : 0.5 : 1, anal crossvein at an angle of thirty degrees, evanescent at its tip, anal vein very weak, vanishing at its base.

Type, Chicago, Illinois, August 10, 1901 (Melander). Another specimen from Niagara Falls, New York, submitted by M. C. Van Duzee, has the arista wholly white and the third vein a little less recurved.

(1) *Platypalpus tenuis*, nov. sp. — Male. Length 2.2 mm. Black, legs entirely pale yellow except the black outer two-thirds of the last tarsal joint, base of antennæ yellow, veins and bristles whitish, disk of mesonotum, polished, front and pleurae pollinose, crossveins approximate and transverse. Hairs of the occiput long sparse and white; face narrower than the front, gray; epistome shining; proboscis brown, palpi small and yellow; third antennal joint brown, pubescent, triangular, over twice as long as wide and slightly longer than the closely pubescent black arista. Humeri, scutellum, narrow sides of the mesonotum and pleurae except sternopleura whitish gray pollinose, the usual bristles pale yellow, acrostichal and dorsocentral hairs individually distinct, together forming four rows, the interspaces glabrous, lateral hairs very sparse. Abdomen entirely shining black, nearly bare, last sternite with a conspicuous fringe of pale yellow hairs, pygidium forming a large rounded termination to the abdomen, its valves not fringed. Legs rather slender, front femora not ciliate, middle femora with about twenty fine blackish setulae in the anterior row and about twelve fine yellowish setæ in the posterior row, no other bristles,

- Face as broad as the front; proboscis brown to black; terminal segments of the ♀ abdomen shining; anterior femora more or less thickened; notal hairs short and very sparse. 44.
44. Penultimate segment of ♀ abdomen long and swollen; pygidium enormously enlarged, nearly as bulky as the remainder of the abdomen; proboscis brown; third joint of the antennæ ovate, shorter than the arista; middle tibiæ with a pronounced spur. P. SATYRIACUS, nov. sp. (1).
- Terminal segments of the ♀ abdomen small; proboscis black; middle tibiæ at most with a small spur 45.
45. First posterior cell narrowed apically, about one-third the costal margin of the submarginal cell in width; styles of ♀ abdomen three times as long as wide; incisures not differentiated P. COLLATERALIS, nov. sp. (2).
- First posterior cell not narrowed, apically about one-half the costal side of the submarginal cell; styles five times as long

middle tibiæ five-sixths as long as the femora, spur minute and yellow. Halteres, calypteres and root of wings very pale yellow, wings three times as long as wide, costal sections proportioned 5 : 4 : 3 : 1, first posterior cell ending at tip, its veins nearly straight and parallel, sections of the fourth vein proportioned 1 : 2, of the fifth vein, 3 : 4 : 8, posterior crossvein transverse, slightly in advance of the anterior, anal vein weak but complete.

Female. Fourth tarsal joint apically infuscated. Ovipositor shining except on incisures, styles four times as long as wide.

Male type from Nahcotta, Washington, May 24, 1917; female from Vashon, Washington, May 28, 1917; five paratypes from Shelton, Quilcene and Piedmont, Washington, July (Melander).

(1) **Platypalpus satyriacus**, nov. sp. — Length 2 mm. Readily distinct by the swollen genitalia of both sexes; related to *lateralis*, but differing in the broader face, longer arista, pronounced tibial spur and sparse short notal hairs. Black, with a castaneous tinge, head opaque gray pollinose except the shining epistome; face short and about as broad as the front; palpi yellow, proboscis dark castaneous; antennæ yellowish at the base, the third joint darker toward the tip, conical, two to two and one-half times as long as broad, not hairy, and shorter than the rather thick arista; vertical hairs yellow. Center of the mesonotum broadly shining, sides and pleuræ whitish pollinose, sternopleuræ largely shining, bristles yellow. Abdomen of male with long hairs, especially apically, the genitalia greatly enlarged, in bulk about equal to the remainder of the abdomen, the penis visible as a strong black hook, the left valve strongly emarginate on the right-hand edge and apically fringed; hairs of female abdomen normal, incisures ochraceous, last tergite elongate, as long as two or three of the preceding segments, overlapping the ovipositor and shining blackish brown, styles short, brown. Legs yellow, anterior femora moderately robust, about sixteen black setulae in the posterior row of the middle femora, middle tibiæ five-sixths as long as their femora, with an acute yellow spur slightly shorter than the diameter of the tibia, last tarsal joint darkened. Wings very long, veins yellow, third and fourth veins gently converging apically, second and third sections of the costa of the male somewhat thickened, the second, third and fourth sections proportioned 1.2 : 1 : 0.2, sections of the fifth vein 1 : 1.2 : 2.2, marginal cilia of male equal to the anterior crossvein in length, of the female short.

Four specimens, Ungava Bay of Hudson Straits, Northern Labrador, collected by L. M. Turner: a pair deposited in the U. S. National Museum.

(2) **Platypalpus collateralis**, nov. sp. — Female. Length 2 mm. Closely related to *lateralis* Loew, but the latter differs in having a narrowed face and front, yellow proboscis, the last two segments of the female abdomen ochraceous, the femora more slender, no tibial spur, and long mesonotal hairs. Black shining, sides of the thorax white pollinose, legs yellow, tibial spur short, tarsi becoming dusky broad at tip, crossveins meeting and transverse; sides of the front parallel, face quite as wide as the front, proboscis black, mesonotum with sparse short whitish hairs, abdomen entirely shining black, fourth vein gently converging toward the third. Front, face and occiput opaque gray pollinose, the face very short, epistome shining; palpi white; base of the antennæ brown (the last joint lacking in the specimens before me); vertical and ocellar bristles long and whitish. Humeri and sides of the mesonotum whitish pollinose like the pleuræ, the sternopleuræ largely polished, bristles whitish; styles about three times as long as broad, black. Anterior femora somewhat thickened, the middle pair with about fifteen black setulae in the posterior row. Calypteres with a yellow fringe. Wings nearly hyaline, with pale veins, second section of the costa somewhat swollen, the second, third and fourth sections proportioned 5 : 3 : 1, third vein straight, ending at the wing-tip, sections of the fifth vein proportioned 1 : 1.5 : 2.5, marginal cilia shorter than the anterior crossvein.

Alaska, several places: Harriman Expedition.

as wide; incisures of the abdominal segments ochraceous;	
proboscis and palpi obliquely porrect; robust species . . .	P. PRORSUS, nov. sp. (1).
46. Pleurae entirely shining, only the pectus above the front coxae pollinose; base of the antennae fuscous (<i>montanus</i> , Melander).	P. SPLENDENS, new name.
Pollen conspicuous along the notopleural suture and on the propleurae; base of the antennae yellow (<i>gilvipes</i> , Coquillett).	P. XANTHOPODUS, new name.
47. Femora more or less marked with black, the hind femora always dark toward the apex; tarsi not annulate	48.
Legs yellow and usually slender	53.
48. Front femora with a round black apical spot, hind femora dark toward apex, legs otherwise yellow; middle femora with regular long black bristles below; tibiae without spur. Front femora without round black spot; middle femora with long black bristles below; tibiae without spur; middle femora, anterior tibiae and all tarsi black	P. APICALIS, Loew.
Front femora without round black spot, middle femora at most with weak pale bristles beneath; legs otherwise colored	P. GESTICULOR, nov. sp. (2).
49. Front rather narrow and subpollinose; anterior femora blackish above; third antennal joint oval	49.
Front shining and usually broad; spur of the middle tibiae minute or wanting, anterior femora not blackish.	50.
	51.

(1) **Platypalpus prorsus**, nov. sp. — Female. Length 3 mm. Related to *lateralis* Loew, but robust, with broad face, black proboscis, incisures of the abdomen narrowly ochraceous and with short sparse notal hair. Head opaque gray pollinose, except the shining epistome; sides of the front nearly parallel, face very short; proboscis geniculate, bending forward below, although this may be accidental in this specimen, the yellowish palpi also porrect; bristles yellowish; base of the antennae reddish, the third joint conical, brown, hairy, about two and one-half times as long as wide and subequal to the arista. The hairs of the third joint are shorter and the arista is thinner than in *lateralis*. Thorax shining black, the sides, encroaching above the humeri, whitish pollinose, like the pleurae, sternopleurae largely shining, bristles whitish, three pairs of scutellar bristles; terminal segments of the abdomen shining, the styles slender and hairy, about five times as long as wide. Legs yellow, the tarsi very weakly annulate, middle femora robust, about seventeen black setulae in the posterior row, middle tibiae five-sixths as long as their femora, no tibial spur but the flexor end sharply truncated. Veins brownish, the second, third and fourth sections of the costa proportioned 4.5 : 2.3 : 1, third vein ending before the wing-tip, nearly parallel with the fourth, sections of the fifth vein proportioned 1 : 1.2 : 2.2, marginal cilia shorter than the anterior crossvein.

One specimen, in the U. S. National Museum, collected by L. M. Turner, at Ungava Bay of Hudson Straits, Northern Labrador.

(2) **Platypalpus gesticular**, nov. sp. — Male. Length 2.3 mm. Body black, notum mostly shining, legs variegated, spur of middle tibiae lacking, antennae black, third antennal joint very small. Front broad, olivaceous, face gray, epistome shining, proboscis black, palpi yellow, rather prominent, cephalic bristles very small; third antennal joint ovate, about one-fourth the length of the slender arista. Mesonotum with double but contiguous acrostical row and dorsocentral patches of pubescence, notopleural suture broadly pollinose, pleurae mostly bare and polished; bristles reduced, yellowish, no humeral. Abdomen shining, with loose yellowish hairs, pygidium globular forming a rather large rounded ending to the abdomen. Coxae pale yellow, all tarsi black, front femora luteous, front tibiae black except toward knees, middle femora black except toward base, with fourteen strong black setae in the posterior row, middle tibiae black, the apical third with matted golden pubescence, hind femora and tibiae each dark on the distal half. Calypteres, fringe and halteres yellow. Wings hyaline, three times as long as wide, veins thin and brown, costal sections proportioned 6 : 4 : 2.3 : 1, third vein ending at the middle of the wing-tip, parallel with the fourth, crossveins located near one-third the wing-length, the distance between them less than the length of the anterior crossvein, posterior crossvein nearly transverse, outer sections of the fifth vein proportioned 3 : 5, anal crossvein reflexed at an angle of 50 degrees, anal vein faint.

Female. Color of legs much less intensive.

Foot of Lake McDonald, Glacier National Park, Montana; August 14, 1916 (Melander).

50. Spur of the middle tibiae as long as the diameter of the tibiae,
base of the hind femora pale, anterior femora with flexor
cilia P. DIVERSIPES, Coquillett.
- Middle tibiae without spur; hind femora entirely blackish,
femora not ciliate P. DIREPTOR, nov. sp.
51. Middle femora without flexor setæ and with 15 setulæ in
posterior row; front narrow; pleuræ shining; third antennal
joint slightly shorter than arista; bristles yellow P. NITIDIPLURA, nov sp. (1).
- Middle femora with flexor setæ and with about 20 or more
posterior setulæ; front broad 52.
52. Third joint of the antennæ three times as long as broad,
subequal to its arista; bristles brown; pygidium rather
large and vertical; upper pleuræ pollinose P. BICORNIS, nov. sp (2).
- Third antennal joint oval, much shorter than its arista; bristles
black; pygidium minute and globular; pleuræ shining P. PUDENS, nov. sp. (3).

(1) **Platypalpus nitidiplura**, nov. sp. — Female. Length 2 mm. Entirely polished black, legs yellow, the hind femora apically dark; crossveins contiguous and transverse; antennæ black, the third antennal joint long, subequal to the arista. Head globular, polished, vertical bristles black; front narrower than the width of the basal antennal joint, its sides parallel; face narrow, its lower half gray pruinose, epistome shiny; third antennal joint two and one half times as long as broad; proboscis black, palpi white, ovoid, rather pointed, with a pair of marginal white setæ. Notum nearly bare, its bristles yellowish, only a faint trace of pollen along notopleural suture and on metanotum. Abdominal hairs fine and very sparse; terminal segments not dusted. Front tarsi uniformly dusky, posterior tarsi dusky beyond base; about fifteen pronounced setulæ in posterior row of middle femora, but no flexor setæ; middle tibia three-fourths length of femora, the apical spur not developed. Halteres white. Wings hyaline, veins yellow at base, otherwise black and thin, costa rather pronounced beyond end of first vein, third and fourth veins straight, slightly diverging, third vein ending distinctly before wing-tip, sections of third vein 1 : 25, of the fourth vein, 2 : 3 : 6, cilia of hind margin short.

A single specimen taken in a cottonwood thicket two miles south of Gardiner, Montana, in the Yellowstone Park, August 17, 1919 (Melander). It is related to *bicornis* and *pudens*, but is quite distinct.

(2) **Platypalpus bicornis**, nov. sp. — Length 2 mm. Black, shining, the upper half of the pleuræ sericeous, legs brownish, spur of the middle tibiae minute, front broad, antennæ porrect, crossveins meeting. Head wider than deep, occiput and face subshining, front and epistome polished; palpi elongate oval, yellowish, with a single apical seta; basal joint of the antennæ minute, third joint subulate, over three times as long as broad, slightly longer than the pale-tipped arista; one pair each of black vertical and ocellar bristles. Mesonotum nearly bare, only four pairs of weak acrostichal setulæ, notopleural suture, scutellum, metathorax, and upper half of the pleuræ sericeous; one pair each of the usual thoracic bristles; abdomen shining, including the genitalia, ovipositor compressed, pygidium rather large and erect. Base of posterior coxae blackish, posterior femora infuscated apically, the hind ones darkest, front femora not ciliate beneath, but densely and finely pubescent instead, middle femora not bristly in front, posteriorly with four long yellow bristles, tarsi blackish apically. Veins blackish, costa slightly swollen at the end of the first vein, its second, third and fourth sections proportioned 4 : 2.6 : 1, third and fourth veins straight and parallel, crossveins meeting on the basal third of the wing, nearly transverse, sections of the fifth vein proportioned 0.3 : 0.5 : 1, anal crossvein forming an angle of fifth degress, base of the anal vein wanting, marginal cilia short.

Twenty-eight specimens, type from Bellingham, Washington, July 29. Paratypes from Brinnon, Everett, Granite Falls, Index, Lake Cushman, Lilliwaup, Quilcene, Tacoma and Tulalip, all in Washington (Melander).

(3) **Platypalpus pudens**, nov. sp. — Male. Length 1.5 mm. Black, shining, bristles black, legs dark testaceous, posterior coxae and distal two-thirds of the hind femora blackish, tibial spur minute, front very broad, third antennal joint short ovate, crossveins meeting. Head wider than deep, occiput and face subshining, front and epistome polished; palpi small, elliptical, blackish, with one apical dark seta, proboscis short; antennæ two-jointed, the outer joint scarcely longer than broad, less than one-half as long as the arista; two pairs of vertical and one of ocellar bristles, long. Mesonotum nearly bare, but three pairs of acrostichal setulæ present, humeral bristles small, one pair each of notopleural, supraalar, postalar, dorsocentral and scutellar bristles; pleuræ polished, the metapleuræ and scutellum alone sub-pollinose; abdomen shining, pygidium minute. Femora not ciliate, middle femora with four stiff black setæ in back of the second row of setulæ, last tarsal joints blackish. Veins pale, costa

slightly thickened at the end of the first vein, its second, third and fourth sections proportioned $3.5 : 2.2 : 1$, third and fourth veins straight and parallel, posterior crossvein nearly transverse, arising a little before the short anterior crossvein, sections of the fifth vein proportioned $0.3 : 0.5 : 1$, anal crossvein weak, forming an angle of fifty degrees, anal vein wanting, marginal cilia short.

Type from Mount Constitution, Orcas Island, Washington, July 31, 1909; paratype from Bovill, Idaho, June 17, 1911 (Melander).

(1) ***Platypalpus inops*, var. *equicornis*, nov. var.** — The typical form of *inops* originally described from Wyoming, occurs also in Glacier National Park, Montana. At this place also the writer took a female *Platypalpus* on August 14, 1916, which conforms most closely with *P. inops*, but differs in having the third antennal joint subequal to the arista instead of about one-third as long. The anal crossvein is also more nearly perpendicular, forming an angle of about 70 degrees with the fifth vein. The anal crossvein of *inops* is more reflexed, making an angle of about 50 degrees.

(2) ***Platypalpus alumnus***, nov. sp. — Male. Length 2 mm. Body black, notum largely polished, sternopleura with large shining spot, hypopluræ entirely pollinose, notal hairs seriate; pygidium large, globose, shining; legs yellow, tarsi not annulate, tibial spur very small; sides of front diverging, antennæ black, the base brownish, third joint conical, arista black, one-half longer than the third joint, palpi small and dusky; crossveins contiguous and transverse, third vein ending just before wing-tip, anal vein almost complete but weak. Head cinereous pollinose, occipital hairs yellow and inconspicuous, vertical bristles brown, face cinereous, nearly twice as long as wide, epistome shining, proboscis black. Sides of notum cinereous, bristles small and brownish, central scutellars moderately long, a bare stripe on each side of the double acrostichal row of pale hairs. Abdomen shining, quite bare, the two large pygidial valves golden fimbriate at edge. Seventeen short dark setæ in posterior flexor row of middle femora and nineteen minute denticles in anterior row, last two joints of tarsi slightly brownish. Calypteres, fringe and halteres pale yellow. Wings hyaline, veins pale brown, anal angle reduced, the anal vein close to margin, first posterior cell a little the widest at three-fifths its length, sections of fifth vein proportioned 3 : 4 : 9.

One male and two females, taken in the alpine meadow at the Continental Divide near Isa Lake, 8200 feet altitude, Yellowstone Park, Wyoming, 8 Aug. 1918 (Melander).

- oval; third vein ending beyond wing-tip; hypopleuræ largely polished P. GLACIALIS, nov. sp. (1).
58. Front pollinose; mesonotal pubescence conspicuous; palpi brownish yellow P. PUBESCENS, nov. sp. (2).
- Front shining; mesonotal pubescence usually short; palpi pale yellow. 59.
59. Middle femora with 8 yellow flexor bristles, front femora with 17 short pale flexor cilia; palpi very large, nearly as long as the proboscis; halteres yellow P. PECTINATOR, Melander.
- Middle femora without conspicuous bristles; front femora not ciliate; palpi normal; halteres tipped with black. P. SIMPLICIPES, nov. sp. (3).

(1) **Platypalpus glacialis**, nov. sp. — Male. Length 2.3 mm. Body black, legs including the coxæ yellow, notum not pollinose but finely pubescent between broad dorsocentral glabrous rows, spur of middle tibiæ not developed, tarsi not annulate, antennæ black, the third joint short, ovate, about one-fourth the arista in length, palpi whitish and large, pygidium large, globose and subshining, crossveins approximate, the posterior nearly transverse. Occiput with olivaceous pollen, its upper hairs golden, ocellar bristles piceous, front lightly dusted, short, its sides diverging, face very narrow, gray, epistome polished, proboscis short, piceous: first and second antennal joints fused, third as long as deep. Notal pubescence evident but fine and rather sparse, humeri and sides of notum with light pollen, lateral bristles yellow, the humeral short, sternopleuræ polished. Abdomen shining, nearly bare, right dorsal valve of the pygidium deeply bifid, left valve small and triangular. Legs slender, front femora weakly ciliate, middle femora with about twenty black setulæ in the anterior row and about sixteen longer setulæ in the posterior row, no yellow bristles, middle tibiæ about seven-eighths as long as the femora, tarsi vaguely becoming dusky apically. Calypteres, fringe and halteres yellow. Wings hyaline, two and a half times as long as broad, veins thin and blackish, costa black at end of first vein, costal sections proportioned 7 : 5 : 3.5 : 1, third vein ending at wingtip, very slightly converging toward fourth, crossveins separated by less than the length of the anterior and by one-half the length of the posterior, located at two-fifths the wing-length, sections of the fifth vein proportioned 2 : 3 : 5, anal crossvein straight, evanescent apically and reflexed at an angle of fifty degrees, anal vein faintly impressed, anal angle weak.

Female. Last two segments ochraceous, styles slender.

Two specimens, Glacier National Park, Montana, August 14, 1916 (Melander).

(2) **Platypalpus pubescens**, nov. sp. — Male. Length 3 mm. Body black, the venter yellowish, legs including the coxæ yellow, notum not pollinose but uniformly and rather densely pubescent, spur of middle tibiæ undeveloped, tarsi not annulate, antennæ black, the third joint one-fourth as long as the arista, palpi brownish, pygidium large, globose and pollinose, crossveins approximate, the posterior nearly transverse. Occiput finely dusted, its upper hairs dusky, vertical and ocellar bristles black, front finely pollinose, two and a half times as long as wide at the middle, face and epistome shining black, proboscis short and brown, palpi brownish yellow; antennæ three jointed, the third joint conical, scarcely longer than wide. Extreme sides of the mesonotum and the pleuræ white pollinose, disc of notum uniformly, closely and conspicuously white-pubescent, lateral bristles yellow, no humeral, one pair of rather distant scutellars; sternopleuræ polished except the extreme upper part. Abdomen not tapering, shining, its hairs rather dense, dorsal valve large, circular and not fringed. Trochanters tipped with black, front femora not ciliate, middle femora robust, with about thirty minute black setulæ in the anterior flexor row and twenty-one setulæ in the posterior row, the latter flanked by a few yellow setæ, middle tibiæ almost three-fourths the femora, only an indication of the spur present, tarsi gradually becoming dusky toward the tip. Calypteres and knob of halteres whitish, fringe and root of halteres luteous. Wings hyaline, two and a fourth times as long as wide, veins narrow and piceous, sections of the costa proportioned 6 : 3.5 : 2.5 : 1, third vein gently recurved and ending at the wing-tip, first posterior cell widest at the apex, last two sections of the fifth vein subequal, anal crossvein straight, reflexed at an angle of fifty degrees, ends of the anal vein evanescent.

Female. End of abdomen not ochraceous, the styles tumid.

Seven specimens, Union Flat, near Pullman, Washington, July 16, 1916 (Melander).

(3) **Platypalpus simplicipes**, nov. sp. — Male. Length 2 mm. Blackish, largely shining, legs yellow, without evident setæ, crossveins meeting. Head globular, occiput lightly dusted, subshining, front shining, moderately wide, broadening above, face short, black; palpi elongate oval, as long as the face, yellow, with a few marginal white setulæ, proboscis brown; antennæ three-jointed, the last joint short-ovate, scarcely longer than broad, one-third as long as the rather pubescent arista; vertical bristles small. Mesonotum uniformly pubescent, no discal setulæ, no humeral bristle, one notopleural, one supraalar, one prescutellar, four scutellar, the outer ones small; upper half of the pleuræ lightly pruinose, sternopleuræ and hypopleuræ shining; abdomen shining, the pygidium globular and puberulent, the fringe of the upper valve long but fine. Legs slender, front femora

60. Mesonotal hairs sparse; two single rows of acrostichal hairs	61.
Mesonotal hairs dense, the acrostichal area covered with irregular hairs; ovipositor tipped with pollen; veins blackish	P. JUVENIS, var. HYENOIDES, nov. var. (1).
61. Ovipositor tipped with pollen; mesonotum without lateral hairs; wings nearly three times as long as broad, veins yellowish; last sternite of ♂ abdomen with a fringe of yellow bristles.	P. JUVENIS, var. JUVENIS, s. str. (2).
Ovipositor completely shining; mesonotum with sparse lateral hairs; wings broader, veins brown; abdomen not fringed.	P. JUVENIS, var. PUEBINUS, nov. var. (3).
62. Disk of thorax uniformly shining; front tibiæ incrassate	63.
Thorax with two shining vittæ more or less connected with a supra-alar shining spot; tibiæ not swollen (<i>P. vittatus</i> , nov. sp.)	66.

not ciliate, middle femora not setose, biserately black-setulose as usual with seventeen setulæ to each row, middle tibiæ two-thirds as long as their femora, the apical spur minute. Halteres tipped with a small black spot. Veins thin and brown, the second, third and fourth sections of the costa proportioned 3 : 2.5 : 1, third vein recurved, ending slightly beyond the tip of the wing, fourth vein more nearly straight, crossveins at two-fifths the wing-length, meeting, the posterior slightly oblique, twice as long as the anterior, sections of the fifth vein proportioned 0.4 : 0.6 : 1, anal crossvein at an angle of fifty-five degrees, anal vein weak, vanishing at its base; marginal cilia shorter than the anterior crossvein.

One specimen, Tacoma, Washington, August 27, 1911.

(1) *Platypalpus juvinis*, var. *hyenooides*, nov. var. — A female from Vashon, Washington, May 28, 1917, differs from the following varieties in having the mesonotal hairs more abundant, thus resembling the species *P. glacialis*, the hairs between the dorsocentral glabrous stripes irregularly placed. Third antennal joint equal to the arista; vertical bristles blackish; end of ovipositor opaque; wings three times as long as broad, veins blackish, stigmal thickening pronounced.

(2) *Platypalpus juvinis*, var. *juvinis*, s. str. — Length 2 mm. Black, disk of the mesothorax shining and nearly bare of hairs, legs yellow, tibial spur minute, tarsi not annulate, antennæ black, the last joint conical and subequal to the arista, front and face densely cinereous, crossveins meeting. Occiput densely cinereous pollinose, sides of the rather broad front diverging above; palpi broadly oval of moderate size, yellow, silky in front and with a single white apical seta; antennæ two-jointed, the basal joint sometimes brown, the outer joint two times as long as wide and slightly shorter than the arista; two pairs of relatively strong yellow vertical bristles and one pair of ocellars. Humeri, sides and hind margin of the mesonotum and pleuræ except a large sternopleural spot, cinereous pruinose, one pair each of the usual bristles yellow, two rows of acrostichal hairs, dorsocentral stripes bare, sides of notum with a few fine white hairs; abdomen shining black, last sternite ♂ with a dense fringe of yellow setæ, pygidium large, its dorsal valve produced backwards as a rather slender curved, nearly smooth process which bears a yellow fringe on its right side, apex of the ovipositor pollinose. Base of the posterior coxæ a little darkened, cilia of the front femora very short and sparse, middle femora with fourteen black setæ in the postero-flexor row and without additional yellow setæ, middle tibiæ five-sixths as long as their femora. Wings nearly three times as long as wide, veins yellowish, costa not thickened, its second, third and fourth sections proportioned 4 : 2.5 : 1, third and fourth veins parallel, the third vein ending at the tip of the wing, posterior crossvein nearly transverse, twice as long as the anterior, located at three-sevenths of the wing-length, sections of the fifth vein proportioned 0.4 : 0.6 : 1, anal crossvein forming an angle of sixty degrees, anal vein almost wanting, marginal cilia short.

Nine specimens, Kokanee Mountains, 7000 feet, British Columbia, collected by R. P. Currie for the U. S. National Museum; Dubois, Wyoming, September, collected by W. M. Wheeler; and Mount Rainier, Washington (Melander). A specimen from San Diego, California (E. P. Van Duzee) has the pygidium more globular.

(3) *Platypalpus juvinis*, var. *puerinus*, nov. var. — A series of specimens from the Kokanee Mountains in British Columbia and also from Mount Katahdin, Maine, differs as follows: the third antennal joint is usually slightly longer than the arista, the vertical bristles are brown, the end of the abdomen ♂ lacks the conspicuous fringe, the dorsal pygidial valve is more blunt, more or less rugulose and not elaborately fringed, the ovipositor is completely shining, the sides of the mesonotum almost devoid of hairs and the wings are broader, with stronger, brownish veins and with the end of the first vein thickened.

63. Tarsi black; bristles, coxae and veins blackish; tibial spur moderate and stout; mesonotum pollinose on sides and rear P. VERPUS, nov. sp. (1).
- Tarsi strongly annulate; bristles, coxae and veins yellow; tibial spur acute 64.
64. Mesonotum mostly shining, not pollinose in front of the scutellum; last section of the fifth vein distinct (*P. politus*, nov. sp.) 65.
- Center of the mesonotum shining; outer angle of the second basal cell rounded, last section of the fifth vein obsolete; base of the antennæ yellow P. ENERVATUS, nov. sp. (2).
65. Fourth vein very perceptibly sinuous; base of the antennæ yellow P. POLITUS, nov. sp. (3).

(1) **Platypalpus verpus**, nov. sp. — Male. Length 2 mm. Black, disk of mesonotum polished, legs brownish yellow, front tibiae swollen, veins, bristles and antennæ black, crossveins separated. Occipital hairs silky and white, two vertical bristles erect, blackish; face with parallel sides, white, front narrow, gray pollinose; ground color of the palpi piceous, the setæ whitish, proboscis black; third antennal joint pyriform, two-thirds as long as the arista. Humeri, notopleural sutures, rear of mesonotum and scutellum gray pollinose; acrostichal and dorsocentral rows and lateral patches of setulæ present; one humeral, three prealar, one posterior dorsocentral and the central scutellar bristles evident; pollen of the pleuræ silvery gray and dense, about two-thirds of the sternopleuræ bare. Abdomen shining, its hairs sparse and dusky, pygidium large and obliquely ascending and rounded, fringe of the left dorsal valve long and dark. Base of the coxae blackish, front femora stout, with twelve pale flexor setæ, middle femora with about eighteen short black setulæ in each flexor row, middle tibiae five-sixths as long as their femora, the terminal spur rather short, stout and black, hind knees infuscated. Halteres with white knobs; calypteres dusky. Wings three times as long as wide, anal angle rather pronounced, veins strong and blackish, costal sections proportioned nearly 4 : 3 : 2 : 1, third vein nearly straight, fourth vein curving and parallel with the third on the outer half, posterior crossvein rather oblique, sections of the fifth vein, 2 : 3 : 2.5, base and apex of anal vein lacking.

Holotype, Union Flat, near Pullman, Washington, June 3, 1918 (Melander).

(2) **Platypalpus enervatus**, nov. sp. — Male. Length 2 mm. Black, sides and back of the mesonotum densely cinereous pollinose, the center of the disk polished, bristles yellow. Head cinereous, face and front rather narrow, their sides parallel, epistome shining; antennæ two-jointed, the basal joint yellowish, the outer joint black, ovate, nearly twice as long as wide and three-fourths as long as the arista; palpi oval, with white setæ, proboscis black. Thorax nearly bare, but with a group of setæ about the notopleural bristle; pleuræ densely whitish pruinose, the sternopleural spot of moderate size; abdomen polished, pygidium more or less triangular in outline and rather small. Legs including the coxae light yellow, the tarsi strongly annulate, front femora as thick as the middle ones, with twelve stiff pale cilia beneath, front tibiae swollen, middle femora with a posterior row of twelve pale setæ, the middle tibiae nearly as long as their femora and with a strong black-tipped spur. Veins pale, third vein ending at the wing-tip, straight, subparallel with the fourth, second, third and fourth sections of the costa proportioned 3.2 : 2.3 : 1, the costa slightly thickened beyond the first vein, posterior crossvein rather oblique, slightly before the middle of the wing, much beyond the anterior crossvein, rounding into the fifth vein, the last section of which, together with the anal crossvein and the anal vein, obsolete, marginal cilia very short.

One specimen. San Diego, California, May, collected by E. P. Van Duzee.

(3) **Platypalpus politus**, nov. sp. — Male. Length 2.6 mm. Robust, black, dorsum shining, legs light yellow, femora ciliate, tibial spur strong, tarsi annulate, antennæ largely yellow, posterior crossvein at the middle of the wing. Occiput finely cinereous dusted, front rather narrow, its sides parallel, densely whitish pruinose, face white, epistome shining; palpi golden yellow, elongate oval, with whitish setæ and long apical bristle, proboscis short, black; antennæ two-jointed, the outer joint short-ovate, its tip black, the arista twice as long as the outer joint; vertical bristles yellow and short, one pair of ocellar and one of vertical bristles. Humeri, notopleural sutures and pleuræ densely whitish pruinose, the glabrous spot of the sternopleuræ moderate in size, scutellum and metanotum subshining; the usual bristles yellow, setulæ very sparse and fine; abdomen shining, pygidium small. Legs robust, front tibiae inflated, larger than the others, middle tibiae nearly as long as their femora. Veins brown, costa a little thickened between the first and second veins, its second, third and fourth sections proportioned 4.5 : 2.4 : 1, third vein straight, fourth vein with a perceptible backward curvature and then turning forward and parallel with the third at the tip, the first posterior cell widest at its middle and ending at the apex of the wing, posterior crossvein nearly transverse, one-third longer than the anterior, the distance between

Third and fourth veins subparallel; antennæ black	P. POLITUS, var. NITENS, nov. var. (1).
66. Legs reddish yellow, tarsi becoming piceous apically	P. VITTATUS, nov. sp. (2).
Middle femora above and hind femora ringed preapically with black, tarsi black except at base	P. VITTATUS, var. PERIMERUS, nov. var. (3).
67. Spur of the middle tibiæ shorter than the diameter of the tibia. Spur of the middle tibiæ at least as long as the diameter of the tibia	68.
68. Crossveins meeting and transverse; spur of the middle tibiæ minute or wanting	69.
Posterior crossvein more or less oblique and located beyond the anterior; at least base of the antennæ yellow, third joint ovate, less than twice as long as wide; bristles yellow; spur of the middle tibiæ small but evident, middle femora ciliate beneath; anal cell incomplete	76.
69. Abdomen more or less pollinose; femora not thickened, the middle femora with 12 setulæ in the posterior row and no bristles; base of the antennæ yellow, the third joint ovate; pleurae entirely pruinose	P. CANUS, Melander.

the crossveins equal to the anterior crossvein, sections of the fifth vein proportioned 1 : 1.5 : 1, anal crossvein at an angle of seventy-five degrees, evanescent toward its end, anal vein indicated only by a very faint fold beyond the anal crossvein, marginal cilia very short.

One specimen, presented by J. M. Aldrich, who collected it at Sierra Morena Mountains, California, April 3, 1906.

(1) *Platypalpus politus*, var. *nitens*, nov. var. — Several specimens differ from the preceding form in having the antennæ black and the third and fourth veins nearly parallel. They represent the following localities: Redwood City, California (J. M. Aldrich); Kerr County, California (W. M. Wheeler); Atherton, Missouri (C. F. Adams), and Washington, D. C. (Knab and Malloch).

(2) *Platypalpus vittatus*, nov. sp. — Length 2.9 mm. Front cinereous, its sides subparallel, face and epistome white-pollinose; palpi black, the exterior side overlaid with white, oval, one-third as long as the black proboscis; antennæ black, three-jointed, the base of the second joint shining, the third joint elongate-ovate, nearly three times as long as broad, short-pubescent beneath, the arista two-fifths longer than the third joint; one pair each of dusky vertical and ocellar bristles. Thorax with a shining black stripe on each side of the middle between the acrostichal and dorsocentral rows, expanding laterally on the posterior part of the notum towards the base of the wings, elsewhere with cinereous pollen on a black ground, the pleurae whiter, sternopleural spot large; abdomen polished black, sparsely hairy, pygidial valves inconspicuous, subdorsal pits prominent. Legs testaceous, base of the posterior coxae black, front femora biseriately ciliate beneath, front tibiae as thick as the middle ones, middle femora yellow-setose in back of the rather coarse black setulae, middle tibiae two-thirds as long as their femora, their spur strong and black, tarsi blackish beyond the base. Veins reddish brown, costa not thickened, its second, third and fourth sections proportioned 5 : 3 : 1, third and fourth veins slightly curving, the first posterior cell widest near the middle, posterior crossvein at the middle of the wing, rather oblique, twice as long as the anterior which is equal in length to the space between the crossveins, sections of the fifth vein proportioned 0.8 : 1.1 : 1, base and apex of the anal vein wanting, the anal crossvein forming an angle of seventy degrees, evanescent, marginal cilia as long as the anterior crossvein.

Five specimens, Converse County, Wyoming, received from Professor Wheeler, and Omak, Washington (Melander). In two of the specimens the shining vittæ of the mesonotum are separated by a pollinose stripe from the supra-alar spot.

(3) *Platypalpus vittatus*, var. *perimerus*, nov. var. — Seven specimens from Chin and Bow Slope, Alberta, collected by Walter Carter from alfalfa, May 30 — June 5, 1923, differ from the preceding in having the legs marked with blackish. This darkening is variable, in its extreme extent forming a dorsal streak on the anterior femora, a broad preapical ring on the posterior femora, the hind tibiæ piceous and all the tarsi black. The lighter colored individuals lack the dorsal streak of the front femora, the preapical ring of the middle femora, and their hind tibiæ are blackened only at the extremity. Types in Canadian National Museum and author's collection.

Abdomen shining, or if pollinose, the species disagrees with the other characters	70.
70. Middle femora with flexor bristles in addition to two rows of setulæ	71.
Middle femora without bristles; antennæ black; mouth- opening not unusually wide	73.
71. Antennæ, palpi and femoral bristles black; mouth-opening normal; coxæ darkened; epistome polished	P. LUCTATOR, nov. sp. (1).
Base of antennæ yellow; palpi and femoral bristles yellow; mouth-opening wide; coxæ yellow (<i>P. hians</i> , Melander)	72.
72. Face broader than long, epistome pollinose; thoracic bristles yellow; halteres yellow	P. HIANS, Melander, s. str.
Face as broad as long, epistome shining; thoracic bristles black; halteres blackish	var. FUSCOHALTERATUS, Melander.
73. Third antennal joint pubescent, subulate, over three times as long as wide, longer than its arista; coxæ more or less blackish at the base, middle femora stout and usually with a few bristles on the anterior side; costa with a strong thickening at the end of the first vein	74.
Third antennal joint ovate, not more than two times as long as wide and shorter than its arista; coxæ and legs yellow; costa not strongly thickened	P. VENATICUS, nov. sp. (2).

(1) **Platypalpus luctator**, nov. sp. — Male. Length 2 mm. Related to *P. hians*, but having black antennæ, palpi and bristles. Occipital setæ dark, arranged in two sets, two pairs of vertical bristles, ocelli black, front two and a half times as long as broad, face twice as long as broad, its ground-color black, epistome polished; mouth-opening normal, proboscis black, palpi of moderate size, decumbent, black and with one apical black hair; outer antennal joint ovate, one-half longer than broad and three-fourths the length of the shortpubescent arista. Mesonotum with olivaceous pollen, its setulæ short and not restricted to the central rows, four scutellar bristles; sternopleuræ largely polished; abdomen wholly polished, the scattered pubescence pale, genitalia coarse and rounding the end of the abdomen, the right and left claspers enclosing an open space between them. Legs uniformly luteous, the coxæ darkened basally, front femora rather stout, middle femora with twenty-four small black setulæ in each row, posterior to which are eight black pronounced bristles, middle tibiae three-fourths their femora, the terminal spur minute. Calypteres, fringe and halteres blackish. Veins light piceous, the second, third and fourth sections of the costa proportioned about 4 : 3 : 1, third vein ending at the wing-tip, nearly parallel with the fourth except at the tip where it converges toward the fourth, crossveins at two-fifths the wing-length, meeting, sections of the fifth vein equal, anal vein uniformly weak and located close to the hind margin, cilia short.

Holotype, Priest Lake, Idaho; August 1, 1916 (Melander).

(2) **Platypalpus venaticus**, nov. sp. — Female. Length 1.8 mm. Cinereous dusted, legs yellow and not setose, crossveins meeting. Head slightly higher than wide, the lower part of the eyes bulging forward, occiput cinereous, front narrow, whitish, face and epistome white; palpi white, oval, one-fourth as long as the black proboscis, with two basal and one apical white setæ; antennæ two-jointed, the basal joint brown, the outer joint black, ovate, not twice as long as wide and three-fourths as long as its arista; two pairs of brown vertical bristles. Thoracic bristles yellow, two humeral, one notopleural, one supraalar, one postalar, two dorsocentral and two scutellar bristles, setulæ very sparse; sternopleural spot large; abdomen shining black. Middle femora short, the tibiae three-fourths as long as their femora, their apical spur minute. Veins thin and pale, costa not thickened, its second, third and fourth sections proportioned 3 : 2.3 : 1, third and fourth veins nearly straight and parallel, the first posterior cell a little the widest at its apex, crossveins before the basal two-fifths of the wing, meeting, the posterior but slightly oblique, more than twice as long as the anterior, sections of the fifth vein proportioned 0.4 : 0.4 : 1, anal crossvein forming an angle of forty-five degrees, anal vein very weak, vanishing at its base, marginal cilia as long as the anterior crossvein.

Type from Pullman, Washington, June 16, 1912. Paratypes from Ilwaco, Washington; Avon and Moscow Mountain, Idaho (Melander) and from Kaslo and South Fork, British Columbia (R. P. Currie).

- | | |
|--|--|
| 74. Thoracic bristles black; tarsi piceous beyond the base, middle femora with anterior black bristles and setose beneath on the postero-flexor edge (<i>P. porrectus</i> , Melander) | 75. |
| Thoracic bristles yellow; last three tarsal joints strongly annulate, middle femora without black bristles and biserrately setulose beneath | P. VELOX, nov. sp. (1). |
| 75. Abdomen completely shining. | P. PORRECTUS, Melander, s. str. |
| Base of the abdominal segments pollinose. | P. PORRECTUS,
var. SUFFASCIATUS, nov. var. (2). |
| 76. Abdomen pollinose and red; front tibiae swollen and ciliate; front tarsi annulate; thorax white-pollinose | P. VIERECKI, Melander. |
| Abdomen black and rarely with pollen; tibiae not swollen | 77. |
| 77. Proboscis yellow; tarsi yellow; white-pollinose species | P. TENELLUS, Melander. |
| Proboscis black | 78. |
| 78. Mesonotum densely golden pollinose, contrasting with the silvery pleuræ; palpi orbicular, not hairy; costa not swollen; tarsi strongly annulate; abdomen with lateral pollinose triangular spots | P. BALLUCATUS, nov. sp. (3). |

(1) ***Platypalpus velox*, nov. sp.** — Male. Length 2 mm. Black, cinereous, bristles yellow, legs yellow, the last three tarsal joints strongly annulate, tibial spur small, femora not setose or ciliate, third antennal joint elongate, arista short, crossveins meeting. Occiput, front and face cinereous, front rather narrow, its sides diverging above, epistome black; antennæ two-jointed, black, the outer joint subulate, over three times as long as wide and nearly twice as long as the arista. One pair each of the usual thoracic bristles, setulæ very sparse, the pollinose coating whitish, sternopleuræ glabrous except the front and upper edge; pygidium globose and small. Posterior coxae slightly brownish at the base, front femora nearly as thick as the middle ones, beneath with fifteen short pale cilia, middle femora not bristly, beneath biserately setulose, the setulæ of the posterior row, twenty in number, scarcely longer than those of the anterior row, middle tibiae nearly as long as their femora. Veins brown, costa thickened at the end of the first vein, its second, third and fourth sections proportioned 3 : 2 : 1, crossveins nearly transverse, located at three-sevenths the length of the wing, the anterior three-fifths as long as the posterior, third and fourth veins subparallel, the first posterior cell a little the widest at two-thirds its length, ending at the apex of the wing, sections of the fifth vein proportioned 0.4 : 0.6 : 1, anal crossvein thin, at an angle of sixty-five degrees, anal vein faint, marginal cilia short.

One specimen, Kettle Falls, Washington, May 3, 1912.

(2) ***Platypalpus porrectus*, var. *suffusolatus*, nov. var.** — Differing from *porrectus* Melander in that the sides of the first segment of the abdomen and the sides and bases of the subsequent segments are cinereous pollinose, leaving the lateral pits conspicuously shining.

Boulder, Colorado (Cockerell); Alma, California (Aldrich).

(3) ***Platypalpus ballucatus*, nov. sp.** — Female. Length 2.5 mm. Mesonotum densely covered with golden pollen, sides of the abdomen pollinose, antennæ reddish yellow. Front densely golden pollinose, its sides parallel, face and epistome silvery, palpi yellow, broadly orbicular, pubescent only at the edge, proboscis black; third joint of the antennæ ovate, scarcely longer than wide, its arista black; one pair of vertical and one of ocellar bristles yellow, lower occipital hairs dense and whitish. One humeral, one notopleural, one supraalar, two scutellar and two dorsocentral bristles yellow, setulæ small and sparse; pollen of the pleuræ dense and white, a small rounded sternopleural glabrous space; abdominal segments with lateral triangular pollinose marks which are broad in front and taper obliquely behind, the middle segments with two prominent lateral pits, venter pollinose. Legs including the coxæ yellow, the apices of the tarsal joints narrowly blackish, front femora with about thirteen pale cilia in each of two flexor rows, black setulæ of middle legs minute, posterior-flexor edge of the middle femora with ten yellow setæ, the anterior edge loosely setose, middle tibiae three-fourths as long as their femora, their spur shorter than their diameter, acute, and tipped with black. Veins pale, crossveins at the middle of the wing, the posterior crossvein somewhat oblique, located beyond the anterior crossvein the length of the latter, costa not tickened, its second, third and fourth sections proportioned $3.8 : 2.4 : 1$, third vein straight, ending just before the tip of the wing, subparallel with the slightly undulating fourth vein, underside of the second basal cell twice as long as the last section of the fifth vein, anal crossvein at two-fifths the length of the basal cell, attenuated apically, forming an angle of eighty degrees, base of the anal vein wanting, marginal cilia short.

Two specimens, Spokane, Washington and Chicago, Illinois (Melander).

Mesonotum white- or cinereous-pollinose, not contrasting with the pleuræ; abdomen not pollinose	79.
79. Crossveins located before the middle of the wing; costa not swollen (<i>P. crassifemoris</i> , Fitch)	80.
Crossveins at or beyond middle of wing, basal cells elongate; costa swollen between first and third veins.	P. CELLARIUS, nov. sp. (1).
80. Base of antennæ reddish-yellow	81.
Antennæ entirely black.	P. CRASSIFEMORIS, var. MELANOCERUS, nov. var. (2).
81. Epistome white pruinose; palpi elongate oval and with two apical setæ	var. CRASSIFEMORIS, s. str.
Epistome shining black.	82.
82. Palpi broadly oval, without apical setæ; tarsi yellow, the last joint black	var. MOLLIS, nov. var. (3).
Palpi elongate oval; tarsi annulate	var. DEBILIS, LOEW.
83. Posterior crossvein transverse, nearly or quite meeting the anterior, if oblique located before the anterior	84.
Posterior crossvein oblique, located beyond the anterior and usually before the middle of the wing	94.
84. Crossveins at or beyond the middle of the wing.	85.
Crossveins distinctly before the middle of the wing; third and fourth veins subparallel	89.
85. Third vein straight, ending before the apex of the wing, fourth vein lightly sinuous, at the end diverging from the third	

(1) **Platypalpus cellararius**, nov. sp. — Male. Length 2.1 mm. Occiput and notum olivaceous gray pollinose, abdomen shining black, legs entirely yellowish, hairs and bristles yellow. Antennæ yellow, the third joint ovate, one-half longer than broad, the black arista twice as long as the third joint; face and front narrow, white pollinose as is also the epistome; palpi whitish, proboscis black. Thoracic setulæ long, one humeral, three notopleural, four scutellar bristles; pleuræ whitish pollinose, with a sternopleural glabrous space; pygidium nearly glabrous, its fringes short. Front femora with twelve yellow flexor setæ, front tibiae as thick as the others, not setose, setulæ of the middle legs minute, a few of the hairs of the front side of the middle femora rather erect, the postero-flexor edge with seven yellow setæ, spur of the middle tibiae small and tipped with black, not as long as the diameter of the tibiae, tarsi apically more or less dusky. Veins pale, hind crossvein somewhat oblique, located a little beyond the anterior and also beyond the middle of the wing, the basal cells therefore long, second and third sections of the costa thickened, subequal and two and one-half times as long as the third, third and fourth veins parallel, straight, the third ending before the wing-tip, the underside of the second basal cell two and six-tenths times as long as the last section of the fifth vein, base of the anal vein wanting, cilia of the hind margin three-fifths as long as the posterior crossvein.

Female. Differs in that the third antennal joint is blackish, the thoracic setulæ are smaller, the dorsum is cinereous pollinose, the tarsi are dusky from the tip of the metatarsus, the crossveins are located at the middle of the wing and the marginal cilia measure one-half the length of the posterior crossvein.

One male and one female, Keremeos, British Columbia (Melander), one female from Lusk, Wyoming, the last collected by W. M. Wheeler.

(2) **Platypalpus crassifemoris**, var. **melanocerus**, nov. var. — Differing from the other forms of *crassifemoris* in having the antennæ entirely black. Palpi clothed with pure white pubescence; epistome white pubescent like the face; tarsi annulate.

Pullman, Wash. (Melander).

(3) **Platypalpus crassifemoris**, var. **mollis**, nov. var. — Antennæ red at base; epistome shining black; palpi broadly oval, without apical setæ; tarsi yellow, the last joint black.

Chester County, Pennsylvania (J. C. Bradley); Milwaukee, Wisconsin (C. T. Brues), Avon, Idaho (Melander) and Quilcene, Washington (Melander).

- vein, first posterior cell widest at its apex, costa thickened from the first to the third vein. P. *BALLISTRARIUS*, nov. sp. (1).
- Third vein curved, converging with the fourth at the end, the first posterior cell widest at the middle, costa not greatly thickened. 86.
86. Third and fourth veins greatly bowed, the first posterior cell at its middle occupying much more than one-third of the wing P. *LYRISTES*, nov. sp. (2).
- Third and fourth veins gently but distinctly curved, the first posterior cell at its middle occupying not more than one-third of wing, third vein ending at or beyond tip of wing 87.
87. Anterior femora with broad blackish annulus, hind femora dark apically; epistome shining; tarsi with dark tip; third antennal joint subequal to arista P. *VALGUS*, nov. sp. (3).

(1) *Platypalpus ballistrarius*, nov. sp. — Male. Length 2.5 mm. Mesonotum moderately dusted with olive-gray pollen, its bristles yellow; wings with long basal cells and costa thickened from the first to the third vein. Front and face white pollinose, the sides of the front parallel, epistome shining black; palpi small, brown but white pollinose and with sparse white setæ, proboscis black; ocellar bristles small, one pair of yellow vertical bristles, one small humeral, three notopleural, one small postalar, four scutellar, three moderate posterior dorsocentrals, four irregular rows of acrostichals; pleurae white pruinose and with a sternopleural glabrous spot; abdomen shining black. Legs including the coxae testaceous, the tarsi darkened apically especially on the distal part of each joint, front femora with a dozen short yellow flexor setæ, middle femora with a few irregular setæ in front and a row in back, their tibiae more than two-thirds the length of the femora, the spur strong and black. Second, third and fourth sections of the costa proportioned 2 4 : 2 5 : 1, crossveins located beyond the middle of the wing, the posterior oblique but placed before the anterior, third vein straight, the fourth vein slightly undulating, the first posterior cell a little the widest at its end which occupies the apex of the wing, sections of the fifth vein proportioned 0.8 : 1 : 1, base of the anal vein weak, anal crossvein forming an angle of 60°, marginal cilia as long as the anterior crossvein.

One specimen, Falls Church, Virginia, May 10, received from Nathan Banks.

(2) *Platypalpus lyristes*, nov. sp. — Male. Length 2.5 mm. Mesonotum with rather dense cinereous pollen, the bristles yellow. Face and front white-pruinose, the latter with parallel sides, epistome black; palpi oval, one-half as long as the proboscis, white, and with white hairs, proboscis black; antennæ two-jointed, black, the outer joint elongate lanceolate, nearly four times as long as wide and a little longer than the arista; ocellar bristles small, two pairs of vertical bristles. One humeral bristle, notopleurals small, one supraalar, one postalar, two dorsocentrals, lateral scutellars small, acrostichals biseriate; sternopleural spot moderately large, pentagonal; abdomen including the venter shining black, base of the segments with indications of pollen, hairs short and sparse above, subdorsal pits prominent, upper valves of the pygidium broad, densely fringed on the left side with long golden cilia. Legs including the coxae yellow, tarsi scarcely darker apically, anterior femora loosely ciliate below, middle femora with but about sixteen setæ in the posterior flexor row, setulae of middle legs minute, middle tibiae five-sixths as long as the femur, the spur longer than the diameter of the tibia and tipped with black. Veins pale at the base, darker distally, the costa brown beyond the first vein and not swollen, crossveins meeting, both long, subequal and nearly transverse, third and fourth veins mutually bowed, the basal and first posterior cells unusually wide, occupying more than one-third of the wing, sections of the fifth vein proportioned 1 : 2 : 1, marginal cilia short.

One specimen, Bright Angel, in the Colorado Canon, Arizona, elevation 3700 feet, collected by H. S. Barbor for the U. S. National Museum.

(3) *Platypalpus valgus*, nov. sp. — Male. Length 2.6 mm. Black, legs mostly yellow, the anterior femora broadly annulate, the hind femora dark on the outer half, tibial spur strong, mesonotum with golden pubescence, crossveins at the middle of the wing, meeting and transverse. Occiput olivaceous pollinose, lower hairs silky white, two black verticals and one ocellar; front similarly pollinose, face white, epistome bare; palpi small, brownish in ground color, proboscis black; antennæ black, the third joint hairy below, slender, two and a half times as long as wide and two-thirds the length of the arista. Lateral thoracic bristles brown, two dorsocentrals, acrostichal and dorsocentral setulae in single rows, pleurae gray pollinose, half of the sternopleuræ shining. Abdomen shining, base of the second segment laterally lightly pollinose, hairs rather long and pale, pygidium obliquely ascending, left dorsal valve triangular, pointed, its fringe dense and yellow. Anterior legs

Femora luteous; arista longer than third antennal joint	88.
88. Epistome pollinose; tarsi annulate; middle section of fifth vein longer than the ultimate; ♀ abdomen with pollinose fasciae.	P. TENAX, nov. sp. (1).
Epistome shining; tarsi gradually darker toward tip; last two sections of fifth vein subequal; abdomen shining	P. RECURVUS, nov. sp. (2).
89. Bristles black; coxae dark, middle femora robust	90.
Bristles yellow; arista slender; legs yellow	91.
90. Third antennal joint much longer than arista; middle femora with five black thorns on anterior face	P. VULNIFICUS, nov. sp. (3).

robust, front femora ciliate as usual, front tibiæ stouter than the hind ones, middle femora with about eighteen setulæ in each flexor row, the hairs rather outstanding but none distinctly setiform, middle tibiæ five-sixths the length of their femora, tarsi apically dusky. Halteres, calypteres and root of wings yellow, veins thin and brown, wings three times as long as wide, basal cells large, third vein uniformly curved ending beyond the tip of the wing, fourth vein sinuous, first posterior cell widest at its middle where it occupies somewhat more than one-third the width of the wing, sections of the costa proportioned 10 : 6 : 4 : 1, of the fourth vein, 1 : 1, of the fifth vein, 2 : 4 : 3, anal crossvein almost perpendicular, base and tip of the anal vein lacking.

Four specimens, from shrubbery, near Adna, Washington, July 10, 1917 (Melander).

(1) **Platypalpus tenax**, nov. sp. — Male. Length 2.3 mm. Black, legs yellow, mesonotum completely covered with golden pollen, bristles black, crossveins transverse, contiguous, located at the middle of the wing, first posterior cell widest at the middle. Upper occiput with olivaceous pollen and a double row of dark setæ, lower occiput with gray pollen and hairs, front olivaceous pollinose, face and epistome white; proboscis black, palpi rather small, oval and yellow; antennæ black, the third joint hairy, lanceolate, over twice as long as wide and three-fourths the length of the arista. Humeral bristle small, posterior dorsocentral and middle scutellar bristles strong, dorsocentral setulæ distinct. Abdomen shining, the hairs fine, long and pale, pygidium not much larger than the diameter of the abdomen, its left dorsal valve with a close fringe of yellowish hairs. Legs robust, including the coxae yellow, the tarsi narrowly annulate, front femora with a dozen fine pale flexor cilia, about twenty-four setulæ in each flexor row of the middle femora, posterior to which are about a dozen irregular yellow setæ, middle tibiæ setulose within, five-sixths as long as their femora, the black-tipped acute spur long. Roots of halteres and of wings, and the calypteres yellow, wings three times as long as wide, veins thin and blackish, sections of the costa proportioned about 7 : 5 : 3 : 1, of the fourth vein, 7 : 8, of the fifth vein, 3 : 4 : 3, third vein gently bowed, ending just beyond the wing-tip, fourth vein conversely bowed, the first posterior cell at its middle occupying about one-third the width of the wing, anal crossvein straight, almost perpendicular, anal vein faint, anal angle moderate.

Female. Abdominal segments with faint basal bands of yellow pollen.

One pair, Ilwaco, Washington, July 15, 1917 (Melander).

(2) **Platypalpus recurvus**, nov. sp. — Female. Length 3 mm. Front cinereous, its sides slightly diverging above, face white-pollinose, epistome black, palpi ovate, half as long as the proboscis, yellow, the exterior side white pruinose, a single apical white seta, proboscis black; antennæ three-jointed, black (in one specimen the antennæ are yellowish at the base), the last joint elongate-oval, twice as long as wide, its arista nearly twice as long as this joint; one pair each of black ocellar and vertical bristles. Pollen of the mesonotum rather dense, yellowish, nearly coating the black ground-color, bristles yellow, the humeral small, one notopleural, one supraalar, one postalar, two dorsocentrals, four scutellars; pleuræ more densely white pruinose, with a large sternopleural spot; abdomen shining black, with visible subdorsal pits. Legs including the coxae testaceous, front femora with twelve flexor cilia, middle femora setose in front and back, middle tibiæ three-fourths as long as their femora, their spur strong, acute, black, tarsi dusky, more pronounced at the apex of the joints. Wings a little dusky, veins strong, costa swollen at the end of the first vein, its second, third and fourth sections proportioned 5 : 3 : 1, third vein uniformly recurved, terminating beyond the tip of the wing, first posterior cell widest at its middle, posterior crossvein oblique, more than three times as long as the anterior, located just before the middle of the wing and half the length of the anterior crossvein beyond that vein, sections of the fifth vein proportioned 0.8 : 1.2 : 1, anal crossvein nearly perpendicular, base of the anal vein indicated only as a slight fold.

Three specimens, Palo Alto, California, April 3, 1906, collected by Dr. J. M. Aldrich. Another female, from Eagle Creek Forest Reserve, near Portland, Oregon (Melander), differs in having the posterior crossvein transverse and meeting the anterior.

(3) **Platypalpus vulnificus**, nov. sp. — Female. Length 3.2 mm. Body black, legs yellow, antennæ elongate, mesonotum pollinose, bristles black, tibial spur strong, middle femora with spinous bristles on the anterior face, crossveins meeting. Front pollinose, rather broad, its sides divergent above, face twice as long as broad, with dark pollen, epistome shining, proboscis black, palpi rather short and stout; antennæ black, the outer joint tapering, four times as long as wide and one-third longer than the arista. Mesonotum with olivaceous pollen, with black setulæ

- Third antennal joint much shorter than the rather stout arista;
middle femora dark, and unarmed on anterior surface . . . *P. PILATUS*, nov. sp. (1).
91. Epistome silvery white pruinose; third antennal joint three-fourths as long as the arista; tarsi strongly annulate; sections of the fifth vein proportioned $0.5 : 0.8 : 1$ *P. EQUALIS*, Loew.
Epistome shining black; antennæ shorter and arista longer;
tarsi not strongly annulate 92.
92. Palpi yellow; base of the antennæ yellow, the third joint with
a straight upper edge 93.
- Palpi blackish inside; antennæ black, the third joint ovate;
last section of the fifth vein subequal to or longer than the
preceding *P. SUTOR*, Melander.
93. Basal cells relatively large, sections of the fifth vein proportioned $0.8 : 1.5 : 1$, first posterior cell ending beyond
the tip of the wing; pollen of the mesonotum golden . . . *P. CREPIDARIUS*, nov. sp. (2).

in four equidistant rows, the usual bristles all present, strong and black, humeral strong; sternopleuræ polished. Sides of the first abdominal segment lightly pollinose, remainder of the abdomen shining except the dull apex, hairs very short, pale and sparse. Posterior coxae black, legs very robust, front femora with spinous black bristles near the knee, the pair on the front face pronounced, middle femora armed with six black spinous thorns in a flexor row before the setulae and with about five more slender black outstanding bristles forming a longitudinal row down the middle of the anterior face, twenty-one setulae in the anterior flexor row and seventeen stout setæ in the posterior row, tibial spur as long as the diameter of the tibiae, tipped with black, all the tarsi weakly annulate. Calypteres and root of halteres brownish, fringe and knob of halteres yellow. Wings hyaline, two and a half times as long as broad, the anal angle full, veins thin and blackish, sections of the costa proportioned $5 : 3 : 2 : 1$, first posterior cell ending at the wing-tip, its veins straight and parallel, crossveins at three-sevenths the wing-length, the posterior before the anterior and nearly transverse, outer sections of fifth vein proportioned $2 : 3$, anal crossvein straight, long, forming an angle of forty-five degrees, anal vein uniformly weak.

One specimen of this distinct species from the southern end of Lake McDonald, Glacier National Park, Montana, August 14, 1916 (Melander).

(1) *Platypalpus pilatus*, nov. sp. — Female. Length 2.2 mm. Black, pollinose, bristles black, legs yellowish, posterior coxae blackish, tibial spur strong, crossveins meeting before the middle of the wing. Head globular, cinereous, front and face short and of moderate breadth, epistome shining black; palpi blackish, spatulate, the anterior surface yellowish pollinose and with a few setæ; antennæ short, three-jointed, the third joint pyriform, one-half longer than wide and two-thirds as long as the rather thick arista; two pairs each of ocellar and vertical bristles. Pollen of the dorsum moderately dense, of a yellowish gray color, two humeral bristles and one pair each of notopleural, supraalar, postalar, dorsocentral and scutellar bristles; pleurae cinereous pollinose, the sternopleuræ largely glabrous; abdomen shining, the pits prominent. Front femora with yellow flexor cilia, middle femora stout, brown, with a few yellow setæ in front and a series of longer ones behind, middle tibiae two thirds as long as their femora, tarsi not annulate. Veins firm, first vein slightly thickened on entering the costa, the second, third and fourth sections of the costa proportioned $4.5 : 2.5 : 1$, third and fourth veins nearly straight and parallel, crossveins meeting at two-fifths the length of the wing, nearly transverse, the posterior three times as long as the anterior, sections of the fifth vein proportioned $0.4 : 0.6 : 1$, marginal cilia short.

One specimen, collected at Friday Harbor, Washington, July 4, 1905, by Dr. J. M. Aldrich.

(2) *Platypalpus crepidarius*, nov. var. — Female. Length 2 mm. Black, dorsum moderately covered with yellow-gray pollen, legs and bristles black. Occiput densely cinereous pollinose, one pair each of ocellar and vertical bristles; front cinereous, its sides parallel, face white pruinose, epistome shining black; palpi oval, yellow, with white apical setæ, proboscis black; antennæ three-jointed, the basal joints yellow, the third joint conical, about one-third as long as the thickened arista. Humeral bristle small, one pair each of notopleural, supraalar, small postalar, dorsocentral and scutellar bristles; pollen of the pleurae denser and white, the glabrous spot occupying one-half of the sternopleuræ; abdomen shining black. Legs including the coxae yellow, tarsi a little dusky towards the apex, front femora ciliate beneath, middle femora ciliate in front and setose behind, tibial spur strong. Veins yellowish, costa not swollen, its second, third and fourth sections proportioned $5 : 3 : 1$, third vein ending at the tip of the wing, crossveins almost meeting, the posterior but little oblique, sections of the fifth vein proportioned $0.8 : 1.5 : 1$, anal crossvein forming an angle of fifty-five degrees, anal vein uniformly faint, marginal cilia shorter than the anterior crossvein.

One specimen, Woodland, Washington, May 23, 1910 (Melander).

Basal cells small, sections of the fifth vein proportioned 0.4 :	
0.4 : 1, first posterior cell ending at the tip of the wing;	
pollen of the mesonotum cinereous	P. SOCCATUS, nov. sp. (1).
94. Pleuræ entirely densely white-pollinose; tarsi yellow	95.
Sternopleura with a glabrous shining spot; front rather narrow, its sides nearly parallel	99.
95. Third antennal joint ovate, two-thirds as long as the arista; bristles of the thorax yellow	96.
Antennæ black, the third joint oval, one-third as long as the arista; bristles black	P. CALIGATUS, Melander.
96. Base of abdominal segments fasciate with cinereous pollen; third vein nearly straight	P. PULVERULENTUS, nov. sp. (2).
Abdomen completely shining, or sides of segments with basal pollinose triangles.	97.
97. Abdomen shining yellow; epistome black; third and fourth veins straight and parallel; slender species less than 2 mm. in length.	98.
Abdomen shining black, with lateral pollinose triangles; epistome white pruinose; base of antennæ yellow; first posterior cell narrowed; robust species measuring 3 to 4 mm.	P. HOLOSERICUS, Melander.
98. Antennæ black; femoral setulæ black	P. RUFIVENTRIS, Melander.

(1) *Platypalpus, soccatus*, nov. var. — Male. Length 1.4 mm. Black, cinereous pollinose, legs entirely yellow, tibial spur equal in length to the diameter of the tibiæ, base of the antennæ yellow, crossveins meeting, basal cells small. Head cinereous, its bristles yellow, front rather broad, its sides diverging, together with the face white pruinose; epistome and proboscis black, palpi broadly spatulate, pale yellow, fringed but not setose. Thoracic bristles yellow, one pair each of the usual bristles, but the scutellum with a small additional lateral pair; sternopleurae glabrous except the upper fourth; abdomen shining black, pygidium large, rotund, shining, not fimbriate. Front femora with short flexor cilia, middle femora with twenty setulæ in the anterior flexor row and twelve black setæ in the posterior row, behind which are some short yellow setæ, middle tibiæ nearly as long as their femora. Veins yellowish, costa but slightly thickened beyond the first vein, its second, third and fourth sections proportioned 3.5 : 3 : 1, third and fourth veins straight, parallel, crossveins meeting at the basal third of the wing, nearly transverse, sections of the fifth vein proportioned 0.4 : 0.4 : 1, anal crossvein at an angle of eighty degrees, anal vein entire, distinct, but very thin, marginal cilia as long as the anterior crossvein.

Female. Last two segments of the abdomen dull gray.

Type, Moscow Mountain, Idaho, July 6, 1912 (Melander). Paratype from same locality, July 4, 1915; and two others from Craig's Mountain, Idaho, June 9, 1918 (Melander).

(2) *Platypalpus pulvorientus*, nov. var. — Length 2.3 mm. Black, thorax cinereous pollinose including the entire pleuræ, abdomen with pollinose fasciæ, legs entirely yellow, tibial spur strong, costa not thickened, crossveins separated, the posterior oblique. Front narrow, cinereous, face white pollinose over a yellowish ground, epistome white over a black ground color, palpi broadly oval, yellow, bare except for two apical white setæ, proboscis short, black; antennæ short, two-jointed, black the base rufous, last joint elongate ovate, twice as long as wide and nearly as long as the arista; one ocellar and two pairs of vertical bristles black. Bristles of the mesonotum blackish, including one humeral, several notopleural, one supraalar, one postalar, two dorsocentral, and four scutellar, acrostichals biseriate, but minute, other setulæ very sparse; pollen of the pleuræ whiter; the pollen of the abdomen including the sides and the bases of the second and following segments. Legs including the coxæ yellow, the posterior coxæ pollinose laterally, lower posterior side of the middle femora with a row of yellow setæ, middle tibiæ two-thirds as long as their femora. Second, third and fourth sections of the costa proportioned 3 : 1.5 : 1, third vein straight, fourth vein undulate, parallel with the third at its very tip, first posterior cell widest at its middle, distance between the crossveins equal to the anterior, which is one-half as long as the posterior, sections of the fifth vein proportioned 0.7 : 0.9 : 1, anal crossvein gradually attenuated, forming an angle of fifty degrees, anal vein uniformly weak, marginal cilia short.

Two specimens, San Diego County, California, March, 8, 1897, collected by W. M. Wheeler.

- Antennæ yellow at base; femoral setulæ yellow; front unusually long. P. OCULEUS, nov. sp. (1).
99. Thoracic bristles black; tarsi not annulate 100.
- Thoracic bristles yellow; tarsi usually more or less annulate. 102.
100. Third antennal joint lanceolate, twice as long as wide and shorter than the arista; middle tibiae two-thirds as long as the femora, the middle femora without yellow setæ in front. P. TALARIS, nov. sp. (2).
- Third antennal joint slender, over three times as long as wide, longer than its arista; tarsi piceous beyond the base, middle tibiae three-fourths as long as their femora 101.
101. Abdominal segments with pollinose fasciæ; base of the anal vein indicated, but weak P. INCURVUS, Melander.
- Abdomen uniformly black; base of the anal vein wanting, veins very coarse P. GRAVIDUS, Melander.
102. Crossveins at middle of wing; tarsi yellow and not or scarcely annulate; costa more or less thickened from first to third veins, base of anal vein wanting; mesonotum with golden pollen; third antennal joint about one-third the arista. 103.
- Crossveins before middle of wing; tarsi usually more or less annulate; costa not thickened 104.

(1) *Platypalpus oculeus*, nov. sp. — Length 1.7 mm. Black, densely cinereous pollinose, even the sternopleuræ covered except for a minute central spot, abdomen shining, testaceous, legs and antennæ yellow, tibial spur equal to the diameter of the tibia. Head obliquely drawn out below, the antennæ inserted well down so that the face is only one-half as long as the front, epistome not pruinose; palpi yellow, proboscis brown; vertex with white setæ. Thoracic bristles yellow, no humeral, three small notopleurals, one supraalar, one postalar, seven dorsocentrals, four scutellars the lateral ones small. Front femora biseriately ciliate beneath, middle femora not thicker than the front ones, their usual two rows of flexor setulæ yellow, posteriorly with six yellow setæ, middle tibiae three-fourths as long as their femora. Veins pale yellow, costa not thickened, third and fourth veins straight, nearly parallel, the first posterior cell widest at apex, second, third and fourth sections of the costa proportioned 2.4 : 2.4 : 1, posterior crossvein before the middle of the wing, nearly transverse, separated from the anterior by the length of the latter, sections of the fifth vein proportioned 0.5 : 0.8 : 1, anal crossvein perpendicular, base of the very weak anal vein entirely wanting, marginal cilia shorter than the anterior crossvein.

One specimen, Philadelphia, Pennsylvania (Brues). The pollinose pleuræ, reddish abdomen, pronounced dorsocentrals, yellow femoral setulæ and slanting head make a combination of characters readily distinguishing this species.

(2) *Platypalpus talaris*, nov. sp. — Length 2.5 mm. Black; front cinerous, rather broad, widening above, face narrow and white, epistome black; palpi oval, yellow, not hairy and with a single apical seta; proboscis black; antennæ three-jointed, the last joint elongate ovate, apparently twice as long as wide and about one-half as long as the arista; one pair of black vertical bristles. Thorax moderately covered with olivaceous-yellow pollen, bristles black, humeral small, one notopleural, one supraalar, one postalar, one dorsocentral, two scutellar, acrostichal setulæ biseriate, sparse; pollen of the pleuræ yellowish, sternopleural spot large; abdomen shining black, pygidium very small. Legs including the coxae testaceous, tarsi very slightly dusky towards the tip, not annulate, front femora with thirteen yellow flexor cilia, middle femora with yellow setæ in back, middle tibiae two-thirds as long as their femora, their spur large. Veins blackish, first vein thickened on entering the costa, the second, third and fourth costal sections proportioned 4 : 2.3 : 1, third vein straight, fourth vein slightly undulating, crossveins slightly before the middle of the wing, nearly meeting, the posterior somewhat oblique, sections of the fifth vein proportioned 0.5 : 1 : 1, anal crossvein perpendicular, anal vein faint, marginal cilia short.

Type ♂ from Mount Constitution, Orcas Island, Washington, July 31, 1908. Additional specimens from Bellingham, Lynden (Melander) and Friday Harbor (Aldrich), Washington; Milwaukee, Wisconsin (Wheeler); Waubamie, Ontario (H. S. Parish), and Montreal, Quebec (Beaulieu).

103. Abdomen strongly fasciate with bands of pollen; first posterior cell widest at middle, its veins recurved *P. FASCIVENTRIS*, nov. sp. (1).
- Abdomen shining; third and fourth vein parallel *P. POSTPOSITUS*, nov. sp. (2).
104. Third antennal joint short-ovate, one-third to one-half as long as its arista; crossveins separated by more than the length of the anterior crossvein, base of the anal vein present but weak; epistome pruinose; front tibiæ with bristles on extensor side. 105.
- Third antennal joint long-ovate, more than half the length of arista; crossveins separated by less than the length of the anterior crossvein, base of the anal vein wanting; front tibiæ with weak or no extensor bristles. 106.
105. Antennæ black; tarsi strongly annulate and not dilated; front tibiæ with evident black extensor setæ, middle femora with yellow setæ in front *P. TRIVIALIS*, Loew.
- Antennæ red with black tip; middle tarsi ♂ black, the metatarsi dilated into a flat disk, tarsi ♀ weakly annulate, the middle pair blackish; extensor bristles of front tibiæ ♂ yellow, of ♀ black; middle femora without yellow setæ in front; palpi yellow *P. DISCIFER*, Loew.

(1) *Platypalpus fasciventris*, nov. sp. — Male. Length 3 mm. Body black, legs yellow, notum closely golden pollinose, bristles yellow, abdomen with pollinose fasciæ, lower third of sternopleuræ polished, tibial spur large, tarsi weakly annulate, front tibiæ without bristles, posterior crossvein oblique and located at the middle of the wing. Occipital hairs abundant and white, vertical bristles short and yellow, front nearly four times as long as broad, closely gray pollinose, its sides little diverging, face gray, epistome bare, proboscis black, palpi yellow, with conspicuous white hairs; antennæ black, the outer joint pyriform, twice as long as deep and less than one-half as long as the rather strong arista. Four yellow setæ grouped around the supra-alar bristle, humeral small, two small lateral scutellars in addition to the central pair, setulæ in four irregular rows, pollen of pleuræ gray. Abdominal fasciæ of gray pollen occupying the basal half of each segment and expanding laterally, seventh tergite entirely pollinose, ventral segments shining, abdominal hairs pale and scattering, pygidium not globose, small, the dorsal valves projecting backward, the left dorsal valve with a close fringe of downward-directed, nearly straight, yellow hairs. Coxæ yellow, front femora of nearly same diameter as the middle pair, weakly ciliate beneath, middle femora with about eighteen setulæ in each flexor row, those behind scarcely longer, the row flanked by a few weak outstanding yellow setæ, middle tibiæ three-fourths the length of the femur, the spur longer than the diameter of the tibiæ and tipped with black, extreme tips of the tarsal joints slightly browned. Calypteres, fringe and halteres yellow. Wings hyaline, anal angle well developed so that the anal vein lies midway between the fifth vein and the margin, third vein gently recurved to end at the wing-tip, first posterior cell widest at the middle, fourth vein sinuous, sections of the fifth vein 2 : 3 : 2, distance between the crossveins nearly equal to the anterior crossvein, anal crossvein almost perpendicular, its outer half abruptly faint, anal vein vanishing basally.

Holotype, Klickitat River near Glenwood, Washington, June 27, 1917 (Melander).

(2) *Platypalpus postpositus*, nov. sp. — Male. Length 1.8 mm. Mesonotum almost completely covered with golden pollen, the black ground color hardly showing through, legs yellow, the tarsi a little dusky apically, tibial spur long and slender, posterior crossvein beyond the middle of the wing, oblique, costa thickened between the first and third veins. Front cinereous, rather narrow, face white-pollinose, epistome shining black; palpi white and with white bristles, proboscis black; antennæ black, two-jointed, the last joint apparently short-ovate, the arista three times as long as this joint; one pair each of vertical and ocellar bristles yellow, hairs of lower occiput moderate and white. Thoracic bristles yellow, one humeral, two notopleural, one supraalar, one dorsocentral, one pair of long and one of short scutellar bristles, two definite rows of acrostichal setulæ; pleuræ silky white, a small oval sternopleural space glistening; abdomen subshining, its hairs white, long but rather sparse, fringes of the pygidium small Coxæ pale yellow, flexor hairs of the anterior femora longer, the middle femora with a posterior row of about fourteen yellow setæ in addition to the two rows of minute setulæ, spur of the middle tibiæ a little longer than the diameter of the tibia, acute and tipped with black. Veins yellow at the base, darker apically, crossveins separated by the length of the anterior crossvein, third and fourth veins parallel, sections of the fifth vein proportioned 0.8 : 1 : 1, base of anal vein wanting, marginal cilia equal in length of the anterior crossvein.

One specimen, Almota, Washington, June 24, 1911 (Melander). This species resembles *cellaris* but has black antennæ, a longer tibial spur, and less setulose thorax.

106. Antennæ and palpi yellow; tarsi entirely yellow; abdomen with interrupted pollinose fasciæ; first posterior cell greatly narrowed at tip P. HARPIGER, Melander.
- Antennæ and palpi black; tarsi more or less annulate; abdomen not pollinose 107.
107. Middle femora without yellow setæ in front, coxæ and legs yellow, tarsi weakly annulate; cilia of hind margin of the wing ♂ nearly as long as the posterior crossvein; arista longer than third antennal joint P. HASTATUS, Melander.
- Middle femora with conspicuous yellow setæ in front; cilia of the hind margin of the wing short; palpi blackish inside 108.
108. Epistome shining; tarsi weakly annulate; third vein straight; middle section of fifth vein shorter than the last section 109.
- Epistome heavily pruinose; tarsi strongly annulate 110.
109. Body black; middle femora with at most a spot of brown P. SUTOR, Melander.
- Body castaneous; all the femora annulate with brown or black P. ARCTICUS, nov. sp. (1).
110. Posterior coxæ dark at base; femora, especially the middle pair, with dark spot or ring; arista obviously longer than third antennal joint, middle section of fifth vein longer than the others P. ARMILLATUS, nov. sp. (2).
- Coxæ and femora yellow; arista subequal to third joint; last two sections of fifth vein subequal P. AEQUALIS, Loew.

(1) *Platypalpus arcticus*, nov. sp. — Female. Length 2 mm. Blackish brown, thorax dusted, sternopleuræ partly shining, bristles yellow, legs yellow, the femora broadly ringed with brown, tibial spur long, third antennal joint elongate oval, posterior crossvein oblique and located beyond the anterior. Occiput and front with yellow pollen, face whitish, nearly as broad as the front, the latter with nearly parallel sides, epistome shining; proboscis black, nearly as long as the head height, palpi blackish within, coated outside with yellowish; antennæ rather short, black, the third joint ovate, one-half longer than broad, the arista twice as long as the antenna. Mesonotum shining through the yellowish coating, bristles yellow, pleuræ whitish tomentose, the sternopleura largely shining, the coating thinner in the center of the mesopleura. Abdomen shining, the last two segments small and opaque yellow, styles brown. Base of the posterior coxæ brown, the dark band of the middle femora strongest, that of the hind femora placed beyond the middle, front and middle femora distinctly biseriately setose beneath, the setæ of the front femora fine, front tibiae short-ciliate outwardly, slightly thicker than the middle pair, middle tibiae three-fourths as long as the femora, the apical spur longer than the diameter of the tibia, slender and acute, tarsi weakly annulate, almost dusky toward the tip. Halteres yellow, calypteres with pale fringe. Wings with a yellowish tinge, veins light brown, anal crossvein forming an angle of sixty degrees, sections of the fifth vein proportioned 1 : 1 : 1.3, third and fourth veins nearly parallel, marginal cilia short.

Four cotype specimens, received from the U. S. National Museum; Fort Chimo, Ungava Bay, Labrador; L. M. Turner, collector.

(2) *Platypalpus armillatus*, nov. sp. Length 2.6 mm. Black; front and face of equal width, with parallel sides, front cinereous, face and epistome white pruinose; palpi elongate oval, black inside, white pruinose outside and with two apical setæ, proboscis black; antennæ black, apparently two-jointed, the last joint elongate oval, slightly more than twice as long as wide and three-fourths as long as the arista, cephalic bristles yellow, one pair each of vertical and ocellar bristles. Mesonotum moderately densely covered with cinereous pollen, its bristles yellow, the humeral bristle small, several notopleurals, one supraalar, one postalar, two dorsocentral and four scutellar bristles, acrostichals biseriate and weak; pollen of pleuræ whitish, anterior half of the sternopleuræ glabrous; abdomen shining black, pygidium moderately large, the dorsal valve with a yellow fringe. Base of the posterior coxæ black, remainder of the legs testaceous, the tarsi rather strongly annulate, posterior femora with a more or less developed blackish ring at the outer two-thirds, in the ♀ much less distinct, sometimes wanting, front femora with long cilia beneath, middle femora with some setæ in back, middle tibiae two-thirds as long as their femora, the spur strong. Veins coarse and black, first vein thickened on entering the costa, costal sections two to four proportioned 6 : 2.3 : 1, third and fourth veins slightly incurved, posterior crossvein oblique, twice as long as the anterior, the space between the crossveins equal to the anterior crossvein, sections of the fifth vein proportioned 0.8 : 1.3 : 1, anal crossvein at an angle of 80 degrees, anal vein very weak, vanishing at both ends, marginal cilia equal in length to the anterior crossvein.

Seventeen specimens, type from Almota, Washington, elevation 600 feet. Paratypes from Pullman, Clarkston, Kettle Falls, Oroville and Omak, Washington; Kendrick, Idaho (Melander) and Bozeman, Montana (R. A. Cooley).

Geographical distribution.

1. *P. aeneicollis*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3008 [1849] (*Tachydromia*); C. & N. Europe.
Strobl, Progr. Seitenstetten, Vol. 14, p. 7 (1880); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*).
2. *P. aeneus*, Macquart, Mém. Soc. Sc. Lille, p. 153 [1823] (*Tachydromia*); W. Europe.
Dipt. N. France, Vol. 3, p. 95 (1827); Hist. Nat. Dipt. Vol. 1, p. 352 (1834); Meigen, Syst. Beschr. Vol. 7, p. 97 [1838] (*Tachydromia*).
3. *P. aequalis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 88: Cent. 5, No. 75 (1864); North America.
Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 222 (1902).
4. *P. agilis*, Meigen, Syst. Beschr. Vol. 3, p. 80 [1822] (*Tachydromia*); Europe.
Zetterstedt, Dipt. Scand. Vol. 1, p. 275 (1842); ibidem, Vol. 8, p. 3004 [1849] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 123 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 91 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 121 [1893] (*Tachydromia*); Ver. Naturw. Hermannst. Vol. 46, p. 24 [1897] (*Tachydromia*); Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 314 [1906] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 411 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 311 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 80 [1913] (*Tachydromia*).
var. *hybridus*, Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 411 [1907] (*Tachydromia*). Finland.
5. *P. ? albens*, Pallas, in Wiedemann, Zool. Mag. Vol. 1 (2), p. 26 [1818] Tauria. (*Empis*).
6. *P. albicornis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 279 [1842] (*Tachydromia*); Europe.
Walker, Ins. Brit. Dipt. Vol. 1, p. 124 (1851); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 112 (1893) (*Tachydromia*); ibidem, Vol. 46, p. 81 [1909] (*Tachydromia*); Wien. Ent. Zeit. Vol. 18, p. 79 [1899] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 308 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 92 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 76 [1913] (*Tachydromia*).
7. *P. albifrons*, Macquart, Mém. Soc. Sc. Lille, 1823, p. 154 [1823] (*Tachydromia*) France.
8. *P. albipennis*, Perris, Ann. Soc. Linn. Lyon. Vol. 5, p. 200 [1852] (*Tachydromia*) France.
9. *P. albiseta*. Panzer, Fauna Ins. Germ. Vol. 103, pl. 17 [1806] (*Tachydromia*); Europe, N. Africa, East Indies.
Meigen, Syst. Beschr. Vol. 3, p. 73 [1822] (*Tachydromia*); ibidem, Vol. 6, p. 342 [1830] (*Tachydromia*); Macquart, Hist. Nat. Dipt. Vol. 1, p. 352 (1834); Loew, Progr. Posen, p. 22 [1840] (*Tachydromia*); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 310 [1842] (*Tachydromia*); Boitard, Man. Vol. 3, p. 322 (1843); Neuhaus, Dipt. March. p. 75 (1886); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 109 [1893] (*Tachydromia*); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 152 [1899] (*Tachydromia*); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 114 [1907] (*Tachydromia*); Frey, ibidem, Vol. 7, p. 413 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34 p. 21 [1908] (*Tachydromia*); Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 180 [1909] (*Tachydromia*); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 79 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica Vol. 3, p. 321 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Bezzi, Ann.

- Mus. Hungar. Vol. 10, p. 492 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 88 [1913] (*Tachydromia*).
? albicornis, Rossi, Fauna Etrusca Mantissa, Vol. 2, p. 77 [1794] (*Empis*); Bezzii, Bull. Soc. Ent. Ital. Vol. 30, p. 152 [1899] (*Tachydromia*).
albocapillatus, Fallen, Dipt. Suec. Empid. p. 9, pt. [1815] (*Tachydromia*); Zetterstedt, ibidem, Vol. 1, p. 309, pt. (1842); ibidem, Vol. 8, p. 3010, pt. [1849] (*Tachydromia*).
castanipes, Meigen, Syst. Beschr. Vol. 3, p. 79 [1822] (*Tachydromia*); Walker, List, Vol. 3, p. 508 (1849)? Ins. Brit. Dipt. Vol. 1, p. 131 (1851); Schiner, Fauna Austr. Vol. 1, p. 90 (1862).
fuscimanus, Zetterstedt, Dipt. Scand. Vol. 1, p. 292 [1842] (*Tachydromia*); A. Costa, Giamb. Vico Napoli, Vol. 2, p. 458 [1857] (*Tachydromia*).
vividus, Meigen, Syst. Beschr. Vol. 7, p. 97 [1838] (*Tachydromia*).
var. *brunnipes*, Strobl, Mem. Soc. Hist. Nat. Esp. Vol. 3, p. 312 [1906] (*Tachydromia*); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 79 [1909] (*Tachydromia*). Spain.
10. *P. albocapillatus*, Fallen, Dipt. Suec. Empid. p. 9, part [1815] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 79 [1822] (*Tachydromia*); Loew, Progr. Posen, p. 22 [1840] (*Tachydromia*); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 309, part [1842] (*Tachydromia*); Walker, List Dipt. Vol. 3, p. 508 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3010, part [1849] (*Tachydromia*); Scholz, Ent Zeitschr. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 152 [1861] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 90 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 109 [1893] (*Tachydromia*)? Wien. Ent. Zeit. Vol. 12, p. 38 [1893] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 413 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Naturw. Steiermark, Graz, Vol. 46, p. 79 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 322 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 87 [1913] (*Tachydromia*). Europe.
- geniculatus*, Fallen, Dipt. Suec. Empid. p. 7, part [1815] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 90, note (1862).
- . *P. albomicans*, Becker in Bezzii, Atti Soc. Veneto-Trent. Sc. Nat. (2), Vol. 1, p. 57 [1893] (*Tachydromia*), no description.
11. *P. albomicans*, Oldenberg, Ent. Mitteil. Deutsche Ent. Mus. Vol. 1, p. 214 Alps. [1912] (*Corynetea*).
12. *P. alexippus*, Walker, List Dipt. Vol. 3, p. 510 (1849). North America.
13. *P. algirus*, Macquart, Explor. Algér. Zool. Vol. 3, p. 445, pl. 3, f. 5 Algeria. (1849); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 119, note [1907] (*Tachydromia*).
14. *P. alpigenus*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, C. Europe. p. 117 [1893] (*Tachydromia*); ibidem, Vol. 34, p. 211, var. [1898] (*Tachydromia*); ibidem, Vol. 46, p. 83 [1909] (*Tachydromia*).
griseiceps, Becker, in litt. in Bezzii, Atti Soc. Veneto-Trent. Sc. Nat. (2), Vol. 1, p. 57 [1893] (*Tachydromia*).
15. *P. alumnus*, nov. sp. Wyoming.
16. *P. analis*, Meigen, Syst. Beschr. Vol. 6, p. 343 [1830] (*Tachydromia*); C. Europe. Schiner, Fauna Austr. Dipt. Vol. 1, p. 91 (1862).
luteus, Zetterstedt, Dipt. Scand. Vol. 1, p. 294, part [1842] (*Tachydromia*).
17. *P. andalusiacus*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 82 [1899] (*Tachydromia*). Spain.
18. *P. annularis*, Bezzii, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 404 [1909] Peru. (*Tachydromia*).

19. *P. annulatus*, Fallen, Empid. 7, part [1815] (*Tachydromia*); Zetterstedt, Ins. Lappon. p. 552, part [1838] (*Tachydromia*); Dipt. Scand. Vol. I, p. 301 [1842] (*Tachydromia*); Walker, List Dipt. Vol. 3, p. 508 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3009 [1849] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. I, p. 131 (1851). C. & N. Europe.
20. *P. annulipes*, Meigen, Syst. Beschr. Vol. 3, p. 77 [1822] (*Tachydromia*); Macquart, Hist. Nat. Dipt. Vol. I, p. 353 (1834); Boitard, Man. Ent. Vol. 3, p. 322 (1843); Schiner, Fauna Austr. Dipt. Vol. I, p. 89 (1862); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 154 [1899] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 297 [1910] (*Tachydromia*). Europe.
- ? *coxatus*, Zetterstedt, Dipt. Scand. Vol. I, p. 281 [1842] (*Tachydromia*); ibidem, Vol. 8, p. 3006 [1849] (*Tachydromia*). var. *obscurior*, Strobl, Progr. Seitenstetten, Vol. 14, p. 7 (1880). C. Europe.
21. *P. anomalicerus*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 45 [1902] (*Tachydromia*). Egypt.
22. *P. apicalis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 90 : Cent. 5, No. 79 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 218 (1902). United States.
- . *P. apicalis*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 265 (1829), no description.
23. *P. apicatus*, new name. Algeria.
apicalis, Becker (not Loew), Zeitschr. Hym. Dipt. Vol. 7, p. 115 [1907] (*Tachydromia*).
24. *P. approximatus*, Becker, Mitteil. Zool. Mus. Berl. Vol. 2, p. 45 [1902] N. Africa.
(*Tachydromia*); Bezzi, Bull. Soc. Ent. Ital. Vol. 37, p. 295 [1905]
(*Tachydromia*); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 117 [1907]
(*Tachydromia*).
25. *P. arcticus*, nov. sp. Labrador.
26. *P. argenteomicans*, Becker, Mitteil. Zool. Mus. Berl. Vol. 4, p. 40 [1908] Canary Islands.
(*Tachydromia*).
27. *P. argenticeps*, Meijere, Tijd. Ent. Vol. 56, Suppl. p. 77 [1914] (*Tachydromia*). Java.
28. *P. armillatus*, new. sp. W. United States.
29. *P. articulatoides*, Frey, Act. Soc. Sc. Fenn. Helsingfors, Vol. 46, p. 2, 11 [1918] (*Tachydromia*). N. Russia.
30. *P. articulatus*, Macquart, Dipt. N. France, Vol. 3, p. 98 (1827); Hist. Nat. Dipt. Vol. I, p. 354 (1834); Meigen, Syst. Beschr. Vol. 8, p. 98 [1838] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. I, p. 284 [1842] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 324 (1843); Walker, List Dipt. Vol. 3, p. 510 (1849); Scholz, Breslau Ent. Zeitschr. Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. I, p. 125 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 4988 [1859] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. I, p. 91 (1862); Strobl, Progr. Seitenstetten, Vol. 14, p. 8 (1880); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 113 [1893] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 408 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 300 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 92 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 77 (1913); Edwards, Ent. Mag. Vol. 50, p. 59 figs. [1914] (*Tachydromia*). Europe.
- maculimanus*, Zetterstedt, Dipt. Scand. Vol. I, p. 248 [1842] (*Tachydromia*).
31. *P. ater*, Wahlberg, Ofvers. Akad. Foerhandl. p. 106 [1844] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3007 [1849] (*Tachydromia*); Strobl, Europe.

Mem. Soc. Hist. Nat. Espan. Vol. 3, p. 315 [1906] (*Tachydromia*); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 84 [1909] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 81 [1913] (*Tachydromia*).

sumelanus, Mik, Wien. Ent. Zeit. Vol. 3, p. 82 [1884] (*Tachydromia*); Becker, Fauna Hernst. Vol. 2 (2), p. 61, fig. 9, 10 [1885] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 120 [1892] (*Tachydromia*); ibidem, Vol. 34, p. 212 [1898] (*Tachydromia*).

montanus, Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 411 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*).

32. *P. baldensis*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 81 [1899] (*Tachydromia*). S. Europe.
- nigrogeniculatus*, Bezz, Bull. Soc. Ent. Ital. Vol. 30, p. 81 [1899] (*Tachydromia*).
var. *nigrifemor*, Strobl, Wien. Ent. Zeit. Vol. Vol. 18, p. 81 [1899] (*Tachydromia*). Spain.
33. *P. ballistrarius*, nov. sp. Virginia.
34. *P. ballucatus*, nov. sp. Washington, Illinois.
35. *P. Beckeri*, Mik, Wien. Ent. Zeit. Vol. 13, p. 166 [1894] (*Tachydromia*). C. Europe, Alps.
- lateralis*, Becker (not Loew), Berl. Ent. Zeit. Vol. 31, p. 137 [1887] (*Tachydromia*).
36. *P. bicolor*, Meigen, Klassif. Beschr. Eur. Zweifl. Ins. Vol. 1, p. 237 [1804] Europe, N. Africa.
(*Tachydromia*); Fabricius, Syst. Antl. p. 143 [1805] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 82 [1822] (*Tachydromia*); Macquart, Mém. Soc. Sc. Lille, p. 152 [1823] (*Tachydromia*); Ins. Dipt. N. France, Vol. 3, p. 98, pl. 2, f. 4 (1827); Hist. Nat. p. 355 (1834); Zetterstedt, Ins. Lappon. p. 549 [1838] (*Tachydromia*); Loew, Progr. Posen, p. 22 [1840] (*Tachydromia*); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 276 [1842] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 324 (1843); Walker, List Dipt. Vol. 3, p. 509 (1849); Scholz, Ent. Zeitschr. Breslau, Vol. 5, p. 19, 57 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 123 (1851); Bonsdorff, Finl. tråv. Ins. Vol. 1, p. 148 [1861] (*Tachydromia*); Schiner, Fauna Austr. Ins. Vol. 1, p. 88 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 381, 384 [1866] (*Tachydromia*); Leunis, Synopsis, Vol. 2, p. 401 (1886); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 113 [1893] (*Tachydromia*); Glasn. Mus. Bosn. Vol. 14, p. 469, var. [1902] (*Tachydromia*); Mitteil. Bosn. Herzeg. Vol. 9, p. 527, var. [1904] (*Tachydromia*); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 117 [1907] (*Tachydromia*); Frey, ibidem, Vol. 7, p. 408 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 298, f. 135 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 77 [1913] (*Tachydromia*).
- calceatus*, Bonsdorff, Finl. tråv. Ins. Vol. 1, p. 149 [1861] (*Tachydromia*).
dickroa, Meigen, Syst. Beschr. Vol. 3, p. 83 [1822] (*Tachydromia*); Walker, List Dipt. Vol. 3, p. 509 (1849); Ins. Brit. Dipt. Vol. 1, p. 127 (1851).
flavipes, Fallen, Empid. p. 6, part [1815] (*Tachydromia*).
pallidiventris, Bonsdorff, Finl. tråv. Ins. Dipt. Vol. 1, p. 148 [1861] (*Tachydromia*).
37. *P. bicornis*, nov. sp. Washington.
38. *P. bivittatus*, Macquart, Dipt. N. France, Vol. 3, p. 97 (1827); Hist. Nat. France.
Ins. Vol. 1, p. 354 (1834); Meigen, Syst. Beschr. Vol. 7, p. 97 [1838] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 328 (1843).
39. *P. brevicornis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 293 [1842] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*). N. Europe.

- Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 76 [1913] (*Tachydromia*).
var. *subbrevis*, Frey, ibidem, Vol. 37 (3), p. 76 [1913] (*Tachydromia*).
brevicornis, Frey (not Zetterstedt), Zeitschr. Hym. Dipt. Vol. 7, p. 410 [1907]
(*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908]
(*Tachydromia*).
40. *P. Brunettii*, new name. W. Himalayas.
flavipes, Brunetti, Rec. India Mus. Vol. 9, p. 23 [1913] (*Howlettia*); Fauna
Brit. India Dipt. Vol. 1, p. 361 [1920] (*Howlettia*).
41. *P. brunneitibia*, Strobl, Wien. Ent. Zeitschr. Vol. 18, p. 78 [1899] (*Tachydromia*); C. & S. Europe.
dromia; Bezzī, Bull. Soc. Ent. Ital. Vol. 30, p. 153 [1899] (*Tachydromia*).
brachystyla, Becker, in litt. in Bezzī, Atti. Soc. Veneto-Trent. Sc. Nat. (2)
Vol. 1, p. 57 [1893] (*Tachydromia*).
42. *P. brunnipes*, Gimmerthal, Bull. Soc. Imp. Nat. Moscou, Vol. 15, p. 667 Russia.
[1842] (*Tachydromia*).
43. *P. calceatus*, Meigen, Syst. Beschr. Vol. 3, p. 87 [1822] (*Tachydromia*); C. & N. Europe.
Macquart, Dipt. N. France, Vol. 3, p. 100 (1827); Hist. Nat. Ins.
Vol. 1, p. 356 (1834); Zetterstedt, Ins. Lappon. p. 550 [1838]
(*Tachydromia*); Loew, Progr. Posen, p. 23 (1840); Isis, Vol. 7,
p. 552 [1840] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 282
[1842] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 324 (1843);
Gimmerthal, Bull. Soc. Imp. Nat. Moscou, Vol. 20 (2), p. 165
[1847] (*Tachydromia*); Walker, List Dipt. Vol. 3, p. 510 (1849);
Zetterstedt, Dipt. Scand. Vol. 8, p. 3006 [1849] (*Tachydromia*);
Scholz, Zeitschr. Ent. Breslau, Vol. 5(19), p. 57 [1851] (*Tachydromia*);
Walker, Ins. Brit. Dipt. Vol. 1, p. 124 (1851); Zetterstedt, Dipt.
Scand. Vol. 13, p. 4988 [1859] (*Tachydromia*); Schiner, Fauna Austr.
Vol. 1, p. 88 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 137
[1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark,
Graz, Vol. 29, p. 113 [1893] (*Tachydromia*); Frey, Hym. Dipt.
Vol. 7, p. 408 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Nat. Ver.
Steiermark, Graz, Vol. 46, p. 81 [1909] (*Tachydromia*); Lundbeck,
Dipt. Danica, Vol. 3, p. 299 [1910] (*Tachydromia*); Wahlgren, Ent.
Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Frey, Acta Soc. Sc.
Fenn. Helsingfors, Vol. 37 (3), p. 77 [1913] (*Tachydromia*).
ecalceatus, Zetterstedt, Ins. Lappon. p. 550 [1838] (*Tachydromia*); Dipt. Sc.
Vol. 1, p. 283 [1842] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1,
p. 124 (1851).
flavipes, Fallen, Empid. p. 6, pt. [1815] (*Tachydromia*).
44. *P. caligaris*, new name. Mexico.
caligatus, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 217, var. (1902).
trivialis, Wheeler and Melander, Biol. C. Amer. Dipt. Vol. 1, p. 375, var.
(1901).
45. *P. caligatus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 216, part, Mexico.
f. 27, 35, 46 (1902); Bezzī, Nova Acta Akad. Naturf. Halle, Vol. 91,
p. 402 [1909] (*Tachydromia*).
trivialis, Wheeler and Melander, part, Biol. C. Amer. Dipt. Vol. 1, p. 375
(1901).
46. *P. callithrix*, nov. sp. Illinois.
47. *P. candicans*, Fallen, Dipt. Suec. Emp. p. 10 [1815] (*Tachydromia*); Europe.
Meigen, Syst. Beschr. Vol. 3, p. 85 [1822] (*Tachydromia*); Zetterstedt,
Ins. Lappon. p. 550, part [1838] (*Tachydromia*); Dipt. Scand. Vol. 1,
p. 285, part [1842] (*Tachydromia*); Walker, List Dipt. Vol. 3, p. 509

- (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3006 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 125 (1851); Pipping, Not. Sællsk. Fenn. Förh. Vol. 4, p. 114 [1858] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 13, p. 4989 [1859] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 149 [1861] (*Tachydromia*); Schiner, Fauna Austr. Vol. 1, p. 89 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 114 [1893] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 409 [1907] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 303, f. 138 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Dahl, Fauna Chorin, p. 466 (1912); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 77 [1913] (*Tachydromia*).
cursitans, Frey (not Fabricius), Zeitschr. Hym. Dipt. Vol. 7, p. 409 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 78 [1913] (*Tachydromia*).
major, Bonsdorff (not Zetterstedt), Finl. tvåv. Ins. Dipt. Vol. 1, p. 149 [1861] (*Tachydromia*).
ventralis, Meigen, Syst. Beschr. Vol. 3, p. 85 [1822] (*Tachydromia*); Walker, Dipt. Brit. Mus. Vol. 3, p. 509 (1849); Schiner, Fauna Austr. Dipt. Vol. 1, p. 88 (1862).
48. *P. candidiseta*, Bezz, Ann. Mus. Hungar. Vol. 10, p. 492 [1912] (*Coryneta*). Formosa.
49. *P. canus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 220, f. 29, 40, California.
 45 (1902); Ent. News, Vol. 17, p. 372 [1906] (*Symballopithalmus*).
50. *P. celer*, Meigen, Syst. Beschr. Vol. 3, p. 80 [1822] (*Tachydromia*). C. Europe.
51. *P. cellarius*, nov. sp. W. North America.
52. *P. chilensis*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 767 (1865); Chile.
 Bezz, Ann. Mus. Hungar. Vol. 3, p. 459 [1905] (*Tachydromia*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 403 [1909] (*Tachydromia*).
53. *P. chionochata*, Bezz, Ann. Mus. Hungar. Vol. 2, p. 359 (1904). New Guinea.
54. *P. ciliaris*, Fallen, Empid. p. 33 [1816] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 86 [1822] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 289 [1842] (*Tachydromia*); Walker, List Dipt. Vol. 3, p. 509 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3006 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 128 (1851); Schiner, Fauna Austr. Vol. 1, p. 90 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 3 [1893] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 412 [1907] (*Tachydromia ciliata*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 316 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 83, f. 26 [1913] (*Tachydromia*). C. & N. Europe.
- pygmaeus*, Zetterstedt, Ins. Lappon. p. 551 [1838] (*Tachydromia*).
55. *P. cinereovittatus*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 78 [1899] (*Tachydromia*). Spain.
56. *P. cingulatus*, Loew, Progr. Posen, p. 23 (1840); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Schiner, Fauna Austr. Vol. 1, p. 91 (1862). C. Europe.
57. *P. collaris*, Meigen, Syst. Beschr. Vol. 7, p. 99 [1838] (*Tachydromia*); Schiner, Fauna Austr. Vol. 1, p. 91 (1862). C. Europe.
58. *P. collateralis*, nov. sp. *lateralis*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 421 (1900).
59. *P. commendatus*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 123 [1914] E. Africa. (*Coryneta*).

60. *P. commiles*, Walker, Ins. Brit. Dipt. Vol. 1, p. 128 (1851). England.
61. *P. commutatus*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 111 [1893] (*Tachydromia*); ibidem, Vol. 46, p. 80 [1909] (*Tachydromia*). C. Europe.
62. *P. comptus*, Walker, Ent. Mag. Vol. 4, p. 228 (1837); List, Vol. 3, p. 508 (1849); Ins. Brit. Dipt. Vol. 1, p. 132 (1851); Schiner, Fauna Austr. Vol. 1, p. 91 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 109 (1893); ibidem, Vol. 34, p. 210 (1898); ibidem, Vol. 46, p. 79 [1909] (*Tachydromia*). C. and N. Europe.
63. *P. compungens*, Walker, Ins. Brit. Dipt. Vol. 1, p. 128 (1851). England.
64. *P. *concitatus*, Meunier, Ann. Sc. Nat. Zool. Vol. 7, p. 89, 100, pl. 5, f. 6, 7 (1908). Baltic Amber.
65. *P. confinis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 307 [1842] (*Tachydromia*); ibidem, Vol. 8, p. 3010 [1849] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 85, f. 28 [1913] (*Tachydromia*). N. Europe.
- . *P. consortus*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 265 [1829] (*Tachydromia*), no description.
66. *P. contiguus*, nov. sp. British Columbia.
67. *P. Coquilletti*, Melander, Occ. Papers Bost. Soc. Hist. Nat. Vol. 5, p. 83 (1824). E. United States.
- trivialis, Melander (not Loew), Trans. Amer. Ent. Soc. Vol. 28, p. 216, f. 26, 34, 43 (1902).
68. *P. cothurnatus*, Macquart, Dipt. N. France, Vol. 3, p. 100 (1827); Hist. Nat. Dipt. Vol. 1, p. 356 (1834); Meigen, Syst. Beschr. Vol. 7, p. 98 [1838] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 285 [1842] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 324 (1843); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 57 [1851] (*Tachydromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 88 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 116 [1893] (*Tachydromia*); ibidem, Vol. 34, p. 211 [1898] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 409 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Acta Soc. Fenn. Helsingfors, Vol. 31, p. 9 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 294 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 79 [1913] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*). N. & C. Europe.
- socculus, Zetterstedt, Ins. Lappon. p. 550 [1838] (*Tachydromia*).
69. *P. crassifemoris*, Fitch, 1 and 2 Rept. N. Y., p. 301 [1856] (*Oscinis*); United States. Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 223 (1902).
- var. *debilis*, Loew, Ent. Zeitschr. Vol. 7, p. 20: Cent. 3, No. 37 (1863); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1896). United States.
- var. *melanocerus*, nov. var. Washington.
- var. *mollis*, nov. var. United States.
70. *P. crassiseta*, Strobl, Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 314 [1906] (*Tachydromia*); Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 180 [1909] (*Tachydromia*). Spain.
71. *P. crepidarius*, nov. sp. Washington.
72. *P. cryptospina*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 8 [1908] (*Tachydromia*); ibidem, Vol. 37 (3), p. 79 [1913] (*Tachydromia*). Finland.
73. *P. cuneipennis*, Melander, Occ. Pap. Bost. Soc. Nat. Hist. Vol. 5, p. 83 (1924). Vermont.

74. *P. cursitans*, Fabricius, Syst. Ent. p. 782 [1775] (*Musca*); Spec. Ins. Vol. 2, Europe.
 p. 447 [1781] (*Musca*); Mantissa, Ins. Vol. 2, p. 349 [1787] (*Musca*);
 Gmelin, Syst. Nat. Vol. 5, p. 2852 [1790] (*Musca*); Fabricius, Ent. Syst. Vol. 4, p. 339 [1794] (*Musca*); Schrank, Fauna Boica, Vol. 3, p. 122 [1803] (*Musca*); Fabricius, Syst. Antl. p. 143 [1805] (*Tachydromia*); Olivier, Encyc. Méth. Vol. 8, p. 30 [1811] (*Musca*); Macquart, Dipt. N. France, Vol. 3, p. 99 (1827); Hist. Nat. Dipt. Vol. 1, p. 355 (1834); Loew, Progr. Posen, p. 22 (1840); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 280 [1842] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 324 (1843); Gimmerthal, Bull. Moscou, Vol. 20 (2), p. 165 [1847] (*Tachydromia*); Walker, List, Vol. 3, p. 509 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3005 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 124 (1851); Schiner, Fauna Austr. Dipt. Vol. 1, p. 88 (1862); Leunis, Synop. Natur. Vol. 2, p. 401 (1886); Neuhaus, Dipt. March. p. 75 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 136 [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 114 [1893] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 303, f. 137 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*).
flavipes, Thunberg, Nova Acta Upsalensis, Vol. 4, p. 26, var. b, part [1784] (*Empis*).
 var. *chrysonotum*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 80 [1899] (*Tachydromia*); Spain.
 Mém. Soc. Esp. N. Hist. Vol. 3, p. 313 [1906] (*Tachydromia*); Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 180 [1909] (*Tachydromia*).
 var. *denominatus*, Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 409 [1907] (*Tachydromia*). Finland.
 var. *hispanicus*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 79 [1899] (*Tachydromia*). Spain.
 var. *minor*, Strobl, ibidem, Vol. 18, p. 79 [1899] (*Tachydromia*). Spain.
75. *P. curticornis*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3004, note [1849] (*Tachydromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 91 (1862). C. Europe.
76. *P. dalmatinus*, Strobl, Zem. Mus. Bosn. Herceg. Vol. 14, p. 469 [1902] (*Tachydromia*); Wiss. Mitteil. Mus. Bosn. Herceg. Vol. 9, p. 527 [1904] (*Tachydromia*). S. Europe.
77. *P. decolor*, nov. sp. Montana.
78. *P. desertorum*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 117 [1907] (*Tachydromia*). Algeria.
79. *P. difficilis*, Frey, ibidem, Vol. 7, p. 410 [1907] (*Tachydromia*); Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 79, f. 21 [1913] (*Tachydromia*). Finland.
80. *P. dilatatovittatus*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 77 [1909] (*Tachydromia*). C. Europe.
81. *P. direptor*, nov. sp. British Columbia.
82. *P. discifer*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 20 : Cent. 3, No. 36 (1863); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 222, 342 (1902). E. United States.
83. *P. dissimilipes*, nov. sp. W. North America.
84. *P. diversipes*, Coquillett, Proc. Wash. Acad. Sc. Vol. 2, p. 422 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 218 (1902). Alaska.
85. *P. divisus*, Walker, Ins. Brit. Dipt. Vol. 1, p. 127 (1851). England.
86. *P. dubius*, ibidem, Vol. 1, p. 132 (1851). Britain.
87. *P. enervatus*, nov. sp. California.
88. *P. engadinicus*, Mik, Wien. Ent. Zeit. Vol. 15, p. 106 [1896] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 77 [1909] (*Tachydromia*). C. Europe.
- testaceus*, Becker (not Philippi), Berl. Ent. Zeitschr. Vol. 31, p. 135 [1887]

- (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 108 [1892].
- var. *Beckeri*, Strobl, ibidem, Vol. 34, p. 210 [1898] (*Tachydromia*). C. Europe.
- Stroblii*, Mik, Wien. Ent. Zeit. Vol. 19, p. 148 [1900] (*Tachydromia*).
89. *P. eumerus*, Bezz, Nova Acta Akad. Naturf. Halle Vol. 91, p. 404 [1909] (*Tachydromia*). Peru.
90. *P. *eversoris*, Meunier, Ann. Sc. Nat. Zool. Vol. 7, p. 90, 102, pl. 6, f. 2 [1908]. Baltic Amber.
91. *P. excisus*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 114 [1907] (*Tachydromia*); Mitteil. Zool. Mus. Berl. Vol. 4, p. 39 [1908] (*Tachydromia*). Europe, Africa, Asia.
92. *P. exiguum*, Meigen, Syst. Beschr. Vol. 3, p. 81 [1822] (*Tachydromia*); N. & C. Europe.
- Walker, List Dipt. Vol. 3, p. 508 (1849); Ins. Brit. Dipt. Vol. 1, p. 132 (1851); Schiner, Fauna Austr. Dipt. Vol. 1, p. 89 (1862); Strobl, Mitteil. Steiermark, Graz, Vol. 29, p. 118 [1892] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 310 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*).
- femoralis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 299 [1842] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 90 (1862); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 120 [1892] (*Tachydromia*).
- geniculatus*, Fallen, Empid. 7, part [1815] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 90, note (1862).
- var. *nigrofemoratus*, Strobl (not 1906), Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*). C. Europe.
93. *P. exilis*, Meigen, Syst. Beschr. Vol. 3, p. 90 [1822] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 90 (1862); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 7, 8 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 313 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 81 [1913] (*Tachydromia*). C. Europe.
- var. *nigroterminatus*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 77 [1909] (*Tachydromia*). C. Europe.
94. *P. fagorum*, Dahl, Fauna Chorin, p. 466 (1912). Germany.
95. *P. fasciatus*, Meigen, Syst. Beschr. Vol. 3, p. 86, pl. 23, f. 22 [1822] (*Tachydromia*); Walker, List Vol. 3, p. 509 (1849); Ins. Brit. Dipt. Vol. 1, p. 126, pl. 5, f. 2 (1851); Kawall, Stett. Ent. Zeit. Vol. 16, p. 229 [1855] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 89 (1862); Glover, Manuscr. notes, 54, pl. 11, f. 8 [1874] (*Tachydromia*); Strobl, Mitteil. Ver. Steiermark, Graz, Vol. 29, p. 115 [1892] (*Tachydromia*); Jahrb. Mus. Kärnten. Vol. 47, p. 204 (1901); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 118 [1907] (*Tachydromia*); Frey, ibidem, Vol. 7, p. 409 [1907] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 305, f. 139 [1910] (*Tachydromia*). Europe, N. Africa.
- candidans*, Zetterstedt, part, Ins. Lappon. p. 550, part [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 285, part [1842] (*Tachydromia*).
- pallidiventris*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 138, part [1887] (*Tachydromia*).
96. *P. fascipes*, Meigen, Syst. Beschr. Vol. 3, p. 78 [1822] (*Tachydromia*); Europe.
- Vol. 6, p. 342 [1830] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 126 (1851); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 411 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*); Frey, Acta Sc. Soc. Fenn. Helsingfors, Vol. 37 (3), p. 79 [1913] (*Tachydromia*).
- confinis*, Bonsdorff, Finl. tvåv. Ins. Vol. 1, p. 152 [1861] (*Tachydromia*).
- notatus*, Meigen, Syst. Beschr. Vol. 3, p. 78 [1822] (*Tachydromia*); Loew,

- Progr. Posen, p. 22 (1840) : Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 275 [1842] (*Tachydromia*); Gimmerthal, Bull. Soc. Mosc. Vol. 20, p. 164 [1844] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3004 [1849] (*Tachydromia*); ibidem, Vol. 13, p. 4986 [1859] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 148 [1861] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 89 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 381 [1866] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 122 [1892] (*Tachydromia*); Jahrb. Mus. Kärnten, Vol. 47, p. 205 (1901); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 84 [1909] (*Tachydromia*).
- var. *nigrofemoratus*, Strobl, Mem. Soc. Esp. Nat. Hist. Vol. 3, p. 315 [1906] (*Tachydromia*). Spain.
- var. *obscurior*, Strobl, Glasnik Mus. Bosn. Herceg. Vol. 14, p. 469 [1902] (*Tachydromia*); Mitteil. Bosn. Herzeg. Vol. 9, p. 528 [1904] (*Tachydromia*). S. Europe.
- var. *pallidicoxa*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 80 [1913] (*Tachydromia*). Finland.
- strigifrons*, Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 410 [1907] (*Tachydromia*).
97. *P. fasciventris*, nov. sp. Washington.
98. *P. ferrugineus*, Brunetti, Rec. Ind. Mus. Vol. 9, p. 40 (1913); Fauna Brit. Ind. Dipt. Vol. 1, p. 376 [1920] (*Tachydromia*). India.
99. *P. flammifer*, Melander, Occ. Pap. Bost. Soc. Nat. Hist. Vol. 5, p. 84 E. United States.
100. *P. ?flavescens*, Rossi, Fauna Etrusca, Mantissa, Vol. 2, p. 77 [1794] (*Empis*); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 152 [1899] (? *Tachydromia*). S. Europe.
101. *P. flavicornis*, Macquart (not Meigen), Mem. Soc. Sc. Lille, p. 152 [1823] (*Tachydromia*). France.
102. *P. flavicornis*, Meigen, Syst. Beschr. Vol. 3, p. 83 [1822] (*Tachydromia*); Macquart, Dipt. N. France, Vol. 3, p. 99 (1827); Hist. Nat. Dipt. Vol. 1, p. 355 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 278 [1842] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 324 (1843); Walker, List. Vol. 3, p. 509 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3005 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 [1851] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 13, p. 4987 [1859] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 149 [1861] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 88 (1862); Strobl, Jahrb. Land. Mus. Kärnt. Vol. 47, p. 203 (1901); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 117 [1907] (*Tachydromia*); Frey, ibidem, Vol. 7, p. 408 [1907] (*Tachydromia*); Med. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Dan. Vol. 3, p. 307 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 92 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 76 [1913] (*Tachydromia*).
- flavipes*, Thunberg, Nova Acta Upsalensis, Vol. 4, p. 26, var. b, part (1784) (*Empis*).
- ruficornis*, Macquart, Dipt. Exot. Suppl. Vol. 4, p. 97 (1849).
103. *P. flavicoxis*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 116 [1907] (*Tachydromia*). Algeria.
104. *P. flavipennis*, Walker, Ins. Brit. Dipt. Vol. 1, p. 130 (1851). England.
105. *P. flavipes*, Fabricius, Ent. Syst. Vol. 4, p. 406 [1794] (*Empis*); Meigen, Klassif. Vol. 1, p. 233 [1804] (*Empis*); Fabricius, Syst. Antl. p. 142 [1805] (*Tachydromia*); Fallen, Empid. p. 6, part [1815] (*Tachydromia*); Billberg, Enumeratio, p. 119 [1820] (*Sicus*); Meigen, Syst. Beschr. Vol. 3, p. 81 [1822] (*Tachydromia*); Macquart, Mém. Soc. Lille, 1823, p. 152 [1823] (*Tachydromia*); Ins. N. France, Vol. 3, p. 97 (1827); Hist. Nat. Dipt. Vol. 1, p. 354, pl. 8, f. 10 (1834); Zetterstedt, Ins. Lappon. p. 549 [1838] (*Tachydromia*); Loew, Progr. Posen, p. 22 (1840); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 273 [1842] (*Tachydromia*); Boitard,

Nouv. Man. Ent. Vol. 3, p. 324 (1843); Walker, List, Vol. 3, p. 508 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3004 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 122, pl. 5, f. 2 (1851); Zetterstedt, Dipt. Scand. Vol. 12, p. 4600 [1855] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 148 [1861] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 87 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 399 [1866] (*Tachydromia*); Walker, Ent. Vol. 5, p. 273 (1871); Jaroschewsky, Trudy Khark. Vol. 11, p. 352 (1877); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 116 [1892] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 409 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3 (6), p. 290 f. 132 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Becker, Ann. Mus. Zool. St-Petersburg, Vol. 17, p. 597 [1913] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 78 [1913] (*Tachydromia*).

vulgaris, Meigen, Klassif. Vol. 1, p. 237 [1804] (*Tachydromia*).

var. *fulvipes*, Meigen, Syst. Beschr. Vol. 3, p. 78 [1822] (*Tachydromia*); Walker, Europe. List. Vol. 3, p. 508 (1849); Ins. Brit. Dipt. Vol. 1, p. 125 (1851); Schiner, Fauna Austr. Dipt. Vol. 1, p. 89 (1862); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 409 [1907] (*Tachydromia*); Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 9 [1908] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 295 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 78 [1913] (*Tachydromia*).

fascipes, Strobl (not Meigen), Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 120 [1892] (*Tachydromia*); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 158 [1899] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 84 [1909] (*Tachydromia*).

var. *gracilis*, Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 57 [1851] (*Tachydromia*). C. Europe.
var. *melanochætus*, Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 158 [1899] (*Tachydromia*). S. Europe.
var. *pseudofulvipes*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31, p. 9 (1908); Finland. Vol. 37 (3), p. 79 [1913] (*Tachydromia*).

106. *P. flavipes*, Scopoli, Ent. Carniolica, p. 366 [1763] (*Asilus*); Thunberg, Nova Acta Upsalensis, Vol. 4, p. 26, var. part [1784] (*Empis*); Olivier, Encyclop. Method. Vol. 6, p. 390 [1791] (*Empis*); Schiner, Verh. Zool-bot. Ges. Wien. Vol. 6, p. 421 [1856] (*Tachydromia*). C. Europe.

107. *P. flavirostris*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 90 : Cent. 5, No. 80 (1864); Coquillett, Proc. U. S. Mus. Vol. 18, p. 438 (1895); Proc. Wash. Acad. Sc. Vol. 2, p. 422 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 219, 342 (1902). New Hampshire.

var. *dilutior*, nov. var. Washington.
var. *microcerus*, nov. var. Idaho.
var. *vittiger*, nov. var. Alaska, Montana.

108. *P. flavus*, Macquart, Mém. Soc. Sc. Lille, p. 153 [1823] (*Tachydromia*).

109. *P. formalis*, Walker, Ins. Brit. Dipt. Vol. 1, p. 130 (1851). France.
110. *P. fulcratus*, Scopoli, Ent. Carniolica, p. 366 [1763] (*Asilus*); Olivier, England.

Encycl. Meth. Vol. 6, p. 390 [1791] (*Empis*); Schiner, Verh. Zool-bot. Ver. Wien. Vol. 6, p. 421 [1856] (*Tachydromia*). C. Europe.

111. *P. fuscicornis*, Zetterstedt, Dipt. Scand. Vol. 1, p. 291 [1842] (*Tachydromia*); ibidem, Vol. 8, p. 3006 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 6 (19), p. 58 [1851] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 90 (1862); Strobl, Wien. Ent. Zeit. Vol. 18, p. 78 [1899] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 316 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. N. & C. Europe.

- Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 83 [1913] (*Tachydromia*).
pallipes, Meigen (not Fallen), Syst. Beschr. Vol. 3, p. 74 [1822] (*Tachydromia*).
112. *P. fuscitarsis*, Zetterstedt, Dipt. Scand. Vol. 13, p. 4990 [1859] (*Tachydromia*); N. Europe.
Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*).
113. *P. geniculatus*, Meigen (not Fallen), Syst. Beschr. Vol. 3, p. 75 [1822] C. Europe.
(*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 1, p. 311, note [1842] (*Tachydromia*).
114. *P. gentilis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 40 [1913]; Fauna Brit. Himalayas.
India Dipt. Vol. 1, p. 377 [1920] (*Tachydromia*).
115. *P. gesticulator*, nov. sp. Montana.
116. *P. gilvipes*, Meigen, Syst. Beschr. Vol. 3, p. 87 [1822] (*Tachydromia*); C. Europe.
Scholz, Zeitschr. Ent. Breslau. Vol. 5 (19), p. 59 [1851] (*Tachydromia*); Loew, Zeitschr. Ges. Naturw. Vol. 8, p. 101 [1857] (*Tachydromia*).
flavipes, Meigen (not Fabricius), Klassif. Beschr. Eur. Zweif. Ins. Vol. 1, p. 238 [1804] (*Tachydromia*).
117. *P. glaber*, Meigen, Syst. Beschr. Vol. 3, p. 89 [1822] (*Tachydromia*); C. Europe.
Walker, Ins. Brit. Dipt. Vol. 1, p. 129 (1851).
118. *P. glabratus*, Meigen, Syst. Beschr. Vol. 7, p. 99 [1838] (*Tachydromia*); C. Europe.
Schiner, Fauna Austr. Dipt. Vol. 1, p. 92 (1862).
119. *P. glacialis*, nov. sp. Montana, Washington.
120. *P. gravidus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 221, pl. 5, California.
f. 25; pl. 6, f. 42 (1902).
121. *P. harpiger*, Melander, Occ. Pap. Bost. Soc. Nat. Hist. Vol. 5, p. 84 Massachusetts.
(1924).
122. *P. hastatus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 222, f. 30 United States.
(1902).
123. *P. hians*, Melander, ibidem, Vol. 28, p. 220, f. 32, 36, 37 (1902); Ent. North America.
News, Vol. 17, p. 372 [1906] (*Symballophtalmus*).
var. *fuscohalteratus*, Melander, Occ. Pap. Bost. Soc. Nat. Hist. Vol. 5, p. 85 E. United States.
(1924).
124. *P. holosericus*, Melander, ibidem, Vol. 5, p. 85 (1924). E. North America.
125. *P. immaculatus*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 45 [1902] Egypt.
(*Tachydromia*).
126. *P. impexus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 219 (1902). United States.
127. *P. incurvus*, Melander, ibidem, Vol. 28, p. 221, f. 31, 33, 44 (1902). California.
128. *P. inferialis*, nov. sp. W. United States.
129. *P. infuscatus*, Meigen, Syst. Beschr. Vol. 3, p. 84 [1822] (*Tachydromia*); C. Europe.
Macquart, Dipt. N. France, Vol. 3, p. 99 (1827); Hist. Nat. Dipt. Vol. 1, p. 355 (1834); Boitard, Man. Ent. Vol. 3, p. 324 (1843); Walker, Ins. Brit. Dipt. Vol. 1, p. 126 (1851); Schiner, Fauna Austr. Dipt. Vol. 1, p. 88 (1862); Neuhaus, Dipt. Marchica, p. 74 (1886); Leunis, Synop. Vol. 2, p. 401 (1886); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 115 [1892] (*Tachydromia*).
130. *P. inops*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 220 (1902); United States.
Ent. News, Vol. 17, p. 372 [1906] (*Symballophtalmus*).
var. *aquicornis*, nov. var. Montana.
131. *P. *interactoris*, Meunier, Ann. Sc. Nat. Zool. Vol. 7, p. 89, 102, pl. 5, Baltic Amber.
f. 12, pl. 6, f. 1 (1908).
132. *P. interjectus*, Lundbeck, Dipt. Danica, Vol. 3, p. 295 [1910] (*Tachydromia*). Denmark.
133. *P. juvenis*, nov. sp. W. North America.
var. *puerinus*, nov. var. North America.
var. *hyænoides*, nov. var. Washington.

134. *P. lacertosus*, nov. sp. Idaho.
135. *P. lacteiseta*, Collin, Ent. M. Mag. London, 699, p. 188 [1922] (*Tachydromia*). Seychelles.
136. *P. læstadianorum*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 83, f. 27 [1913] (*Tachydromia*); Ent. Tidskr. 1914, p. 80 (1914). Finland, Sweden.
fuscicornis, Frey (not Zetterstedt), Zeitschr. Hym. Dipt. Vol. 7, p. 412 [1907] (*Tachydromia*); Medd. Soc. Sc. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*).
137. *P. latabilis*, nov. sp. New York.
138. *P. latetus*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 91 : Cent. 5, No. 81, (1864); Coquillett, Proc. U. S. Mus. Vol. 18, p. 438 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 219, 341 (1902). New Hampshire.
139. *P. lateralis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 89 : Cent. 5, No. 78 (1864); Coquillett, Proc. U. S. Mus. Vol. 18, p. 438 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 222 (1902). E. United States.
140. *P. latericia*, Becker, Ann. Soc. Ent. France, Vol. 83, p. 124 [1914] (Coryneta). E. Africa.
141. *P. laticinctus*, Walker, Ins. Brit. Dipt. Vol. 1, p. 127 (1851). England.
142. *P. lessinensis*, Strobl, Wien. Ent. Zeit. Vol. 12, p. 38 [1893] (*Tachydromia*). S. Europe.
143. *P. leucocephalus*, Roser, Correspondenzbl. Landw. Württemb. Vol. 1, p. 54 [1840] (*Tachydromia*). C. Europe.
144. *P. leucochætus*, Becker, Berl. Ent. Zeitschr. Vol. 33, p. 345 [1889] (*Tachydromia*); ? Strobl, Wien. Ent. Zeit. Vol. 12, p. 38 [1893] (*Tachydromia*); Bezzi, Bull. Soc. Ent. Ital. Vol. 30, p. 152 [1899] (*Tachydromia*). C. & S. Europe.
145. *P. leucothrix*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 81 [1909] (*Tachydromia*). C. Europe.
146. *P. lineatus*, Meigen, Klassif. Vol. 1, p. 238, pl. 12, f. 5 [1804] (*Tachydromia*); Syst. Beschr. Vol. 3, p. 88 [1822] (*Tachydromia*). C. Europe.
147. *P. longicornis*, Meigen, ibidem, Vol. 3, p. 73, pl. 23, f. 17 [1822] (*Tachydromia*); Walker, List, Vol. 3, p. 507 (1849); Ins. Brit. Dipt. Vol. 1, p. 133 (1851); Neuhaus, Dipt. March. p. 75 (1886); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 413 (1907); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 320, f. 141 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 87 [1913] (*Tachydromia*). Europe.
- pubicornis*, Zetterstedt, Ins. Lappon. p. 553 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 304 [1842] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 151 [1861] (*Tachydromia*); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 137 [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, p. 210 [1898] (*Tachydromia*); Wien. Ent. Zeitschr. Vol. 18, p. 78 [1899] (*Tachydromia*); Jahrb. Kärnten. Vol. 47, p. 203 (1901); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 79 (1909).
148. *P. longimanus*, Strobl, ibidem, Vol. 46, p. 79 (1909). C. Europe.
149. *P. longirostris*, Bezz, Ann. Mus. Hungar. Vol. 10, p. 491 [1912] (Coryneta). Formosa.
var. *xanthopus*, Bezz, Suppl. Ent. Berlin, Vol. 3, p. 78 [1914] (Coryneta). Formosa,
150. *P. luctator*, nov. sp. Idaho.
151. *P. lupatus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 340 (1902). New Mexico.
152. *P. luteicornis*, Meigen, Syst. Beschr. Vol. 7, p. 97 [1838] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 91 (1862). C. Europe.
153. *P. luteipalpis*, Macquart, Hist. Nat. Dipt. Vol. 1, p. 579, Errata (1834). France.
flavipalpus, Macquart (not Meigen), Dipt. N. France, Vol. 3; p. 96 (1827); Hist. Nat. Vol. 1, p. 353 (1834); Boitard, Man. Ent. Vol. 3, p. 322 (1843).
fuscipennis, Macquart, Hist. Nat. Dipt. Vol. 2, p. 659 (1835).

154. *P. luteus*, Meigen, Klassif. Vol. 1, p. 238 [1804] (*Tachydromia*); Fallen, Europe.

Empid. p. 10 [1815] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 89 [1822] (*Tachydromia*); Macquart, Dipt. N. France, Vol. 3, p. 100 (1827); Hist. Nat. Dipt. Vol. 1, p. 356 (1834); Zetterstedt, Fauna Ins. Lappon. p. 552 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 294 part [1842] (*Tachydromia*); Walker, List, Vol. 3, p. 510 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3007 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 129 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 150 [1861] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 91 (1862); Neuhaus, Dipt. March. p. 75 (1885); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 108 [1892] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 411 (1907); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 8 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 312 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94, f. 19 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 81 [1913] (*Tachydromia*).

glaberr. Zetterstedt (not Meigen), Dipt. Scand. Vol. 1, p. 295 [1842] (*Tachydromia*); ibidem, Vol. 8, p. 3007 [1849] (*Tachydromia*).

pallidus, Meigen, Syst. Beschr. Vol. 3, p. 90 [1822] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5, p. 59 [1851] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 91 (1862).

155. *P. lyristes*, nov. sp.

Arizona.

156. *P. macropalpus*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 80 [1899] (*Tachydromia*).

Spain.

157. *P. maculifemoratus*, nov. sp. (1).

Java.

158. *P. maculifemur*, Meijere, Tijdschr. v. Ent. Vol. 56, Suppl. p. 78 [1914] (*Tachydromia*).

Java.

159. *P. maculipes*, Meigen, Syst. Beschr. Vol. 3, p. 79 [1822] (*Tachydromia*);

Europe.

Zetterstedt, Dipt. Scand. Vol. 1, p. 274 [1842] (*Tachydromia*); Walker, List, Vol. 3, p. 508 (1849); Ins. Brit. Dipt. Vol. 1, p. 122 [1851] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 12, p. 4600 [1855] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 148 [1861] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 87 (1862); Neuhaus, Dipt. March. p. 74 (1886); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 117 [1892] (*Tachydromia*); Mem. Soc. Esp. Vol. 3, p. 313 [1906] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 409 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Verh.

(1) *Platypalpus maculifemoratus*, nov. sp. — Length 2 mm. Black, thorax dusky, legs including the coxae yellow, tarsi not annulate, middle femora marked with a black spot at the outer two-thirds of the anterior face, tibial spur sharp and yellow, crossveins touching. Head rather densely golden pollinose, front short and broad, face much widened below and silvery pruinose, palpi small and yellow, proboscis two-thirds as long as the head, reddish in front, otherwise black, antennae black, the third joint defective in the type specimens, vertical bristles strong and black. Thorax moderately covered with yellowish pollen, slightly shining, its bristles black, pleurae whitish pruinose, sternum and hypopleurae largely polished. Abdomen shining; the pygidium moderately small, globular, closed, not hairy; styles of the ovipositor slender, yellowish. Front tibiae as thick as the others, with three extensor black setae, middle femora moderately thickened, the pale flexor setae long on both sides, patella yellow, middle tibiae two-thirds as long as their femora, the spur as long as the diameter of the tibia, tarsi becoming dark apically. Halteres yellow. Wings hyaline, veins pale brown, third vein slightly converging toward the fourth near the tip, crossveins forming a continuous line, located at two-fifths the length of the wing, second, third and fourth sections of the costa proportioned 1 : 0.7 : 0.2, sections of the fifth vein, 0.3 : 0.6 : 1, fringe of the hind margin as long as the anterior crossvein.

Two specimens, Tjibodas, Java, belonging to the U. S. National Museum.

- Zool.-bot. Ges. Wien, Vol. 59, p. 180 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 291, f. 133 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 79 [1913] (*Tachydromia*).
nigrosetosus, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 117 [1892] (*Tachydromia*); Vol. 34, p. 211 [1898] (*Tachydromia*); Vol. 46, p. 83 [1909] (*Tachydromia*).
160. *P. maculus*, Zetterstedt, Dipt. Scand. Vol. 1, p. 289 [1842] (*Tachydromia*); C. & N. Europe.
Wahlberg, Ofvers. K. Vet. Akad. Förhandl. Vol. 2, p. 254 [1845] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 8, p. 3006 [1849] (*Tachydromia*); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 134 [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 80 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 315 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 83; f. 25 [1913] (*Tachydromia*); Collin, Ent. M. Mag. (2), Vol. 24, p. 130 [1913] (*Tachydromia*).
flavipalpis, Zetterstedt (not Meigen), Ins. Lappon. p. 551, var. [1838] (*Tachydromia*).
161. *P. major*, Zetterstedt, Dipt. Scand. Vol. 1, p. 287 [1842] (*Tachydromia*); C. & N. Europe.
Vol. 7, p. 3006 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 88 (1862); Jaroszewski, Trudy Khark. Vol. 11, p. 352 (1877); Beling, Verh. Zool.-bot. Ges. Wien, Vol. 38, p. 2 (1888); Strobl, Verh. Mitteil. Siebenbürg. Ver. Hermannst. Vol. 46, p. 23 [1896] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 408 [1907] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 81 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 306, f. 140 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 77 [1913] (*Tachydromia*).
cursitans, Meigen (not Fabricius), Klassif. Vol. 1, p. 237 [1804] (*Tachydromia*); Syst. Beschr. Vol. 3, p. 83, pl. 23, f. 23 [1822] (*Tachydromia*).
162. *P. Meigenianus*, Gimmerthal, Bull. Soc. Imp. Nat. Moscou, Vol. 7, p. 117 [1834] (*Tachydromia Meigenia*); Vol. 15, p. 667 [1842] (*Tachydromia Meigenia*); Beazzi, Katal. pal. Dipt. Vol. 2, p. 285 (1903) correction. Russia.
163. *P. melanogaster*, nov. sp. Labrador.
164. *P. melleus*, nov. sp. New York.
165. *P. mesogrammus*, Loew, Berl. Ent. Zeitschr. Vol. 7, p. 21 (1863); Cent. 3, No. 38; Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 219 (1902). E. United States.
166. *P. Mikii*, Becker, Wien. Ent. Zeit. Vol. 9, p. 67 [1890] (*Tachydromia*); C. Europe.
Strobl, Verh. Mitteil. Siebenbürg. Hermannst. Vol. 46, p. 23 [1896] (*Tachydromia*).
167. *P. mimus*, nov. sp. New York.
168. *P. minutissima*, Strobl, Wien. Ent. Zeitschr. Vol. 18, p. 82 [1899] (*Tachydromia*); Mém. Soc. Esp. Hist. Nat. Vol. 3, p. 317 [1906] (*Tachydromia*). Spain.
169. *P. minutus*, Meigen, Klassif. Vol. 1, p. 238 [1804] (*Tachydromia*); Syst. Beschr. Vol. 3, p. 76 [1822] (*Tachydromia*); Macquart, Dipt. N. France, Vol. 3, p. 95 (1827); Hist. Nat. Vol. 1, p. 353 [1834]; Zetterstedt, Dipt. Scand. Vol. 1, p. 303 [1842] (*Tachydromia*); Europe.

Boitard, Man. Ent. Vol. 3, p. 322 (1843); Walker, List, Vol. 3, p. 508 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 309 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 131 (1851); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 151 [1861] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 89 (1862); Siebke, Nyt Mag. Naturvid. Vol. 14, p. 379 (1866); Strobl, Progr. Seitenstetten, Vol. 14, p. 7 (1880); Neuhaus, Dipt. March. p. 75 (1886); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 134 [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 118 [1892] (*Tachydromia*); Wien. Ent. Zeit. Vol. 18, p. 82 [1899] (*Tachydromia*); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 309 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*); Frey, Acta Sc. Fenn. Helsingfors, Vol. 37 (3), p. 80 [1913] (*Tachydromia*).

annulatus, Fallen, Dipt. Suec. Empid. 7, part [1815] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 77 [1822] (*Tachydromia*); Macquart, Dipt. N. France, Vol. 3, p. 96 (1827); Hist. Nat. Vol. 1, p. 353 [1834]; Zetterstedt, Ins. Lappon. p. 552, part [1838] (*Tachydromia*); Loew, Progr. Posen, p. 22 (1840); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 322 (1843); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 151 [1861] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 411 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 80 [1913] (*Tachydromia*).

femoralis, Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 150 [1861] (*Tachydromia*).

var. *obscuripes*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 82 [1899] (*Tachydromia*); Verh. Ges. Wien, Vol. 59, p. 181 [1909] (*Tachydromia*). Spain.

170. *P. montanus*, Becker, Berl. Ent. Zeitschr. Vol. 31, p. 138 [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 119 [1892] (*Tachydromia*); Bezzì, Bull. Soc. Ent. Ital. Vol. 30, p. 158 [1899] (*Tachydromia*); Strobl, Wien. Ent. Zeit. Vol. 18, p. 82 [1899] (*Tachydromia*); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 84 [1909] (*Tachydromia*). Europe.
171. *P. monticola*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 217, f. 41 (1902). Colorado.
172. *P. multisetosus*, Bezzì, Bull. Soc. Ent. Ital. Vol. 30, p. 155 [1899] (*Tachydromia*). Italy.
173. *P. mundus*, Walker, Ent. Mag. Vol. 4, p. 228 (1837); Ins. Brit. Dipt. Vol. 1, p. 129 (1851). England.
174. *P. nepalensis*, Brunetti, Fauna Brit. Indian Dipt. Brachyc. Vol. 1, p. 377 [1920] (*Tachydromia*). India.
- longicornis*, Brunetti (not Meigen), Rec. India Mus. Vol. 9, p. 22 [1913] (*Brevios*).
175. *P. niger*, Macquart, Mém. Soc. Sc. Lille, p. 154 [1823] (*Tachydromia*); Boitard, Man. Ent. Vol. 3, p. 322 (1843). France.
176. *P. niger*, Meigen, Klassif. Vol. 1, p. 238 [1804] (*Tachydromia*); Syst. Beschr. Vol. 3, p. 75 [1822] (*Tachydromia*); Macquart, Dipt. Nord France, Vol. 3, p. 94 (1827); Hist. Nat. Vol. 1, p. 352 (1834); Loew, Progr. Posen, p. 22 (1840); Isis, Vol. 7, p. 551 [1840] (*Tachydromia*); Walker, List. Vol. 3, p. 508 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Neuhaus, Dipt. March. p. 75 (1886). C. Europe.
177. *P. nigricoxus*, Mik, Wien. Ent. Zeit. Vol. 3, p. 82 [1884] (*Tachydromia*); Becker, Fauna Hernstein, Vol. 2, p. 60, f. 7, 8 [1885] (*Tachydromia*); C. Europe.

- Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*).
 178. *P. nigrifemoratus*, Macquart, Hist. Nat. Vol. 1, p. 354 (1834); Boitard, S. Europe.
 Man. Ent. Vol. 3, p. 324 (1843); Bezzi, Bull. Soc. Ent. Ital. Vol. 24,
 p. 116 (1892).
 179. *P. nigrimanus*, Roser, Correspondenzbl. Landw. Würtemb. Vol. 1, p. 54 C. Europe.
 [1840] (*Tachydromia*).
 180. *P. nigrimanus*, Strobl, Progr. Seitenstetten, Vol. 14, p. 8 (1880); Mik, S. & C. Europe.
 Verh. Ges. Wien. Vol. 31, p. 348 [1881] (*Tachydromia*); Strobl,
 Glasnik Zem. Mus. Bosn. Herceg. Vol. 10, p. 401 [1898] (*Tachydromia*);
 Mitteil. Mus. Bosn. Herceg. Vol. 7, p. 563 [1900] (*Tachydromia*);
 Jahrb. Mus. Kärnten, Vol. 47, p. 204 (1901); Mitteil. Naturw. Ver.
 Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*).
 181. *P. nigrinus*, Meigen, Syst. Beschr. Vol. 3, p. 76 [1822] (*Tachydromia*); C. & N. Europe.
 Walker, List, Vol. 3, p. 508 (1849); Becker, Berl. Ent. Zeitschr.
 Vol. 31, p. 138, var. [1887] (*Tachydromia*); Strobl, Mitteil. Naturw.
 Ver. Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*).
 182. *P. nigrinus*, Schiner, Fauna Austr. Dipt. Vol. 1, p. 89 (1862); Strobl, C. Europe.
 Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*).
 183. *P. nigripalpis*, Bigot, Ann. Soc. Ent. France (6), Vol. 1, p. 365 (1881). Alps.
 184. *P. nigripes*, Meigen, Syst. Beschr. Vol. 6, p. 343 [1830] (*Tachydromia*). C. Europe.
 185. *P. nigritarsis*, Fallen, Empid. p. 34 [1816] (*Tachydromia*); Meigen,
 Syst. Beschr. Vol. 3, p. 74 [1822] (*Tachydromia*); Zetterstedt, Fauna
 Ins. Lappon. p. 553 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1,
 p. 307 [1842] (*Tachydromia*); Walker, List, Vol. 3, p. 507 (1849);
 Zetterstedt, Dipt. Scand. Vol. 8, p. 3010 [1849] (*Tachydromia*); Scholz,
 Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*);
 Walker, Ins. Brit. Dipt. Vol. 1, p. 133 (1851); Bonsdorff, Finl. tvåv.
 Ins. Dipt. Vol. 1, p. 152 [1861] (*Tachydromia*); Schiner, Fauna
 Austr. Dipt. Vol. 1, p. 91 (1862); Strobl, Mitteil. Naturw. Ver.
 Steiermark, Graz, Vol. 29, p. 110 [1892] (*Tachydromia*); Wien. Ent.
 Zeit. Vol. 18, p. 78 [1899] (*Tachydromia*); Mem. Soc. Esp. Hist.
 Nat. Vol. 3, p. 313 [1906] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt.
 Vol. 7, p. 413 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors,
 Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver.
 Steiermark, Graz, Vol. 46, p. 79 [1909] (*Tachydromia*); Lundbeck,
 Dipt. Danica, Vol. 3, p. 319 [1910] (*Tachydromia*); Wahlgren, Ent.
 Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc.
 Fenn. Helsingfors, Vol. 37 (3), p. 87 [1913] (*Tachydromia*).
 186. *P. nigromaculatus*, Roser, Correspondenzbl. Landw. Würtemb. Vol. 1, C. Europe.
 p. 54 [1840] (*Tachydromia*).
 187. *P. nitidipleura*, nov. sp. Montana.
 188. *P. nitidus*, Macquart, Dipt. N. France, Vol. 3, p. 95 (1827); Hist. Nat.
 Vol. 1, p. 352 (1834); Meigen, Syst. Beschr. Vol. 7, p. 97 [1838]
 (*Tachydromia*). France.
 189. *P. niveiseta*, Zetterstedt, Dipt. Scand. Vol. 1, p. 311 [1842] (*Tachydromia*); N. Europe.
 Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*).
 190. *P. Novakii*, Strobl, Wien. Ent. Zeit. Vol. 12, p. 37 [1893] (*Tachydromia*). S. Europe.
 — *P. ochraceus*, Becker, in Bezzi, Atti Soc. Veneto-Trent. (2), Vol. 1, p. 57
 [1893] (*Tachydromia*), no description.
 191. *P. ochricollis*, nov. sp. Washington.
 192. *P. oculatus*, nov. sp. Pennsylvania.

193. *P. aedicnemus*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, C. & S. Europe.
p. 211 [1898] (*Tachydromia*); Jarb. Mus. Kärnten, Vol. 47, p. 204
(1901).
flaviventris, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, p. 211
[1898] (*Tachydromia*).
varius, Bezzi (not Walker), Bull. Soc. Ent. Ital. Vol. 24, p. 116 (1892).
194. *P. oriens*, nov. sp. (1). Java.
195. *P. orientalis*, Brunetti, Rec. Indian Mus. Vol. 9, p. 39 (1913); Fauna India.
Brit. India Dipt. Vol. 1, p. 378 (1920).
196. *P. ostiorum*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 44 [1902] N. Africa.
(*Tachydromia*); Zeitschr. Hym. Dipt. Vol. 7, p. 115 [1907] (*Tachydromia osteriorum*).
197. *P. pachycnemus*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 89 : Cent. 5, E. United States.
No. 77 (1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438
(1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 217 (1902).
198. *P. pallidiventris*, Meigen, Syst. Beschr. Vol. 3, p. 82 [1822] (*Tachydromia*); Europe, N. Africa.
Macquart, Dipt. N. France, Vol. 3, p. 98 (1827); Hist. Nat. Dipt. Vol. 1, p. 355 (1834); Zetterstedt, Ins. Lappon. p. 549 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 277 [1842] (*Tachydromia*); Walker, Ins. Dipt. Vol. 1, p. 123 (1851); Zetterstedt, Dipt. Scand. Vol. 13, p. 4987 [1859] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 88 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 138, part [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 113 [1892] (*Tachydromia*); Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 117 [1907] (*Tachydromia*); Frey, ibidem, Vol. 7, p. 408 [1907] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 81, 83 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 301, f. 136 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 93 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 77 [1913] (*Tachydromia*).
flavipes, Fallen, Empid. p. 6, part [1815] (*Tachydromia*).
longiseta, Zetterstedt, Dipt. Scand. Vol. 1, p. 278 [1842] (*Tachydromia*); ibidem, Vol. 13, p. 4987 [1859] (*Tachydromia*); Strobl, Jahrb. Kärnten, Vol. 47, p. 204 (1901); Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 83 [1909] (*Tachydromia*).
199. *P. pallipes*, Fallen, Empid. p. 8 [1815] (*Tachydromia*); Zetterstedt, Fauna Europe.
Ins. Lappon. p. 553 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 305 [1842] (*Tachydromia*); Walker, List, Vol. 3, p. 507 (1849); Zetterstedt, Dipt. Scand. Vol. 8, p. 3009 [1849] (*Tachydromia*); Schiner, Fauna Austr. Dipt. Vol. 1, p. 90 (1862); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 412 [1907] (*Tachydromia*); Medd. Soc. Fenn.

(1) *Platypalpus oriens*, nov. sp. — Length 2.6 mm. Very close to *albisetata* Panz. Body black, mesonotum polished except narrowly along the sides; pleurae uniformly pollinose except largely on the sternopleurae. Palpi yellow, base of the proboscis reddish. Antennae elongate, the white arista longer than the antennae. Pygidium globular, more rotund than in *albisetata* but terminating in a sharp claw, the dorsal valve hairy and with a blunt backward prolongation. Legs yellow, the front femora with nearly uniform outstanding yellow hairs, front tibiae without basal thickening. Wings nearly hyaline, neuration as in *albisetata*.

Several specimens, Tjibodas, Mt. Gede, 4,500 to 9,000 feet altitude, Java, from the Bryant and Palmer collection in the U. S. National Museum. The entirely pruinose mesopleurae, lack of prominent central bristles of the front femora, the yellow palpi and blunt process of the dorsal valve are distinguishing characters, which differ from those of *albisetata*. In that species the dorsal valve is not hairy and is projected backward as a long acuminate process. *P. subulifer* Meijere is also closely related but has the arista equal to the antenna, the base of the front tibia swollen, spindle-like, the proboscis black and the legs brown.

- Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 318 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 83, f. 24 [1913] (*Tachydromia*).
- flavipalpis*, Meigen, Syst. Beschr. Vol. 3, p. 74 [1822] (*Tachydromia*); Macquart, Hist. Nat. Vol. 1, p. 352 (1834); Zetterstedt, Ins. Lappon. p. 551, part [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, p. 288 [1842] (*Tachydromia*); ibidem, Vol. 8, p. 3006 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Zetterstedt, Dipt. Scand. Vol. 13, p. 4990 [1856] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 150 [1861] (*Tachydromia*); Schiner, Fauna Austr. Vol. 1, p. 90 (1862); Strobl, Mitteil. Bosn. Herceg. Vol. 7, p. 563 [1900] (*Tachydromia*); Dahl, Fauna Chorin, p. 466 (1912); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 81 [1909] (*Tachydromia*).
200. *P. paludosus*, Perris, Ann. Soc. Linn. Lyon, Vol. 5, p. 200 [1852] (*Tachydromia*). France.
201. *P. parvicornis*, Zetterstedt, Dipt. Scand. Vol. 13, p. 4992 [1859] (*Tachydromia*). N. Europe.
202. *P. Paulseni*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 768 (1865); Bezzi, Ann. Mus. Hungar, Vol. 3, p. 459 [1905] (*Tachydromia*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 402 [1909] (*Tachydromia*).
203. *P. pectinator*, Melander, Occ. Pap. Boston Soc. Nat. Hist. Vol. 5, p. 85 (1924). North America.
204. *P. pectoralis*, Fallen, Empid. Vol. 9 [1815] (*Tachydromia*); Meigen, Syst. Beschr. Vol. 3, p. 87 [1822] (*Tachydromia*); Macquart, Hist. Nat. Vol. 1 p. 356 (1834); Zetterstedt, Dipt. Scand. Vol. 1, p. 295 [1842] (*Tachydromia*); Walker, List, Vol. 3, p. 510 (1849); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 129 (1851); Loew, Zeitschr. Ges. Naturwiss. Vol. 10, p. 101 (1857); Schiner, Fauna Dipt. Austr. Vol. 1, p. 91 (1862); Verrall, Ent. Vol. 23, p. 153 [1890] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 108 [1892] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 411 [1907] (*Tachydromia*); Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 8 [1908] (*Tachydromia*); Medd. Soc. Sc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 77 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 314 [1910] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 81 [1913] (*Tachydromia*). C. & N. Europe.
- var. *nonstriatus*, Strobl, Jahrb. Mus. Kärnten, Vol. 47, p. 203 (1901). C. Europe.
- var. *stramineipes*, Zetterstedt, Dipt. Scand. Vol. 1, p. 296 [1842] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Walker, Ins. Brit. Dipt. Vol. 1, p. 130 [1851] (*stramineipes*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 7, 8 [1908] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 81 [1913] (*Tachydromia*). C. & N. Europe.
205. *P. pedestris*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 118 [1907] (*Tachydromia*). Algeria.
206. *P. picipes*, Zetterstedt, Dipt. Scand. Vol. 1, p. 298 [1842] (*Tachydromia*). N. Europe.
207. *P. pictipennis*, Bezz, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 402, f. 19 [1909] (*Tachydromia*). Peru, Costa Rica.
208. *P. pictitarsis*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 2, p. 44 [1902] (*Tachydromia*). Egypt.
209. *P. pilatus*, nov. sp. Washington.

210. *P. pluto*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 217, f. 39 [1902]; Pacific United States.
Bezzi, Ann. Mus. Hungar. Vol. 3, p. 456, 459 [1905] (*Tachydromia*);
Nova Acta Akad. Naturf. Halle, Vol. 91, p. 402 [1909] (*Tachydromia*).
211. *P. podocarpi*, Becker, Ann. Soc. Ent. France, Vol. 88, p. 123 [1914] E. Africa.
(*Coryneta*).
212. *P. politus*, nov. sp. California.
var. *nitens*, nov. var. United States.
213. *P. pollinosus*, Strobl, Glasnik. Zem. Mus. Bosn. Herceg. Vol. 10, p. 400 S. Europe.
[1898] (*Tachydromia*); Mitteil. Mus. Bosn. Herceg. Vol. 7, p. 562 [1900] (*Tachydromia*).
fallidus, Strobl (not Meigen), Wien. Ent. Zeit. Vol. 12, p. 38 [1893] (*Tachydromia*).
214. *P. Popplesi*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (10), p. 11 Finland.
[1913] (*Tachydromia*).
215. *P. porrectus*, Melander, Occ. Pap. Bost. Soc. Nat. Hist. Vol. 5, p. 86 (1924). W. United States.
var. *suffasciatus*, nov. var. W. United States.
216. *P. postpositus*, nov. sp. Washington.
217. *P. *? predatoris*, Meunier, Ann. Sc. Nat. (Zool.) Vol. 7, p. 89, 101, pl. 5, Baltic Amber.
f. 8-11 (1908).
218. *P. prorsus*, nov. sp. Labrador.
219. *P. proserpina*, Bezzi, Nova Acta Akad. Naturf. Halle, Vol. 91, p. 405 Peru.
[1909] (*Tachydromia*).
220. *P. pseudobicolor*, Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, C. Europe.
p. 82 [1909] (*Tachydromia*).
221. *P. pseudociliaris*, Strobl, ibidem, Vol. 46, p. 80 [1909] (*Tachydromia*). C. Europe.
222. *P. pseudoexiguus*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 181 Spain.
[1909] (*Tachydromia*).
223. *P. pseudomaculipes*, Strobl, Wien. Ent. Zeit. Vol. 18, p. 80 [1899] (*Tachydromia*). Spain.
224. *P. pseudounguiculatus*, Strobl, Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 181 Spain.
[1909] (*Tachydromia*).
225. *P. pubescens*, nov. sp. Washington.
226. *P. pudens*, nov. sp. W. United States.
227. *P. pulchellus*, Walker, Ins. Brit. Dipt. Vol. 1, p. 130 (1851). England.
228. *P. pulicarius*, Meigen, Syst. Beschr. Vol. 6, p. 343 [1830] (*Tachydromia*). C. Europe.
229. *P. pulverulentus*, nov. sp. California.
230. *P. pygmaeus*, Macquart, Mém. Soc. Sc. Lille, p. 154 [1823] (*Tachydromia*). France.
231. *P. pygmaeus*, Meigen (not Macquart), Syst. Beschr. Vol. 7, p. 99 [1838] (*Tachydromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 92 (1862). C. Europe.
232. *P. rapidus*, Meigen, Syst. Beschr. Vol. 3, p. 81 [1822] (*Tachydromia*). C. Europe.
233. *P. recurvus*, nov. sp. California.
234. *P. robustus*, Walker, Ent. Mag. Vol. 4, p. 228 (1837); Ins. Brit. Dipt. Vol. 1, p. 128 (1851). England.
235. *P. rondonensis*, Strobl, Mém. Soc. Esp. Hist. Nat. Vol. 3, p. 316 [1906] Spain.
(*Tachydromia*).
236. *P. rubefactus*, nov. sp. Illinois, Maryland.
237. *P. ruficornis*, Roser, Correspondenzbl. Land. Würtemb. Vol. 1, p. 54 C. Europe.
[1840] (*Tachydromia*).
238. *P. rufipes*, Meigen, Syst. Beschr. Vol. 7, p. 99 [1838] (*Tachydromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 89 (1862). C. Europe.
239. *P. rufiventris*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 341 (1902). New Mexico.
240. *P. sahlbergi*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 31 (9), p. 7, 8 Finland, Sweden.
(1908); ibidem, Vol. 37 (3), p. 81, f. 22 (1913); Ent. Tidskr. 1914,
p. 80 (1914).
- straminipes*, Strobl (not Zetterstedt), Mitteil. Naturw. Ver. Steiermark, Graz,

- Vol. 29, p. 109 [1892] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 412 [1907] (*Tachydromia stramineipes*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 77 (1909).
var. *nigricollis*, Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 82 [1913] Finland, Sweden.
(*Tachydromia*).
241. *P. sanguineus*, Arribalzaga, Natural. Argent. Vol. 1, p. 294 [1878] (*Tachydromia*); Bezzi, Ann. Mus. Hungar, Vol. 3, p. 459 [1905] (*Tachydromia*); Nova Acta Akad. Naturf. Halle, Vol. 91, p. 402 (1909). Argentina.
242. *P. sanguinolentus*, new name. Java.
sanguineus, Meijere (not Arribalzaga), Tijd. Ent. Vol. 56, Suppl. p. 76 [1914]
(*Tachydromia*).
243. *P. satyriacus*, nov. sp. Labrador.
244. *P. semihyalipennis*, Grimmerthal, Bull. Soc. Imp. Moscou, Vol. 20 (2), p. 166 [1847] (*Tachydromia*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 92 [1862] (*semihyalinus*). Russia.
245. *P. sericatus*, nov. sp. British Columbia.
246. *P. serviceus*, Macquart, Mém. Soc. Sc. Lille, p. 153 [1823] (*Tachydromia*). France.
247. *P. simplicipes*, nov. sp. Washington.
248. *P. soccatus*, nov. sp. Idaho.
249. *P. sordidus*, Zetterstedt, Ins. Lappon. p. 552 [1838] (*Tachydromia*); Dipt. Scand. Vol. 1, 300 [1842] (*Tachydromia*); ibidem, Vol. 8, p. 3007 [1849] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 120 [1892] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 410 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 94 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 79 [1913] (*Tachydromia*). C. & N. Europe.
250. *P. spinosus*, nov. sp. California.
251. *P. splendens*, new name. Colorado.
montanus, Melander (not Becker), Trans. Amer. Soc. Vol. 28, p. 213, f. 7, 8 [1902] (*Elaphropeza*); Ent. News, Vol. 17, p. 372 [1906] (*Symbalophthalmus*).
252. *P. stigmatellus*, Zetterstedt, Dipt. Scand. Vol. 1, p. 306 [1842] (*Tachydromia*); ibidem, Vol. 8, p. 3010 [1849] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19) p. 58 [1851] (*Tachydromia*); Bonsdorff, Finl. tvåv. Ins. Dipt. Vol. 1, p. 151 [1861] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 110 [1892] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 412 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 79 [1909] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 87, f. 29 [1913] (*Tachydromia*). C. & N. Europe.
pallipes, Zetterstedt, part, Ins. Lappon. p. 553, part [1838] (*Tachydromia*).
253. *P. strigifrons*, Zetterstedt, Dipt. Scand. Vol. 8, p. 3005 [1849] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 293, f. 134 [1910] (*Tachydromia*); Becker, Ann. Mus. Zool. St. Petersbourg, Vol. 17, p. 597 [1913] (*Tachydromia*); Frey, Acta Soc. Sc. Helsingfors, Vol. 37 (3), p. 79 [1913] (*Tachydromia*). N. Europe, Persia.
254. *P. Stroblianus*, new name. C. Europe.
diversipes, Strobl (not Coquillett) Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 84 [1909] (*Tachydromia*).
255. *P. subulifer*, Meijere, Tijd. Ent. Vol. 56, Suppl. p. 78 [1914] (*Tachydromia*). Java.
256. *P. sutor*, Melander. Occ. Pap. Boston, Soc. Nat. Hist. Vol. 5, p. 87 (1924). North America.

257. *P. tachistiformis*, nov. sp. Montana.
258. *P. tenuatus*, Meigen, Syst. Beschr. Vol. 3, p. 88 [1822] (*Tachydromia*); C. Europe.
Strobl. Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 78 [1909]
(*Tachydromia*).
259. *P. talaris*, nov. sp. North America.
260. *P. tectifrons*, Becker, Ann. Mus. Zool. St. Petersbourg, Vol. 12, p. 314 Tibet.
[1907] (*Tachydromia*).
261. *P. tenax*, nov. sp. Washington.
262. *P. tenellus*, Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 223, f. 28, 30 United States.
(1902).
263. *P. teneriffensis*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 41 [1908] Canary Islands.
(*Tachydromia*).
264. *P. tenuis*, nov. sp. Washington.
265. *P. tergestinus*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 345 (1860). S. Europe.
266. *P. tersus*, Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 439 (1895); E. United States.
Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 219 (1902).
267. *P. testaceus*, Philippi, Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 767 (1865); Chile.
Bezzi, Ann. Mus. Hungar. Vol. 3, p. 459 [1905] (*Tachydromia*);
Nova Acta Akad. Naturf. Halle, Vol. 91, p. 402 [1909] (*Tachydromia*).
268. *P. thoracicus*, Lundbeck, Dipt. Danica, Vol. 3, p. 317 [1910] (*Tachydromia*); Denmark.
Collin, Ent. M. Mag. London (2), Vol. 24, p. 130 [1913]
(*Tachydromia*).
—. *P. tibialis*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 265 [1829] (*Tachydromia*), no description.
269. *P. trivialis*, Loew, Berl. Ent. Zeitschr. Vol. 8, p. 88 : Cent. 5, No. 76 E. United States.
(1864); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1896);
Proc. Ent. Soc. Wash. Vol. 5, p. 265 (1903).
270. *P. turgidus*, Becker, Zeitschr. Hym. Dipt. Vol. 7, p. 118 [1907] (*Tachydromia*). Algeria.
271. *P. uncinatus*, nov. sp. Costa Rica.
272. *P. unguiculatus*, Zetterstedt, Ins. Lappon. p. 551 [1838] (*Tachydromia*); N. & C. Europe.
Dipt. Scand. Vol. 1, p. 291 [1842] (*Tachydromia*); Scholz, Zeitschr. Ent. Breslau, Vol. 5 (19), p. 58 [1851] (*Tachydromia*); Schiner, Fauna Dipt Austr. Vol. 1, p. 90 (1862); Strobl, Progr. Seitenstetten, Vol. 14, p. 8 (1880); Becker, Berl. Ent. Zeitschrift, Vol. 31, p. 134 [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 111 [1892] (*Tachydromia*); Frey, Zeitschr. Hym. Dipt. Vol. 7, p. 412 [1907] (*Tachydromia*); Medd. Soc. Fenn. Helsingfors, Vol. 34, p. 21 [1908] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 80 [1909] (*Tachydromia*); Wahlgren, Ent. Tidskr. Vol. 31, p. 95 [1910] (*Tachydromia*); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 82, f. 23 [1913]
(*Tachydromia*).
273. *P. univittatus*, Loew, Oefv. Kongl. Vet. Akad. Förhandl. Vol. 14, p. 371 S. Africa.
(1857); Dipterent. Südafrik. Vol. 1, p. 269 (1860); Adams, Kansas.
Univ. Sc. Bul. Vol. 3, p. 158 (1905).
274. *P. valens*, nov. sp. (1). Java.

(1) *Platypalpus valens*, nov. sp. — Female. Length 4 mm. Robust, head, thorax and abdomen black, legs yellow, tarsi not annulate, tibial spur very large, antennæ short, posterior crossvein oblique. Front long and narrow, scarcely wider above, face linear; palpi white without, yellowish within, with a few strong white setæ, proboscis one-half the height of the head, black; antennæ three-jointed, the basal joint minute, the third joint ovate, scarcely longer than wide, the arista thin, very sparsely hairy and nearly three times as long as the antennæ. Mesonotum with sparse golden dust, the bristles black, upper pleurae entirely pruinose, fringe of calypteres strong and

275. *P. valgus*, nov. sp. Washington.
 276. *P. varicolor*, Becker, Mitteil. Zool. Mus. Berlin, Vol. 4, p. 39 [1908] Canary Islands.
(Tachydromia).
 277. *P. varipes*, Meigen, Syst. Beschr. Vol. 3, p. 88 [1822] (*Tachydromia*). C. Europe.
 278. *P. varius*, Walker, Ins. Brit. Dipt. Vol. 1, p. 126 (1851); Schiner, Fauna Dipt. Austr. Vol. 1, p. 88 (1862); Becker, Berl. Ent. Zeitschr. Vol. 31, p. 134 [1887] (*Tachydromia*); Strobl, Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 46, p. 81 [1909] (*Tachydromia*); Lundbeck, Dipt. Danica, Vol. 3, p. 302 [1910] (*Tachydromia*). C. Europe.
 279. *P. velox*, nov. sp. Washington.
 280. *P. venaticus*, nov. sp. W. North America.
 —. *P. venosus*, Stephens, Syst. Cat. Brit. Ins. Vol. 2, p. 265 [1829] (*Tachydromia*), no description.
 281. *P. verpus*, nov. sp. Washington.
 282. *P. versipes*, nov. sp. Virginia.
 283. *P. versutus*, Melander, Occ. Pap. Bost. Soc. Nat. Hist. Vol. 5, p. 87 (1924). E. United States.
 284. *P. vicarius*, Walker, Trans. Ent. Soc. London, n. s., Vol. 4, p. 148 (1857); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 438 (1895); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 220 (1902). North America.
 285. *P. Vierecki*, Melander, ibidem, Vol. 28, p. 340 (1902). New Mexico.
 286. *P. Villeneuvei*, Becker, Deut. Ent. Zeitschr. p. 647 (1910). Corsica.
 —. *P. vitripennis*, Becker, Ann. Mus. Zool. St. Petersbourg, Vol. 17, p. 597 [1913] (*Tachydromia*), attributed to Meigen, no description. Persia.
 287. *P. vittatus*, nov. sp. Wyoming.
 var. *perimerus*, nov. var. Alberta.
 288. *P. vulnificus*, nov. sp. Montana.
 289. *P. xanthochiton*, nov. sp. Washington.
 290. *P. xanthopodus*, new name. North America.
gilvipes, Coquillett (not Meigen), Proc. Wash. Acad. Sc. Vol. 2, p. 422 (1900); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 224 (1902).

EXTINCT EMPIDIDÆ : FOSSIL SPECIES AND GENERA

All the described fossil Empididæ come from the Tertiary. Some more recent species of *Empis* and *Rhamphomyia* are known from copal, but have not been described. Two pre-Tertiary species have been assigned to the Empididæ, namely *Empidia Wulpii* and *Hasmona Leo*, but it is not at all certain that they belong to this family. The Baltic amber has produced a great number of specimens of Empididæ; Meunier called attention to fifteen hundred inclusions that he had studied; but the descriptions and figures too often miss the salient characters to be of much value in deciphering phylogeny. It may be stated that the Tertiary Empididæ almost all belong to genera of the present day holarctic fauna. Such genera as are extinct present no startling characters, and could very well produce living species without causing astonishment. The following notes show the status of the extinct genera, which have been assigned to this family.

yellowish. Abdomen shining, the last two small segments opaque yellow, styles of the ovipositor long and narrow. Legs robust, front femora thickened, with two flexor rows of pale short, sparse hairs, front tibiæ outwardly ciliate on the apical half with black setæ and closely biserately black-setulose beneath, patella and tibial spur black, middle tibiæ two-thirds as long as their femora, tarsi uniformly blackish apically. Halteres yellow. Wings hyaline, veins firm and black, first vein ending at the middle of the wing, second, third and fourth sections of the costa proportioned 1 : 0.6 : 0.12, third and fourth veins converging apically, posterior crossvein located beyond the anterior a distance equal to the anterior, sections of the fifth vein proportioned 0.8 : 1 : 1.2, marginal cilia shorter than the anterior crossvein.

One specimen, in the U. S. National Museum, from the Bryant and Palmer collection. Tjibodas, Mt. Gede, Java.

I. GENUS BURMITEMPIS, COCKERELL

Burmitempis, Cockerell, Amer. Journ. Sc. Vol. 44, p. 367 (1917).

Characters. — Head broad, dichoptic, third antennal joint very large, oblong-oval, hairy; with very long simple arista; wings large and broad, first basal cell long, emitting three veins from apex, and one from lower side, second posterior cell somewhat contracted apically, no discal cell, second basal small, narrow and nearly vertical, anal cell large and apically truncate, anal angle of wing rectangular; halteres enormous, with long thick knob; legs long, ordinary; genitalia small and simple.

Dr. Cockerell located this genus in the Empididae, apparently nearest to *Microsania*, but suggestive of the Tachydromiinae in some of its characters. The removal of *Microsania* to the Platypodidae may carry with it this genus. If the vein arising from the lower side of the first basal cell is the anterior branch of the media and the lower outer side of the basal the long and longitudinally placed anterior crossvein then the homologies of the neuration are clear though strangely aberrant. If, however, the small vein at the apex of the first basal cell is the anterior crossvein then the radius has two branches and the media three, a condition not in conformity with expectations among the Diptera. From the data at hand the systematic position of this fossil can not be definitely stated.

Geographical distribution.

1. *B. halteralis*, Cockerell, Amer. Journ. Sc. Vol. 44, p. 367, f. 7 (1917). Burmese Amber,
Miocene.

2. GENUS DRAPETIELLA, MEUNIER

Drapetella, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 86, note 3, p. 97 (1908).

Characters. — This genus was located with *Drapetis*, differing by the ovate style-bearing third joint of the antennae and the greatly inclined second basal cell. As a short style is anomalous in the entire subfamily Tachydromiinae and as the figure shows an impossible venation this genus is far from being certain. It possibly is an Ocydromiinae, with open discal cell, such as is shown by *Euthyneura aperta* to-day.

Geographical distribution.

1. *D. definita*, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 89, 97, pl. 3, f. 16, pl. 4, f. 2 (1908). Baltic Amber.

3. GENUS ELECTROCYRTOMA, COCKERELL

Electrocyrtoma, Cockerell, Ann. Ent. Soc. Amer. Vol. 10, p. 22 (1917).

Characters. — Resembling *Bicellaria*, but with lengthened arista, slender tibiae and tarsi, no intercalary vein, and with the fork of second and third veins much before anterior crossvein. Thorax gibbous, finely hairy; discal cell open but with a slight bend in fourth vein at two-fifths of its last section; base of front femora about two times as thick as apex.

Geographical distribution.

1. *E. burmanica*, Cockerell, Ann. Ent. Soc. Amer. Vol. 10, p. 22, f. 5 (1917). Burmese Amber,
Miocene?

4. GENUS EMPIDIA, WEYENBERGH

Empidia, Weyenbergh, Arch. Mus. Teyl, Haarlem, Vol. 2, p. 258 (1869).

Characters. — Handlirsch, who has examined the type, states that it is not all at clear that this fossil belongs to the Empididæ. The length of the specimen is about nineteen millimeters. The *Empis* wing pictured with the fossil has nothing to do with the insect.

Geographical distribution.

1. *E. Wulpi*, Weyenbergh, Arch. Mus. Teyl, Haarlem, Vol. 2, p. 258, pl. 34, f. 5, 5a (1869); Tijdschr. v. Ent. (2), Vol. 4, p. 237 (1869); Period. Zool. Vol. 1, p. 89 (1874); Scudder, Bull. U. S. Geol. Surv. Vol. 31, p. 88 (1886); Meunier, Ann. Soc. Sc. Bruxelles, Vol. 19, p. 178 (1895); Handlirsch, Foss. Ins. p. 634 (1909).

5. GENUS EUTHYNEURELLA, MEUNIER

Euthyneurella, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 113 (1908).

Characters. — « This genus differs from *Euthyneura* Macquart by some small peculiarities in the wings and by the scutellar hairs not being vertical. » From the figures of the wing and of the antenna the insect seems to be a *Trichina*.

Geographical distribution.

1. *E. longirostris*, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 91, 113, pl. 8, f. 8, 9; Vol. 9, p. 1 (1908).

6. GENUS HASMONA, GIEBEL

Hasmona, Giebel, Ins. Vorwelt, p. 209 (1856).

Characters. — This species is apparently Nematocerous, according to Handlirsch. The distinctive feature of the genus is the neuration which Giebel states is unlike that of living Empids. There is a strong simple vein extending parallel to the costa to the wing-tip, followed by three simple veins united before the middle by an oblique crossvein. The insect measures two millimeters.

Geographical distribution.

1. *H. leo*, Giebel, Brodie, Foss. Ins. p. 34, 121, pl. 3, f. 11 [1845] (*Empidæ*); Vale of Warbour, Giebel, Ins. Vorwelt, p. 209 (1856); Handlisch, Foss. Ins. p. 634, pl. 51, f. 14 (1908).

7. GENUS HILARITES, HEER

Hilarites, Heer, Viertjahresb. Naturf. Ges. Zürich, Vol. 1, p. 38 (1856).

Characters. — Heer stated that this insect apparently belongs to *Hilara*, but since the wings are not preserved it was impossible to make a certain determination.

Geographical distribution.

1. *H. bellus*, Heer, Viertjahresb. Naturf. Ges. Zurich, Vol. 1, p. 38, pl. 2, f. 5 (1856). Aix, France; Ligurian, Lower Oligocene.

8. GENUS MEGHYPERELLA, MEUNIER

Meghyperella, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 112 (1908).

Characters. — Antennæ three-jointed, the last joint conical, a little longer than its distinct arista. Wings rather broad and pointed, the discal cell emitting two posterior veins, the intercalary absent. The genus is apparently related to *Leptopeza*, differing in the possession of a more generalized arista. It may be a predecessor of *Stenoproctus*.

Geographical distribution.

1. *M. porphyropsoides*, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 91, 112, Baltic Amber.
pl. 8, f. 5-7 (1908)

9. GENUS MESOMYITES, COCKERELL

Mesomyites, Cockerell, Proc. U. S. Nat. Mus. Vol. 52, p. 377 (1917).

Characters. — *Mesomyites* was originally described as a new genus of the Empididæ, and the small size, slender wing and general features of the neuration are suggestive of the Clinoceratinæ. However, the second basal cell emits a vein parallel with the discal cell, the second branch of the cubitus. Thus the anal cell is widely open, an impossible condition among the Brachycera. Dr. Cockerell, in Nature, Vol. 103, p. 44 (1919), has referred his genus to the anomalous tipulid genus *Styringomyia*. The species on which *Mesomyites* was based is the following.

Geographical distribution.

1. *M. concinnus*, Cockerell, Proc. U. S. Nat. Mus. Vol. 52, p. 377, pl. 31, Isle of Wight, Oligocene.
f. 7 (1917).

10. GENUS OUSTALETIMYIA, MEUNIER

Oustaletimyia, Meunier, Ann. Soc. Ent. France, Vol. 62, Bull. p. 332 (1893).

Characters. — A provisional genus described as being related to *Edalea* but with simple femora and two-jointed antennæ.

Geographical distribution.

1. *O. succinorum*, Meunier, Ann. Soc. Ent. France, Vol. 62, Bull. p. 332 Baltic Amber.
(1893).

11. GENUS PALÆOEDALEA, MEUNIER

Palaeoedalea, Meunier, Ann. Soc. Sc. Bruxelles, Vol. 26, p. 101 (1902).

Characters. — Related to *Edalea* in neuration and three-jointed elongate antennæ, but with slender simple legs.

Geographical distribution.

1. *P. samlandica*, Meunier, Ann. Soc. Sc. Bruxelles, Vol. 26, p. 101 (1902). Baltic Amber.
elegans, Meunier, ibidem, Vol. 26, f. 5 (1902).

12. GENUS PALÆOLEPTOPEZA, MEUNIER

Palæoleptopeza, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 111 (1908).

Characters. — Near *Leptopeza* but with spinose hind femora.

Geographical distribution.

1. *P. gracilis*, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 91, 111, pl. 7, Baltic Amber.
f. 15, 16, pl. 8, f. 1-4 (1908).

13. GENUS PALÆOPARAMESIA, MEUNIER

Palæoparamesia, Meunier, Ann. Soc. Sc. Bruxelles, Vol. 26, p. 98 (1902); Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 107 (1908).

Characters. — It is impossible to tell to which division of *Clinocera*, in the broad sense, this form belongs.

Geographical distribution.

1. *P. Proosti*, Meunier, Ann. Soc. Sc. Bruxelles, Vol. 26, p. 98, f. 3, 4 (1902); Baltic Amber.
Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 90, 107 (1908).

14. GENUS PARATHALASSIELLA, MEUNIER

ParathalassIELLA, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 90 (1908).

Characters. — Discal cell present, the fourth vein forked, claws and pulvilli long and robust, third joint of the antennæ conical, the arista as long as the antenna. From this information alone it is impossible to determine the position of this fossil fly. Meunier was familiar with the figure of *Parathalassius* in the Entomological News, 1906, and located the genus near this form, but the furcate fourth vein indicates a very different insect.

Geographical distribution.

1. *P. problematica*, Meunier, Ann. Sc. Nat. (Zool.) 9 ser. Vol. 7, p. 90, 106, Baltic Amber.
pl. 6, 17, 18; pl. 7, f. 2 (1908).

15. GENUS PROTOEDEALEA, COCKERELL

Protoedalea, Cockerell, Proc. U. S. Nat. Mus. Vol. 57, p. 252 (1920).

Characters. — A precursor of *Œdalea* and *Anthalia*, its ancestral traits evidenced by the lengthened first and second veins, narrow discal cell, oblique forking of the cubitus, the posterior branch of which squarely terminates the anal cell, simple legs, thorax only moderately gibbous, and large size, being over four mm. in length. Antennæ short the last joint oboviform, style indistinguishable, proboscis retracted, discal cell unusually long, fourth vein abbreviated, venter of the male with sparse long hairs, body black, wings dusky.

Geographical distribution.

1. *P. brachystoma*, Cockerell, Proc. U. S. Nat. Mus. Vol. 57, p. 252, f. 8 Eocene, Green River
(1920). Shales, Colorado.

16. GENUS THIRZA, GIEBEL

Thirza, Giebel, Ins. Vorwelt, p. 209 (1856).

Characters. — Antennæ very short, first joint cylindrical, second and third joints globose; eyes separated; proboscis short and thick. Thorax convex; femora short and slightly thickened at middle, tibiæ twice as long as femora, with two apical bristles, metatarsi half as long as tibiæ. The wings have a peculiar neuration, which is difficult to visualize, and therefore the original description will be quoted : « Die Schulterader verhält sich wie gewöhnlich, aber die äuszere Mittelader entspringt von einer die Schulterader mit der innern Mittelader verbindenden Querader, und zwar sogleich zweiästig, der äuszere Ast endet einfach an der kleinen Querader, welche die Schulterader mit dem innern Ast verbindet; dieser gabelt sich kurz vor der Flügelmitte in demselben Niveau, wo der innerste Ast der schulterader sich gabelt; beide Aeste sind dann später durch eine Querader verbunden, von welcher der gewöhnliche Zwischenlängsast ausgeht. Auch die innere Mittelader entspringt nicht unmittelbar am Grunde, sondern von einer feinen Querader neben diesem. Sie biegt sich vor der Mitte zum Innenrande und sendet an der Biegungsstelle die kleine Querader zum innersten Ast der äuszern Mittelader, von welcher ein Längsast abgeht. Das Analfeld durchziehen zwei von der Randkerbe and stark divergirende Aeste. »

The type and only species is *Th. Naumannii* Giebel. The genus was located in the Hybotidæ by its describer

Geographical distribution.

1. *Th. Naumannii*, Giebel, Ins. Vorwelt, p. 210 (1856).

Baltic Amber, Lower Oligocene.

List of the Fossil Empididæ Belonging to Existing Genera.

References to these species occur in the lists of species of the various genera.

Brachystoma spinulosum Loew.

Chelifera detestata Meunier.

Chelopoda delicata Meunier, *dolosa* Meunier, *rustica* Meunier (Subgenus *Phyllodromia*) *vaga* Meunier.

Drapetis brevis Meunier, *decolorata* Meunier, *decorata* Meunier, *mortua* Meunier, *vitiosa* Meunier.

Dysaletria diabolica Meunier.

Empis bulbirostris Meunier, *carbonum* Germar, *florissantana* Cockerell, *macilenta* Meunier, *macrophthalma* Förster, *mala* Meunier, *malefica* Meunier, *melia* Heyden, *miocenica* Cockerell, *mordax* Meunier, *morosella*, n. n.; *perdita* Cockerell, *personata* Meunier, *Poeppigi* Giebel, *pulvillata* Loew, *stilicornis* Meunier, *tibialis* Meunier, *tritava*, n. n.

Gloma hirta Loew.

Hemerodromia detestata Meunier.

Hilara exilis Meunier, *Heerii* Meunier, *litigiosa* Meunier, *macilenta* Meunier, *tarda* Meunier.

Hybos exilis Meunier, *tanuis* Meunier.

Leptopeza clavipes Loew, *coccinna* Meunier, *spinigera* Meunier.

Micrempis eocenica Meunier, *suspiciosa* Meunier.

Microphorus defunctus Handlirsch, *eocenicus* Meunier, *putidus* Meunier.

Edalea robusta Meunier.

Platypalpus concitatus Meunier, *eversoris* Meunier, *interfactoris* Meunier, *predatoris* Meunier.

Ragas generosa Meunier.

Rhamphomyia ablata Meunier, *angusta* Meunier, *antipedalis* Loew, *corrupta* Meunier, *crinitarsis* Loew, *distans* Loew, *enena* Cockerell, *errabunda* Meunier, *formosa* Loew, *hypolitha* Cockerell, *insolita* Meunier, *involuta* Meunier, *media* Meunier, *obtusa* Meunier, *adaloides* Meunier, *polymorpha* Meunier, *porrecta* Meunier, *pteropa* Loew, *ptilopa* Loew, *remitarsis* Loew, *sepulta* Cockerell, *ungulina* Loew.

Tachydromia stilpon Meunier.

Tachypeza egelata Meunier, *voracis* Meunier.

Timalphes acuticornis Loew, *gracilis* Meunier, *pahata* Loew.

Trichopeza sucina, n. n.

GENERAL REMOVED FROM THE EMPIDIDÆ (1).

The following genera have been held by various authors as belonging to the Empididæ. The resemblances to members of this family have proved to be superficial and the genera have one by one been removed to other families. The forms here placed in the Bombyliidæ are highly specialized in that family, with neuration reduced in accompaniment with their small size. They can hardly be regarded as annectent forms, linking the Bombyliidæ with the Empididæ, because they are more highly specialized in other respects than the Hybotinæ with which they connect. Their resemblances therefore are to be regarded as parallel developments. The status of the genera in question is given herewith.

I. GENUS HILARIMORPHA, SCHINER

Hilarimorpha, Schiner, Wien. Ent. Monatschr. Vol. 4, p. 54 (1860); Fauna Dipt. Austr. Vol. 1, p. 116 (1862); Mik, Verh. Zool.-bot. Ges. Wien. Vol. 31, p. 329 (1881); Williston, Psyche, 1888, p. 99 (1888); Bigot, Bull. Soc. Zool. France, 1891, p. 15 (1891); Osten-Sacken, Berl. Ent. Zeitschr. Vol. 35, p. 303 (1890); Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 388 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 336 [1902] (*Hilaromorpha*); Kertész, Cat. Dipt. Vol. 3, p. 333 (1908); Verrall, Brit. Flies, Vol. 5, p. 243 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 552 (1910).

Characters. — This genus contains a few small blackish bristleless species that run to the Empididæ in existing keys of Diptera. The general opinion is that *Hilarimorpha* belongs to the Leptidae, in which family it is aberrant in having minute empodia, reduced tibial spurs and no discal cell. Five species have been described.

Type species : *H. singularis*, Egger.

Geographical distribution.

1. *H. maculata*, Matsumura, Addit. Tokyo, Vol. 2, p. 357, pl. 21, f. 19 (1916). Japan.
2. *H. Mikii*, Williston, Psyche, 1888, p. 100 (1888); Melander, Trans. Amer. Illinois. Ent. Soc. Vol. 28, p. 337 (1902).
3. *H. obscura*, Bigot, Bull. Soc. Ent. France (6), Vol. 7, p. 141 (1887); Ann. Soc. Ent. France (6), Vol. 9, p. 129 (1889); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 337 (1902).
4. *H. singularis*. Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 346 (1860); Europe. Schiner, Fauna Dipt. Austr. Vol. 1, p. 117 (1862); Verh. Zool.-bot. Ges. Wien, Vol. 18, p. 909 (1868); Mik ibidem, Vol. 31, pl. 16, f. 19-21 (1881); Williston, Stett. Ent. Zeit. Vol. 46, p. 401 (1885); Verrall, Brit. Flies, Vol. 5, p. 243, f. 178 (1909).
5. *H. tristis*, Egger, Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 347 (1860); C. Europe. Schiner, Fauna Dipt. Austr. Vol. 1, p. 117 (1862); Mik, Verh. Zool.-bot. Ges. Wien, Vol. 31, pl. 16, f. 22 (1881)

(1) A. L. Tonnoir, Rec. Austr. Mus. 1925, transfers *Ironymia* (vide antea, p. 48) to the Platypezidæ. P. Tillyard, Insects of Australia, 1926, p. 366, transfers *Sciadocera* (antea, p. 48) to the Phoridae, notwithstanding its different antennal structure.

2. GENUS MICROSANIA, ZETTERSTEDT

Microsania, Zetterstedt, Isis, Vol. 1, p. 30 (1837); Fauna Ins. Lappon., p. 534 (1838); Dipt. Scand. Vol. 1, p. 333 (1842); Bigot, Ann. Soc. Ent. France (6), Vol. 9, p. 123 (1889); Melander, Man. N. Amer. Dipt. p. 225 (1908); Lundbeck, Dipt. Danica, Vol. 3, p. 18 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 43, 47 (1910); Melander, Psyche, Vol. 29, p. 43-48 (1922).

Microcyrta, Bigot, Ann. Soc. Ent. France (3), Vol. 10, p. 557, 564 (1857); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253 (1903).

Pachyzeza, Lioy (not Serville, Coleoptera, 1835), Atti. Inst. Sc. Veneto, Venezia (3), Vol. 9, p. 723 (1863); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 255 (1903).

Platytelma, Rondani, Dipt. Ital. Vol. 1, p. 138 (1856); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 256, 261 (1903).

Characters. — The species of *Microsania* look much like *Bicellaria* with which genus they have repeatedly been associated. *Microsania* departs from all Empid genera in possessing a large and inflexed hypopygium and a single median row of acrostichal hairs. The costa is setulose and is interrupted at the end of the first vein, the anal cell is pointed, the proboscis is short and fleshy, and the middle tibiae possess apical bristles.

The genus belongs to the Platypezidae, as discussed by the writer in *Psyche* for April, 1922, being closely related to *Opetia* and *Platycnema*. Loew recognized the Platypezid relationships when he described one of the American species as *Platycnema imperfecta*. The other European writers have all located *Microsania* with the Empididae. Almost none of the Empididae are common to both Europe and North America, but two of the three species of *Microsania* occur on both continents.

Type species : *M. stigmatica*, Zetterstedt.

Geographical distribution.

1. *M. pallipes*, Meigen, Syst. Beschr. Vol. 6, p. 356 [1830] (*Cyrtoma*); Macquart, C. Europe. Hist. Nat. Dipt. Vol. 1, p. 360 [1834] (*Cyrtoma*); Rondani, Dipt. Ital. Vol. 1, p. 138 [1856] (*Platytelma*); Bigot, Ann. Soc. Ent. France (3), Vol. 5, p. 557, 564 [1857] (*Microcyrta*); Schiner, Fauna Dipt. Austr. Vol. 1, p. 76 [1862] (*Cyrtoma*); Lioy, Atti. Inst. Veneto, Sc. Venezia (3), Vol. 9, p. 723 [1863] (*Pachyzeza*).
2. *M. pectinipennis*, Meigen, Syst. Beschr. Vol. 6, p. 356, pl. 66, f. 15, 16 [1830] Europe, North America. (*Cyrtoma*); Zetterstedt, Dipt. Scand. Vol. 1, p. 335 [1842] (*pectinipennis*); ibidem, Vol. 8, p. 3013 [1849] (*pectinipes*); Wahlgren, Ent. Tidskr. Vol. 31, p. 47 [1910] (*pectinipennis*); Melander, Psyche, Vol. 29, p. 46 (1922).
3. *M. stigmatica*, Zetterstedt, Fauna Ins. Lappon. p. 534 [1838] (*Cyrtoma*); Dipt. Scand. Vol. 1, p. 334 (1842); ibidem, Vol. 8, p. 3013 (1849); ibidem, Vol. 13, p. 4098 (1859); Bonsdorff, Finl. tváv. Ins. Dipt. Vol. 1, p. 157 (1861); Lundbeck, Dipt. Dan. Vol. 3, p. 19, f. 3 (1910); Wahlgren, Ent. Tidskr. Vol. 31, p. 47, f. 3 (1910); Frey, Acta Soc. Sc. Fenn. Helsingfors, Vol. 37 (3), p. 8 (1913); Melander, Psyche, Vol. 29, p. 46 (1922).
imperfecta, Loew, Berl. Ent. Zeitschr. Vol. 9: Cent. 6, No. 82 [1865] (*Platycnema*); Aldrich, Cat. Dipt. N. Amer. p. 342 [1905] (? *Platycnema*).

3. GENUS MALTHACOTRICA, BECKER

Heterotropus, Loew, Beschr. Eur. Dipt. Vol. 3, p. 180 (1873).

Malthacotrica, Becker, Ann. Mus. Zool. Acad. Sc. St. Pétersb. Vol. 12, p. 312, pl. 2, f. 18 (1907); Kertész, Cat. Dipt. Vol. 6, p. 14 (1909); Becker, Ann. Mus. Zool. Acad. Sc. St. Pétersb. 1912, Vol. 17, p. 482 (1913).

Characters. — Becker, in 1907, described *Malthacotrica glauca* as a new genus of Empididæ, locating it in the Hybotinæ because of the lengthened anal cell and the horizontal proboscis, even though the insect has the third vein forked and the discal cell emitting three posterior veins. Later, in 1913, Becker assigned his genus to the Bombyliidæ as synonymous with the older *Heterotropus* Loew. The genus occurs in Central and Western Asia, with two known species. Apparently it is closely related to the American *Prorates* Melander, which see on the following page.

Geographical distribution of the species of *Heterotropus*.

1. *H. albidiennis*, Loew, Beschr. Eur. Dipt. Vol. 3, p. 180 (1873); Becker, Transcasplia. Ann. Mus. Zool. Acad. Sc. St. Pétersb. Vol. 17, p. 586 (1913).
2. *H. glaucus*, Becker, ibidem. Vol. 12, p. 313, pl. 2, f. 13 [1907] (*Malthaco-* Turkestan. *tricha*); ibidem, Vol. 17, p. 482, f. 25 (1913).

4. GENUS MYTHICOMYIA, COQUILLETT

Mythicomyia, Coquillett, Ent. News Philad. Vol. 4, p. 209 (1893); Proc. U. S. Nat. Mus. Vol. 18, p. 409 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 337 (1902); Coquillett, Proc. Ent. Soc. Wash. Vol. 5, p. 253 (1903); Williston, Man. N. Amer. Dipt. p. 211, 218 (1908); Kertész, Cat. Dipt. Vol. 3, p. 333 (1908); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 573 (1910); Cresson, Ent. News Philad. Vol. 26, p. 448-456 (1915); Greene, Proc. Ent. Soc. Wash. Vol. 26, p. 60-64 (1924).

Heterhybos, Brethes, Revista Chilena Hist. Nat. Vol. 23, p. 40 (1919).

Characters. — Coquillett erected the genus *Mythicomyia* in 1893, placing it in the Empididæ. Since that time the group has been shifted here and there by various writers. Aldrich and Kertész, in their respective catalogs, placed it in the Leptidæ, the writer in 1902, then unacquainted with specimens, left it in the Empididæ but segregated it in a subfamily, the Mythicomyiinæ, Cresson located it in the Empididæ, and Williston and Greene probably in the Bombyliidæ.

Mythicomyia is represented by eight described species, restricted to the Western United States, where the minute flies are found hovering over bush-flowers, such as ocean spray. In addition, *Heterhybos hyalinipennis* Brethes was described as a new genus of Empididæ from Rio Blanco, South America. This genus is completely synonymous with *Mythicomyia*. These species are closely related to the European *Glabellula* Bezz, 1902, redescribed by Greene, 1924, as *Pachyneres*, and belong to the subfamily Cyrtosiinæ of the Bombyliidæ, as defined by Becker, Ann. Mus. Zool. Acad. Sc. St. Pétersb. Vol. 17, p. 472 (1913). *Glabellula* has the second basal cell fused with the discal cell and the femora more or less enlarged. In *Mythicomyia* the discal cell is complete, the two basals weakly separated and the femora are slender.

Type species : *M. Rileyi*, Coquillett, the original species.

Geographical distribution.

- | | |
|--|----------------------|
| 1. <i>M. armata</i> , Cresson, Ent. News Philad. Vol. 26, p. 455, f. 3 (1915). | W. United States. |
| 2. <i>M. armipes</i> , Cresson, ibidem, Vol. 26, p. 454, f. 5, 6 (1915). | New Mexico. |
| 3. <i>M. atra</i> , Cresson, Ent. News Philad. Vol. 26, p. 456 (1915). | New Mexico. |
| 4. <i>M. californica</i> , Greene, Proc. Ent. Soc. Wash. Vol. 26, p. 61 (1924). | California. |
| 5. <i>M. flavipes</i> , Cresson, Ent. News Philad. Vol. 26, p. 452, f. 1, 2 (1915). | W. United States. |
| 6. <i>M. hyalinipennis</i> , Brethes, Revista Chilena. Hist. Nat. Vol. 23, p. 40 [1920] (Heterhybos). | Chile. |
| 7. <i>M. minuta</i> , Greene, Proc. Ent. Soc. Wash. Vol. 26, p. 62 (1924). | New Mexico. |
| 8. <i>M. pictipes</i> , Coquillett, Proc. U. S. Nat. Mus. Vol. 25, p. 103 (1902); Cresson, Ent. News Philad. Vol. 26, p. 453 (1915). | S. W. United States. |
| 9. <i>M. Rileyi</i> , Coquillett, Ent. News Philad. Vol. 4, p. 209 (1893); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 338 (1902); Cresson, Ent. News Vol. 26, p. 454 (1915). | S. W. United States. |
| 10. <i>M. scutellata</i> , Coquillett, Proc. U. S. Nat. Mus. Vol. 25, p. 102 (1902); Cresson, Ent. News Philad. Vol. 26, p. 451, f. 4 (1915). | W. United States. |
| 11. <i>M. tibialis</i> , Coquillett, Proc. U. S. Nat. Mus. Vol. 18, p. 409 (1896); Melander, Trans. Amer. Ent. Soc. Vol. 28, p. 338 (1902); Cresson, Ent. News Philad. Vol. 26, p. 453 (1915). | W. United States. |

5. GENUS PRORATES, MELANDER

Prorates, Melander, Ent. News Philad. Vol. 17, p. 372 (1906); Williston's Man. N. Amer. Dipt. p. 224 (1908); Kertész, Cat. Dipt. Vol. 6, p. 3 (1909); Coquillett, Proc. U. S. Nat. Mus. Vol. 37, p. 595 (1910).

Characters. — Small opaque gray bristleless species, somewhat resembling a Therevid, stigma strong, third and fourth veins both forked, alula well developed, legs simple, third antennal joint with a microscopic style. Head globose, broader than long, occiput concave, the face extremely short and retracted, eyes of the male contiguous from the antennæ to the vertex, the facets of the upper two-thirds enlarged, of the female broadly separated on the front, the facets uniform, front of the female with parallel sides, no ocellar bristles; eyes reduced in back beneath, the lower part of the head made up of the occiput, mouth opening large, with rounding edges; proboscis rigid, porrect, slightly shorter than the head, palpi retracted; antennæ three-jointed, the basal joints similar and small, the third joint compressed, conical, tipped with a microscopic bristle. Thorax but little convex, nearly bare, bristles very weak, one notopleural, one postalar and one pair of scutellars present, mesopleuræ bearing weak hairs. Abdomen cylindrical, scarcely tapering in either sex, pygidium terminal, not enlarging the abdomen, entirely included within a pair of lateral valves, no ovipositor. Legs slender, simple, nowhere thickened, devoid of all bristles, the hairy covering very short, pulvilli minute. Wings rather broad, anal angle rectangular, no basal bristle, costal hairs microscopically small, costa continuing to the posterior branch of the third vein, auxiliary vein distinct from the first and gently curving forward at its end to meet the costa, a distinct elliptical stigma surrounding the end of the first vein which is not far beyond the middle of the wing, second vein rather short, ending nearer the first than to the tip of the wing, pedicel of the second and third veins arising over the middle of the second basal cell, anterior crossvein at the middle of the discal cell, the first basal therefore longer than the second, discal cell pointed apically, the posterior crossvein nearly parallel with the hind margin of the wing, petiole of the second posterior cell very short, none of the posterior veins attaining the hind margin, anal cell larger than usual, its outer angle acute, anal vein reaching the hind margin, alula well developed in both sexes but not margined, calypteres large for an Empid, membranous and with a thickened fringed margin.

Type species : *P. claripennis*, Melander, the only species known. Many specimens of this Insect were taken by Mr. H. L. Viereck in 1902 at Alamogordo, New Mexico, at which time and place he took also the two species of the recently described genus *Cænotus* Cole. *Cænotus*, which is strikingly like *Prorates*, although generically quite distinct, was assigned by Cole to the Therevidæ, in which family it is aberrant in lacking the « small crossvein » at the apex of the second basal cell and in being bristleless.

Prorates has been regarded as a primitive but anomalous Empid. Judging by the structure of its proboscis it probably is anthophilous. In appearance it suggests a minute Therevid or Bombyliid with reduced neuration, or a Scenopinid. According to existing tabulations to the families of Diptera the species would be located in the Empididæ in the subfamily Hybotinæ. From all the members of this group, however, it departs in having the costa stopping at the third vein, the auxiliary vein distinct and ending in the costa, the antennal style microscopic, and the mesopleuræ furnished with some pubescence.

Meghyperus, which is more clearly Hybotine than *Prorates* is, shows many parallel characters, such as the distinct though evanescent auxiliary vein, presence of alula, dichopticism of female, and bristleless legs. These important characters, all at variance with other Hybotinæ, perhaps indicate primitive traits retained from a common ancestor, but more likely are the result of a parallel evolution among insects having similar flower-frequenting habits.

It is not yet time to remove *Meghyperus* from the Hybotinæ, but *Prorates* may properly be placed in the Bombyliidæ, where with *Mallacotricha* (= *Heterotropus*) and also with *Cænotus* it will comprise the subfamily Heterotropinæ. The three genera may be separated as follows :

- | | |
|---|-----------------------------------|
| 1. <i>Proboscis porrect; first basal cell much longer than the second.</i> | 2. |
| <i>Proboscis very short, not projecting; first basal cell not greatly longer than the second.</i> | Genus <i>Cænotus</i> , Cole. |
| 2. <i>Pygidium deeper than long; discal cell wider than the second basal, emitting three posterior veins; third antennal joint subulate, with bristle-tipped style.</i> | Genus <i>HETEROTROPUS</i> , Loew: |
| <i>Pygidium longer than deep; discal cell narrower than the second basal, fourth vein petiolate; third antennal joint conical, with microscopic style . . .</i> | Genus <i>PRORATES</i> , Melander. |

Geographical distribution.

1. *P. claripennis*, Melander, Ent. News Philad. Vol. 17, p. 373, f. (1906); New Mexico.
Williston's Man. N. Amer. Dipt. p. 219, f. 5 (1908).

ADDENDA : RECENTLY DESCRIBED EMPIDIDÆ

GENUS ACARTERUS, LOEW

- A. varipes*, Senior-White, Spol. Zeylan. Colombo, Vol. 12, p. 394 (1924). Ceylon.

GENUS ARIASELLA, GIL

- Ariassel**a, Gil, Bol. Soc. Esp. Hist. Nat. Vol. 23, p. 151 (1923) (1).

- A. semiaaptera*, Gil, Bol. Soc. Esp. Hist. Nat. Vol. 23, p. 151, figs. (1923). Spain.

(1) **Ariassel**a, a genus related to *Pieltainia*, Arias, but characterized by having vestigial wings. Type *A. semiaaptera*, Gil, a species slightly under three millimeters in length. Thorax compressed, body blackish, legs especially the hind femora and tibiæ, robust, one notopleural, two scutellar bristles; face narrow, one pair ocellar bristles, third antennal joint circular, with dorsal arista; wings vestigial, reaching the fifth abdominal segment, the basal two-thirds consisting of a strap-like pedicel, with only the costal and a single longitudinal vein; halteres absent.

GENUS BICELLARIA, MACQUART

- B. nigrita*, Collin, Ent. Mo. Mag. Vol. 62, p. 190 (1926). England.
B. subpilosa, Collin, ibidem, Vol. 62, p. 190 (1926). England.

GENUS BRACHYSTOMA, MEIGEN

- B. Hamiltoni*, Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 451, f. 38-41 New Zealand. (1923).

GENUS CHELIFERA, MACQUART

- C. angusta*, Collin, Ent. Mo. Mag. Vol. 63, p. 95 (1927). England.
C. aperticauda, Collin, ibidem, Vol. 63, p. 95 (1927). England.
C. astigma, Collin, ibidem, Vol. 63, p. 94 (1927). England.
C. concinnicauda, Collin, ibidem, Vol. 63, p. 95 (1927) *monostigma*, Zetterstedt, not Meigen. N. W. Europe.
precatoria, var. B, Zetterstedt
stigmatica, Lundbeck, not Schiner.
C. diversicauda, Collin, Ent. Mo. Mag. Vol. 63, p. 95 (1927). British Isles.
C. erecta, Collin, ibidem, Vol. 63, p. 94 (1927). British Isles.
C. fontanalis, Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 422, fig. [1923] New Zealand (*Hemerodromia*).
C. fectinicauda, Collin, Ent. Mo. Mag. Vol. 63, p. 95 (1927). England.

GENUS CHELIPODA, MACQUART

- C. oliraensis*, Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 443, f. 11-19 New Zealand. [1923] (*Litanomyia*).

GENUS CLINOCERA, MEIGEN

- C. hirtiloba*, Speiser, Schr. Physik-Ökonom. Ges. Königsberg, Vol. 64, p. 13 Prussia. [1924] (*Atalanta*, *Euceltidia*).

GENUS DRAPETIS, MEIGEN

- D. (Elaphropeza) abdominalis*, Senior-White, Mem. Dept. Agric. India, Ceylon. Vol. 7, p. 151 (1922).
D. convergens, Collin, Ent. Mo. Mag. Vol. 62, p. 149 (1926). England.
D. (Elaphropeza) distincta, Senior-White, Mem. Dept. Agric. India, Vol. 7, Ceylon. p. 152 (1922).
D. incompleta, Collin, Ent. Mo. Mag. Vol. 62, p. 148 (1926). England.
exilis, Loew and Lundbeck, not Meigen.
D. (Elaphropeza) nigropunctata, Senior-White, Mem. Dept. Agric. India, Vol. 7, India. p. 152 (1922).
D. (Elaphropeza) notatithorax, Senior-White, ibidem, Vol. 7, p. 151, pl. 14, Ceylon. f. 10 (1922).
D. parilis, Collin, Ent. Mo. Mag. Vol. 62, p. 148 (1926). England.
D. (Elaphropeza) plumicornia, Senior-White, Mem. Dept. Agric. India, Vol. 7, Ceylon. p. 150, pl. 14, f. 9 (1922).

GENUS EMPIMORPHA, COQUILLETT

- E. torrentalis*, Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 463, f. 65-68 (1923). New Zealand.

GENUS EMPIS, LINNÆUS

Kritempis, Collin, Ent. Mo. Mag. Vol. 62, p. 236 (1926) (1).

Leptempis, Collin, ibidem, Vol. 62, p. 235 (1926) (2).

Rhadinempis, Collin, ibidem, Vol. 62, p. 236 (1926) (3).

<i>E. (Rhadinempis) Bazini</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 236 (1926).	France.
<i>E. (Pterempis) cyrenaica</i> , Beazzi, Boll. Mus. Zool. Torino, Vol. 39, N. S. (18), p. 12 (1924).	Cyrenaica.
<i>E. dalmatica</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 320 (1925).	Dalmatia.
<i>E. (Xanthempis) laetabilis</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 237 (1926).	England.
<i>E. lamelligornis</i> , Becker (antea, p. 159).	
subsp. <i>planetica</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 23 (1927).	England.
subsp. <i>protarsalis</i> , Collin, ibidem, Vol. 63, p. 24 (1927).	England.
<i>E. limata</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 22 (1927).	England.
<i>E. Lindneri</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 317 (1925).	Dalmatia.
<i>E. pennaria</i> , Fallen (antea, p. 167).	
subsp. <i>bicuspidata</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 22 (1927).	England.
subsp. <i>laminata</i> , Collin, ibidem, Vol. 63, p. 22 (1927).	England.
<i>E. prævia</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 23 (1927).	England.
<i>E. subbrunipennis</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 319 (1925).	Dalmatia.
<i>E. subinfumata</i> , Malloch, No. Am. Fauna, No. 46, p. 184 (1923).	Pribilof Islands.
<i>E. (Anachrostichus) Verralli</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 21 (1927).	England.
<i>E. (s. str.) Woodi</i> , Collin, ibidem, Vol. 53, p. 21 (1927).	England.

GENUS EUTHYNEURA, MACQUART

E. Halidayi, Collin, Ent. Mo. Mag. Vol. 62, p. 214 (1926). England.

GENUS FRAUDATOR, HUTTON (4)

Fraudator, Hutton, Trans. N. Zeal. Inst. 1900, Art. 1, p. 23 (1900); Tillyard, Ins. Austr. and N. Zealand, p. 365 [1926] (ref. to Empididæ).

F. perspicuus, Hutton, Trans. N. Zeal. Inst. 1900, Art. 1, p. 23 (1900); New Zealand. Tillyard, Ins. Austr. N. Zeal. p. 365 (1926).

(1) **Kritempis**, subgenus of *Empis*. Type *E. algira*, Macquart: includes *livida*, Linnæus, *Maquarti*, Becker (= *geniculata*, Macquart, not Zetterstedt), *macropalpa*, Egger, *nigrimana*, Becker, *sibillina*, Beazzi. Anal vein faint, not reaching margin, discal cell not truncate, with the first or the first and second posterior veins not reaching the margin, cubital fork rather acute; a small bristle in comb at tip of hind tibiae behind; male facets scarcely enlarged, no rows of long bristles beneath the middle femora, upper lamellæ of pygidium often very large, always much larger than the side lamellæ, in *livida* composing almost the whole of the pygidium; species never very small.

(2) **Leptempis**, subgenus of *Empis*. Type *E. grisea*, Fallen. Rather large, narrow-bodied, long-legged, wings ample, never blackish, sometimes maculated, anal vein distinct to margin; hypopygium with side lamellæ long and narrow, upper lamellæ rather prominent and bilobed, penis long and usually very undulated; female legs extensively yellow.

(3) **Rhadinempis**, subgenus of *Empis*. Type *E. Bazini*, Collin, the only assigned species. Resembles the *Holoclera* group of *Rhamphomyia*, but has the third vein forked. Antennæ short, the third joint short and pointed, about as long as the basal two together, and half as long as the microscopically pubescent arista; acrostichals absent, dorsocentral bristles uniserial; axillary excision of wings greater than a right angle.

(4) Originally assigned to the Bombyliidæ, but superficially resembling *Timaphes fumosa*, Hutton Type, and only species, *F. perspicuus*, Hutton. Eyes contiguous, upper facets coarser; third antennal joint swollen at base and gradually tapering to a long point; proboscis short; thorax black, humeri brown, scutellum yellowish, pleurae silvery; abdomen linear, brown; wings smoky, stigma large, third vein forked, third basal cell closed and petiolate, fourth vein arched; 7 mm. in length.

GENUS HALPOMERA, MACQUART

H. chiloensis, Brethes, Revista Chilena Hist. Nat. Vol. 28, p. 105 (1924). Chile.

GENUS HEMERODROMIA, MEIGEN

<i>H. (s. str.) adulatoria</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 96 (1927).	England.
<i>H. (s. str.) baetica</i> , Collin, ibidem, Vol. 63, p. 96 (1927).	England.
<i>H. (s. str.) laudatoria</i> , Collin, ibidem, Vol. 63, p. 96 (1927).	England.

GENUS HILARA, MEIGEN

<i>H. apta</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 66 (1927).	England.
<i>H. auripila</i> , Curran, Can. Ent. Vol. 58, p. 247 (1926).	Alberta.
<i>H. biseta</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 64 (1927).	England.
<i>H. brevistyla</i> , Collin, ibidem, Vol. 63, p. 28 (1927).	England.
<i>pinetorum</i> , of British lists, not Zetterstedt.	
<i>H. Crickmayi</i> , Curran, Can. Ent. Vol. 58, p. 246 (1926).	N. W. Canada.
<i>H. Curtisi</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 26 (1927).	England.
<i>H. dracophylli</i> , Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 456, f. 50 (1923).	New Zealand.
<i>H. flavinceris</i> , Miller, ibidem, Vol. 54, p. 453, f. 42, 43 (1923); Tillyard, Ins. Austr. N. Zeal. p. 365, f. W 56 (1926).	New Zealand.
<i>H. fossalis</i> , Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 460, f. 54-56 (1923).	New Zealand.
<i>H. galactoptera</i> , Strobl (antea, p. 124).	
<i>vallium</i> , Verrall-Collin, Ent. Mo. Mag. Vol. 63, p. 62 [1927] (no description).	
<i>H. Garretti</i> , Curran, Can. Ent. Vol. 58, p. 248 (1926).	Alberta.
<i>H. granditarsis</i> , Curran, ibidem, Vol. 58, p. 248 (1926).	Alberta.
<i>H. griseifrons</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 65 (1927).	England.
<i>H. hirtella</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 64 (1927).	England.
<i>H. hirtipes</i> , Collin, ibidem, Vol. 63, p. 65 (1927).	England.
<i>H. hyperborea</i> , Frey, Mem. Acad. Sc. Russ. Vol. 29 (10), p. 12 (1915).	N. Siberia.
<i>H. implicata</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 66 (1927).	England.
<i>H. littoralis</i> , Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 454, f. 44-47 (1923).	New Zealand.
<i>H. media</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 63 (1927).	England.
<i>H. merula</i> , Collin, ibidem, Vol. 63, p. 63 (1927).	England.
<i>H. monedula</i> , Collin, ibidein, Vol. 63, p. 63 (1927).	England.
<i>H. morata</i> , Collin, ibidem, Vol. 63, p. 67 (1927).	England.
<i>H. nigrokirta</i> , Collin, ibidem, Vol. 63, p. 65 (1927).	England.
<i>H. platyura</i> , Loew (antea, p. 118).	
<i>obesa</i> , Verrall-Collin, Ent. Mo. Mag. Vol. 63, p. 61 [1927] (no description).	
<i>H. primula</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 64 (1927).	England.
<i>H. quadriseta</i> , Collin, ibidem, Vol. 63, p. 64 (1927).	England.
<i>H. rejecta</i> , Collin, ibidem, Vol. 63, p. 26, 66 (1927).	England.
<i>cinereomicans</i> , of British lists, not Strobl.	
<i>H. rufopunctata</i> , Curran, Can. Ent. Vol. 58, p. 245 (1926).	Alberta.
<i>H. setosa</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 66 (1927).	England.
<i>H. subaura</i> , Collin, ibidem, Vol. 63, p. 63 (1927).	England.
<i>H. subpollinosa</i> , Collin, ibidem, Vol. 63, p. 26 (1927).	England.
<i>quadrifaria</i> , Collin, ibidem, Vol. 49, p. 106 (1913), not Strobl.	
<i>H. vector</i> , Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 454, f. 48, 49 (1923).	New Zealand.
<i>H. Woodi</i> , Collin, Ent. Mo. Mag. Vol. 63, p. 63 (1927).	England.

GENUS HILAREMPIS, BEZZI

- H. insularis*, Brethes, Revista Chilena Hist. Nat. Vol. 28, p. 106 (1924). Chile.
H. Moreirai, Brethes, ibidem, Vol. 28, p. 106 (1924). Chile.
H. nigra, Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 462, f. 63, 64 (1923). New Zealand.

GENUS ITEAPHILA, MACQUART

- I. curva*, Curran, Can. Ent. Vol. 57, p. 24 (1925). Labrador.

GENUS LEPTOPEZA, MEIGEN

- L. fulvescens*, Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 448 (1923). New Zealand.
flavescens, Miller, ibidem, Vol. 54, f. 34-36 (1923).
L. nigra, Miller, ibidem, Vol. 54, p. 447, f. 29-33 (1923). New Zealand.

GENUS OROPEZELLA, COLLIN

- Oropezella**, Collin, Ent. Mo. Mag. Vol. 62, p. 214 (1926).
Leptometopiella, nov. gen., antea, p. 70.

GENUS PLATYPALPUS, MACQUART

- P. agilellus*, Collin, Ent. Mo. Mag. Vol. 62, p. 153 [1926] (*Tachydromia*). England.
P. albifacies, Collin, ibidem, Vol. 62, p. 156 [1926] (*Tachydromia*). England.
P. aristatus, Collin, ibidem, Vol. 62, p. 157 [1926] (*Tachydromia*). England.
P. aurantiacus, Collin, ibidem, Vol. 62, p. 152 [1926] (*Tachydromia*). England.
P. calcaratus, Collin, ibidem, Vol. 62, p. 155 [1926] (*Tachydromia*). England.
P. Carteri, Collin, ibidem, Vol. 62, p. 158 [1926] (*Tachydromia*). England.
P. ceylonensis, nov. nom. Ceylon.
monticolus, Senior-White (not Melander), Spol. Zeyl. Colombo, Vol. 12, p. 399 [1924] (*Tachydromia*).
P. clarandus, Collin, Ent. Mo. Mag. Vol. 62, p. 157 [1926] (*Tachydromia*). England.
P. coarctatus, Collin, ibidem, Vol. 62, p. 158 [1926] (*Tachydromia*). N. Europe.
flavipes, Zetterstedt and Lundbeck, part.
P. engadinicus, Mik, var. *infuscatus*, Oldenberg (not Meigen), Entom. Mitteil. Berlin, Vol. 13, p. 88 [1924] (*Tachydromia*). Alps.
P. eximus, Oldenberg, ibidem, Vol. 13, p. 84 [1924] (*Tachydromia*). C. Europe.
P. extricatus, Collin, Ent. Mo. Mag. Vol. 62, p. 188 [1926] (*Tachydromia*). England.
P. incertus, Collin, ibidem, Vol. 62, p. 152 [1926] (*Tachydromia*). England.
P. infectus, Collin, ibidem, Vol. 62, p. 157 [1926] (*Tachydromia*). England.
P. ingenuus, Collin, ibidem, Vol. 62, p. 155 [1926] (*Tachydromia*). England.
P. interstinctus, Collin, ibidem, Vol. 62, p. 158 [1926] (*Tachydromia*). N. Europe.
flavipes, Zetterstedt and Lundbeck, part.
P. latistrigatus, Meijere, Tijdschr. v. Ent. Vol. 67, Suppl. p. 16 [1924] (*Tachydromia*). Sumatra.
P. luteolus, Collin, Ent. Mo. Mag. Vol. 62, p. 186 [1926] (*Tachydromia*). England.
P. nanus, Oldenberg, Deutsche Ent. Zeitschr. 1924, p. 234 [1924] (*Tachydromia*). C. Europe.
P. optivus, Collin, Ent. Mo. Mag. Vol. 62, p. 157 [1926] (*Tachydromia*). England.
P. pallidicornis, Collin, ibidem, Vol. 62, p. 186 [1926] (*Tachydromia*). N. W. Europe.
albicornis, Zetterstedt, part, and Lundbeck.

DIPTERA

<i>P. parvicauda</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 154 [1926] (<i>Tachydromia</i>).	England.
<i>P. politus</i> , Collin (not <i>politus</i> , nov. sp., antea, p. 333 (1); ibidem, Vol. 62, p. 155 [1926] (<i>Tachydromia</i>).	England.
<i>P. praecinctus</i> , Collin, ibidem, Vol. 62, p. 159 [1926] (<i>Tachydromia</i>).	England.
<i>P. stigma</i> , Collin, ibidem, Vol. 62, p. 188 [1926] (<i>Tachydromia</i>).	England.
<i>P. subtilis</i> , Collin, ibidem, Vol. 62, p. 188 [1926] (<i>Tachydromia</i>).	England.
<i>P. sylvicola</i> , Collin, ibidem, Vol. 62, p. 153 [1926] (<i>Tachydromia</i>).	England.
<i>P. tantalus</i> , Collin, ibidem, Vol. 62, p. 158 [1926] (<i>Tachydromia</i>).	England.
<i>P. Verralli</i> , Collin, ibidem, Vol. 62, p. 185 [1926] (<i>Tachydromia</i>).	England.
<i>P. zeylanicus</i> , Senior-White, Mem. Dept. Agric. India, Vol. 7, p. 148 [1922]	Ceylon.

GENUS RAGAS, WALKER

R. setosa, Beazzi, Boll. Mus. Zool. Torino, Vol. 39, N. S. (18), p. 14 (1924). Cyrenaica.

GENUS RHAMPHOMYIA, MEIGEN

Aclonempis, Collin, Ent. Mo. Mag. Vol. 62, p. 216 (1926) (2).

Amydroneura, Collin, ibidem, Vol. 62, p. 216 (1926) (3).

<i>R. (Pararhamphomyia) albatarsis</i> , Collin, ibidem, Vol. 62, p. 216 (1926).	England.
<i>R. (Aclonempis) albohirta</i> , Collin, ibidem, Vol. 62, p. 218 (1926).	N. W. Europe.
? <i>eupterota</i> , Frey, not Loew.	
<i>R. armipes</i> , Sack, Rept. Norweg. Exped. Nov. Zeml. No. 15, p. 6, fig. (1923).	Nova Zemlya.
<i>R. brevipila</i> , Oldenberg, Deutsche Ent. Zeitschr. 1922, p. 342 (1922).	Alps.
<i>R. (Lundstroemiella) brevistylata</i> , Oldenberg, Konowia, Vol. 6, p. 12 (1927).	C. Europe.
<i>R. (Holoclera) caliginosa</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 218 (1926).	England.
<i>R. claripennis</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 339 (1922).	Alps.
<i>R. (Pararhamphomyia) cibrata</i> , Oldenberg, Konowia, Vol. 6, p. 20 (1927).	Lapland.
<i>R. (Lundstroemiella) dalmatica</i> , Oldenberg, ibidem, Vol. 6, p. 10 (1927).	Dalmatia.
<i>R. (Lundstroemiella) Dudai</i> , Oldenberg, ibidem, Vol. 6, p. 16 (1927).	S. Europe.
<i>R. engadinica</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 344 (1922).	Alps.
<i>R. (s. str.) hercynica</i> , Oldenberg, Konowia, Vol. 6, p. 28 (1927).	S. Europe.
<i>R. hirsuta</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 346 (1922).	Alps.
<i>R. (Amydroneura) hirsutipes</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 219 (1926).	England.
<i>R. hirtimana</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 340 (1922).	Alps.
<i>R. (Lundstroemiella) Kertessi</i> , Oldenberg, Konowia, Vol. 6, p. 18 (1927).	Hungary.
<i>R. (Holoclera) lamellata</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 218 (1926).	England.
<i>R. (Pararhamphomyia) micropyga</i> , Collin, ibidem, Vol. 62, p. 217 (1926).	England.
<i>R. minor</i> , Oldenberg, Deutsche Ent. Zeitschr. p. 340 (1922).	Alps.
<i>R. (Pararhamphomyia) murina</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 216 (1926).	England.
<i>R. (Pararhamphomyia) nudipes</i> , Oldenberg, Konowia, Vol. 6, p. 25 (1927).	Germany.
<i>R. opacithorax</i> , Malloch, No. Am. Fauna, No. 46, p. 185 (1923).	Pribilof Islands.

(1) *P. politus*, nov. sp., antea, p. 333, may be renamed *politellus*, nov. sp.

(2) **Aclonempis**, Collin, subgenus of *Rhamphomyia*. Type *R. albohirta*, Collin; including *R. eupterota*, Loew, *galactoptera*, Strobl, *longipes*, Meigen, *minor*, Oldenberg, *nox*, Oldenberg, and *umbripes*, Becker. Labella long, narrow and very short-haired; both sides of prosternum and on episterna of prothorax with hairs; no distinct bristle in comb at tip of hind tibiae behind.

(3) **Amydroneura**, Collin, subgenus of *Rhamphomyia*. Type *R. erythrophthalma*, Meigen; including *R. bipila*, Strobl, *crassicauda*, Strobl, *gibba*, Fallen, *hirsutipes*, Collin, *pseudogibba*, Strobl and *serotina*, Oldenberg. Discal cell elongate, with the vein forming its anterior margin very indistinct, often visible only as a slight depression on the wing-surface.

<i>R. Porteri</i> , Brethes, Revista Chilena Hist. Nat. Vol. 28, p. 106 (1924).	Chile.
<i>R. (Lundstroemiella) rupestris</i> , Oldenberg, Konowia, Vol. 6, p. 4 (1927).	Alps.
<i>R. (Lundstroemiella) subalpina</i> , Oldenberg, ibidem, Vol. 6, p. 11 (1927).	S. Europe.
<i>R. (s. str.) subcinerascens</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 218 (1926).	England.
<i>R. (s. str.) sulcatella</i> , Collin, ibidem, Vol. 62, p. 217 (1926). <i>sulcata</i> , Meijere, Tijd. 1918, p. 135.	N. W. Europe.
<i>R. (Holoclera) trigemina</i> , Oldenberg, Konowia, Vol. 6, p. 19 (1927).	Germany.

GENUS STILPON, LOEW

<i>S. nubila</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 149 (1926).	N. W. Europe.
<i>lunata</i> , Walker, pl. 5.	
<i>graminum</i> , var. B, Fallen and Zetterstedt.	

GENUS SYNECHES, WALKER

<i>S. Jardinei</i> , Senior-White, Spol. Zeyl. Colombo, Vol. 12, p. 395 (1924).	Ceylon.
<i>S. maculithorax</i> , Senior-White, ibidem, Vol. 12, p. 397 (1924).	Ceylon.
<i>S. peradeniyæ</i> , Senior-White, ibidem, Vol. 12, p. 396 (1924).	Ceylon.

GENUS TACHYDROMIA, MEIGEN

<i>T. Halidayi</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 150 [1926] (<i>Tachista</i>). <i>morio</i> , Walker, not Zetterstedt.	England.
<i>T. halterata</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 151 [1926] (<i>Tachista</i>).	England.
<i>T. Woodi</i> , Collin, ibidem, Vol. 62, p. 151 [1926] (<i>Tachista</i>).	England.

GENUS TRICHINA, MEIGEN

<i>T. bilobata</i> , Collin, Ent. Mo. Mag. Vol. 62, p. 213 (1926).	British Isles; C. Europe.
--	---------------------------

GENUS TRICHOPEZA, RONDANI

<i>T. longipenna</i> , Miller, Trans. Proc. N. Zeal. Inst. Vol. 54, p. 446, f. 20-28 (1923).	New Zealand.
--	--------------

CORRIGENDA

Oldenberg, 1924, published the results of an examination of von Roser's collection, which alters the nomenclature of various species, as follows :

Hilara albiventris Roser is not *heterogastra* Nowicki, but is the species later described by Wood as *albocingulata*.

Hilara ferruginea Roser (nom. nudum) is *thoracica* Macquart, which is stated to be the same as *flava* Schiner.

Hilara geniculata Roser becomes *litorea* Fallen.

Hilara albitarsis Roser is the same as *Braueri* Strobl.

Hilara albipennis Roser is the same as *niveipennis* Zetterstedt, which it antedates.

Empis pallens Roser is not *æqualis* Loew, but is *parvula* Egger, which it antedates.

Bicellaria aterrima Roser in part is probably *spuria* Fallen, and in part is *Atelestus pulicarius* Fallen, of the Platypezidæ.

Bicellaria tibialis Roser is *Atelestus pulicarius* Fallen.

Drapetis pygmaea Roser is not *assimilis* Fallen, but is probably *pusilla* Loew.

Tachydromia nigromaculata Roser becomes *Drapetis (Elaphropeza) ephippiata* Fallen.

Tachyptesa costalis Roser belongs to the genus *Tachydromia* (*Tachista*).

Tachyptesa brevipennis Roser belongs to *Tachydromia*, and is the same as the later-described *microptera* Loew.

Tachyptesa obscura Roser is not *Wirthemi* Zetterstedt, but belongs to *Platypalpus*, being synonymous with *leucochæta* Becker.

Platypalpus ruficornis Roser is the same as *thoracicus* Lundbeck.

Collin, in the Entomologists' Monthly Magazine, 1926-1927, is issuing a cursory review of British Empididæ, which contains many changes in nomenclature, largely the result of examinations of the type collections of Fallen, Meigen, Zetterstedt, Curtis, Walker and Lundbeck. Collin's review is presumably not completed at the time the final proofs of this fascicle are being returned, so the student of this family is referred to forthcoming issues of the magazine for notes on the sub-families Clinoceratinæ and Hybotinæ. A summary of Collin's findings so far as published is presented herewith:

Platypalpus ambiguus Macquart, on which Bigot founded the genus *Crossopalpus*, is the same species as *Drapetis flexuosa* Loew, instead of being *Drapetis flavipes* Macquart, as generally supposed. The subgenus *Eudrapetis*, Melander, therefore, falls into synonymy with *Crossopalpus* as a subgenus of *Drapetis*.

Drapetis (Crossopalpus) curvipes Meigen is not *assimilis* Fallen, but includes *aterrima* Curtis, *nigra* Fallen and Zetterstedt, but not Meigen, *moriella* Zetterstedt, *picipes* Zetterstedt and *setigera* Lundbeck, but not Loew.

Drapetis (Crossopalpus) nigritella Zetterstedt is not *aterrima* Curtis, but includes *aterrima* Lundbeck and *nervosa* Loew.

Drapetis (Crossopalpus) minima Meigen is a valid species, not the same as *assimilis* Fallen, but *minima* Zetterstedt is *assimilis* Fallen, as is also *exilis* Zetterstedt of the Insecta Lapponica.

Drapetis exilis Meigen includes *pusilla* Loew and the female of *nigripes* Zetterstedt.

Platypalpus minutus Meigen is the same as *exiguus* Meigen, but the *exiguus* of the British lists is *niger* Meigen.

Platypalpus fasciatus Lundbeck, not Meigen, is *cursitans* Fabricius, and *cursitans* Lundbeck is *laticinctus* Walker.

Platypalpus glaber Meigen is the same as *luteus* Meigen.

Platypalpus infuscatus Meigen becomes *annulipes* Meigen.

Platypalpus ecalceatus Zetterstedt is a valid species, larger than *calceatus* Meigen.

Bicellaria melana Haliday is synonymous with *spuria* Fallen, but *sulcata* Zetterstedt is distinct from *spuria* Fallen.

Œdalea pallipes Zetterstedt is transferred to the genus *Trichina*.

Euthyneura Myrtilli Macquart includes *consobrina* Zetterstedt and *rostrata* Zetterstedt, which were placed in *Anthepiscopus*, ante a, p. 107.

Microphorus crassipes Macquart is distinct from *anomalus* Meigen.

Rhamphomyia sulcata Meigen, *rugicollis* Meigen, and *propinqua* Meijere are the same.

Empis barbata Macquart is a *Rhamphomyia*, and is the same as *bicolor* Macquart 1823, not 1827, and *pennata* Macquart.

Rhamphomyia plumipes Frey, Zetterstedt, Lundbeck, etc., is *geniculata* Meigen.

Rhamphomyia aethiops Zetterstedt is distinct from *caudata* Zetterstedt, but *longestylata* Frey is the same as *caudata* Zetterstedt.

Rhamphomyia attenuata Frey is *ignobilis* Zetterstedt.

Rhamphomyia villosa Zetterstedt, *morio* Zetterstedt, and *fumipennis* Zetterstedt are one species.

Empis vernalis of Meigen, Zetterstedt, etc. is *pennaria* Fallen, but the *pennaria* of the British lists is *nuntia* Meigen.

Empis (Pachymeria) picipes Meigen includes *brevicornis* Loew and *maculipes* Zetterstedt.

Empis (Pachymeria) scotica Curtis (1824) is the same as *palparis* Egger (1860).

Empis (Pachymeria) Erberi Nowicki becomes *tumida* Meigen.

Empis lamellicornis ♂ described by Lundbeck is *gymnopoda* Bezzi.

Empis dasythrix Meijere is *plumipes* Zetterstedt, as is also *decora* of Lundbeck.

Empis confluenta Becker is *albibennis* Meigen.

Hilara lasiopyga Lundbeck is *manicata* Meigen.

Hilara carinthiaca Becker is *fuscipes* Fabricius.

Hilara intermedia Fallen is not the same as *fuscipes* Fabricius, but is the species generally known as *pubipes* Loew.

Hilara pinetorum Zetterstedt is *clypeata* Meigen.

Hilara bivittata Strobl becomes *longevittata* Zetterstedt.

Hilara galactoptera Strobl is a distinct species from *pseudosartrix* Strobl.

Chelifera melanocephala Haliday (1833) and Mik is the same as *trapezina* Zetterstedt (1838), but Lundbeck's *melanocephala* is *flarella* Zetterstedt.

Hemerodromia albicornis of Gercke, and possibly of Walker, is *raptoria* Meigen.

BIBLIOGRAPHY

- Adams, C. F., Diptera Africana. Part 1 (Kansas Univ. Sc. Bull. Vol. 3, p. 147-208 [1905]).
 Adams, F. C., *Rhamphomyia tenuirostris* Fallen, taken in New Forest (Ent. M. Mag. London (2), Vol. 16, p. 94 [1905]).
 Adolph, E., Die Dipterenflügel, ihr Schema und ihr Ableitung (Nova Acta Akad. Naturf. Halle, Vol. 47, p. 271-314, pl. 1-4 [1885]).
 Aldrich, J. M., Catalogue of North American Diptera. Empididæ, p. 310-333 (Smithsonian Miscell. Coll. Vol. 46, publ. 1444, 680 p. [1905]).
 Aldrich, J. M. & Turley, L. A., A balloon-making fly (Amer. Naturalist, 1889, p. 809-812 [1889]).
 Arias, J., Description preliminar de un nuevo Empido de Espana (Bol. Soc. Esp. Hist. Nat. Vol. 19, p. 479-481 [1919]).
 Arribalzaga, F. L., Argentine Empididæ (El Naturalista Argentino, Buenos Aires, Vol. 1, p. 292-294 [1879]).
 Barbut, J., Genera Insectorum of Linnæus (London, 1781, éd. 2 [1783]).
 Becker, E., Zur Kenntnis der Mundtheile der Dipteren (Denkschr. Akad. Wiss. Wien, Vol. 45, p. 123-162, pl. 4 [1882]).
 Becker, Th. Beiträge zur Kenntnis der Dipteren-Fauna von St. Moritz (Berl. Ent. Zeitschr. Vol. 31, p. 93-141 [1887]).
Hilara sartor, nov. sp., und ihr Schlier (Berl. Ent. Zeitschr. Vol. 32, p. 7-12 [1888]).
 Neue Dipteren aus Dalmatien (Berl. Ent. Zeitschr. Vol. 33, p. 335-346 [1889]).
 Altes und Neues aus des Schweiz (Wien. Ent. Zeit. Vol. 8, p. 73-84, pl. 1 [1889]).
 Berichtigung : *Symbalophthalmus*, n. n. for *Macroptera* (Wien. Ent. Zeit. Vol. 8, p. 285 [1889]).

- Einige Bemerkungen zu Herrn J. M. F. Bigot's Classificirung der Empiden (Wien. Ent. Zeit. Vol. 9, p. 32-35 [1890]).
- Altes und Neues aus Tyrol und Salzburg (Wien. Ent. Zeit. Vol. 9, p. 65-70 [1890]).
- Neues aus Süd-Tyrol und Steiermark (Wien. Ent. Zeit. Vol. 10, p. 281-288, pl. 3 [1891]).
- Neues aus der Schweiz (Wien. Ent. Zeit. Vol. 10, p. 289-296 [1891]).
- Berichtigung zu meinen Dipterologischen Beiträge im Heft 9, 1891, der Wiener Entomologischen Zeitung : Neues aus Süd-Tyrol und Steiermark (Wien. Ent. Zeit. Vol. 11, p. 125, 126 [1892]).
- Bemerkung zu *Hilara longicornis* Strobl (Wien. Ent. Zeit. Vol. 13, p. 156-159 [1894]).
- Beiträge zur Dipteren-Fauna Siberiens (Acta Soc. Sc. Fenn. Helsingfors, Vol. 26 (9), p. 1-66, 2 pl. [1900]).
- Aegyptische Dipteren (Mitteil. Zool. Mus. Berlin, Vol. 2, p. 1-66 [1902]).
- Zur Kenntnis der Dipteren von Central-Asien (Ann. Mus. Zool. Acad. Sc. St. Petersb. Vol. 12, p. 253-317 [1907]).
- Die Ergebnisse meiner Dipterologischen Frühjahrstrreise nach Algier und Tunis, 1906 (Zeitschr. Hym. Dipt. Vol. 7, p. 33-61, 97-128, 225-256, 369-407 [1907]).
- Dipteren der Kanarischen Inseln (Mitteil. Zool. Mus. Berlin, Vol. 4, p. 1-180, 4 pl. [1908]).
- Dipteren der Insel Madeira (Mitteil. Zool. Mus. Berlin, Vol. 4, p. 181-206 [1908]).
- Microphorus* Macquart und seine nächsten Verwandten (Wien. Ent. Zeit. Vol. 28, p. 25-28 [1909]).
- Collections recueillies par M. Maurice de Rothschild dans l'Afrique orientale anglaise (Bull. Mus. Hist. Nat. Paris, 1909, p. 113-121 [1909]).
- Voyage de M. Maurice de Rothschild en Ethiopie et dans l'Afrique centrale (Ann. Soc. Ent. France, Vol. 79, p. 22-30 [1910]).
- Dipterologische Sammelreise nach Korsika (Deutsche Ent. Zeitschr. 1910, p. 635-665 [1910]).
- Die Loew'schen Typen in der Rosenhauerschen Dipteren-Sammlung (Wien. Ent. Zeit. Vol. 30, p. 71-76 [1911]).
- Genera Bombyliidarum (Ann. Mus. Zool. Acad. Sc. St. Petersb. Vol. 17, p. 421-502 [1913]).
- Persische Dipteren von den Expeditionen des Herrn N. Zarudny, 1898 und 1901 (Ann. Mus. Zool. Acad. Sc. St. Petersb. Vol. 17, p. 503-654 [1913]).
- Dipteren aus Marokko (Ann. Mus. Zool. Acad. Sc. St. Petersb. Vol. 18, p. 62-95 [1913]).
- Diptères nouveaux d'Afrique orientale (Ann. Soc. Ent. France Vol. 83, p. 120-130 [1914]).
- Zur Kenntnis der Dipteren vom Arktischen West-Siberien (Mém. Acad. Sc. Petrograd (8), Vol. 28 (7), p. 1-67, pl. 11-13 [1915]).
- Voyage de Ch. Alluaud et R. Jeannel en Afrique Orientale 1911-1912. Résultats scientifiques. Insectes Diptères, 5, Diptera Brachycera (1), Paris, 147 and 190 p. (1915).
- Diptères, Brachycères (Mission de l'Arc Méridien en Amérique du Sud, Paris, Vol. 10, p. 163-215, pl. 14-17 [1919]).
- Voyage de M. le Baron Maurice de Rothschild en Ethiopie et en Afrique Orientale Anglaise. Diptera, p. 796-836, 2 pl., Paris (1922).
- Beling, T., Beitrag zur Metamorphose zweiflügiger Insekten (Arch. Naturg. Berlin, Vol. 41, Pt. 1, p. 31-57 [1875]).
- Beitrag zur Metamorphose zweiflügiger Insekten aus der Familien Tabanidæ, Leptidæ, Asilidæ, Empidæ, Dolichopidæ und Syrphidæ (Arch. Naturg. Berlin, Vol. 48, p. 186-240 [1882]).
- Beitrag zur Metamorphose einiger zweiflügiger Insekten aus der Familien Tabanidæ, Empidæ und Syrphidæ (Verh. Zool.-bot. Ges. Wien, Vol. 38, p. 1-4 [1888]).

- Bellardi, L., Saggio di Ditterologia Messicana, parte 2 (Mem. Accad. Sc. Torin. Vol. 21, p. 103-225, 3 pl. [1861]).
- Berendt, G. C., Die im Bernstein befindlichen organischen Reste der Vorwelt (Berlin, 2 vols., 1845 and 1856).
- Bethune, C. J. S., Insects of the Northern parts of British America (Canad. Ent. Vol. 13, p. 165 [1881]).
- Beutenmueller, W., Types of Diptera in the Collection of the American Museum of Natural History (Bull. Amer. Mus. Nat. Hist. Vol. 20, p. 87-99 [1904]).
- A new Empid from the Black Mountains, North Carolina (Insecutor Inscitiae Menstruus, Vol. 1, p. 130 [1913]).
- Bezzi, M., I Ditteri del Trentino, saggio di un elenco delle specie di Ditteri osservati nel Trentino (Atti Soc. Veneto-Trentina Sci. Nat. Padova (2) Vol. 1, fasc. 1, Empididae, p. 257-263 [1892]). (Lists 75 Empididae, including several undescribed species named by Becker.)
- Contribuzioni alla fauna ditterologica italiana, 1, Ditteri della Calabria (Bull. Soc. Ent. Ital. Vol. 27, p. 39-78 [1895]).
- Rhamphomyia heterochroma*, nova Dipterorum species ex Hungaria (Termes. Fuzetek, Budapest, Vol. 21, p. 439, 440 [1898]).
- Contribuzioni alla fauna ditterologica italiana, 2, Ditteri delle Marche e degli Abruzzi (Bull. Soc. Ent. Ital. Vol. 30, p. 121-164 [1899]).
- De nova dipterorum specie faunæ hungaricæ pertinente (Termes. Fuzetek, Budapest, Vol. 23, p. 251, 252 [1900]).
- Contribuzioni alla fauna ditterologica italiana, 2, Ditteri della Marche e degli Abruzzi, Seconda continuazione (Bull. Soc. Ent. Ital. Vol. 32, p. 77-102 [1900]).
- Sulla presenza del genere *Chionea* in Italia, e la riduzione delle ali nei ditteri (Rend. Instit. Lombardo di Scienze, Milano (2), Vol. 33, p. 16 [1900]).
- Neue Namen für einige Dipteren-Gattungen (Zeitschr. Hym. Dipt. Vol. 3, p. 191, 192 [1902]).
- Katalog der Palæarctischen Dipteren, Vol. 2; Orthorrhapha Brachycera. Empididae, p. 218-289 (Budapest [1903]).
- Alcune notizie sui ditteri cavernicoli (Rivista Ital. Speleologia, Vol. 1, p. 8-16 [1903]).
- Empididae novæ palæarcticæ ex Museo nationali hungarico (Ann. Mus. Hungar. Vol. 2, p. 198-202 [1904]).
- Empididae Indo-australiani raccolti dal Signor L. Biro (Ann. Mus. Hungar. Vol. 2, p. 320-361 [1904]).
- Verzeichniss des bis jetzt bekannten Arten der dipterengattung *Drapetis* Meigen (Wien. Ent. Zeit. Vol. 23, p. 143-146 [1904]).
- Clinoceræ tres novæ ex Europa (Ann. Mus. Hungar. Vol. 3, p. 362-365 [1905]).
- Empididae neotropicæ musei nationalis hungarici (Ann. Mus. Hungar. Vol. 3, p. 424-460 [1905]).
- Ditteri Eritrei, Vol. 1 (Catalogue of African Empididae, p. 294, 295) (Bull. Soc. Ent. Ital. Vol. 37, p. 195-304 [1906]).
- Leptidae et Empididae in Insula Formosa a clar. H. Sauter collectæ (Ann. Mus. Hungar. Vol. 5, p. 564-568 [1907]).
- Empididae : in L. Schultze, Zool. u. anthrop. Ergebnisse e. Forschungsreise in Südafrika, Bd. 1, Lfg. 1 (Denkschr. Med. Nat. Ges. Jena, Vol. 13, p. 179-201 [1908]).
- Diagnoses d'espèces nouvelles de Diptères d'Afrique (Ann. Soc. Ent. Belg. Vol. 52, p. 374-388 [1908]).

- Rhagionidæ et Empididæ palæarcticæ novæ ex Museo Nationali Hungarico (Ann. Mus. Hungar. Vol. 6, p. 389-396 [1908]).
- Einige neue palæarktische Empis-Arten (Deutsche Ent. Zeitschr. 1909, Beiheft, p. 85-103 [1909]).
- Eine neu Empidide aus Paraguay (Wien. Ent. Zeit. Vol. 28, p. 319-322 [1909]).
- Diptera syriaca et ægyptia a cl. P. Béraud S. J. collecta (Broteria Lisboa : Revista S. Fiel, Vol. 8, p. 37-67, pl. 9 [1909]).
- Beiträge zur Kenntniss der südamerikanischen Dipterenfauna, fam. Empididæ (Nova Acta Akad. Naturf. Halle, Vol. 91, p. 293-407, pl. 1, 1909 [1910]).
- Biospeologica, 20; Diptères, première série, suivi d'un Appendice sur les Diptères cavernicoles des Balkans (Arch. Zool. Expér. Gén. Paris (5), Vol. 8, p. 1-87 [1911]).
- Rhagionidæ et Empididæ ex insula Formosa a clar. H. Sauter missæ (Ann. Nat. Hungar. Vol. 10, p. 442-496 [1912]).
- H. Sauter's Formosa-Ausbeute. Rhagionidæ et Empididæ (Suppl. Ent. Berlin, Vol. 3, p. 65-78 [1914]).
- Studi sulla ditterofauna nivale della Alpi italiane (Mem. Soc. Ital. Sc. Nat. Milan, Vol. 9, fasc. 1, p. 164 [1918]).
- Ditteri di Cirenaica (Atti Soc. Ital. Sc. Nat. Milano, Vol. 60, p. 432-443 [1921]).
- Materiali per lo studio della Fauna Tunisina (Ann. Mus. Civ. Storia Nat. Genova (3) Vol. 10, p. 97-139 [1922]).
- Ditteri di Cirenaica raccolti dal Prof. Alessandro Ghigi durante l'escursione organizzata dal Touring Club Italiano nel mese d'aprile 1920 (Atti Ital. Sc. Nat. Mus. Civ. Storia Nat. Milano, Vol. 60, p. 432-443 [1922]).
- Missione del Dr. E. Festa in Cirenaica, Pt. 11, Ditteri di Cirenaica (Boll. Mus. Zool. Anat. Comp. Torino, Vol 39 (1924). N. S., no. 18, 26 p. [1925]).
- Materiali per una Fauna dell'arcipelago Toscano, Pt. 17, Ditteri del Giglio (Ann. Mus. Civ. Storia Nat. Genova (3), Vol. 10, Pt. L, p. 291-354 [1925]).
- Bigot, J. M. F., Essai d'une classification générale et synoptique de l'ordre des Insectes Diptères (Ann. Soc. Ent. France (3), Vol. 5, p. 517-564 [1857]).
- Dipterorum aliquot nova genera (Revue Mag. Zool. Paris, Vol. 11, p. 305-315 [1859]).
2. *Enoplemnis*; 3. *Megacyllarus* (Bull. Soc. Ent. France (5), Vol. 10, p. 47 [1880]).
- Diptères nouveaux ou peu connus, XXVI., 17 (Ann. Soc. Ent. France (6), Vol. 1, p. 363-371 [1881]).
- Enoplemnis cinerea* (Ann. Soc. Ent. France (6), Vol. 2, Bull. p. 91 (1882); Bull. Soc. Ent. France, 1882 (9), p. 112 [1882]).
- Diptères nouveaux ou peu connus. Leptides (Bull. Soc. Ent. France, Vol. 12, p. 97-118 [1887]).
- Diagnoses de quelques espèces nouvelles de Diptères (Bull. Soc. Ent. France (6), Vol. 7, p. 139-142 [1887]).
- Diptères (Mission scient. du Cap Horn, Zool. Vol. 6, p. 1-45, pl. 1-4 [1888]).
- Note : *Heleodromia ochracea* referred to *Hilara* (Wien. Ent. Zeit. Vol. 7, p. 109 (1888); Ann. Soc. Ent. France (6), Vol. 8, Bull. p. 30 [1888]).
- Note rectificative concernant quelques Diptères du Cap Horn (Bull. Soc. Zool. France, Vol. 13, p. 101 [1888]).
- Diptères nouveaux ou peu connus, 3^{me} partie, XLII. Empidi (Ann. Soc. Ent. France (6), Vol. 9, p. 111-134 [1889]).
- Voyage de M. Ch. Allaud aux îles Canaries. Diptères (Bull. Soc. Zool. France, Vol. 16, p. 275-279 [1891]).
- Hilarimorpha nota Leptid (Bull. Soc. Zool. France, 1891, p. 15 [1891]).

- Billberg, G. J., *Enumeratio Insectorum in Museo Billberg* (Holmiæ. Gadel. 138 p. [1820]).
- Blanchard, E., *Histoire naturelle des Insectes*, Paris, 3 vol. (1840).
- Gay : *Historia física y política de Chile*, Zool. Dipt. in vol. 7 (1852).
- Bohemian, C. H., *Entomologiska antekningar under en resa i Södra Sverige*, 1851 (Vetensk. Acad. Handl. Stockholm, 1851, p. 53-210 [1852]).
- Entomologiska antekningar under en resa i norra Skåne och södra Halland år 1862* (Vetensk. Acad. Handl. Stockholm, Vol. 20, p. 57-85 [1863]).
- Spetsbergens Insekt-fauna (Vetensk. Akad. Handl. Stockholm, 1865, p. 563-577 [1865]).
- Boie, Fr., *Zur Entwicklungsgeschichte mehrerer Trypeta-Arten* (Stett. Ent. Zeit. Vol. 8, p. 326-331 [1847]).
- Boitard, P., *Manuel d'Entomologie*, 2 vol. Atlas avec 110 pl. Paris (1828).
- Nouveau Manuel complet d'Entomologie, 3 vol. et atlas, Paris (1843).
- Bonsdorff, E. J., *Finlands tvåvingade Insekter, forteckn. och beskr.* Helsingfors, 2 vol. (1861).
- Bouché, P. Fr., *Naturgeschichte der Insecten besonders in Hinsicht ihrer ersten Zustände als Larven und Puppen*. Berlin, 216 p. 10 pl. (1834).
- Brauer, Fr., *Kurze Characteristik der Dipteren-Larven* (Verh. Zool.-bot. Ges. Wien, Vol. 19, p. 843-852 [1869]).
- Vergleichenden Untersuchungen des Flügelgeäders der Dipteren nach Adolph's Theorie (Denkschr. Wissensch. Akad. Wien, Vol. 44, p. 59-110 [1882]).
- Systematische Studien auf Grundlage der Dipteren-Larven nebst einer Zusammenstellung von Beispielen aus der Literatur über dieselben und Beschreibung neuer Formen (Denkschr. Wiss. Akad. Wien, Vol. 47, p. 1-100, 5 pl. [1883]).
- Brethes, J., *Catalogo de los Dipteros de las Republicas del Plata* (Anal. Mus. Hist. Nat. Buenos-Aires, Vol. 16, p. 277-305 [1907]).
- Dipteros é Himenopteros de Mendoza (Anales Mus. Hist. Nat. Buenos-Aires, Vol. 19, p. 85-105 [1909]).
- Un nouvel Empididæ du Chili (Revista Chilena de Hist. Nat. Santiago, Vol. 20, p. 79, 80 [1916]).
- Cueillette d'Insectes du Rio Blanco (Revista Chilena de Hist. Nat. Santiago, Vol. 23, p. 40-44 [1920]).
- Sur quelques diptères Chiliens (Revista Chilena Hist. Nat. Buenos-Ayres, Vol. 28, p. 104-111 [1924]).
- Britton, W. E., *Check list of the insects of Connecticut* (Conn. Geol. & Nat. Hist. Surv. Bull. 31, Hartford, 397 p. [1920]).
- Brocher, F., Quelques observations sur un Diptère amphibia d'eau douce (*Clinocera barbatula* Mik) (Bull. Soc. Zool. Genève, Vol. 1, p. 115-117 [1908]).
- Métamorphoses de l'*Hemerodromia prectoria* Fallen (Ann. Biol. Lacustre, Bruxelles, Vol. 4, p. 44, 45 [1909]).
- Brullé, A., *Expédition Scientifique de Morée sous la direction de M. Bory de Saint-Vincent*, Paris, Vol. 3 (1832).
- Brunetti, E., *New Indian Empidæ* (Rec. Indian Mus. Vol. 9, Pt. 1, No. 2, p. 11-45 [1913]).
- Fauna of British India, etc., Vol. 1, Diptera Brachycera, London, 401 p. 4 pl. (1920).
- Carter, A. E. J., *Hybos culiciformis*, Fab., in Scotland (Ent. Monthly Mag. London, Vol. 47, p. 161 [1911]).
- On the *Hybos grossipes*, L., of the British List (Ent. Monthly Mag. London, Vol. 48, p. 59, 60 [1912]).

- Cockerell, T. D. A., Notes and observations : *Hilara viridis* (The Entomologist, London, Vol. 36, p. 49 [1903]).
- Miocene fossil insects (Proc. Acad. Nat. Sc. Philad. Vol. 46, p. 634-648 [1914]).
- Two Diptera of the Genus *Rhamphomyia* from Colorado (Canad. Ent. Vol. 48, p. 123 [1916]).
- Some American fossil insects (Proc. U. S. Nat. Mus. Vol. 51, No. 2146, p. 89-106 [1916]).
- Arthropods in Burmese amber (Amer. Journ. of Sc. Vol. 44, p. 360-368 [1917]).
- New tertiary insects (Proc. U. S. Nat. Mus. Vol. 52, p. 373-384, pl. 31 [1917]).
- Fossil insects (Ann. Ent. Soc. Amer. Vol. 10, p. 1-22 [1917]).
- The oldest mosquitoes (Nature, London, Vol. 103, p. 44 [1919]).
- Eocene insects from the Rocky Mountains (Proc. U. S. Nat. Mus. Vol. 57, p. 233-260 [1920]).
- Cole, F. R., Some Diptera of Laguna Beach (Report Laguna Marine Labor. Vol. 1, p. 150-162 [1912]).
- A revision of the North American two-winged flies of the family Therevidæ (Proc. U. S. Nat. Mus. Vol. 62, Art. 4, 140 p. 13 pl. [1923]).
- Notes on the Diptera of Laguna Beach, California (Jour. Ent. Zool., Claremont, 1925, p. 55-60 [1925]).
- Cole, F. R. & Lovett, A., L. An annotated list of the diptera of Oregon (Proc. Calif. Acad. Sc. San Francisco (4), Vol. 11, No. 15, p. 197-344 [1921]).
- Collin, J. E., Thirty additions to the list of British Diptera (Ent. Monthly Mag. London, Vol. 49, p. 104-106; 130-135 [1913]).
- Hormopeza oblitterata* Zetterstedt associated with *Melanophila acuminata* De Geer on burning pines in Berkshire (Ent. Monthly Mag. London, Vol. 54, p. 278 [1918]).
- Empidæ from the Seychelles (Ent. Monthly Mag. London, No. 699, p. 184-189 [1922]).
- Notes on the Empididæ (Diptera) with additions and corrections to the British List (Ent. Mo. Mag. Vol. 62, p. 146-159, 185-190, 213-219, 231-237 (1926); Vol. 63, p. 20-29, 61-67, 93-96 [1927]).
- Coquebert de Montbret, A. J., Illustrata Iconographa Insectorum, quae in Musaeis parisiniis observavit et lucem edidit Joh. Christ Fabricius, praemissis eiusdem descriptionibus, Parisiis (1790-1804).
- Coquillett, D. W., An Anomalous Empid (*Mythicomyia rileyi*) (Ent. News, Philad. 1893, p. 208-209 [1893]).
- Revision of the North American Empidæ (Proc. U. S. Nat. Mus. Vol. 18, 1895, p. 387-440 [1896]).
- Arthropoda of the Commander Islands : Diptera (Fur Seals and Fur-Seal Islands, Vol. 4, p. 341-346 [1899]).
- Papers from the Harriman Alaska Expedition, Vol. 9, Entomological Results (3), Diptera (Proc. Wash. Acad. Sc. Vol. 2, p. 389-464 (1900); Reprinted, Doubleday, New-York (1904); Reprinted, Smithsonian Publication, No. 1996 [1910]).
- New Diptera in the U. S. National Museum (Proc. U. S. Nat. Mus. Vol. 23, p. 593-618 [1901]).
- Roederoides juncta* (in Needham & Betten, Aquatic Insects of the Adirondacks, p. 585, 586, pl. 15, f. 5-8 [1901]).
- New Diptera from North America (Proc. U. S. Nat. Mus. Vol. 25, p. 83-126 [1902]).
- New Orthorrhaphous Diptera from Mexico and Texas (Journ. New York Ent. Soc. Vol. 10, p. 136-141 [1902]).
- The genera of the Dipterous family Empididæ, with notes and new species (Proc. Ent. Soc. Wash. Vol. 5, p. 245-272 [1903]).

- The genera of the Dipterous family Empididæ, addenda (Proc. Ent. Soc. Wash. Vol. 6, p. 51-52 [1904]).
- Reports on Californian and Nevadan Diptera (Invertebrata Pacifica, Diptera, Vol. 1, p. 17-39 [1905]).
- New genera and species of North American Diptera (Proc. Ent. Soc. Wash. Vol. 12, p. 124-131 [1910]).
- The Type-species of the North American Genera of Diptera (Proc. U. S. Nat. Mus. Vol. 37, p. 409-647 [1910]).
- Corti, E., Eine neue Art der Dipterengattung *Tachydromia* (Meigen), Loew (Wien. Ent. Zeit. Vol. 26, p. 101, 102 [1907]).
- Costa, A., Frammenti di Entomologia Napolitana, Articolo I, Nuove specie di Ditteri (Annal. Scientif. Napoli, Vol. 1, p. 76 [1854]).
- Contribuzione alla Fauna Ditterologica Italiana (Il Giambatt. Vico Napoli, Vol. 2, p. 438-460 [1857]).
- Dipterologische Mittheilungen (Wien. Ent. Monatschr. Vol. 3, p. 63, 64 [1859]).
- Crampton, G. C., Genitalia of male Diptera and Mecoptera (Trans. Amer. Ent. Soc. Vol. 48, p. 207-225, pl. 8-10 [1922]).
- Cresson, E. T. Jr., A revision of the genus *Mythicomyia* (Ent. News Philad. Vol. 26, p. 448-456 [1915]).
- Curran, C. H., Three new Diptera from Labrador (Can. Ent. Vol. 57, p. 24-26 [1925]).
- The species of *Hilara* occurring in Banff and vicinity (Can. Ent. Vol. 58, p. 245-249 [1926]).
- Curtis, J., British Entomology, London, 16 vol. 770 pl. (1823-1840) (Empididæ, Vol. 8, 1824).
- A guide to an arrangement of British Insects. London, 256 columns (1829) (118 spp. Empids).
- Observations on the natural history and economy of various Insects, etc., affecting the corn-crops, including their parasitic enemies (Journ. Agric. Soc. London, Vol. 6, p. 498, pl. N, f. 5 [1845]).
- Cuvier, G. L. C. D., Description de deux espèces nouvelles d'insectes (Millim. Magaz. Encyclop. Vol. 1, p. 205-207, pl. [1795]).
- Czizek, K., Neue Beiträge zur Dipterenfauna Mährens (Zeitschr. Mähr. Landesmus. Brünn. Vol. 7, p. 157-177 [1907]).
- Dahl, F. Die Gatting *Limosina* und die biocönotische Forschung (Sitzber. Ges. Naturf. Freunde, Berlin [1909]).
- Ueber die Fauna des Plagefenngebietes (Beiträge zur Naturdenkmalpflege, Vol. 3; Das Plagefenn bei Chorin. Berlin, p. 341-638 [1912]).
- Dahlbom, A. G., Anteckningar öfver Insekter, observerade på Gotland och i en del af Calmar Län. (Vet. Akad. Handl. Stockholm, 1850, p. 155-229 [1851]).
- Dale, C. W., History of Glanville's Wootton, Diptera, p. 239-293 (1878).
- De Geer, C., Mémoires pour servir à l'Histoire des Insectes, Stockholm, 7 vol. (1752-1778).
- De Villers, Ch. J., Caroli Linnæi Entomologia, London, 4 vol. (1789).
- Dufour, L., Recherches anatomiques et physiologiques sur les Diptères (Mém. Acad. Sc. Mathémat. des Savants étrangers, Paris, Vol. 11, p. 171-360, 11 pl. [1850]).
- Edwards, F. W., A remarkable case of venational teratology in Diptera (Ent. Monthly Mag. London, Vol. 50, p. 59, fig. [1914]).
- Egger, J., Dipterologische Beiträge (Verh. Zool.-bot. Ges. Wien, Vol. 10, p. 339-358 [1860]).
- Enderlein, G., Die Insekten des Antarkto-Archipalata-Gebietes (Vet. Akad. Handl. Stockholm, Vol. 48, No. 3, 170 p. [1912]).

- Engel, E., Ueber einige Dipteren, deren Vorkommen in der Mark nicht oder wenig bekannt ist, No. 4, *Hilara lugubris* (Ent. Nachr. Berlin, Vol. 12, p. 45-47 [1886]).
- Das Dipterengenuss *Atalanta* Meigen (*Clinocera* ol.) (Deutsche Ent. Zeitschr. 1918, p. 1-80, 197-268 [1918]).
- Eschscholtz, J. Fr., Entomographien, Berlin, 128 p. (1822).
- Eversmann, E., Diptera Wolgam fluvium inter et montes Uralensis observata (Bull. Soc. Nat. Moscou, Vol. 7, p. 420-432 [1834]).
- Fabricius, J. Chr., Systema Entomologiae, Flensburgi et Lipsiae (1775).
- Species Insectorum, Hamburg et Kiel, 2 vol. (1781).
 - Mantissa Insectorum, Hafniæ, 2 vol. (1787).
 - Entomologia Systematica, Hafniæ, 4 vol. (1792-1794).
 - Systema Antliatorum, Brunsvigae (1805).
- Fabricius, O., Fauna Groenlandica, Hafniæ et Lipsiae (1780).
- Fallen, C. F., Diptera Sueciæ, Empidiae sueciæ, p. 1-16 (1815); p. 17-34 (1816). Lundæ, 2 vol. (1814-1825).
- Fitch, A., First and second report on the noxious, beneficial and other Insects of the State of New York, Albany (1856).
- Förster, B., Die Insekten des plattigen Steinmergels von Brunstatt (Abh. Geol. Specialkarte von Elsass-Lothr. Vol. 3 [1891]).
- Fourcroy, A. Fr., Entomologia parisiensis, Paris, 2 vol. 544 p. (1785).
- Frey, R., För Finlands fauna nya Dipterer (Medd. Soc. Fauna Flora Fenn. Helsingfors, Vol. 32, p. 107-109 [1906]).
- Beiträge zur Kenntnis der Dipteren-Fauna Finlands, 1, *Hilara barbipes*, nov. sp. (Medd. Soc. Fauna Flora Fenn. Helsingfors, Vol. 33, p. 67, 68 [1907]).
- Uebersicht der finnischen Arten der Gattung *Tachydromia* Meigen (= *Platypalpus* Macquart) (Zeitschr. Hymen. Dipt. Vol. 7, p. 407-413 [1907]).
- Finlands Tachydromia-Arter (Medd. Soc. Fauna Flora Fenn. Helsingfors, Vol. 34, p. 20-21 [1908]).
- Mittheilungen über finnändische Dipteren, 1, Neue und seltene Arten; 2, Uebersicht der Gruppen der Gattung *Rhamphomyia* Meigen (Acta Soc. Fauna Flora Fenn. Helsingfors, Vol. 31 (9), p. 1-24 [1908]).
- Zur Kenntnis der Dipterenfauna Finlands, 2, Empididae (Acta Soc. Fauna Flora Fenn. Helsingfors, Vol. 37 (3), p. 1-89, pl. 3 [1913]).
- Beitrag zur Kenntnis der Dipterenfauna des Nördl. Europäischen Russlands (Acta Soc. Fauna Flora Fenn. Helsingfors, Vol. 37, No. 10, p. 6-20 [1913]).
- Verzeichnis von Dr. E. Strand in Norwegen gesammelter Diptera Brachycera (Nyt Mag. Naturvidensk. Vol. 51, p. 320-322 [1914]).
- Nya svenska Empidider (Ent. Tidskr. 1914, p. 78-80 [1914]).
- Dipteren Brachycera aus dem arktischen Kustengegenden Sibiriens (Mém. Acad. Sci. Russie, St. Petersburg, Phys.-Math. (8), Vol. 29, No. 10, 35 p. 2 pl. [1915]).
- Dipteren aus dem Sarekgebiet. Diptera brachycera (Naturwiss. Unters. des Sarekgebirges in Schwedisch-Lappland gel. von. Dr. Axel Hamberg (with B. Poppius & C. Lundström). Bd. 4, Zoologie, Lief 6, Stockholm [1916]).
- Ein Beitrag zur Kenntnis der Dipterenfauna Ceylons (Oefv. Vet. Soc. Stockholm, Vol. 59, A, No. 20, p. 1-36 [1917]).

- Beitrag zur Kenntnis der Dipterenfauna des Nördl-Europäischen Russlands, 2, Dipteren aus Archangelsk. (Acta Soc. Fauna Flora Fenn. Helsingfors, Vol. 46, No. 2, 32 p. [1918]).
- Vorarbeiten zu einer Monographie der Gattung *Rhamphomyia* Meigen (Notulæ Entomol. Vol. 2, p. 1-10, 34-45, 65-77 [1922]).
- Alfons Dampf : Zur Kenntnis der Estländischen Moorfauna [2 Beitrag] (Sitzber. Naturf. Ges. Univ. Dorpat, Vol. 31 [1924]).
- Geoffroy, E. L., Histoire abrégée des Insectes qui se trouvent aux environs de Paris, Paris (1762).
- Gercke, G., Dipterologische Miscellaneen (Wien. Ent. Zeit. Vol. 5, p. 161-168, pl. 2 [1886]).
- Germar, E. F., Fauna Insectorum Europæ, fasc. 3-24. Empididæ, in fasc. 12, Halæ (1829).
Insectorum protogaeæ specimen sistens insecta carbonum fossilium cura E. F. Germar, in fasc. 19 (1837).
- Giard, A., Un insecte imitateur du *Bibio marci* (*Empis ciliata*) (Bull. Scientif Dépt du Nord, Lille, Vol. 5, p. 192-194 [1873]).
- Giard, M., Traité élémentaire d'Entomologie, Paris, Vol. 3, 1110 p. (1885).
- Giebel, Ch. G. A., Ein systematisches Verzeichniss aller in Deutschland und den angrenzenden Ländern verkommenden Petrefacten, Leipzig, 706 p. (1852).
- Die Insecten und Spinnen der Vorwelt, mit steter Berücksichtigung der lebenden Insecten und Spinnen. Leipzig, 529 p. (1856).
- Giglio-Tos, E., Mission scientifique de M. Ch. Alluaud aux Iles Séchelles. Diptères (Ann. Soc. Ent. France (7), Vol. 5, p. 353-368 [1895]).
- Gil, J., Estudio de un nuevo Taquidromino de España (Bol. Soc. Esp. Hist. Nat. Vol. 23, p. 150-154 [1923]).
- Gimmerthal, B. A., Observations de quelques nouvelles espèces de Diptères, accompagnées des recherches sur la métamorphose de quelques autres (Bull. Soc. Nat. Moscou, Vol. 7, p. 98-121 [1832]).
- Uebersicht der Zweiflügler Livlands und Curlands (Bull. Soc. Nat. Moscou, Vol. 15, p. 639-686 [1842]).
- Vierter Beitrag zu einer künftig bearbeitenden Dipterologie Russlands (Bull. Soc. Nat. Moscou, Vol. 20 (2), p. 140-208 [1847]).
- Girschner, E., Zur Biologie von *Hilara* (Ent. Nachr. Berlin, Vol. 15, p. 220-222 [1889]).
Beiträge zur Biologie von *Hilara* (Ent. Nachr. Berlin, Vol. 20, p. 61-64 [1894]).
Entgegnung auf Herrn Prof. Mik's Artikel in Nr. 10, Jahrg. 1894, dieser Zeitschrift (Ent. Nachr. Berlin, Vol. 20, p. 241-244 [1894]).
- Glover, T., Manuscript notes from my Journal. Diptera. Washington (1874).
- Gmelin, J. F., Linné : Systema naturæ, ed. 13, aucta, reformata cura Joa. Frid. Gmelin. Lipsiæ. 10 vol. (1788-1793) (Empididæ in Vol. 5).
- Greene, C. T., New species of *Mythicomyia* and its relationship with a new genus (Proc. Ent. Soc. Wash. Vol. 26, p. 60-64 [1924]).
- Grünberg, K., Die blutsaugenden Dipteren. Jena (1907).
Die Süßwasserfauna Deutschlands, herausgegeben von A. Brauer. Heft 2a. Diptera. Jena (1910).
- Guérin-Maneville, F. Ed., Iconographie du Règne Animal de G. Cuvier, Paris, 7 vol. Insecta, in Vol. 7 (1835).
- Haliday, A. H., Catalogue of diptera occurring about Holywood in Downshire (Ent. Monthly Mag. London, Vol. 1, p. 147-180 [1833]).

- Sendschreiben an C. A. Dohrn über die Dipteren der in London befindlichen Linneischen Sammlung (Stett. Ent. Zeit. Vol. 12, p. 131-145 [1851]).
- Hamm, A. H., Observations on *Empis livida* L. (Ent. Monthly Mag. London, Vol. 44, p. 181-184 [1908]).
- Observations on *Empis opaca* F. (Ent. Monthly Mag. London, Vol. 45, p. 132-134 [1909]).
- Further observations on the Empinæ (Ent. Monthly Mag. Vol. 45, p. 157-162 [1909]).
- Handlirsch, A., Beitrag zur Kenntnis des Gespinnstes von *Hilara sartrix* Becker (Verh. Zool-bot. Ges. Wien, Vol. 39, p. 623-626 [1889]).
- Die fossilen Insekten und die Phylogenie der rezenten Formen. Leipzig, 1470 p. 51 pl. (1906-1908).
- Canadian fossil Insects. Insects from the tertiary lake deposits of the southern interior of British Columbia (Canad. Geol. Surv. Contrib. to Canad. Palaeont. Ottawa, Vol. 2, Pt. 3, p. 93-129 [1910]).
- Hansen, H. J., Faunula Insectorum Færoeensis (Naturhist. Tidskr. (3), Vol. 13 [1880]).
- Harris, M., An exposition of English Insects. Edit. 3, 166 p. 50 pl. (1782).
- Heeger, E., Beiträge sur Naturgeschichte der Insecten, 6. Fortsetzung (Sitzber. Math. Nat. Cl. Akad. Wiss. Wien, Vol. 9, p. 774-781 [1852]).
- Heer, O., Ueber die fossilen Insekten von Aix in der Provence (Vierteljahrsschr. Naturf. Gesell. Zürich, Vol. 1, p. 1-40 [1856]).
- Herbst, J. Fr. W., Kurze Einleitung zur Kenntniss der Insecten für Ungeübte und Anfänger, Berlin, 3 Bde. 144 pl. (1784-87). Also as Vol. 6-8 of Borowski's Gemeinnützige Naturgeschichte des Thiereichs.
- Heyden, L. (von), Fossil Diptera of Rott (Palaeontographica, Vol. 17 [1870]).
- Holmgren, A. E., Bidrag till kännedomen om Beerens Eilands och Spetsbergens insekta fauna (Svensk. Akad. Handl. Stockholm Vol. 8 (5), p. 1-55 [1869]).
- Insekten fra Nordgrönland, samlade af Prof. A. E. Nordenskiöld år 1870 (Oefv. Vet. Akad. Förhandl. Stockholm, 1872, p. 97-105 [1872]).
- Illustrissimo viro A. E. Nordenskiöldio in patriam reduci salutem dicit plurimum nova species insectorum cura et labore A. E. Nordenskiöldii e Novaia Semlia coactorum descripsit. Holmiæ, 1880, 24 p. (1881).
- Insecta a viris doctissimis Nördenskiöld illum ducem sequentibus in insula Waigatsch et Novaja Semlia anno 1873 collecta (Ent. Tidskr. Vol. 4, p. 139-194 [1883]).
- Howard, L. O., A contribution to the study of the fauna of human excrement (Proc. Wash. Acad. Sc. Vol. 2, p. 541-604 [1900]).
- The Insect Book, New York, 429 p. (1901).
- Howlett, M., Note on the coupling of *Empis borealis* (Ent. Monthly Mag. London, Vol. 43, p. 229-232 [1907]).
- Hutton, F. W., Synopsis of the Diptera Brachycera of New Zealand (Trans. New Zeal. Inst. Wellington, Vol. 33, p. 1-95 [1901]).
- Jacobsen, Frz. V. S., *Rhamphomyia marginata* Meigen (Kröyer, Tidskr. Vol. 4, p. 211 [1842]).
- Jaroschewsky, W. A., Verzeichniss der zweiflügler welche vorzüglich in Charkow u. Umgebung gesammelt sind (Trudy Naturf. Ges. Kharkoff, Vol. 10, p. 1-49 [1876]).
- Zusätze zu der Verzeichniss der Dipteren Kharkoffs, nebst Verbreitung im Gebiete der europäischen Russlands (Trudy Naturf. Ges. Kharkoff, Vol. 11, p. 317-454 (1877); ibidem, Vol. 12 [1880]).

- Johnson, C. W., Catalog of the Insects of New Jersey, Diptera (Rept. N. J. State Bd. Agric. Vol. 29, suppl. 617-699 [1899]; Rept. N. J. State Mus. 1909, p. 703-814 [1910]), new ed. Insects of Florida (Bull. Am. Mus. Nat. Hist. Vol. 32, Art. 3, p. 37-90 [1913]). A revised list of the Diptera of Jamaica (Bull. Am. Mus. Nat. Hist. Vol. 41, Art. 8, p. 421-449 [1919]).
- Fauna of New England, Pt. 15, List of the Diptera (Occ. Pap. Boston Soc. Nat. Hist. Vol. 7, p. 1-326 [1925]).
- Kawall, H., Entomologische Notizen aus Kurland (Stettin. Ent. Zeit. Vol. 16, p. 227-232 [1855]).
- Kertész, K., Verzeichniss einiger, von L. Biro in Neu-Guinea und am Malayischen Archipel gesammelten Dipteren (Termes. Fuzetek. Vol. 22, p. 173-195 [1899]). Note on spinning Empidæ (Rovartani Lapok, Vol. 8, p. 43 [1901]). Catalogus dipterorum hucusque descriptorum (Budapest, Empididæ, Vol. 6, p. 1-169 [1909]). Egy különös élétmodú légynemröl. Subgenera of *Clinocera* and habits of the species (Rovartani Lapok, Budapest, Vol. 18, p. 65-68 [1911]).
- Kieffer, J. J., Beiträge zur Biologie und Morphologie der Dipteren (Illustr. Zeitschr. Ent. Vol. 5, p. 131-133 [1900]).
- Kirby, W., Fauna boreali-Americana or the Zoology of the Northern Parts of British Columbia, Vol. 4, London (1837).
- Kleine, R., Einige Bemerkungen über *Rhamphomyia sulcata* Fall. und *cinerascens* Meigen (Soc. Ent. Steglitz, Vol. 24, p. 65 [1909]). Zur Kenntnis der Diptera. 4. Larva of *Rhamphomyia platyptera* (Zeitschr. f. Naturw. Leipzig, Vol. 81, 1909, p. 188-196 [1910]).
- Kowarz, F., Beschreibung sechs neuer Dipteren-Arten (Verh. Zool.-bot. Ges. Wien, Vol. 17, p. 319-324 [1868]). Verzeichniss der in den Umgebungen von Herculesbad und Orsova gesammelten Dipteren (Verh. Zool.-bot. Ges. Wien, Vol. 23, p. 456, 457 [1873]).
- Kunckel d'Herculaïs, J., Recherches sur l'organisation et le développement des Diptères. Paris, 2 Vol. (1875-1881).
- Kuntze, A., Tabelle zum Bestimmen der Arten der Gattung *Empis* L. (Zeitschr. Hymen. Dipt. Vol. 6, p. 209-216; 297-304 [1906]; Vol. 7, p. 25-32; 155-160 [1907]). Dipterologische Sammelreise in Korsika (Deutsche Ent. Zeitschr. 1913, p. 544-552 [1913]).
- Lamarck, J., Histoire naturelle des Animaux sans Vertèbres. Paris, 7 Vol. (1815-1822).
- Latreille, P. A., in Dictionnaire d'Histoire naturelle de Déterville, Vol. 24. Paris (1804). Histoire naturelle, générale et particulière, des Crustacés et des Insectes. Paris, 14 vol. 1802-1805 (Empididæ in Vol. 14 [1804]).
- Genera Crustaceorum et Insectorum. Paris, 4 vol. 1806-1809 (Empididæ in Vol. 4 [1809]). Considérations générales sur l'ordre naturel des Animaux composant les classes des Crustacés, des Arachnides et des Insectes. Paris, 444 p. (1810).
- Lefebvre, A., Note sur l'*Empis platyptera*, extrait d'une lettre adressée à M. Guérin (Ann. Soc. Ent. France (2), Vol. 9, p. 125-130, f. [1851]).
- Leonardi, G., Gli insetti nocivi ai nostri orti, campi, frutteti e boschi, all'uomo ed agli animali domestici. Napoli, Vol. 3, Imenotteri e Ditteri (1900).
- Lepelletier de Saint-Fargeau & Serville, J. G., L'Encyclopédie méthodique. Insectes, Vol. 10. Paris (1825).
- Leunis, J., Synopsis der Thierkunde, Hannover, 2 vol. 3. ed. (1886).

- Lichtwardt, B., Dipterologische Bemerkungen (Zeitschr. Hymen. Dipt. Vol. 5, p. 309-311 [1905]).
- Linnæus, C., Fauna Suecica. Holmiæ, Ed. 2 (1761).
- Systema Naturæ. Holmiæ, Ed. 10 (1758). Ed. 12, 3 parts (1766-1768).
- Lioy, P., Sopra una straordinaria invasione di Ditteri della famiglia degli Empiti (Atti Soc. Ital. Sc. Nat. Milano, Vol. 6, p. 380-384 [1864]).
- I Ditteri distributti secondo un nuovo metodo di classificazione naturale (Atti Instituto Veneto Sc. Venezia (3), Vol. 9. Empiti in p. 598-604; p. 719-773 [1864]).
- Loew, H., Bemerkungen über die in der Posener Gegend einheimischen Arten mehrerer zweiflügler Gattungen (Program Posen, 1840, 40 p. 1 pl. [1840]).
- Ueber die in Grossherzogthum Posen aufgefundenen Zweiflügler (Oken's Isis, Vol. 7, p. 512-584 [1840]). Repetition of preceding.
- Ueber die europäischen Raubfliegen (Linnæa Ent. Vol. 4, p. 1-155 [1849]).
- Ueber den Bernstein und die Bernsteinfauna (Dipteren) (Berlin, Progr. Meseritz, 44 p. [1850]).
- Meghyperus* und *Arthropeas*, zwei neue Dipterengattungen (Stett. Ent. Zeit. Vol. 11, p. 302-308 [1850]).
- In Rosenhauer's : Die Thiere Andalusiens. Erlangen (1856).
- Review of Walker's Insecta Brittanica, Diptera. (Natural History Review, Lond. Vol. 12, p. 67-97 [1856]).
- Neue Beiträge zur Kenntniss der Dipteren. Part 4. Berlin, 57 p. (1856).
- Eine dipterologische Razzia auf dem Gebiete des naturwissenschaftlichen Vereins für Sachsen und Thüringen (Zeitschr. ges. Naturwiss. Jena. Vol. 10, p. 97-112 [1857]).
- Zehn neue Dipteren (Wien. Ent. Monatsschr. Vol. 2, p. 7-15 [1858]).
- Ueber die Arten der Gattung *Clinocera* (Wien. Ent. Monatsschr. Vol. 2, p. 238-253, 257-262 [1858]).
- Noch eine neue *Clinocera* (Wien. Ent. Monatsschr. Vol. 2, p. 386, 387 [1858]).
- Synamphotera pallida*, nov. sp. et gen. (Zeitschr. Ges. Naturwiss. Jena, Vol. 11, p. 453-456 [1858]).
- Bidrag till kännedomen om Afrikas Diptera (Oefv. Vet. Acad. Förhandl. Stockholm, Vol. 13, p. 255-264 (1856); ibidem, Vol. 14, p. 337-383 (1857); ibidem, Vol. 15, p. 335-341 [1858]).
- Neue Beiträge zur Kenntnis der Dipteren. Part 6 (Progr. Meseritz, 50 p. [1859]).
- Ueber der schlesisch Arten der Gattung *Tachypora* (Zeitsch. Ent. Breslau, Vol. 14, p. 1-32 [1860]).
- Ueber die schlesischen Arten der Gattung *Microphorus* (Zeitschr. Ent. Breslau, Vol. 14, p. 33-50 [1860]).
- Diptera americana ab Osten-Sackenio collecta, decas prima (Wien. Ent. Monatsschr. Vol. 4, p. 79-84 [1860]).
- Die Dipteren-Fauna Süd-Afrikas (Abhandl. Nat. Ver. Sachsen u. Thüringen, Vol. 2, p. 57-402 (1860); Reprint, Berlin [1860]).
- Diptera aliquot in insula Cuba collecta (Wien. Ent. Monatsschr. Vol. 5, p. 33-43 [1861]).
- Beschreibung einiger neuer europäischen Dipteren (Wien. Ent. Monatsschr. Vol. 5, p. 348-353 [1861]).
- Ueber einige bei Varna gefangene Dipteren (Wien. Ent. Monatsschr. Vol. 6, p. 160-175 [1862]).
- Diptera Americæ septentrionalis indigena (Berl. Ent. Zeitschr. Vol. 5, Centuria 1 (1861); Vol. 6, Centuria 2 (1862); Vol. 8, Centuria 5 (1864); Vol. 13, Centuria 8 (1869); Vol. 16, Centuria 10 [1872]).
- Diptera common to Europe and America (Silliman, Journal Sc. Arts, Vol. 37 [1864]).
- Die Europäischen Tipula-Arten deren Weibchen verkümmerte Flügel haben (Wien. Ent. Monatsschr. Vol. 8, p. 120-128 [1864]).

- Die oesterreichischen *Hemerodromia*-Arten (Wien. Ent. Monatsschr. Vol. 8, p. 237-255 [1864]).
 Ueber *Empis cothurnata* Brull. und *Empis hispanica* Loew (Wien. Entom. Monatsschr. Vol. 8, p. 255-258 [1864]).
Clinocera bivittata, nov. sp. (Wien. Ent. Monatsschr. Vol. 8, p. 258-260 [1864]).
 Ueber die *Pachymeria*-Arten aus dem Verwandtschaftskreise der *P. femorata* Fabricius (Wien. Ent. Monatsschr. Vol. 8, p. 353-366 [1864]).
 Ueber einige bei Kutais in Imeretien gefangene Dipteren (Berl. Ent. Zeitschr. Vol. 9, p. 234-242 [1865]).
 Ueber *Empis ciliata* Fabricius und über die ihr zunächst verwandten Arten (Berl. Ent. Zeitschr. Vol. 11, p. 1-10 [1867]).
 Ueber den Verwandtschaftskreise der *Empis stercorea* Linn. (Berl. Ent. Zeitschr. Vol. 11, p. 11-24 [1867]).
 Ueber diejenige mit *Empis chioptera* Meigen verwandten Arten, welche dunkle Schwinger haben (Berl. Ent. Zeitschr. Vol. 11, p. 25-62 [1867]).
 Nachträgliche Bemerkungen zu den *Empis*-Arten aus den Verwandschaftskreisen der *E. stercorea* und *chionoptera* (Berl. Ent. Zeitschr. Vol. 11, p. 157-166 [1867]).
 Ueber *Empis albicans* Meigen und eine derselben nahe verwandte Art (Berl. Ent. Zeitschr. Vol. 12, p. 168-175 [1868]).
 Ueber *Empis nitida* Meigen und die ihr verwandten Arten (Berlin. Ent. Zeitschr. Vol. 12, p. 231-240 [1868]).
 Nachträgliches über den Verwandtschafts-Kreis von *Empis albicans* (Berl. Ent. Zeitschr. Vol. 12, p. 387-393 [1868]).
 Ueber Dypteren der Augsburger Umgegend (20. Bericht. Naturh. Verein, Augsburg [1869]).
 Beschreibungen europäischer Dipteren, 3 vol., Halle, Vol. 1 (1869); Vol. 2 (1871); Vol. 3 (1873).
 Ueber einige *Empis*-Arten, welche zu den im Vol. 11 Bande besprochenen Verwandschaftskreisen gehören (Berl. Ent. Zeitschr. Vol. 13, p. 65-94 [1869]).
 Ueber die bisher auf der Galizischen Seite des Tetragebirges beobachteten Dipteren (Jahrb. Gel. Ges. Krakau, Vol. 42, p. 163-181 [1871]).
 Diptera Nova, in Pannonia inferiori et in confinibus Daciæ regionibus a Fern. Kowarzio capta (Berl. Ent. Zeitschr. Vol. 17, p. 33-52 [1873]).
 Diptera nova a Hug. Theod. Christopho collecta (Zeitschr. Ges. Naturwiss. Jena, Vol. 43, p. 413-420 [1874]).
 Beschreibungen neuer amerikanischer Dipteren (Zeitschr. Ges. Naturwiss. Jena, Vol. 48, p. 317-340 [1876]).
 Lucas, H., Note sur *Rhamphomyia platyptera* Panzer (Ann. Soc. Ent. France (3), Vol. 7, Bull. p. 243 [1859]).
 Lundbeck, W., Diptera Groenlandica (Videnskab. Meddel. Naturhist. Forening. Kjöbenhavn, 1898, p. 236-314 (1898); 1900, p. 281-316 [1900]).
 Diptera Danica, genera and species of flies hitherto found in Denmark. Vol. 3, Empididæ, with 141 fig. Copenhagen-London, p. 1-329 (1910).
 A new species of *Hilara* (Nathist. Meddel. Kjöbenhavn, Vol. 64, p. 325-327 [1913]).
 Conspectus faunæ groenlandicæ, pars 2, Landarthropoder (with K. L. Hendriksen) (Kjöbenhavn, Meddel. Grönl. Vol. 22 (2), p. 481-821 [1918]).
 Macquart, J., Monographie des Insectes Diptères de la famille des Empides, observés dans le Nord-Ouest de la France (Mém. Soc. Sc. Lille, 1822, p. 137-165 [1823]).

DIPTERA

- Insectes Diptères du Nord de la France, Lille, 5 vol. (Empids, Vol. 3, 1826 [1827]).
 Histoire naturelle des Insectes Diptères (suites à Buffon). Paris, 2 vol. (1834-1835).
 Description d'un nouveau genre d'Insectes Diptères de la famille des Tanystomes (*Euthyneura Myrtilli*) (Ann. Soc. Ent. France (1), Vol. 5, p. 517-520 [1836]).
 Diptères exotiques ou peu connus (Mém. Soc. Sc. Lille, 11 parts and 5 suppl. [1838-1855]).
 Explorations scientifiques de l'Algérie, Zoologie, Vol. 3 (1849).
- Malloch, J. R., *Empis hyalipennis* Fln. in Dumbartonshire (Ent. Monthly Mag. London, Vol. 42, p. 257 [1906]).
 A preliminary classification of Diptera, exclusive of Pupipara, based upon larval and pupal characters (Illinois Lab. Nat. Hist. Bull. Urbana, Vol. 12, Art. 3, p. 161-409, pl. 28-57 [1917]).
 The Diptera collected by the Canadian Expedition 1913-1918, excluding the Tipulidae and Culicidae (Rept. Canad. Arctic Exped. 1913-18, Vol. 3, C, 90 p. [1919]).
 Diptera of the Pribilof Islands, Alaska (No. American Fauna, No. 46, p. 170-227, pl. 12-15 [1923]).
 A new Empid from the Eastern United States (Ent. News Philad. Vol. 34, p. 5 [1924]).
- Matsumura, S., Erster Beitrag zur Insekten-Fauna Sachalin (Jour. Coll. Agric. Sapporo, Vol. 4, p. 1-145 [1911]).
 Thousand insects of Japan. Diptera (Additamenta, Vol. 2, Tokyo, p. 185-473, pl. 16-25 [1916]).
- McAtee, W. L., Some habits of Empidinae (Ent. News Philad. Vol. 20, p. 359-361 [1909]).
 Meigen, J. W., Nouvelle classification des Mouches à deux ailes. Paris, 40 p. (1800). Reprinted with notes by Hendel : Verh. Zool.-bot. Ges. Wien, 1908, p. 43-69 [1908]).
 Versuch einer neuen Gattungseintheilung der europäischen zweiflüglichen Insekten (Illiger's Magaz. Ins. Vol. 2, p. 259-281 [1803]).
 Klassification und Beschreibung der europäischen zweiflüglichen Insekten. Braunschweig, 1 vol. in 2 parts (1804).
 Systematische Beschreibung der bekannten europäischen zweiflüglichen Insekten. Aachen, Vol. 1-2; Hamm, Vol. 3-7 (1818-1838).
 Neue Arten von Diptera aus der Umgegend von München (Gistl's Faunus, Vol. 2, p. 66-72 [1835]).
- Meijere, J. C. H. (de), Diptera, Nova Guinea, Résultats de l'expédition scientifique néerlandaise à la Nouvelle-Guinée en 1902. Leide (E.-J. Brill), Vol. 5; Livr. 1, p. 67-99 (1906).
 Eerste Supplement op de Nieuwe Naamlijst van Nederlandsche Diptera (Tijdschr. v. Ent. Vol. 50, p. 151-195 [1907]).
 Studien über südostasiatische Dipteren. 1 (Tijdschr. v. Ent. Vol. 50, p. 196-264 [1907]).
 Studien über südostasiatische Dipteren. 4. Die neue Dipterfauna von Krakatau. (Tijdschr. v. Ent. Vol. 53, p. 58-194 [1910]).
 Die Dipteren der arktischen Inseln : in Fauna Arctica, Vol. 5, Jena, p. 13-72 (1910).
 Studien über südostasiatische Dipteren. 6 (Tijdschr. v. Ent. Vol. 54, p. 258-432 [1911]).
 Dipteren von Ceram und Waigeu (Bijdr. Dierk. Amsterdam, Vol. 19, p. 45-67 [1913]).
 Studien über südostasiatische Dipteren. 8 (Tijdschr. v. Ent. Vol. 56, suppl. p. 1-99 [1914]).
 Fauna simbarensis, Diptera (Tijdschr. v. Ent. Vol. 58, 1915, Suppl. p. 1-63 [1916]).
 Neue holländische Dipteren (Tijdschr. v. Ent. Vol. 61, p. 128-141, pl. 8 [1918]).
 Beitrag zur Kenntnis der Sumatraniischen Dipteren (Bijdr. Dierk. Amsterdam, Vol. 21, p. 13-39, pl. [1919]).

- Studien über süd-ostasiatische Dipteren, Pt. 15, dritter Beitrag zur Kenntnis der sumatranischen Dipteren (Tijdschr. v. Ent. Vol. 67, suppl. p. 1-64 [1924]).
- Meinert, F., Fluernes Munddele. Kjöbenhavn. 91 p. (1881).
- Melander, A. L., Gynandromorphism in a new species of *Hilara* (Psyche, Vol. 9, p. 213-215 [1901]).
Monograph of North American Empididæ, part 1 (Trans. Amer. Ent. Soc. Vol. 28, p. 195-367, pl. 5-9 [1902]).
- Some new or little-known genera of Empididæ (Ent. News, Philad. Vol. 17, p. 370-379 [1906]).
- Empididæ : in Williston's Manual of North American Diptera, 3. ed. p. 218-227 (1908).
- The genus *Tachydromia* (Psyche, Vol. 17, p. 41-62, pl. [1910]).
- The dipterous genus *Drapetis* Meigen (Ann. Ent. Soc. Amer. Vol. 11, p. 183-221 [1918]).
- Microsania*, a genus of the Platypezidæ (Psyche, Vol. 29, p. 43-48 [1922]).
- New species of *Platypalpus* occurring in New England (Occ. Papers Boston Soc. Nat. Hist. Vol. 5, p. 83-87 [1924]).
- Menzbier, M. A., Ueber das Kopfskelet und die Mundwerkzeuge der Zweiflügler (Bull. Soc. Nat. Moscou, Vol. 55, Pt. 1, p. 8-70 [1880]).
- Meunier, F., Note sur quelques Diptères fossiles de l'ambre tertiaire (Bull. Soc. Ent. France, 1893, p. 332-334 [1893]).
- An undescribed species of *Phyllodromia* from tertiary amber (Bull. Soc. Ent. France (6), Vol. 15, p. 13 [1895]).
- Revision des Diptères fossiles types de Loew conservés au Musée provincial de Königsberg (Miscell. Ent. Vol. 7, p. 161-165; 169-182 [1899]).
- Etude de quelques Diptères de l'ambre (Ann. Sc. Nat. (Zool.) Vol. 16, p. 395-406 [1902]).
- Description de quelques Diptères de l'ambre (Ann. Soc. Scient. Bruxelles, Vol. 26, p. 96-104 [1902]).
- Les Empidæ de l'ambre de la Baltique (Comptes rendus Acad. Sc. Paris, Vol. 145, p. 146-147 [1907]).
- Monographie des Empidæ de l'ambre de la Baltique et catalogue bibliographique complet des diptères de cette résine (Ann. Sc. Nat. (Zool.) (9), Vol. 7, p. 81-135 [1908]).
- Nouvelles recherches sur quelques Insectes des plâtrières d'Aix-en-Provence (Verh. Akad. Amsterdam, Wetensch. Sect. 10, Vol. 28, No. 5, 18 p. 5 pl. [1915]).
- Mik, J., Beiträge zur Dipteren-Fauna Oesterreichs (Verh. Zool.-bot. Ges. Wien, Vol. 19, p. 19-36, pl. 4 [1869]).
- Dipterologische Untersuchungen (Jahresber. Akad. Gymnas. Wien, 1878, 26 p. 1 pl. [1878]).
- Beschreibung neuer Dipteren. 1. Eilf neue europäische *Clinocera*-Arten (Verh. Zool.-bot. Ges. Wien, Vol. 30, p. 347-353 [1880]).
- Dipterologische Mitteilungen. Vol. 5. Die Gattung *Clinocera* Meigen (Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 320-329, pl. 16 [1881]).
- Einige Worte über P. Gabriel Strobl's « Dipterologische Funde um Seitenstettin » (Verh. Zool.-bot. Ges. Wien, Vol. 31, p. 345-352 [1881]).
- Ueber die Dipteren-Arten *Hemerodromia precatoria* Fallen und *Hemerodromia melanocephala* Hal. (Wien. Ent. Zeit. Vol. 1, p. 39-42 [1882]).
- Note on feeding of *Empis punctata* (Wien. Ent. Zeit. Vol. 1, p. 203 [1882]).
- Verschiedene Nahrung der Männchen und Weibchen mancher Insecten (Ent. Nachr. Berlin, Vol. 8, p. 116-119 [1882]).
- Eine neue Dipteren-Art aus Niederösterreich (Wien. Ent. Zeit. Vol. 3, p. 4-6 [1884]).
- Vier neue Dipteren aus Nieder-Oesterreich (Wien Ent. Zeit. Vol. 3, p. 81, 82 [1884]).

- Diptera des Gebietes von Hernstein in Nieder-Oesterreich und der weiteren Umgebung (in G. Beck, Fauna von Hernstein in Nieder-Oesterreich. [1885]).
- Eine neue Dipteren-Arten aus Süd-Tirol (Wien. Ent. Zeit. Vol. 5, p. 22-24 [1886]).
- Dipterologische Miscellen, ser. 1, Pt. 2, No. 13 (Wien. Ent. Zeit. Vol. 5, p. 278 [1886]).
- Die Dipteren-genera Paolo Lioy's (Ent. Nachr. Berlin, Vol. 12, p. 321-328 [1886]).
- Ueber einige Empiden aus Kärnten (Wien. Ent. Zeit. Vol. 6, p. 99-103 [1887]).
- Diagnosen neuer Dipteren, 1. Zwei neue Arten aus dem Alten Genus *Clinocera* Meigen (Wien. Ent. Zeit. Vol. 6, p. 161, 162 [1887]).
- Dipterologische Miscellen, ser. 1, Part 13, No. 63 (Wien. Ent. Zeit. Vol. 7, p. 299 [1888]).
- Dipterologische Miscellen, ser. 1, Part 14, No. 68 (Wien. Ent. Zeit. Vol. 7, p. 327 [1888]).
- Ein spinnendes Dipteron (Verh. Zool.-bot. Ges. Wien, Vol. 38, Sitzber. p. 97 [1888]).
- Antwort auf Herrn Dr. J. Schnabl's « Entgegnung » auf meine Kritik seiner « Contributions à la faune diptérologique » (Ent. Nachr. Berlin, Vol. 14, p. 41-45 [1888]).
- Eine neue schweizerische Art aus der alten Gattung *Clinocera* (Wien. Ent. Zeit. Vol. 8, p. 71, 72 [1889]).
- Eine neue aus den Beskiden stammende Art der alten Gattung *Clinocera* Meigen (Wien. Ent. Zeit. Vol. 8, p. 150-152 [1889]).
- Vorläufige Notiz über *Parathalassius blasigii*, ein neues Dipteren aus Venedig (Wien. Ent. Zeit. Vol. 10, p. 216, 217 [1891]).
- Dipterologische Miscellen, ser. 1, Part. 20, No. 99 (Wien. Ent. Zeit. Vol. 11, p. 55 [1892]).
- Zur Kenntniss der Dipteren Gattung *Hilara* (Wien. Ent. Zeit. Vol. 11, p. 78-85 [1892]).
- Dipterologische Miscellen, 4 (Wien. Ent. Zeit. Vol. 13, p. 49 [1894]).
- Dipterologische Miscellen, 5, No. 33 (Wien. Ent. Zeit. Vol. 13, p. 164-165 [1894]).
- Zur Verständigung (in Sachen der *Hilara sartor*) (Wien. Ent. Zeit. Vol. 13, p. 197-199 [1894]).
- Ein Beitrag zur Biologie einiger Dipteren, 2, Ueber spinnende Hilaren (Wien. Ent. Zeit. Vol. 13, p. 271-284 [1894]); (Jahresber. Akad. Gymnas. Wien, 1893-1894, p. 10-20 [1894]).
- Einige Worte über *Hilara sartor* (Ent. Nachr. Berlin, Vol. 20, p. 49-53 [1894]).
- Einige Worte zu Herrn Girschner's Artikel in den entomologischen Nachrichten Jahrg. 1894, p. 61, betitelt « Beiträge zur Biologie von *Hilara* » (Ent. Nachr. Berlin, Vol. 20, p. 151-155 [1894]).
- Dipterologische Miscellen, ser. 2, Part. 7 (Wien. Ent. Zeit. Vol. 15, p. 106-114 [1896]).
- Dipterologische Miscellen, ser. 2, Part. 9 (Wien. Ent. Zeit. Vol. 16, p. 34-40 [1897]).
- Dipterologische Miscellen, ser. 3, Part. 1 (Wien. Ent. Zeit. Vol. 19, p. 143-152 [1900]).
- Miller, D., A new species of *Empis* (Trans. New Zeal. Inst. Wellington, Vol. 42, p. 235, 236 [1910]).
- New species of New Zeal. and Empididae (Trans. New Zeal. Instit. Wellington, Vol. 45, p. 198-206, pl. [1913]).
- Material for a monograph on the Diptera Fauna of New Zealand : Part 3 - Family Empididae (Trans. Proc. New Zealand Institute, Vol. 54, p. 437-464, fig. [1923]).
- Müller, H., Die Entwicklung der Blumenthätigkeit der Insekten (Kosmos, Vol. 5 [1881]).
- Mueller, O. Fr., Fauna insectorum Friedrichsdalina, Hafniæ et Lipsiæ, 120 p. (1764).
- Zoologiæ Danicæ prodromus, Hafniæ, 314 p. (1776).
- Needham, J. G. & Betten, C., Aquatic Insects in the Adirondacks (Bull. New York Mus. Nat. Hist. Vol. 47, p. 383-612, 36 pl. [1901]).
- Neuhaus, G. H., Diptera Marchica. Systematisches Verzeichniss der Zweiflügler der Mark Brandenburg mit kurzer Beschreibung und analytischen Bestimmungs-Tabellen, Berlin, 371 p. 4 pl. (1886).

- Nowicki, M., Beschreibung neuer Dipteren (Verh. Naturf. Ver. Brünn. Vol. 6, p. 70-97, pl. 2 [1868]).
 Notes on *Microphorus* (Gel. Ges. Krakau, Vol. 42, p. 72-73 [1871]).
 Beiträge zur Insekten-Fauna Galiziens (Krakau University Press., 52 p. [1873]).
- Nylander, W., Hymenoptera et Diptera från Oesterbotten (Notizer ur Sällsk. Fauna Flore Fenn. Förh. Helsingfors, Vol. 4, p. 245-249 [1858]).
- Oldenberg, L., Einige südeuropäische Empididen (Ann. Mus. Hungar. Vol. 8, p. 344-352 [1910]).
 Drei neue Dipteren aus Tirol (Ent. Mitteil. Deutsche Ent. Mus. Dahlem-Berlin, Vol. 1, p. 209-215 [1912]).
 Ueber einige *Rhamphomyia*-Arten (Arch. Naturg. Berlin, Vol. 80 (1914), p. 69-91 [1915]).
 Veränderlichkeit der Beinfärbung bei *Atalanta* (*Clinocera*) (Arch. Naturg. Berlin, Vol. 80 (1914), p. 92 [1915]).
 Einige neue und alte *Hilara*-Arten (Arch. Naturg. Berlin, Vol. 81 (1915), A (9), p. 166-172 [1916]).
 Vier alpine *Rhamphomyia*-Arten (Arch. Naturg. Berlin, Vol. 82, A (1), p. 153-164 [1916]).
 Die Rhamphomyien des Wiener Hofmuseums (Arch. Naturg. Berlin, Vol. 83 (1917), A (6), p. 14-27 [1919]).
 Bemerkungen zu Engels Arbeit « Das Dipterengenus *Atalanta* » (Deutsche Ent. Zeitschr. 1919, p. 390-392 [1919]).
 Dipteren aus den Alpen (Zool. Jahrb. Vol. 43, Syst. p. 221-234 [1920]).
 Sechs alpine *Rhamphomyia*-Arten (Deutschen Ent. Zeitschr. 1922, p. 339-347 [1922]).
 Ueber einige *Tachydromia*- (*Coryneta*-) Arten (Ent. Mitteil. Berlin, Vol. 13, p. 84-89 [1924]).
 Die Empididen v. Rosers in Stuttgart (Deutsche Ent. Zeitschr. 1924, p. 226-236 [1924]).
 Drei dalmatische *Empis*-Arten (Deutsche Ent. Zeitschr. 1925, p. 317-322 [1925]).
 Beiträge zur Kenntnis der paläarktischen Rhamphomyien (Konowia, Vol. 6, p. 1-29 [1927]).
- Olivier, A. G., et al., Encyclopédie méthodique, dictionnaire des Insectes, Paris, 10 vol. (1789-1825); Diptera, Vol. 6 (1791).
- Osten-Sacken, C. R., A singular habit of *Hilara* (Ent. Monthly Mag. London, Vol. 14, p. 126 [1877]).
 Catalog of the described Diptera of North America (Smithsonian Miscell. Coll. 270, 2 ed. (1878); 1 ed. [1858]).
 Priorität oder Continuität? (Wien. Ent. Zeit., Vol. 1, p. 191-193 [1881]).
 Enumeration of the Diptera of the Malay Archipelago, collected by Prof. O. Beccari, Mr. L. M. d'Albertis and others (Ann. Mus. Stor. Nat. Genova, Vol. 16, p. 393-492 [1881]).
 Diptera from the Philippine Islands, brought home by Carl Semper (Berl. Ent. Zeitschr. Vol. 26, p. 83-120, 187-252 [1882]).
 Eine Beobachtung an *Hilara* (Ent. Nachr. Berlin, Vol. 12, p. 1, 2 [1886]).
 Biologia Centrali-Americanana, Diptera, Vol. 1, p. 129-216, pl. 3 (1887).
Hilarimorpha is a Leptid. (Berl. Ent. Zeitschr. Vol. 35, p. 303 [1890]).
- Palm, J., Beitrag zur Dipterenfauna Tirols (Verh. Zool.-bot. Ges. Wien, Vol. 19, p. 395-454 [1869]).
- Panzer, G. W. F., Fauna Insectorum Germaniae, Nürnberg, (1793-1844) (1829-1844 by H. Schaeffer (Empidæ 1801-1806)).
- Pearce, E. K., Typical flies ; a photographic atlas of Diptera. Cambridge, 45 p. (1915).
- Perris, Ed., Seconde excursion dans les Grandes Landes. Lettre adressée à M. Mulsant (Ann. Soc. Linn. Lyon, Vol. 5, p. 145-216 [1852]).
- Perty, Einige Insekten-Missbildungen (Mitteil. Naturf. Ges. Bern. 1867, p. 298-309 [1867]).
- Philippi, R. A., Aufzählung der Chilenischen Dipteren (Verh. Zool.-bot. Ges. Wien, Vol. 15, p. 595-782, pl. 23-29 [1865]).

- Pipping, List of Finland Empids (Notiser Sällsk. Faun. Fl. Fenn. Förh. Helsingfors, Vol. 4, p. 114 [1858]).
- Pokorny, E., Beitrag zur Dipterenfauna Tirols, Vol. 3 (Verh. Zool.-bot. Ges. Wien, Vol. 37, p. 381-420, pl. 7 [1887]).
- Poppius, B., För Finlands Fauna nya Dipterer (Medd. Soc. Fauna Fl. Fenn. Helsingfors, Vol. 32, p. 107 [1906]).
- Poulton, E. B., Predaceous Insects and their prey (Trans. Ent. Soc. Lond. 1906, p. 323-409 [1906]).
Empidæ and their prey in relation to courtship (Oxford Univ. Gazette, 1913, p. 952-953 (1913); Ent. Monthly Mag. London, Vol. 49, p. 177-180 [1913]).
- Raddatz, A., Uebersicht der in Mecklenburg bis jetzt beobachteten Fliegen (Arch. Ver. Freunde Nat. Mecklenburg, Rostock, Vol. 27, p. 22-131 [1873]).
- Reed, E. C., Catalogo de los Insectos Dipteros de Chile (Ann. Univ. Chile, Vol. 73, 51 p. [1888]).
- Retzius, A. J., De Geer : Genera et Species Insectorum, Lipsiæ, 220 p. (1783).
- Robert, Ch., Description d'un Diptère nouveau du genre *Paramesia* (Ann. Soc. Ent. France, Vol. 5, p. 537-538 [1836]).
- Röder, V. (von), Dipterologisch-synonymische Bemerkungen, No. 3 (Wien. Ent. Zeit. Vol. 3, p. 291 [1884]).
Rhamphomyia argentata, nov. sp. (Wien. Ent. Zeit. Vol. 6, p. 114 [1887]).
- Analytische Tabelle der Hemerodrominæ mit Einschluss der Gattung *Synamphotera* Loew (Wien. Ent. Zeit. Vol. 6, p. 169 [1887]).
- Dipterologische Beiträge (Wien. Ent. Zeit. Vol. 7, p. 95, 96 [1888]).
- Eine neue Diptere aus Kleinasien (Ent. Nachr. Berlin, Vol. 20, p. 202, 203 [1894]).
- Rondani, C., Dipterologiæ Italiciæ prodromus, Parinæ, 8 vol. (1856-1880).
Muscaria exotica Musei civici Januensis observata et distincta, Fragmentum. 3 (Ann. Mus. Stor. Nat. Genova, Vol. 7, p. 421-464 [1875]).
- Roser, C. L. F. (von), Verzeichniss der in Würtemberg vorkommenden zweiflüglichen Insekten (Correspondenzbl. landwirthsch. Ver. Würtemb. Stuttgart, 19 p. (1834). Supplement, 1840, Vol. 1, Pt. 1, p. 49-64 [1840]).
- Rossi, P., Fauna Etrusca, Liburni, 2 vol. (1790), 2 ed. (1807).
- Sack, P., Report of the scientific results of the Norwegian expedition to Novaya Zemlya, 1921, No. 15, Diptera, Christiana, p. 1-10 (1923).
- Say, Th., Descriptions of Dipterous Insects of the United States (Journ. Acad. Nat. Sc. Philad. Vol. 3, p. 9-54, 73-104 (1823); Complete Writings, Vol. 2, p. 38-89 [1859]).
- Keating's narrative of an expedition to the source of St. Peter's River, Lake Winnepeg, under the command of Major Long, Philadelphia (1824) (Complete Writings, Vol. 1, p. 176-258 [1857]).
- Schaeffer, J. Chr., Icones Insectorum circa Ratisbonam indigenorum, 3 vol. (1766-1779).
- Schellenberg, J. R., Gattungen der Fliegen in Vol. 42 Kupfertafeln entworfen und gezeichnet, Zurich, 95 p., 42 pl. (1803).
- Schiner, J. R., Scriptores austriaci rerum dipterologicarum (Verh. Zool.-bot. Ges. Wien, Vol. 6, p. 399-424 [1856]).
- Vorläufiger Commentar zum dipterologischen Theile der Fauna austriaca (Wien. Ent. Monatsschr. Vol. 4, p. 47-55 [1860]), etc.
- Fauna Austriaca, Diptera, Wien, 2 vol. (1862-1864).
- Catalogus systematicus Dipterorum Europæ, Vienna, 125 p. (1864).

- Erwiederung auf wiederholte Angriffe des Herrn H. Loew in Meseritz gegen meine Person und gegen meine Fauna austriaca (Wien. Ent. Monatsschr. Vol. 8, p. 296-301 [1864]).
- On *Hilarimorpha* (Verh. Zool.-bot. Ges. Wien, Vol. 18, p. 909, 910 [1868]).
- Reise der oesterreichischen Fregatte Novara um die Erde. Zoologischer Teil. Diptera, VI. and 388 p. pl. 4 (1868).
- Miscellen : Ueber neue Dipteren (Verh. Zool.-bot. Ges. Wien, Vol. 22, p. 74 [1872]).
- Schiödte, J. G., Beretning om Resultaterne af en i Sommeren 1838 foretagen entomologisk Undersøgelse af det sydlige Sjaelland (Kröyer, Naturhist. Tidskr. Vol. 2, p. 309-395 [1839]).
- Naturhistoriske Tillæg til Rinck : « Grönland geographisk og statistisk beskrevet » (Etzel Groenland, p. 604-620 [1860]).
- Uebersicht der Land-, Süsswasser- und Ufer-Arthropoden Grönlands (German translation of Tillæg) (Berl. Ent. Zeitschr. Vol. 3, 1859, p. 134-157 [1860]).
- Scholz, H., Ueber den Aufenthalt der Dipteren während ihrer ersten Stände (Zeitschr. Ent. Breslau, 1848-1849, p. 1-34 [1849]).
- Beiträge zur Kunde der schlesischen Zweiflüger, Part. 1, p. 35-40 (1850) : Part. 2, p. 41-60 (1851) (Zeitschr. Ent. Breslau, Vol. 5, No. 19 [1851]).
- Schrank, Fr. v. P., Enumeratio Insectorum Austriæ indigenorum, 9 and 548 p. (1781).
Fauna Boica, Nürnberg, 3 vol. (1798-1803).
- Schroeder, G., Beiträge zur Dipteren-Fauna Pommerns, Part. 4 (Stettin, Ent. Zeit. Vol. 73, p. 179-205 [1912]).
- Schummel, T. E., Ueber die von ihm auf dem Altvater-Gebirge gefangenen Insecten (Uebers. Arb. Schles. Ges. Vaterl. Kultur, Breslau, 1843, p. 184-199 [1843]).
- Schwarz, E. A., Feeding habit of a species of Empidæ (Proc. Ent. Soc. Wash. Vol. 2, p. 146, 147 [1891]).
- Scopoli, J. A., Entomologia Carniolica, Vindobonæ (1763).
- Scott-Elliott, G. Fr., Notes on flower haunting diptera (Trans. Ent. Soc. Lond. 1896, p. 117-128 [1896]).
- Scudder, S. H., Nomenclator Zoologicus : Part 1, Supplemental list, Part 2, Universal index (Bull. U. S. Nat. Mus. Vol. 19 [1884]).
- Systematische Uebersicht der fossilen Myriapoden, Arachnoideen und Insekten (Zittel, Handb. Palæontol. Vol. 1, p. 747-831 [1885]).
- Sendel, N., Historia succinorum corporea aliena involventium et naturæ opere pictorum et cælatorum, Lipsiæ, 336 p. 13 pl. (1742); (*Empis*, sp. p. 44, f. 19).
- Senior-White, R., New species of Diptera from the Indian Region (Mem. Dept. Agric. India, Entom. Ser. Vol. 7, p. 107-169 [1922]).
- New Ceylon Diptera, Pt. 3 (Spolia Zeylanica Colombo, Vol. 12, p. 375-406 [1924]).
- Serres, P. M. T. (de), Géognosie des terrains tertiaires du Midi de la France, Montpellier, 92 and 276 p. (1829) (*Empis*, sp. p. 232).
- Sharp, D., Sclater, W. L., et al. The Zoological Record, London, 62 vol. to 1925 (1864-1926).
- Siebke, H., Beretning om en i Sommeren 1861 foretagen entomologisk Reise (Nyt Magaz. Naturv. Vol. 12, p. 105-192 [1864]).
- Entomologisk Reise i Romsdals Amt i Sommeren 1864 (Nyt Magaz. Naturv. Vol. 14, p. 375-388 [1866]).
- Entomologiske Undersøgelser, foretagene i Sommeren 1865 (Nyt Magaz. Naturv. Vol. 14, p. 389-420 [1866]).

- Enumeratio Insectorum Norvegicorum, Vol. 4; Catalogus Dipterorum Norvegiæ, Christianiæ, 255 p. (1877).
- Slosson, A. T., Hunting Empids (Ent. News Philad. Vol. 14, p. 265-269 [1903]).
- Speiser, P., Dipteren aus Deutschlands afrikanischen Kolonien (Berl. Ent. Zeitschr. Vol. 52, 1907, p. 127-149 [1908]).
4. Orthorhapha : in Sjöstedts Kilimandjaro-Meru-Expedition, Stockholm, Vol. 10, Pt. 4, p. 31-112 (1910).
- Vorarbeiten zur einer Dipterenfauna der Provinz Ostpreusen (Schr. Physik-ökonom. Ges. Königsberg, Vol. 64, p. 7-18 [1924]).
- Stæger, C., Critisk Bemærkninger til Antliatslägten *Ocydromia* Meigen (Kröyer, Naturh. Tidsskr. Vol. 4, p. 98-102 [1842]).
- Haustellum hos Slægterne *Empis* og *Rhamphomyia* (Kröyer, Naturh. Tidsskr. Vol. 4, p. 102 [1842]).
- Beskrivelse af Grönlands Antliater (Kröyer, Naturh. Tidsskr. 2 ser. Vol. 1, p. 346-369 [1845]).
- Stein, J. P. E. Fr., Ein Ausflug nach dem Altvater-Gebirge (Stettin, Ent. Zeit. Vol. 34, p. 233-243 [1873]).
- Stein, P., Zwei neue Dipteren (Wien. Ent. Zeit. Vol. 9, p. 108, 109 [1890]).
- Einige Mitteilungen über *Trichopera longicornis* Meigen (Wien. Ent. Zeit. Vol. 22, p. 225-228 [1903]).
- Stellweg, F., Tanzende Fliegen (Nat. Wochenschr. Vol. 32, p. 281-283 [1917]).
- Stephens, J. F., A systematic catalogue of British Insects. London, 2 vol. (1829).
- Strobl, G., Dipterologische Funde um Seitenstettin (Jahresber. Obergymnas. Seitenstett. Progr. Vol. 14 [1880]).
- Zur Synonymie von *Symbalophthalmus pictipes* Becker (Wien. Ent. Zeit. Vol. 10, p. 267 [1891]).
- Die österreichischen Arten der Gattung *Hilara* Meigen, mit Berücksichtigung der Arten Deutschlands und der Schweiz (Verh. Zool.-bot. Ges. Wien, Vol. 42, p. 85-182 [1892]).
- Beiträge zur Dipteren-Fauna des österreichischen Litorale (Wien. Ent. Zeit. Vol. 12, p. 29-42; 74-80; 89-108; 121-136; 160-170 [1893]).
- Die Dipteren von Steiermark (Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 29, p. 1-199, 1892 [1893]; Vol. 30, p. 1-52, 1893 [1894]).
- Hilara longicornis*, nov. sp. (Wien. Ent. Zeit. Vol. 13, p. 59 [1894]).
- Siebenbürgische Zweiflügler (Verh. Siebenbürg. Ver. Naturwiss. Hermannstadt, Vol. 46, 1896 [1897]).
- Die Dipteren von Steiermark, 4. Theil. Nachträge (Mitteil. Naturw. Ver. Steiermark, Graz, Vol. 34, 192-298 [1898]).
- Spanische Dipteren (Wien. Ent. Zeit. Vol. 29 (2), p. 12-27; (3), 77-83; etc. [1899]).
- Dipterenfauna von Bosnien, Hercegovina und Dalmatia (Glasnik Zem. Mus. Bosni i Herzegov. Sarajevo, Vol. 10, p. 387-616 (in Serbian) (1898); Wiss. Mitteil. Bosnien, Hercegov. Sarajevo, Vol. 7, p. 552-670 (in German) [1900]).
- Tief's dipterologischer Nachlass aus Kärnten und Oesterr-Schledien (Jahrb. Mus. Kärnten. Klazentfurt, Vol. 26, p. 171-246 [1901]).
- Neue Beiträge zur Dipterenfauna der Balkanhalbinsel (Glasnik Zem. Mus. Bosni i Herzegov. Sarajevo, Vol. 14, p. 461-517 (in Serbian) (1902); Wiss. Mittel. Bosn. Hercegov. Sarajevo, Vol. 9, p. 519-581 (in German) [1904]).
- Spanische Dipteren, 2. Beitrag (Mem. Soc. Esp. Hist. Nat. Vol. 3, p. 271-422 [1906]).

- Spanische Dipteren, 3. Beitrag (Verh. Zool.-bot. Ges. Wien, Vol. 59, p. 121-301 [1909]).
- Die Dipteren von Steiermark, 2. Nachtrag (Mitteil. Naturw. Steiermark, Graz, Vol. 46, 1909, p. 45-293 [1910]).
- Sulzer, J. H., Abgekürzte Geschichte der Insekten nach dem Linneischen system. Winterthur (1776).
- Thomson, C. G., Kongliga Svenska Fregatten Eugenies Resa omkring Jorden. Vetenskapliga Iakttagelser, Zoologi, Pt. 1. Insekta, Häft 12, Diptera, p. 443-614, 9 pl. (1868).
- Thunberg, C. P., Novæ Insectorum species descriptæ (N. Acta reg. Soc. Upsalensis, Upsala, Vol. 4, p. 1-28 [1784]).
- Tillyard, P., The Insects of Australia and New Zealand, 560 p. Sydney (1926).
- Tonnoir, A. L., Australian Platypezidæ (Rec. Austr. Mus. Sydney, Vol. 14, p. 306-312 [1925]).
- Tucker, E. S., Some results of desultory collecting of Insects in Kansas and Colorado (Kansas. Univ. Sc. Bull. Vol. 4, p. 51-112 [1907]).
- Verhoeff, C., Zur Biologie von *Hilara* (Ent. Nachr. Berlin, Vol. 20, p. 1, 2 [1894]).
- Verrall, G. H., On the species of *Empis* allied to *E. stercorea*, Linn. including one new to science (Ent. Monthly Mag. Vol. 8, p. 281-284 [1872]).
- Notes on Scotch Diptera (Scotish. Nat. Vol. 2, p. 199-202 [1874]).
- Diptera in Arran. (Ent. Monthly Mag. London, Vol. 19, p. 222-226 [1883]).
- A hundred new species of Diptera (Ent. Monthly Mag. Vol. 22, p. 179-182; 199-202; 230-234 [1886]).
- List of British Diptera, Cambridge, 1 ed. (1888), 2 ed. (1901).
- Critical notes on Diptera (The Entom. London, Vol. 23, p. 150-154 [1890]).
- A second hundred new British species of Diptera (Ent. Monthly Mag. London, Vol. 30, p. 76-79; 140-146 [1894]).
- British Flies, Vol. 5. Stratiomyidæ, etc., London, 780 p. (1909).
- Villeneuve, J., Étude sur quelques Diptères, No. 6, *Empis setigera* (Bull. Soc. Ent. France, 1903, p. 212 [1903]).
- Notes synonymiques (Wien. Ent. Zeit. Vol. 31, p. 96-97 [1912]).
- Diptères nouveaux ou intéressants (Feuille Jeunes Natur. Paris, Vol. 43, p. 111-113 [1913]).
- Wahlberg, P., F. Nya Diptera från Lappland (Oefv. K. Vet. Akad. Holm. Förhandl. Stockholm, Vol. 1, p. 64-68; 106-110; 217-219 [1844]).
- Andamålet med Tachydromidernes fortbildung (Oefv. K. Vet. Akad. Holm. Förhandl. Stockholm, Vol. 2, p. 253-254 [1845]).
- Wahlgren, E., Zur Kenntnis swedischer Dipteren, No. 2, *Brachystoma aucta* Zetterstedt und *Paramesia tenella* Wahlberg (Ent. Tidskr. Vol. 31, p. 28-34 [1910]).
- Svensk insekt-fauna, Pt. 11, Diptera, Fam. 24, Empididæ (Ent. Tidskr. Stockholm, Vol. 31, p. 41-95 [1910]); Reprint, Uppsala, p. 63-117 (1910).
- Det ölandska alvarets djurvärld, Pt. 1. (Arkiv. Zool. Svensk. Vetensk. Vol. 9 (19), p. 135 [1915]); Pt. 2. (ibidem, Vol. 11 (1), p. 130 [1917]).
- De europeiska polaröarnas insektafauna, dess sammansättning och härkomst (Ent. Tidskr. Vol. 41 (1), 23 p. [1920]).
- Walker, Fr. Notes on Diptera (Ent. Monthly Mag. London, Vol. 3, p. 178-182 (1835); Vol. 4, p. 226-230 [1836]).
- Descriptions of Diptera collected by Capt. King in the survey of the Straits of Magellan (Trans. Linn. Soc. Lond. Vol. 17, p. 331-359 [1837]).

- List of the specimens of Dipterous Insects in the collection of the British Museum, London, 4 parts and 3 suppl. (1848-1855).
- Insecta Saundersiana, or characters of undescribed insects in the collection of W. W. Saunders, London, 474 p., 8 pl., in 5 parts (1850-1856).
- Insecta Britannica, Diptera. London, 3 vol. (1851-1856).
- Characters of undescribed Diptera in the collection of Wm. Saunders (Trans. Ent. Soc. Lond. n. s. Vol. 4, p. 119-158; 190-235 [1857]; Vol. 5, 268-334 [1858]).
- Catalogue of the Dipterous Insects collected in Amboyna by Mr. A. R. Wallace (Journ. Proc. Linn. Soc. London. Suppl. to Vol. 4, 1860, p. 144-168 [1861]).
- Catalogue of the Dipterous Insects collected in Batchian, Kaisaa and Makian, and at Tidon in Celebes by Mr. A. R. Wallace (Journ. Proc. Linn. Soc. Lond. Vol. 5, p. 270-303 [1861]).
- Catalogue of the Dipterous Insects collected in Waigiou, Mysol and North Ceram by A. R. Wallace (Journ. Proc. Linn. Soc. Lond. Vol. 7, p. 202-238 [1864]).
- Descriptions of new species of the Dipterous Insects of New Guinea (Journ. Proc. Linn. Soc. Lond. Vol. 8, p. 102-130 [1865]).
- List of the Diptera collected in Egypt and Arabia by J. K. Lord, Esq.; with descriptions of the species new to science (The Ent. London, Vol. 5, p. 254-263; 271-275; 339-346 [1871]).
- Wallengren, H. D. J., Anteckningar i Entomologi (Oefv. Vet. Akad. Förhandl. Stockholm, 1870, p. 145-182 [1870]).
- Wasmann, E., *Hilara sartor* Becker (Ber. Offenbach Ver. Naturk. Münster, Vol. 41, p. 436 [1895]).
- Webster, F. M., Notes and observations on several species of Diptera (Canad. Ent. Vol. 30, p. 18-19 [1898]).
- Wesché, W., The labial and maxillary palpi in Diptera (Trans. Linn. Soc. Lond. ser. 2, Vol. 9, p. 219-230, pl. 8-10 [1904]).
- Westwood, J. O., An introduction to the modern classification of Insects. London, 2 vol. (1839-1840).
- Wheeler, M. W., A new Empid with remarkable middle tarsi (Ent. News Philad. Vol. 7, p. 189-191 [1896]).
- Wheeler, W. M. & Melander, A. L., Empididae, in Biologia Centrali-Americana, Diptera, Vol. 1, p. 367-376 (1901).
- White, A., The Diptera brachycera of Tasmania, Part 3 (Pap. Proc. Roy. Soc. Tasmania, Hobart, 1916, p. 148-266 [1917]).
- White, Adam, Nomenclator of British Diptera (List Coll. Brit. Mus. London, Vol. 15, 42 p. [1853]).
- Whitfield, F. G. S., The natural control of the leaf-miner *Phytomyza aconiti* Hendel by *Tachydromia minuta* Meigen (Bull. Ent. Res. Vol. 16, Pt. 1, p. 95-97, 3 fig. [1925]). Review (Rev. Appl. Ent. Vol. 13, A, p. 422 [1925]).
- Wiedemann, C. R. W., Ueber einige neue Fliegengattungen (Zool. Mag. Berlin, Vol. 1, Pt. 1, p. 57-61 [1817]).
- Neue Zweiflüger aus der Gegend um Kiel (Zool. Mag. Berlin, Vol. 1, Pt. 1, p. 61-86 [1817]).
- Aus Pallas dipterologischen Nachlinse (Zool. Mag. Berlin, Vol. 1 (2), p. 1-40 [1818]).
- Analecta Entomologica, Kiliæ (1824).
- Aussereuropäische zweiflügelige Insekten, Hamm, 2 vol. (1828-1830).
- Williston, S. W., Ueber einige Leptiden-Characteren (Stett. Ent. Zeit. Vol. 46, p. 400, 401 [1885]).
- Synopsis of the families and genera of North American Diptera, New Haven, 84 p. [1888]).
- Hilarimorpha and Apiocera (Psyche, 1888, p. 99-102 [1888]).

- New or little-known Diptera (Kansas Univ. Quarterly, Vol. 1, p. 59-78 [1893]).
Manual of the families and genera of North American Diptera, ed. 2, New Haven, 167 p. (1896).
On the Diptera of St. Vincent (West Indies) (Trans. Ent. Soc. Lond. 1896, p. 253-449, pl. 8-14 [1896]).
Manual of North American Diptera, ed. 3, New Haven, 405 p. (1908).
Wood, J. H., *Hilara albocingulata*, nov. sp. (Ent. Monthly Mag. London, Vol. 49, p. 13, 14 [1913]).
Wood, W., Illustrations of the Linnean genera of Insects, London, 2 vol. (1821).
Wulp, F. M. (van der), Eenige Noord-Americaansche Diptera (Tijdschr. v. Ent. 2 ser. Vol. 2, p. 125-164, pl. 3-5 [1867]).
Catalogue of the described Diptera from South Asia (The Hague, Dutch Ent. Soc. 1896, 220 p. [1896]).
Zur Dipteren-Fauna von Ceylon (Termesz. Füzetek, Budapest, Vol. 20, p. 136-144 [1897]).
Verbeteringen en Aanvullingen in den Catalogue of the described Diptera from South Asia (Tijdschr. v. Ent. Vol. 42, p. 41-57 [1899]).
Wulp, F. M. (van der), & Meijere, J. C. H. (de). Nieuwe Naamlijst van Nederlandsche Diptera (Tijdschr. v. Ent. Vol. 41, p. 149 [1898]).
Yerbury, J. W., List of the Diptera of Glamorgan (Trans. Cardiff Naturalists Soc. Vol. 51 (Empids on p. 54 et seq.) [1920]).
Zetterstedt, J. W. Några nya Svenske Insect-arter fundne och beskrifne (Vet. Acad. Handl. Stockholm, 1819, p. 69-86 [1819]).
Resa genom Umeå Lappmarker år 1832, Orebro (1833).
Conspectus familiarum, generum et specierum Dipterorum in Fauna Lapponica descriptorum (Isis, 1837, Vol. 1, p. 28-67 [1837]).
Insecta lapponica descripta, Lipsiæ (1838-1840).
Diptera scandinaviæ disposita et descripta, Lundæ, 14 vol. (1842-1860).

INDEX (*)

	Pages		Pages		Pages
abbreviata Lw. (<i>Empis</i>)	145	æstiva Lw. (<i>Empis</i>)	146	alipes Walk. (<i>Chelifera</i>)	262
abbrevinervis Meij. (<i>Empis</i>)	145	æstiva Scop. (<i>Empis</i>)	146	albisetra Panz. (<i>Platypalpus</i>)	345
abcirus Walk. (<i>Empis</i>)	145	æthiops Zett. (<i>Hilara</i>)	113	albisetra Zett. (<i>Chelipoda</i>)	265
abdita Coq. (<i>Rhamphomyia</i>)	205	æthiops Zett. (<i>Rhamphomyia</i> , not candata)	188-385	albissima Frey (<i>Rhamphomyia</i>)	184
abdominalis Bez. (<i>Rhamphomyia</i>)	184	affinis Egg. (<i>Empis maculata</i>)	162	albitarsis Collin (<i>Rhamphomyia</i>)	382
abdominalis Bonsd. (<i>Hilara obscuritarsis</i>)	123	affinis Lw. (<i>Drapetis assimilis</i>)	308	albitarsis Ros. (<i>Hilara</i>)	113, 383
abdominalis Scholz (<i>Hilara heterogastra</i>)	119	affinis Steph. (<i>Empis</i>)	146	albitarsis Zett. (<i>Tachydromia annulimana</i>)	283
abdominalis Wied. (<i>Drapetis</i>)	312	agasics Walk. (<i>Rhamphomyia</i>)	184	albiventris Ros. (<i>Hilara</i> , not heterogastra)	119, 383
abdominalis Zett. (<i>Hilara</i>)	113	Agastashys Scud. (<i>Stilpon Lw.</i>)	301	albocapillatus Fall. (<i>Platypalpus</i>)	346
abdominenotata White (<i>Drapetis</i>)	378	agasthus Walk. (<i>Empis</i>)	146	albocapillatus Fall. (<i>Platypalpus albiseta</i>)	346
aberrans Bez. (<i>Hilara</i>)	113	Agatachys Meig. (<i>Stilpon Lw.</i>)	301	albocincta Bohem. (<i>Trichopeza</i>)	219
* ablata Meun. (<i>Rhamphomyia</i>)	184	agens Mel. (<i>Tachyempis</i>)	292	albocingulata Wood (<i>Hilara albiventris</i>)	113, 383
abnormis Bez. (<i>Hilara</i>)	113	agilellus Coll. (<i>Platypalpus</i>)	381	albogeniculata Ros. (<i>Rhamphomyia</i>)	185
abrupta Thom. (<i>Empis</i>)	145	agilis Meig. (<i>Platypalpus</i>)	345	albohalteralis Brun. (<i>Empis</i>)	147
Acallomyia (nov. subgen. <i>Empis</i>)	140	alampra Lw. (<i>Empis</i>)	146	albohalterata Stro. (<i>Dolichocephala ocellata</i>)	247
Acanthopeza Beck. (<i>Stenoprocitus Lw.</i>)	72	alata Zett. (<i>Rhamphomyia vesiculosa</i>)	210	albohirta Coll. (<i>Rhamphomyia</i>)	382
Acarterus auct. (<i>Parahybos Kert.</i>)	45	albata Coq. (<i>Rhamphomyia</i>)	184	albomicans Beck. (<i>Platypalpus</i>)	346
Acarterus Lw. (gen.)	38	albens Pallas (<i>Platypalpus</i>)	345	albonotatus Lw. (<i>Syneches</i>)	40
acephala Panz. (<i>Hilara flavipes</i>)	117	albicans Meig. (<i>Empis</i>)	146	albopilosa Coq. (<i>Rhamphomyia</i>)	185
Actonempis Coll. (subgen. <i>Rhamphomyia</i>)	382	albicheta Frey (<i>Rhamphomyia nitens</i>)	200	albopilosus Beck. (<i>Microphorus</i>)	92
acrodactyla, nov. sp. (<i>Drapetis</i>)	312	albicincta Lw. (<i>Empis</i>)	146	albosegmentata Zett. (<i>Rhamphomyia</i>)	185
Acromyia Bon. (<i>Hybos Meig.</i>)	18	albicornis auct. (<i>Hemerodromia raptoria</i>)	254, 385	aldrichi Mel. (<i>Parathalassius</i>)	87
acroptera, nov. sp. (<i>Microphorella</i>)	88	albicornis auct. (<i>Platypalpus pallidicornis</i>)	345, 381	aldrichii Mel. (<i>Empis</i>)	147
* acuticornis Lw. (<i>Timalaphes</i>)	98	albicornis Meig. (<i>Hemerodromia</i>)	254	alexippus Walk. (<i>Platypalpus</i>)	346
adelensis Mill. (<i>Homalocnemis</i>)	14	albicornis Rossi (<i>Platypalpus albisetosus</i>)	346	algecirasensis Stro. (<i>Empis</i>)	147
adulatoria Coll. (<i>Hemerodromia</i>)	380	albicornis Zett. (<i>Platypalpus</i>)	345	algecirasensis Stro. (<i>Hilara</i>)	113
adusta Lw. (<i>Empis</i>)	145	albida Meig. (<i>Hilara fuscipes</i>)	118	algira Macq. (<i>Empis</i>)	147
adversa Coq. (<i>Rhamphomyia</i>)	184	albidipennis Lw. (<i>Heterotropus</i>)	375	algira Macq. (<i>Hilara clypeata</i>)	116
æmula Lw. (<i>Empis</i>)	145	albidiseta Beck. (<i>Empis</i>)	146	algitus Macq. (<i>Platypalpus</i>)	346
æmula Lw. (<i>Tachydromia</i>)	283	albidiventris Str. (<i>Rhamphomyia</i>)	184	alipes Meig. (<i>Rhamphomyia</i>)	185
ænea Walk. (<i>Drapetis</i>)	310	albitacies Coll. (<i>Platypalpus</i>)	381	alternigra Mel. (<i>Drapetis</i>)	307
æneicollis Zett. (<i>Platypalpus</i>)	345	albifrons Bez. (<i>Empis</i>)	146	alterpicta Beck. (<i>Tachydromia</i>)	283
ænescens Wied. (<i>Drapetis</i>)	310	albifrons Macq. (<i>Platypalpus</i>)	345	Allanthalia (nov. gen.)	61
æneus Macq. (<i>Platypalpus</i>)	345	albinervis Meig. (<i>Empis</i>)	146	almériensis Stro. (<i>Hilara</i>)	113
æqualis Lw. (<i>Empis</i>)	145	albipennis Fall. (<i>Rhamphomyia</i>)	184	alpestris Schum. (<i>Rhamphomyia anthracina</i>)	186
æqualis Lw. (<i>Platypalpus</i>)	345	albipennis Meig. (<i>Empis</i>)	146, 385	alpicola Stro. (<i>Empis</i>)	147
æquicornis, nov. var. (<i>Platypalpus inops</i>)	330	albipennis Perris (<i>Platypalpus</i>)	345	alpicola Stro. (<i>Empis chioptera</i>)	151
æripes Mel. (<i>Empis</i>)	145	albipennis Rossi (<i>Hilara</i>)	113, 383	alpigenus Stro. (<i>Platypalpus</i>)	346
aerobatica Mel. (<i>Empis</i>)	146	albipennis Zett. (<i>Empis albimervis</i>)	146		
aeronetha Mik (<i>Hilara</i>)	113	albipennis Zett. (<i>Euthyneura myrtilli</i>)	54		

(*) Names of species and varieties are spelled with lower case letters; names of genera and subgenera are in thick type and begun with a capital; names of families and subfamilies in thick capitals. Extinct forms are indicated by a prefixed asterisk. Species synonyms are in *italics* and have their valid names given in parentheses. Genus synonyms are so indicated by their valid names following in parentheses.

Pages	Pages	Pages			
alpina Bez. (<i>Bicellaria</i>)	76	annulatus Fall. (<i>Platypalpus m-</i>			
alpina Eng. (<i>Wiedemannia</i>)	242	<i>nutus</i>)	360	appendiculata Hansen (<i>Cinocera</i>	
alpina Lw. (<i>Empis</i>)	147	annulimana Meig. (<i>Tachydromia</i>)	283	<i>stagnalis</i>)	238
alpina Lw.-Ost.-Sack. (<i>Hilara</i>		annulipes Bez. (<i>Syneches</i>)	40	appendiculata Macq. (<i>Rhamphomyia</i>	
<i>sartor</i>)	125	annulipes Meig. (<i>Platypalpus</i>)	347, 384	<i>forcipata</i>)	193
alpina Stro. (<i>Rhamphomyia tenui-</i>		annulipes Wh.-Mel. (<i>Empis</i>)	147	appendiculata Zett. (<i>Cinocera</i>)	236
<i>cornis</i>)	195	Anodontina Macq. (<i>Haplomera</i>		approximatus Beck. (<i>Platypalpus</i>)	347
alpina Zett. (<i>Rhamphomyia</i>)	185	Macq.)	130	apraca Steph. (<i>Empis</i>)	147
alumnus, nov. sp. (<i>Platypalpus</i>)	330	anomala Lw. (<i>Hilara pilipes</i>)	123	aprilis White (<i>Rhamphomyia</i>)	186
amaranta Beck. (<i>Hilara</i>)	113	anomala Old. (<i>Rhamphomyia</i>)	185	apta Coll. (<i>Hilara</i>)	380
amarantha Beck. (<i>Clinocera</i>)	238	anomala Scholz (<i>Dolichocephala</i>		apteropus Bez. (<i>Empis opaca</i>)	166
amaurus Beck. (<i>Halsanalotes</i>)	298	<i>irrorata</i>)	247	aquatica Beck. (<i>Clinocera</i>)	236
ambigua Bez. (<i>Empis</i>)	147	Anomalempis (nov. gen.)	14	aquilex Lw. (<i>Wiedemannia</i>)	243
ambigua Macq. (<i>Drapetis</i> , not		anomalicerus Beck. (<i>Platypalpus</i>)	347	aquilus White (<i>Empis</i>)	147
<i>flavipes</i>)	309, 384	anomalina Frey (<i>Rhamphomyia</i>		arctica Zett. (<i>Rhamphomyia alboseg-</i>	
ambigua Phil. (<i>Brachystoma</i>)	13	<i>gufistar</i>)	194	<i>mentata</i>)	185
americana Wied. (<i>Rhamphomyia</i>)	185	anomalina Zett. (<i>Rhamphomyia</i>)	185	arcticus, nov. sp. (<i>Platypalpus</i>)	344
Amlictoides Bez. (<i>Deuteragonista</i>		anomalinervis Frey (<i>Rhamphomyia</i>)	187	arcuata Coq. (<i>Rhamphomyia</i>)	186
Phil.)	132	anomalipennis Meig. (<i>Rhampho-</i>		arcuata Lw. (<i>Drapetis assimilis</i>)	308
amoena Lw. (<i>Rhamphomyia</i>)	185	<i>myia</i>)	186	arcuincta Bez. (<i>Rhamphomyia</i>)	186
amplectens, n. sp. (<i>Proclinoptyga</i>)	221	anomalous Meig. (<i>Microphorus</i>)	92, 384	ardesiaca Wied.-Meig. (<i>Empis</i>)	147
amplicella Coq. (<i>Rhamphomyia</i>)	185	antarctica Walker (<i>Empis</i>)	147	ardopeodes, nov. sp. (<i>Euhybos</i>)	32
amplipedis Coq. (<i>Rhamphomyia</i>)	185	antennalis, nov. sp. (<i>Leptopeza</i>)	66	Ardoptera Macq. (<i>Dolichocephala</i>	
amplitarsis Brun. (<i>Empis</i>)	147	antennata Beck. (<i>Drapetis</i>)	312	Macq.)	245
Amydroneura Coll. (subgen.		antennata Frey (<i>Rhamphomyia</i>)	186	arenaria Hal. (<i>Chersodromia</i>)	296
<i>Rhamphomyia</i>)	382	Anthalia Coq. (<i>Euthyneura</i> Macq.)	52	argentata Phil. (<i>Haplomera</i>)	131
amyti Walk. (<i>Empis</i>)	147	Anthalla Zett. (gen.)	57	argentata Roed. (<i>Rhamphomyia</i>)	186
Anachrostichus Bez. (subgen.		Anthepiscopus Beck (gen.)	104	argentea Big. (<i>Rhamphomyia lim-</i>	
<i>Empis</i>)	140	anthracina Big. (<i>Empis</i>)	147	<i>bata</i>)	196
analis Meig. (<i>Platypalpus</i>)	346	anthracina Meig. (<i>Rhamphomyia</i>)	186	argenteomicans Beck. (<i>Platypalpus</i>)	347
analis Thoms. (<i>Chelifera</i>)	262	anthracina Zett. (<i>Rhamphomyia</i>		argenticeps Meij. (<i>Platypalpus</i>)	347
anaxo Walk. (<i>Rhamphomyia</i>)	185	<i>vesiculos</i>)	210	argentifera Bez. (<i>Hilarempis</i>)	128
andalusiaca Stro. (<i>Rhamphomyia</i>)	185	anthracinella Stro. (<i>Rhamphomyia</i>)	186	argentula Beck. (<i>Hilarempis</i>)	128
andalusiacus Stro. (<i>Platypalpus</i>)	346	anthracodes Coq. (<i>Rhamphomyia</i>)	186	argus, nov. sp. (<i>Dolichocephala</i>)	246
andermattensis Stro. (<i>Hilara longe-</i>		* antipedalis Lw. (<i>Rhamphomyia</i>)	186	Argyrandrus Bez. (subgen.	
<i>vittata</i>)	120	antipodus Bez. (<i>Anthepiscopus</i>)	107	<i>Empis</i>)	141
anfractuosa Bez. (<i>Rhamphomyia</i>)	185	Apalecnemis Phil. (gen.)	100	argyreata Egg. (<i>Empis maculata</i>)	162
anfractuosa Mik (<i>Empis dasypoda</i>)	153	aperta Lw. (<i>Rhamphomyia</i>)	186	argyria, nov. sp. (<i>Euthyneura</i>)	53
anglokanica Lund. (<i>Hilara</i>)	113	aperta Mel. (<i>Euthyneura</i>)	53	argyrina Bez. (<i>Rhamphomyia</i>)	186
angulifera Frey (<i>Rhamphomyia</i>)	185	aperta Zett. (<i>Rhamphomyia dispar</i>)	191	argyriventris Beck. (<i>Atrichopleura</i>)	134
angusta Coll. (<i>Chelifera</i>)	378	aperticauda Coll. (<i>Chelifera</i>)	378	argyropalpa Bez. (<i>Coloboneura</i>)	294
* angusta Meun. (<i>Rhamphomyia</i>)	185	afelbecki Stro. (<i>Empis</i>)	147	argyrosoma Stro. (<i>Hilara braueri</i>)	114
angustata Coll. (<i>Drapetis</i>)	310	apicalis Beck. (<i>Platypalpus apicatus</i>)	347	argyrotarsis Bez. (<i>Rhamphomyia</i>)	186
angustifrons Stro. (<i>Hilara aeronetha</i>)	113	apicalis Bez. (<i>Rhamphomyia</i>)	186	argyrozoa Phil. (<i>Empis</i>)	147
angustifurca, nov. sp. (<i>Bicellaria</i>)	75	apicalis Lw. (<i>Empis</i>)	147	argyrozoa Phil. (<i>Hilarempis</i>)	128
angustipennis Beck. (<i>Rhamphomyia</i>		apicalis Lw. (<i>Edalea</i>)	64	Ariasella Gil (gen.)	377
<i>stigmosa</i>)	206	apicalis Lw. (<i>Platypalpus</i>)	347	aristalis Mel. (<i>Drapetis</i>)	312
angustipennis Bez. (<i>Empis</i>)	147	apicalis Steph. (<i>Platypalpus</i>)	347	aristatus Coll. (<i>Platypalpus</i>)	381
angustipennis Lw. (<i>Rhamphomyia</i>)	185	apicata Staeg. (<i>Tachydromia terri-</i>		armata Beck. (<i>Rhamphomyia</i>)	186
annularis Bez. (<i>Platypalpus</i>)	346	<i>cola</i>)	287	armata Cress. (<i>Mythicomyia</i>)	376
annularis, nov. sp. (<i>Tachypesa</i>)	274	apicatus, n. n. (<i>Platypalpus</i>)	347	armata Eng. (<i>Wiedemannia</i>)	244
annulata Phil. (<i>Empis</i>)	147	apicis Brun. (<i>Hybos</i>)	20	armata Mel. (<i>Drapetis</i>)	310
annulatus Fall. (<i>Platypalpus</i>)	347	apicis Will. (<i>Tachyempis</i>)	292	armatus, nov. sp. (<i>Syneches</i>)	40
				armillatus, nov. sp. (<i>Platypalpus</i>)	344

DIPTERA

Pages	Pages	Pages			
armimana Old. (<i>Rhamphomyia</i>)	186	auripes Brun. (<i>Hybos</i>)	20	benigna O.-S. (<i>Lamprempis</i>)	136
armipes Bez. (<i>Drapetis</i>)	310	auripila Curr. (<i>Hilara</i>)	380	Bergenstammia Mik (subgen.)	
armipes Cress. (<i>Mythicomyia</i>)	376	<i>australis</i> Frey (<i>Rhamphomyia tenuicornis</i>)	195	<i>Clinocera</i>	230
armipes Lw. (<i>Empis</i>)	147	avida Coq. (<i>Empis</i>)	148	bezzii Frey (<i>Rhamphomyia</i>)	187
armipes, nov. sp. (<i>Microphorus</i>)	91	avida Coq. (<i>Rhamphomyia</i>)	187	bezzii Kert. (<i>Hybos</i>)	20
armipes Sack (<i>Rhamphomyia</i>)	382	avida Harr. (<i>Rhamphomyia culicina</i>)	190	bibioniformis Old. (<i>Rhamphomyia</i>)	187
arrognans Linn. (<i>Tachydromia</i>)	283	azteca Wh.-Mel. (<i>Empis</i>)	148	bicaudata, nov. sp. (<i>Drapetis</i>)	313
arrognans Zett. (<i>Tachydromia annulimana</i>)	283	bacis Walk. (<i>Drapetis</i>)	313	Bicellaria Macq. (gen.)	73
arthritica Mel. (<i>Empis</i>)	147	baculifer Mel. (<i>Hilara</i>)	113	bicincta Tuck. (<i>Clinocera</i>)	237
articulatoides Frey (<i>Platypalpus</i>)	347	bætica Coll. (<i>Hemerocephala</i>)	380	bicingulata Bez. (<i>Hilarempis</i>)	129
articulatus Macq. (<i>Platypalpus</i>)	347	bakeri, nov. sp. (<i>Syneches</i>)	45	bicolor Bell. (<i>Empis</i>)	148
asema Mel. (<i>Empis</i>)	147	bakeri Wh.-Mel. (<i>Euhybos</i>)	33	bicolor Bez. (<i>Drapetis</i>)	313
asiliformis Bon. (<i>Hybos grossipes</i>)	22	<i>baldensis</i> Stro. (<i>Empis pennaria</i>)	168	bicolor Big. (<i>Brachystoma</i>)	13
asiliformis Fab. (<i>Syneches muscarius</i>)	43	<i>baldensis</i> Stro. (<i>Platypalpus</i>)	348	bicolor Big. (<i>Itaphila</i>)	103
assimilis Fall. (<i>Drapetis</i>)	307	<i>ballistrarius</i> , nov. sp. (<i>Platypalpus</i>)	338	bicolor Big. (<i>Syneches bigoti</i>)	41
assimilis Stro. (<i>Empis</i>)	148	<i>ballucatus</i> , nov. sp. (<i>Platypalpus</i>)	336	bicolor Lw. (<i>Boreodromia</i>)	216
astigma Coll. (<i>Chelifera</i>)	378	<i>balnearia</i> White (<i>Hilara</i>)	113	bicolor Macq. 1823 (<i>Rhamphomyia</i>)	
astylata, nov. sp. (<i>Edalea</i>)	63	<i>barbata</i> Lw. (<i>Empis</i>)	179	<i>barbata</i> ; 187, 384	
Atalanta auct. (<i>Clinocera</i>)	227	<i>barbata</i> Macq. (<i>Rhamphomyia</i> , not <i>Empis</i>)	148, 384	bicolor Macq. 1827 (<i>Rhamphomyia</i>)	187, 384
ater, nov. sp. (<i>Syneches</i>)	40	<i>barbata</i> Old. (<i>Tachydromia</i>)	284	bicolor Meigen (<i>Platypalpus</i>)	348
ater Wahlb. (<i>Platypalpus</i>)	347	<i>barbatula</i> Bez. (<i>Hilarempis</i>)	128	bicolor Phil. (<i>Chelifera</i>)	262
aterrima Curt. (<i>Drapetis curvipes</i>)	310,	<i>barbatula</i> Mik (<i>Clinocera</i>)	238	bicolor Phil. (<i>Deuteragonista</i>)	132
	384	<i>barbipes</i> Frey (<i>Hilara</i>)	113	bicolor Walk. (<i>Syneches</i>)	44
aterrima Frey (<i>Rhamphomyia</i>)	186	<i>bares</i> Walk. (<i>Hilara</i>)	113	bicolor Wied. (<i>Rhamphomyia</i>)	187
aterrima Lund. (<i>Drapetis nigritella</i>)	310,	<i>baropeodes</i> , nov. sp. (<i>Euhybos</i>)	32	bicoloripes Brun. (<i>Drapetis</i>)	313
	384	<i>barypoda</i> Coq. (<i>Rhamphomyia</i>)	187	bicornis, nov. sp. (<i>Platypalpus</i>)	320
aterrima Meij. (<i>Syndyas</i>)	37	<i>barypodus</i> , nov. sp. (<i>Syneches</i>)	40	<i>bicuspidata</i> Coll. (<i>Empis pennaria</i>)	370
aterrima Ros. (<i>Bicellaria spuria</i>)	76, 384	<i>basalis</i> Bez. (<i>Drapetis</i>)	313	<i>bicuspidata</i> Eng. (<i>Wiedemannia</i>)	242
atra Cress. (<i>Mythicomyia</i>)	376	<i>basalis</i> Coll. (<i>Drapetis</i>)	310	<i>bifasciata</i> Oliv. (<i>Empis</i>)	148
atra Lw. (<i>Hilara</i>)	113	<i>basalis</i> Lw. (<i>Empis</i>)	148	<i>bifasciata</i> Rossi (<i>Tachydromia arro-</i>	
atra Macq. (<i>Empis leptodopus</i>)	160	<i>basalis</i> Lw. (<i>Hilara</i>)	113	<i>gans</i>	284
atra Meig. (<i>Bicellaria spuria</i>)	77	<i>basalis</i> Lw. (<i>Oreogeron</i>)	100	<i>bifilata</i> Coq. (<i>Rhamphomyia</i>)	187
atra Meig. (<i>Rhamphomyia</i>)	186	<i>basalis</i> Lw. (<i>Rhamphomyia</i>)	187	<i>bigoti</i> Bez. (<i>Syneches</i>)	41
atra Walker (<i>Drapetis aterrima</i>)	310	<i>basalis</i> , nov. sp. (<i>Trichina</i>)	55	<i>bigoti</i> Mel. (<i>Empis</i>)	148
atra Wied. (<i>Empis</i>)	148	<i>basilaris</i> Beck. (<i>Empis</i>)	148	<i>bihamata</i> Bez. (<i>Drapetis</i>)	313
atrata Coq. (<i>Rhamphomyia</i>)	186	<i>batylmenensis</i> Frey (<i>Rhamphomyia</i> <i>paradoxa</i>)	201	<i>bihamata</i> Meij. (<i>Drapetis hirsutitibia</i>)	314
atratus Coq. (<i>Microphorus</i>)	93	<i>bazini</i> Coll. (<i>Empis</i>)	379	<i>bilineata</i> Lw. (<i>Empis digramma</i>)	154
atriiceps Bohem. (<i>Dysalitria</i>)	288	<i>beckeri</i> Dahl. (<i>Stilpon graminum</i>)	304	<i>bilineata</i> Meig. (<i>Rhamphomyia</i>)	187
Atrichopleura Bez. (gen.)	134	<i>beckeri</i> Mik (<i>Platypalpus</i>)	348	<i>bilineatus</i> Mel. (<i>Microphorus</i>)	93
atrifemur Wh.-Mel. (<i>Empis</i>)	148	<i>beckeri</i> Mik (<i>Wiedemannia</i>)	242	<i>bilobata</i> Coll. (<i>Trichina</i>)	383
atrifennis Zett. (<i>Rhamphomyia vesiculosus</i>)	210	<i>beckeri</i> , nov. sp. (<i>Chersodromia</i>)	296	<i>bilobata</i> Old. (<i>Wiedemannia</i>)	245
atripes Mel. (<i>Trichina</i>)	55	<i>beckeri</i> Stro. (<i>Hilara</i>)	113	<i>bimaculata</i> Bez. (<i>Leptopeza</i>)	67
atripes Stro. (<i>Empis tessellata</i>)	175	<i>beckeri</i> Stro. (<i>Microphorus</i>)	93	<i>bimaculata</i> Lw. (<i>Tachydromia</i>)	284
attenuata Frey (<i>Rhamphomyia ignobilis</i>)	187, 385	<i>beckeri</i> Stro. (<i>Platypalpus eneadienicus</i>)	353	<i>bimaculata</i> Mel. (<i>Tachydromia maculipennis</i>)	285
aucta Old. (<i>Rhamphomyia</i>)	187	<i>bella</i> Mel. (<i>Hilara</i>)	114	<i>binotata</i> Lw. (<i>Clinocera</i>)	237
aucta Zett. (<i>Clinocera appendiculata</i>)	236	<i>bellatorius</i> White (<i>Empis</i>)	148	<i>binotata</i> , nov. sp. (<i>Tachypesa</i>)	275
aurantiacus Coll. (<i>Platypalpus</i>)	381	* <i>bellus</i> Heer (<i>Hilarites</i>)	369	<i>binotata</i> Meij. (<i>Drapetis</i>)	313
aurata Coq. (<i>Hilara</i>)	113	<i>benhami</i> Mill. (<i>Hilarempis</i>)	129	<i>biplagiata</i> Hal. (<i>Clinocera</i>)	239
aurata Vill. (<i>Empis ciliata</i>)	151			<i>bipila</i> Stro. (<i>Rhamphomyia</i>)	187
				<i>biplagiata</i> Bez. (<i>Leptoptera</i>)	67

Pages	Pages	Pages
bipunctata Schum. (<i>Hemerodromia</i>) 254	brachystyla Beck. (<i>Platypalpus brunneitibia</i>) 349	brunnipes Str. (<i>Platypalpus albisetosus</i>) 346
biroi Bez. (<i>Ptilophylldromia</i>) 268	braueri Mik (<i>Wiedemannia</i>) 245	brusewitzii Holm. (<i>Rhamphomyia</i>) 187
biroi Bez. (<i>Rhamphomyia</i>) 187	braueri Stro. (<i>Hilara albifrons</i>) 114, 383	brussnevi Frey (<i>Rhamphomyia</i>) 187
biseta Coll. (<i>Hilara</i>) 380	brevicornis Frey (<i>Platypalpus sub-brevis</i>) 349	bucinator Mel. (<i>Euthyneura</i>) 53
bisetosus Bez. (<i>Hybos</i>) 20	brevicornis Lw. (<i>Empis picipes</i>) 149, 385	*bulbirostris Meun. (<i>Empis</i>) 150
bispina Mel. (<i>Drapetis</i>) 308	brevicornis Lw. (<i>Hormopeza</i>) 97	bulbosa Mel. (<i>Anthalia</i>) 61
bistortæ Meig. (<i>Empis</i>) 148	brevicornis Meij. (<i>Chersodromia</i>) 296	bullata Bez. (<i>Empis</i>) 150
bistortæ Walk. (<i>Empis</i>) 148	brevicornis Stro. (<i>Œdalea</i>) 64	bullata Mel. (<i>Hormopeza</i>) 97
bistigma Bez. (<i>Tachydromia</i>) 284	brevicornis Zett. (<i>Platypalpus</i>) 348	*burmanica Ckll (<i>Electrocyrtoma</i>) 368
bistigma Curt. (<i>Wiedemannia</i>) 245	brevicula, n. n. (<i>Drapetis</i>) 310	* Burmitempis Ckll. (gen.) 368
bistriata Stro. (<i>Rhamphomyia flava</i>) 192	brevifurca, nov. sp. (<i>Bicellaria</i>) 75	cacuminifer Mel. (<i>Empis</i>) 150
bistriata Zett. (<i>Hilara</i>) 114	brevifurca Stro. (<i>Hilara clypeata</i>) 116	cælebs Beck. (<i>Antheptiscopus</i>) 107
biuncinata, nov. sp. (<i>Drapetis</i>) 313	brevior Brun. (<i>Drapetis</i>) 308	cæligena Mel. (<i>Empis</i>) 150
bivittata Lw. (<i>Clinocera</i>) 236	brevior Meij. (<i>Syndyas</i>) 37	Cænotus Cole (gen.) 377
bivittata Phil. (<i>Hemerodromia</i>) 254	Brevios Brun. (<i>Platypalpus Macq.</i>) 315	cærulescens Old. (<i>Hilara</i>) 114
bivittata Stro. (<i>Hilara longevity-tata</i>) 114, 385	brevipennis Ros. (<i>Tachydromia</i>) 286,	cæsia Wied.-Meig. (<i>Rhamphomyia</i>) 187
bivittata Stro. (<i>Rhamphomyia flaviventris</i>) 192	384	calcarata Bez. (<i>Empis</i>) 150
bivittata Wied. (<i>Empis</i>) 148	brevipennis Zett. (<i>Chersodromia arenaria</i>) 296	calcarata Stro. (<i>Tachydromia</i>) 284
bivittatus Macq. (<i>Platypalpus</i>) 348	brevipes Lw. (<i>Euthyneura</i>) 53	calcaratus Coll. (<i>Platypalpus</i>) 381
bivittatus Macq. (<i>Scelolabes</i>) 71	brevipila Lw. (<i>Hilara</i>) 114	calcarifera Bez. (<i>Drapetis</i>) 313
blasigii Mik (<i>Parathallassius</i>) 87	brevipila Old. (<i>Rhamphomyia</i>) 382	calceana Meig. (<i>Tachydromia</i>) 284
Blepharoprotæ Lw. (<i>Brachystoma</i>) 12	brevirostris Macq. (<i>Empis</i>) 149	calceatus Bonsd. (<i>Platypalpus bicolor</i>) 348
bohemani Zett. (<i>Wiedemannia</i>) 244	brevis Bez. (<i>Drapetis brevicerca</i>) 310	calceatus Meig. (<i>Platypalpus</i>) 349, 384
boja Schrank (<i>Empis ciliata</i>) 151	brevis Lw. (<i>Empis</i>) 149	californica Coq. (<i>Rhamphomyia</i>) 187
boliviana Bez. (<i>Chelifera</i>) 262	brevis Lw. (<i>Rhamphomyia</i>) 187	california Greene (<i>Mythicomyia</i>) 376
boliviana Bez. (<i>Rhamphomyia</i>) 187	*brevis Meun. (<i>Drapetis</i>) 310	caligaris, n. n. (<i>Platypalpus</i>) 349
borealis Linn. (<i>Empis</i>) 148	brevistyla Coll. (<i>Hilara</i>) 380	caligatus Mel. (<i>Platypalpus caligatus</i>) 349
borealis O. Fab. (<i>Rhamphomyia nigrita</i>) 199	brevistylata Old. (<i>Rhamphomyia</i>) 382	caliginosa Coll. (<i>Rhamphomyia</i>) 382
borealis Old. (<i>Hilara</i>) 114	brevitibia, nov. sp. (<i>Clinocera</i>) 235	callithrix, nov. sp. (<i>Platypalpus</i>) 325
borealis Zett. (<i>Leptopeza</i>) 67	breviventris Frey (<i>Rhamphomyia</i>) 187	callositibia Bez. (<i>Drapetis</i>) 313
borealis Zett. (<i>Wiedemannia bistrigata</i>) 245	breviventris Phil. (<i>Apalocnemis</i>) 101	calopoda Bez. (<i>Lampræmpis</i>) 136
Boreodromia Coq. (gen.) 215	breviventris Phil. (<i>Deuteragonista</i>) 132	calva Mel. (<i>Drapetis</i>) 313
bovina Beck. (<i>Hilara</i>) 114	brevivittata Bonsd. (<i>Hilara bistrigata</i>) 114	calva Mel. (<i>Tachyempis</i>) 292
bracata Rond. (<i>Œdalea</i>) 64	brevivittata Bonsd. (<i>Hilara pilipes</i>) 123	calvimontis Ckll. (<i>Rhamphomyia</i>) 187
brachialis Mel. (<i>Tachypeza</i>) 276	brevivittata Macq. (<i>Hilara</i>) 114	Camelopsis Eng. (subgen. <i>Wiedemannia</i>) 241
brachialis, n. sp. (<i>Hemerodromia</i>) 255	Brochella (nov. gen.) 108	cana Bez. (<i>Chelifera</i>) 262
brachialis Rond. (<i>Hybos</i>) 20	brouni Hutt. (<i>Empis</i>) 149	cana Coq. (<i>Hilara</i>) 114
brachygastera Phil. (<i>Haplomera</i>) 131	brunetti, n. n. (<i>Platypalpus</i>) 349	cana Zett. (<i>Rhamphomyia</i>) 187
brachyrrhyncha Thom. (<i>Hilarempis</i>) 129	brunnea Coq. (<i>Empis</i>) 149	canaliculata Macq. (<i>Rhamphomyia</i>) 187
brachysoma Coq. (<i>Empis</i>) 149	brunnea, nov. sp. (<i>Hemerodromia</i>) 255	canaster Mel. (<i>Empis</i>) 150
brachystigma Bez. (<i>Hybos</i>) 20	brunneitibia Stro. (<i>Platypalpus</i>) 349	candicans Fall. (<i>Platypalpus</i>) 349
*brachystoma Ckll. (<i>Protœdalea</i>) 371	brunnipennis Meig. (<i>Empis</i>) 149	candicans Lw. (<i>Rhamphomyia</i>) 187
brachystoma Coq. (<i>Hilara umbrosa</i>) 127	brunnipennis, nov. sp. (<i>Clinocera</i>) 232	candicans Zett. (<i>Platypalpus fasciatus</i>) 353
Brachystoma Meig. (gen.) 12	brunnipes Brun. (<i>Hybos</i>) 20	candida Rossi (<i>Empis</i>) 150
brachystoma Phil. (<i>Hilarempis</i>) 129	brunnipes Gimm. (<i>Platypalpus</i>) 349	candidata Lw. (<i>Empis</i>) 150
BRACHYSTOMATINÆ (subf.) 11	brunnipes Macq. (<i>Drapetis ænescens</i>) 310	candidatus Mel. (<i>Parathallassius</i>) 87
	brunnipes Meig. (<i>Empis</i>) 150	candidiseta Bez. (<i>Platypalpus</i>) 350
	brunnipes Steph. (<i>Hybos</i>) 20	canescens Zett. (<i>Hilara</i>) 114

DIPTERA

Pages	Pages	Pages			
<i>cantabrica</i> Stro. (<i>Empis ciliatopennata</i>)	151	<i>chichimeca</i> Wh.-Mel. (<i>Lamprempis</i>)	136	<i>cinerea</i> Müll. (<i>Empis</i>)	151
<i>cantabrica</i> Stro. (<i>Hilara bistrata</i>)	114	<i>chibinensis</i> Frey (<i>Rhamphomyia</i>)	188	<i>cinerea</i> , nov. sp. (<i>Tachyempis</i>)	290
<i>canus</i> Mel. (<i>Platypalpus</i>)	350	<i>chilensis</i> Bez. (<i>Haplomera</i>)	131	<i>cinerea</i> Zett. (<i>Empis</i>)	151
<i>capensis</i> Schin. (<i>Hilara</i>)	114	<i>chilensis</i> Phil. (<i>Platypalpus</i>)	350	<i>cineromicans</i> auct. (<i>Hilara rejecta</i>)	380
<i>capnodes</i> , nov. sp. (<i>Tachyempis</i>)	289	<i>chiloensis</i> Brethes (<i>Haplomera</i>)	380	<i>cineromicans</i> Stro. (<i>Hilara</i>)	115
<i>capnopterus</i> , nov. sp. (<i>Oreogeton</i>)	99	<i>chionochæta</i> Bez. (<i>Platypalpus</i>)	350	<i>cinerovittatus</i> Stro. (<i>Platypalpus</i>)	350
<i>captus</i> Coq. (<i>Empis</i>)	150	<i>chionoptera</i> Bez. (<i>Rhamphomyia</i>)	188	<i>cingulata</i> Bez. (<i>Apalocnemis</i>)	101
<i>captus</i> Coq. (<i>Hemerodromia</i>)	255	<i>choiptera</i> Beck. (<i>Empis helophila</i>)	158	<i>cingulata</i> Dahlb. (<i>Hilara</i>)	115
<i>carbonaria</i> Brun. (<i>Empis</i>)	150	<i>choiptera</i> Meig. (<i>Empis</i>)	150	<i>cingulata</i> Gimm. (<i>Empis</i>)	152
<i>carbonaria</i> Mel. (<i>Hilara</i>)	114	<i>choiptera</i> Meig. (<i>Empis dasypoda</i>)	153	<i>cingulatus</i> Lw. (<i>Platypalpus</i>)	350
<i>carbonaria</i> Wied.-Meig. (<i>Rhamphomyia</i>)	187	<i>chiragra</i> Bez. (<i>Empis</i>)	151	Cladodremia Bez. (subgen. <i>Chelifera</i>)	259
<i>carbonella</i> Zett. (<i>Hilara</i>)	114	<i>chiragra</i> Bez. (<i>Parahybos</i>)	46	<i>clarandus</i> Coll. (<i>Platypalpus</i>)	381
* <i>carbonum</i> Germ. (<i>Empis</i>)	150	<i>chiragra</i> , nov. sp. (<i>Microphorella</i>)	89	<i>claripennis</i> Mel. (<i>Prorates</i>)	377
<i>carenifera</i> Bez. (<i>Rhamphomyia</i>)	188	<i>chiragrica</i> Speiser (<i>Hilara</i>)	115	<i>claripennis</i> Old. (<i>Rhamphomyia</i>)	382
<i>carinata</i> Bez. (<i>Hilara</i>)	129	Chiromantis Rond. (<i>Chelipoda</i> Macq.)	263	<i>claripennis</i> Stro. (<i>Hybos grossipes</i>)	22
<i>carinthiaca</i> Beck. (<i>Hilara fuscipes</i>)	115, 385	<i>chloropa</i> Bez. (<i>Hoplopesa</i>)	71	<i>clauda</i> Coq. (<i>Empis</i>)	152
<i>carinthiaca</i> Str. (<i>Hilara</i>)	115	Choreodremia Frey (subgen. <i>Rhamphomyia</i>)	180	<i>clauda</i> Coq. (<i>Rhamphomyia</i>)	188
<i>carteri</i> Coll. (<i>Platypalpus</i>)	381	<i>chorica</i> Fall. (<i>Hilara</i>)	115	<i>clauda</i> Schrank (<i>Empis</i>)	152
<i>castanipes</i> Lw. (<i>Hilara</i>)	115	<i>chorica</i> Schin. (<i>Hilara cornicula</i>)	116	<i>clausa</i> Coq. (<i>Empis</i>)	152
<i>castanipes</i> Meig. (<i>Platypalpus albisetosus</i>)	346	<i>chrysonotum</i> Stro. (<i>Platypalpus curvistans</i>)	352	<i>clavator</i> Coq. (<i>Rhamphomyia</i>)	188
<i>castellana</i> Stro. (<i>Empis morio</i>)	164	<i>cilians</i> Frey (<i>Platypalpus ciliaris</i>)	350	<i>clavigera</i> Lw. (<i>Rhamphomyia</i>)	189
<i>catalonica</i> Stro. (<i>Tachydromia</i>)	284	<i>ciliaris</i> Fall. (<i>Platypalpus</i>)	350	<i>clavipes</i> Fab. (<i>Hybos grossipes</i>)	22
<i>cataluna</i> Stro. (<i>Hemerodromia oratoria</i>)	257	<i>ciliata</i> Coq. (<i>Rhamphomyia</i>)	188	<i>clavipes</i> Harris (<i>Hilara ciliipes</i>)	115
<i>caucasica</i> Bez. (<i>Empis</i>)	150	<i>ciliata</i> Fab. (<i>Empis</i>)	151	* <i>clavipes</i> Lw. (<i>Leptopeza</i>)	67
<i>caudata</i> Zett. (<i>Rhamphomyia</i>), 188, 385		<i>ciliata</i> Fall. (<i>Empis plumipes</i>)	169	<i>clavipes</i> Lw. (<i>Tachypeza</i>)	276
<i>caudatula</i> Lw. (<i>Empis</i>)	150	<i>ciliata</i> Macq. (<i>Hilara ciliipes</i>)	115	<i>clavipes</i> Meig. (<i>Trichina</i>)	56
<i>cavaticum</i> Beck. (<i>Lamposoma</i>)	248	<i>ciliata</i> Meig. (<i>Empis pennipes</i>)	168	Cleptodremia Corti (<i>Platypalpus</i>)	315
<i>celer</i> Meig. (<i>Platypalpus</i>)	350	<i>ciliata</i> Mel. (<i>Tachydromia</i>)	285	Clinocera Meig. (gen.)	226
<i>celeripes</i> Meig. (<i>Stilpon</i>)	304	<i>ciliatocosta</i> Bez. (<i>Drapetis</i>)	312	CLINOCERATINÆ (subfam.) 210	
<i>cellarius</i> , nov. sp. (<i>Platypalpus</i>)	337	<i>ciliatopennata</i> Stro. (<i>Empis</i>)	151	Clinocerella Eng. (subgen. <i>Wiedemannia</i>)	241
<i>centralis</i> Brun. (<i>Empis</i>)	150	<i>ciliipes</i> Meig. (<i>Hilara</i>)	115	<i>clypeata</i> Macq. (<i>Rhamphomyia</i>)	189
Cephalodremia Beck. (gen.)	267	<i>ciliipes</i> Say (<i>Rhamphomyia</i>)	188	<i>clypeata</i> Meig. (<i>Hilara</i>)	116, 385
Ceratempis (nov. gen.)	218	<i>ciliipes</i> Schin. (<i>Hilara spinimana</i>)	126	<i>coarctatus</i> Coll. (<i>Platypalpus</i>)	381
Cerathybos Bez (gen.)	36	<i>cimicoides</i> Fab. (<i>Tachydromia arrogans</i>)	284	<i>carnealis</i> Beck. (<i>Hilara scrobiculata</i>)	125
Ceratomerus Phil. (gen.)	214	<i>cimicoides</i> Meig. (<i>Tachydromia connexa</i>)	285	<i>cognata</i> Egg. (<i>Empis</i>)	152
<i>certa</i> Harr. (<i>Empis lutea</i>)	162	<i>cimicoides</i> Walk. (<i>Tachydromia annulimana</i>)	283	<i>cognata</i> Steph. (<i>Empis</i>)	152
<i>certa</i> Walk. (<i>Hilara</i>)	115	<i>cincinnatula</i> Lw. (<i>Empis</i>)	151	Celabria (nov. gen.)	258
<i>cervina</i> Lw. (<i>Hilara</i>)	129	<i>cinefacta</i> Coq. (<i>Rhamphomyia</i>)	188	<i>coleophora</i> , nov. subsp. (<i>Hemero-</i>	
<i>ceylonensis</i> , n. n. (<i>Platypalpus</i>)	381	<i>cineracea</i> Coq. (<i>Rhamphomyia</i>)	188	<i>dromia empiformis</i>)	256
<i>ceylonica</i> Bez. (<i>Empis</i>)	150	<i>cineraria</i> Bez. (<i>Empis velutina</i>)	177	<i>collaris</i> Meig. (<i>Platypalpus</i>)	350
<i>chaetoproctus</i> Bez. (<i>Parahybos</i>)	46	<i>cinerascens</i> Meig. (<i>Rhamphomyia</i>)	188	<i>collateralis</i> , nov. sp. (<i>Platypalpus</i>)	327
Chamaedipsa Mik. (subgen. <i>Wiedemannia</i>)	242	<i>cinerea</i> Big. (<i>Empis bigoti</i>)	148	<i>collina</i> Phil. (<i>Bicellaria</i>)	76
Charadrodremia , nov. gen.	292	<i>cinerea</i> Fab. (<i>Rhamphomyia</i>)	188	<i>collina</i> Phil. (<i>Empis polita</i>)	169
<i>chelana</i> , nov. sp. (<i>Tachydromia</i>)	282	<i>cinerea</i> Macq. (<i>Hilara</i>)	115	<i>collusor</i> Mel. (<i>Chelifera</i>)	263
Chelifera Macq. (gen.)	258	<i>cinerea</i> Meig. (<i>Rhamphomyia tipularia</i>)	208	Coleoboneura Mel. (gen.)	293
Chelipoda Macq. (gen.)	263			<i>colonica</i> Walk. (<i>Empis</i>)	152
Chersodremia Walk. (gen.)	294			<i>colorata</i> Coq. (<i>Rhamphomyia</i>)	189

Pages	Pages	Pages			
comata Mel. (<i>Drapetis</i>)	313	coquilletti Mel. (<i>Platypalpus</i>)	351	* crinitarsis Lw. (<i>Rhamphomyia</i>)	189
comata, nov. sp. (<i>Clinocera</i>)	232	coquilletti, nov. sp. (<i>Euhybos</i>)	27	crocata Coq. (<i>Euthyneura</i>)	53
combinata Beck. (<i>Dolichocephala</i>)	246	coracina Bez. (<i>Empis</i>)	152	Crossopalpus Big. (<i>Drapetis</i>)	
commendatus Beck. (<i>Platypalpus</i>)	350	coracina Old. (<i>Hilara</i>)	116	Meig.)	305
commiles Walk. (<i>Platypalpus</i>)	351	coracina Zett. (<i>Rhamphomyia</i>)	189	cryptospina Frey (<i>Platypalpus</i>)	351
commutatus Stro. (<i>Platypalpus</i>)	351	coracula Lundb. (<i>Hilara</i>)	116	Ctenodrapetis Bez. (subgen.	
compacta Brun. (<i>Hilara</i>)	116	corycrica Bez. (<i>Empis</i>)	152	<i>Drapetis</i>)	307
completa Lw. (<i>Empis</i>)	152	coriacea Big. (<i>Iteaphila</i>)	103	ctenistes, nov. sp. (<i>Clinocera</i>)	234
compta Coq. (<i>Empis</i>)	152	cormus Walk. (<i>Iteaphila</i>)	103	culiciformis Fab. (<i>Hybos</i>)	20
compta Coq. (<i>Leptopeza</i>)	67	cornicula Lw. (<i>Hilara</i>)	116	culiciformis Gmel. (<i>Hybos grossipes</i>)	22
compta Coq. (<i>Rhamphomyia</i>)	189	* corrupta Meun. (<i>Rhamphomyia</i>)	189	culicina Fall. (<i>Rhamphomyia</i>)	189
compta Zett. (<i>Rhamphomyia pen-</i> <i>nata</i>)	202	corticalis Mel. (<i>Tachypeza</i>)	276	cuneata Lw. (<i>Hilara</i>)	116
comptus Walk. (<i>Platypalpus</i>)	351	corvina Lw. (<i>Empis</i>)	152	cuneipennis Bez. (<i>Empis</i>)	153
compungens Walk. (<i>Platypalpus</i>)	351	corvina Lw. (<i>Rhamphomyia</i>)	189	cuneipennis Mel. (<i>Drapetis</i>)	312
* concinna Meun. (<i>Leptopeza</i>)	67	Coryneta auct. (<i>Platypalpus</i>)		cuneipennis Mel. (<i>Platypalpus</i>)	351
concinnaicauda Coll. (<i>Chelifera</i>)	378	Macq.)	315	cursitans Fab. (<i>Platypalpus</i>)	352
* concinnus Ckll. (<i>Mesomyites</i>)	370	Coryneta Meig. (<i>Tachydromia</i>)		cursitans Frey (<i>Platypalpus candi-</i> <i>cans</i>)	350
concisa Lw. (<i>Empis</i>)	152	Meig.)	279	cursitans Lundb. (<i>Platypalpus lati-</i> <i>cinctus</i>)	352, 384
* concitatus Meun. (<i>Platypalpus</i>)	351	costalis Ros. (<i>Tachydromia</i> , not		cursitans Meig. (<i>Platypalpus major</i>)	359
concolor Verr. (<i>Empis</i>)	152	<i>Tachypeza</i>)	276, 384	cursitans Zett. (<i>Chersodromia</i>)	296
confidens Harr. (<i>Empis femorata</i>)	156	costata Zett. (<i>Rhamphomyia</i>)	189	cursoia Scop. (<i>Tachypeza nubila</i>)	277
confinis Bonsd. (<i>Platypalpus fas-</i> <i>cipes</i>)	353	cothurnata Big. (<i>Ocydromia</i>)	69	curta Lw. (<i>Empis</i>)	153
confinis Zett. (<i>Platypalpus</i>)	351	cothurnata Brullé (<i>Empis</i>)	152	curticornis Zett. (<i>Platypalpus</i>)	352
confinis Zett. (<i>Rhamphomyia</i>)	189	cothurnatus Macq. (<i>Platypalpus</i>)	351	curtipes Beck. (<i>Heleodromia</i>)	225
confirmata Walk. (<i>Hilara</i>)	116	cotoxanthus Blanch. (<i>Empis</i>)	152	curtisi Coll. (<i>Hilara</i>)	380
confusa Beck. (<i>Empis albipennis</i>)	152,	coxalis, nov. sp. (<i>Colabris</i>)	258	curva Curr. (<i>Iteaphila</i>)	381
	385	coxalis Ros. (<i>Ocydromia glabricula</i>)	69	curvata Beck (<i>Cephalodromia</i>)	268
conformis Frey (<i>Rhamphomyia stig-</i> <i>mosa</i>)	206	coxalis Thom. (<i>Empis</i>)	152	curvinervis Old. (<i>Rhamphomyia</i>)	190
conformis Kow. (<i>Rhamphomyia la-</i> <i>vipes</i>)	196	coxatus Zett. (<i>Platypalpus annulipes</i>)	347	curvinervis Zett. (<i>Drapetis moriella</i>)	311
confusa Lw. (<i>Empis maculata</i>)	163	crassa Bez. (<i>Atrichopleura</i>)	134	curveurina, nov. sp. (<i>Syneches</i>)	41
congregaria Mel. (<i>Hilara</i>)	116	crassa Lw. (<i>Drapetis anescens</i>)	310	curvipes Coq. (<i>Rhamphomyia</i>)	190
coniatus, nov. sp. (<i>Hybos</i>)	20	crassa Now. (<i>Empis</i>)	152	curvipes Fab. (<i>Syneches</i>)	41
conjuncta Coq. (<i>Iteaphila</i>)	103	crassicauda Stro. (<i>Rhamphomyia</i>)	189	curvipes Lw. (<i>Empis</i>)	153
conjuncta Lw. (<i>Clinocera</i>)	237	crassifemoris Fitch (<i>Platypalpus</i>)	351	curvipes Meig. (<i>Drapetis flavipes</i>)	309
conjuncta Lw. (<i>Rhamphomyia</i>)	189	crassifila Lw. (<i>Empis</i>)	153	curvipes Meig. (<i>Drapetis</i> , not assi- milis)	309, 384
connexa Beck. (<i>Empis</i>)	152	crassimana Stro. (<i>Rhamphomyia</i>)	189	curvipes, nov. sp. (<i>Stilpon</i>)	303
connexa Meig. (<i>Tachydromia</i>)	285	crassinervis Lw. (<i>Rhamphomyia</i>)		curvipes Siebke (<i>Hilara</i>)	116
conservativa Mall. (<i>Rhamphomyia</i>)	189	<i>sordida</i>)	205	curvula Frey (<i>Rhamphomyia</i>)	190
consobrina Zett. (<i>Euthyneura myr-</i> <i>tilli</i> , not <i>Anthepiscopus</i>)	107, 384	crassipes Fab. (<i>Euhybos</i>)	33	cuspidatus, nov. sp. (<i>Euhybos</i>)	32
consortus Steph. (<i>Platypalpus</i>)	351	crassipes Fourer. (<i>Hilara maura</i>)	122	cyanea Bell. (<i>Lampræmpis</i>)	136
constans Harr. (<i>Empis livida</i>)	161	crassipes Macq. (<i>Microphorus</i> , not		cyanescens Bez. (<i>Hilarempis</i>)	129
contigua Lw. (<i>Empis</i>)	152	<i>anomalus</i>)	92, 384	cyanogaster Wh.-Mel. (<i>Rhampho-</i> <i>myia</i>)	190
contiguus, nov. sp. (<i>Platypalpus</i>)	320	crassipes Meig. (<i>Empis</i>)	153	cyanophthalmus Str. (<i>Symbalophthal-</i> <i>mus dissimilis</i>)	300
convergens Coll. (<i>Drapetis</i>)	378	crassipes Schrank (<i>Empis chioptera</i>)	151	cylindrica Fab. (<i>Empis</i>)	153
cophas Walk. (<i>Rhamphomyia</i>)	189	crassirostris Fall. (<i>Rhamphomyia</i>)		cymballista, nov. sp. (<i>Oreogeton</i>)	99
Coptophlebia Bez. (subgen.		<i>nigripes</i>)	199	cyrenaica Bez. (<i>Empis</i>)	379
<i>Empis</i>)	141	crassiseta Stro. (<i>Platypalpus</i>)	351	Cyrtoma Meig. (<i>Bicellaria</i>	
copulifera, nov. sp. (<i>Hormopeza</i>)	96	<i>Macq.</i>)	73		

DIPTERA

	Pages		Pages		Pages
czernyi Bez. (<i>Wiedemannia</i>)	244	dichrous Bez. (<i>Syneches</i>)	41	ditæniata Bez. (<i>Clinocera</i>)	237
czernyi Stro. (<i>Hilara</i>)	116	difficilis Frey (<i>Platypalpus</i>)	352	divergens Bez. (<i>Drapetis distans</i>)	308
czernyi Stro. (<i>Hilara borealis</i>)	114	difficilis Lundb. (<i>Chersodromia</i>)	296	divergens Lw. (<i>Drapetis</i>)	308
dalmatica Old. (<i>Empis</i>)	379	digramma Meig. (<i>Empis</i>)	154	divergens Lw. (<i>Empis</i>)	154
dalmatica Old. (<i>Rhamphomyia</i>)	382	dilatativittatus Stro. (<i>Platypalpus</i>)	352	diversa Coq. (<i>Rhamphomyia</i>)	191
dalmatina Str. (<i>Hilara</i>)	116	dilutata Frey (<i>Tachypeza servicer-palpis</i>)	278	diversa Mel. (<i>Drapetis</i>)	308
dalmatinus Stro. (<i>Microphorus</i>)	93	dilutior, nov. var. (<i>Platypalpus flavirostris</i>)	325	diversicauda Coll. (<i>Chelifera</i>)	378
dalmatinus Stro. (<i>Platypalpus</i>)	352	dilutipes Stro. (<i>Drapetis moriella</i>)	311	diversipennis Beck. (<i>Rhamphomyia</i>)	191
dana Walk. (<i>Rhamphomyia</i>)	190	dimidiata Bell. (<i>Lactistomyia</i>)	35	diversipes Coq. (<i>Platypalpus</i>)	352
daria Walk. (<i>Rhamphomyia</i>)	190	dimidiata Lw. (<i>Clinocera</i>)	239	diversipes Mel. (<i>Drapetis</i>)	310
dasychira Mik (<i>Empis</i>)	153	dimidiata Lw. (<i>Empis locutiana</i>)	161	diversipes Mel. (<i>Tachydromia</i>)	285
dasynota Lw. (<i>Empis</i>)	153	dimidiata Lw. (<i>Rhamphomyia</i>)	190	diversipes Stro. (<i>Hilara</i>)	117
daspoda Egg. (<i>Empis</i>)	153	dimidiata Meig. (<i>Empis</i>)	154	diversipes Stro. (<i>Platypalpus stroblianus</i>)	365
dasyprocta Lw. (<i>Empis</i>)	153	dimidiata Stro. (<i>Hilara</i>)	116	dividua Mel. (<i>Drapetis</i>)	308
Dasyrhamphomyia Frey (sub-gen. <i>Rhamphomyia</i>)	182	dimidiatus Lw. (<i>Euhybos electus</i>)	34	divisa Lw. (<i>Empis</i>)	154
dasythrix Meij. (<i>Empis plumipes</i>)	153, 385	dimidiatus Walk. (<i>Euhybos</i>)	34	divisus Walk. (<i>Platypalpus</i>)	352
debulis Coq. (<i>Syneches</i>)	41	dinopus, nov. sp. (<i>Euhybos</i>)	31	dolabraria Mel. (<i>Empis</i>)	154
debilis Lw. (<i>Platypalpus crassifemoris</i>)	351	dinoscelis Bez. (<i>Syneches</i>)	45	dolicheretma Mel. (<i>Clinocera</i>)	235
debilis Lw. (<i>Rhamphomyia</i>)	190	Dionaea auct. (<i>Rhamphomyia</i> Meig.)	180	Dolichocephala Macq. (gen.)	245
decolor, nov. sp. (<i>Platypalpus</i>)	321	Dionaea Meig. (<i>Empis</i> Linn.)	138	dolichocera Bez. (<i>Rhamphomyia</i>)	191
* decolorata Meun. (<i>Drapetis</i>)	308	Dipsomyia Bez. (gen.)	220	dolichopodina Schin. (<i>Lamprematis</i>)	136
decora Lundb. (<i>Empis plumpipes</i>)	153, 386	direptor, nov. sp. (<i>Platypalpus</i>)	310	dolichoptera Bez. (<i>Rhamphomyia</i>)	191
decora Meig. (<i>Empis</i>)	153	discalis Mel. (<i>Drapetis</i>)	310	dolorosa, nov. sp. (<i>Tachypeza</i>)	275
* decorata Meun. (<i>Drapetis</i>)	310	discifer Lw. (<i>Platypalpus</i>)	352	dolorosa Wh.-Mel. (<i>Empis</i>)	154
decoripes Roder (<i>Empis</i>)	154	discifera, nov. sp. (<i>Tachypeza</i>)	273	* dolosa Meun. (<i>Chelipoda</i>)	266
dedecor Lw. (<i>Empis</i>)	154	discoidalis Beck. (<i>Rhamphomyia</i>)	190	dorsalis Brun. (<i>Hemerodromia</i>)	255
defecta Lw. (<i>Chelifera</i>)	263	discoidalis Bez. (<i>Drapetis</i>)	312	dorsalis Curt. (<i>Empis punctata</i>)	171
defessa Will. (<i>Hemerodromia</i>)	255	discoidalis Lundb. (<i>Hilara</i>)	117	dorsalis Lw. (<i>Syndyas</i>)	37
deficiens Walk. (<i>Syneches</i>)	45	discoidalis Meij. (<i>Hybos</i>)	21	dorsalis Meig. (<i>Ocydromia glabricula</i>)	69
* definita Meun. (<i>Drapetilla</i>)	368	discolor Lw. (<i>Empis</i>)	154	dorsata Beck. (<i>Rhamphomyia</i>)	191
* defunctus Handl. (<i>Microphorus</i>)	93	discolor Stro. (<i>Hilara</i>)	117	dorsata, nov. sp. (<i>Hemerodromia</i>)	255
* delicate Meun. (<i>Chelipoda</i>)	265	disconvenita Mel. (<i>Empis dolabraria</i>)	154	dorsieta Mel. (<i>Drapetis</i>)	310
denominatus Frey (<i>Platypalpus cur-silans</i>)	352	dispar Adams (<i>Drapetis</i>)	313	dracophylli Miller (<i>Hilara</i>)	380
dentata Old. (<i>Rhamphomyia</i>)	190	dispar Curt. (<i>Ocydromia glabricula</i>)	69	Drapetilla Meun. (gen.)	368
denticulata Old. (<i>Tachydromia</i>)	285	dispar Old. (<i>Bicollaria</i>)	76	Drapetis Meig. (gen.)	305
dentipes Wied. (<i>Euhybos</i>)	34	dispar Scholz (<i>Empis</i>)	154	drapetoides Walk. (<i>Microphorus</i>)	93
dentipes Zett. (<i>Rhamphomyia</i>)	190	dispar Zett. (<i>Rhamphomyia</i>)	190	Drymodromia Beck. (gen.)	263
depilis Lw. (<i>Empis</i>)	154	disparilis Coq. (<i>Rhamphomyia</i>)	191	Dryodromia Rond. (<i>Synampho-tera</i> Lw.)	216
derodactylus, nov. sp. (<i>Euhybos</i>)	33	disparilis Mel. (<i>Leptopeza</i>)	67	dubia Schin. (<i>Empis femorata</i>)	156
desertorum Beck. (<i>Platypalpus</i>)	352	dissimilipes, nov. sp. (<i>Platypalpus</i>)	318	dubius Walk. (<i>Platypalpus</i>)	352
* detestata Meun. (<i>Chelifera</i>)	260	dissimilis Fall. (<i>Symbalophthalmus</i>)	300	dudai Old. (<i>Rhamphomyia</i>)	382
* detestata Meun. (<i>Hemerodromia</i>)	255	dissimilis Zett. (<i>Rhamphomyia maculipennis</i>)	197	dumetorum Phil. (<i>Empis</i>)	155
Deuteragonista Phil. (gen.)	132	distans Bez. (<i>Drapetis</i>)	308	dunicola Phil. (<i>Hilara</i>)	129
* diabolica Meun. (<i>Dysalatris</i>)	288	distans Lw. (<i>Empis</i>)	154	duplex, nov. sp. (<i>Syneches</i>)	41
diaphorina O.-S. (<i>Lamprematis</i>)	136	* distans Lw. (<i>Rhamphomyia</i>)	191	duplex Walk. (<i>Euhybos</i>)	34
dichætophorus Bez. (<i>Syneches</i>)	41	distans, nov. sp. (<i>Tachypeza</i>)	273	duplicis Coq. (<i>Rhamphomyia</i>)	191
dichogenus, nov. sp. (<i>Syneches</i>)	41	distincta White (<i>Drapetis</i>)	378	dusmeti Stro. (<i>Empis</i>)	155
dichroa Bez. (<i>Tachyempis</i>)	292	distinguendum Schin. (<i>Brachystoma</i>)	13	Dysalatris Lw. (gen.)	287
dichroa Meig. (<i>Platypalpus bicolor</i>)	348				

Pages	Pages	Pages
ecalceatus Zett. (<i>Platypalpus</i> , not <i>calceatus</i>) 349, 384	*errabunda Meun. (<i>Rhamphomyia</i>) 191	exquisita Mall. (<i>Coloboneura</i>) 294
ecetra Walk. (<i>Rhamphomyia</i>) 191	erythrophthalma Meig. (<i>Rhamphomyia</i>) 191	extricatus Coll. (<i>Platypalpus</i>) 381
echinata White (<i>Hilarempis</i>) 129	erythrophthalma Stro. (<i>Rhamphomyia pusilla</i>) 203	exul O.-S. (<i>Drapetis</i>) 314
ectorus, nov. sp. (<i>Hybos</i>) 21	escheri Zett. (<i>Wiedemannia</i>) 243	facialis Mel. (<i>Drapetis</i>) 310
effera Coq. (<i>Rhamphomyia</i>) 191	escorialensis Stro. (<i>Hilara</i>) 117	fagorum Dahl (<i>Platypalpus</i>) 353
efficiens White (<i>Hilara</i>) 117	Eucelidia Mik (subgen. <i>Wiedemannia</i> Zett.) 241	falcata Mel. (<i>Empis</i>) 155
*egelata Meun. (<i>Tachypeza</i>) 276	eucera, nov. sp. (<i>Niphogenia</i>) 217	fallaciosa Lw. (<i>Wiedemannia</i>) 244
Elaphropeza Macq. (subgen. <i>Drapetis</i>) 307	eudamides Walk. (<i>Empis</i>) 155	fallax Egg. (<i>Empis</i>) 155
* Electrocyrtoma Ckll. (gen.) 368	Eudrapetis Mel. (subgen. <i>Drapetis</i> = <i>Crossopalpus</i> Big.) 307, 384	falleni Meig. (<i>Rhamphomyia vesculosa</i>) 210
electus Mel. (<i>Euhybos</i>) 34	Euhybos Coq. (gen.) 24	fasciata Meig. (<i>Hilara</i>) 117
electus Mel. (<i>Euhybos duplex</i>) 34	eumelanus Mik (<i>Platypalpus ater</i>) 348	fasciata Schum. (<i>Empis</i>) 155
elegans Bez. (<i>Hilarempis</i>) 129	eumera Bez. (<i>Syndyas</i>) 37	fasciatus Lundb. (<i>Platypalpus curvifrons</i>) 353, 384
elegans Big. (<i>Ocydromia</i>) 69	eumera Lw. (<i>Empis</i>) 155	fasciatus Meig. (<i>Platypalpus</i>) 353
elegans Brun. (<i>Empis</i>) 155	eumerus Bez. (<i>Platypalpus</i>) 353	fasciculata Stro. (<i>Empis</i>) 155
*elegans Meun. (<i>Palaoedalea samilandica</i>) 370	eunordquistii Frey (<i>Rhamphomyia obscura</i>) 201	fascifemorata Brun. (<i>Drapetis</i>) 308
elevatus Bez. (<i>Syneches</i>) 42	eupeza Lw. (<i>Empis</i>) 155	fascipes Meig. (<i>Platypalpus</i>) 353
elongata Hal. (<i>Trichina</i>) 56	eupterota Frey (<i>Rhamphomyia albohirta</i>) 382	fascipes Roser (<i>Drapetis flavipes</i>) 309
elongata Meig. (<i>Empis</i>) 155	eupterota Lw. (<i>Rhamphomyia</i>) 191	fascipes Stro. (<i>Platypalpus flavipes</i>) 355
elongata Meij. (<i>Syndyas</i>) 37	eurypterus Bez. (<i>Euhybos</i>) 34	fascipennis Meij. (<i>Chelipoda</i>) 266
elongata Mel. (<i>Chelipoda</i>) 266	eustylatus Big. (<i>Syneches</i>) 45	fascipennis Zett. (<i>Rhamphomyia</i>) 191
* Empidia Weyenb. (gen.) 369	Euthyneura Macq. (gen.) 52	fasciventris, nov. sp. (<i>Platypalpus</i>) 343
EMPIDINÆ (subfam.) 78	* Euthyneurella Meun. (gen.) 369	femoralis Bez. (<i>Chelifera</i>) 262
empidiformis Beck. (<i>Rhamphomyia</i>) 191	eversmanni Lw. (<i>Empis</i>) 155	femoralis Bez. (<i>Drapetis femorata</i>) 308
empiformis Say (<i>Hemerodromia</i>) 255	*eversoris Meun. (<i>Platypalpus</i>) 353	femoralis Bonsd. (<i>Platypalpus minoratus</i>) 360
Empimorpha Coq. (gen.) 178	excisa Lw. (<i>Tachydromia</i>) 285	femoralis Wh.-Mel. (<i>Drapetis</i>) 310
Empis Linn. (gen.) 138	excisa, nov. sp. (<i>Tachypeza</i>) 274	femoralis Zett. (<i>Platypalpus exiguum</i>) 353
enecator Mel. (<i>Tachydromia</i>) 285	excitus Beck. (<i>Platypalpus</i>) 353	femorata Fab. (<i>Empis</i>) 155
*enena Ckll. (<i>Rhamphomyia</i>) 191	exigua Lw. (<i>Rhamphomyia</i>) 191	femorata Heeger (<i>Stilpon lunata</i>) 304
enervatus, nov. sp. (<i>Platypalpus</i>) 333	exiguus auct. (<i>Platypalpus niger</i>) 353, 384	femorata Lw. (<i>Haploxytoma</i>) 78
engadinica Old. (<i>Rhamphomyia</i>) 382	exiguus Meun. (<i>Platypalpus minutus</i>) 353,	femorata Lw. (<i>Hilara</i>) 117
engadinicus Mik (<i>Platypalpus</i>) 352	384	femorata Mel. (<i>Drapetis</i>) 308
Enicopteryx Steph. (<i>Rhamphomyia</i> Meig.) 180	exilis auct. (<i>Drapetis incompleta</i>) 378	femorata, nov. sp. (<i>Anthalia</i>) 59
enodis Mel. (<i>Empis</i>) 155	exilis Coq. (<i>Empis</i>) 155	femorata, nov. sp. (<i>Hemorodromia</i>) 256
Enoplempis Big. (subgen. <i>Empis</i>) 142	exilis Macq. (<i>Drapetis flavipes</i>) 309	femorata Walk. (<i>Empis palparis</i>) 167
*eocenica Meun. (<i>Micrempis</i>) 299	exilis Meig. (<i>Drapetis</i>) 308, 384	femoratus Müll. (<i>Hybos</i>) 21
*eocenicus Meun. (<i>Microphorus</i>) 93	exilis Meig. (<i>Platypalpus</i>) 353	femorella Zett. (<i>Hilara nitidula</i>) 123
ephippiata Fall. (<i>Drapetis</i>) 313	*exilis Meun. (<i>Hilara</i>) 117	fenestrata Say (<i>Tachypeza</i>) 276
ephippium Scholz (<i>Hilara</i>) 117	*exilis Meun. (<i>Hybos</i>) 21	fenestrata Zett. (<i>Tachydromia sabulosa</i>) 286
epibosca, nov. sp. (<i>Tachyempis</i>) 291	exilis Zett. [1838] (<i>Drapetis assimilis</i>) 309, 384	ferruginea Brun. (<i>Drapetis</i>) 314
Epiceia Walk. (<i>Syneches</i> Walk.) 39	eximus Old. (<i>Platypalpus</i>) 381	ferruginea Fab. (<i>Rhamphomyia</i>) 192
erberi Mik (<i>Rhamphomyia</i>) 191	exoleta Hal. (<i>Dolichocephala guttata</i>) 247	ferruginea Meig. (<i>Empis trigramma</i>) 176
erberi Now. (<i>Empis tumida</i>) 155, 385	exotica Wied. (<i>Empis</i>) 155	ferruginea Meig. (<i>Rhamphomyia</i>) 192
erecta Coll. (<i>Chelifera</i>) 378	exorrecta, nov. sp. (<i>Proclinopyga</i>) 222	ferruginea Roser (<i>Hilara thoracica</i>) 127,
eremita Beck. (<i>Schistostomia</i>) 94	expulsa Lw. (<i>Rhamphomyia</i>) 191	383
erinacioides Mall. (<i>Rhamphomyia</i>) 191		ferruginea Scop. (<i>Empis stercorea</i>) 174
Eriogaster Macq. (<i>Empis</i> Linn.) 138		ferrugineus Brun. (<i>Platypalpus</i>) 354
erminea Mik (<i>Wiedemannia</i>) 242		ferrugineus Walk. (<i>Syneches</i>) 45
erosa Lw. (<i>Empis</i>) 155		ficana Walk. (<i>Rhamphomyia</i>) 192

DIPTERA

Pages	Pages	Pages			
filata Lw. (<i>Empis</i>)	156	flavipes Fall. (<i>Platypalpus bicolor</i>)	348	fratellus Brun. (<i>Syneches</i>)	42
filata Zett. (<i>Rhamphomyia</i>)	192	flavipes Fall. (<i>Platypalpus calceatus</i>)	349	fraterna Lw. (<i>Empis</i>)	156
filicauda Lundb. (<i>Rhamphomyia</i>)	192	flavipes Fall. (<i>Platypalpus pallidiventris</i>)	362	fraternellus Zett. (<i>Anthepeiscopus adalinus</i>)	107
fimbria Walk. (<i>Empis</i>)	156	flavipes Macq. (<i>Drapetis</i>)	308	Fraudator Hutton (gen.)	379
fimbriata Coq. (<i>Rhamphomyia</i>)	192	flavipes Mats. (<i>Rhamphomyia</i>)	192	frauscheri Stro. (<i>Empis</i>)	156
fimbriatipes Now. (<i>Rhamphomyia squamigera</i>)	206	flavipes Meig. (<i>Hilara</i>)	117	frigeli Zett. (<i>Chelifera</i>)	260
fistulator, nov. sp. (<i>Proclinoptyga</i>)	222	flavipes Meig. (<i>Hybos femoratus</i>)	21	frontalis Coq. (<i>Empis</i>)	156
fiumana Egg. (<i>Empis</i>)	156	flavipes Meig. (<i>Leptopeza ruficollis</i>)	67	frontalis Lw. (<i>Rhamphomyia</i>)	193
fixa Harr. (<i>Rhamphomyia sulcata</i>)	207	flavipes Meig. (<i>Platypalpus vilvipes</i>)	356	fugeo Harr. (<i>Empis livida</i>)	161
flabilis White (<i>Empis</i>)	156	flavipes Phil. (<i>Chelifera</i>)	262	fulcratus Scop. (<i>Platypalpus</i>)	355
flammifer Mel. (<i>Platypalpus</i>)	354	flavipes Steph. (<i>Rhamphomyia</i>)	192	fuliginella Zett. (<i>Rhamphomyia dispar</i>)	191
flava Bonsd. (<i>Rhamphomyia sciarina</i>)	204	flavipes Thunb. (<i>Platypalpus curvifrons</i>)	352	fuliginosa Meig. (<i>Syneches</i>)	42
flava Coq. (<i>Anthalia</i>)	61	flavipes Thunb. (<i>Platypalpus flavicornis</i>)	354	fulva Macq. (<i>Empis</i>)	156
flava Fall. (<i>Rhamphomyia</i>)	192	flavipes Thunb. (<i>Tachypeza truncorum</i>)	278	fulva Phil. (<i>Hilarempis</i>)	129
flava Müll. (<i>Empis</i>)	156	flavipes Will. (<i>Drapetis xanthopoda</i>)	312	fulva Walk. (<i>Empis walkeri</i>)	178
flava Schin. (<i>Hilara thoracica</i>)	117, 383	flavipes Winth.-Meig. (<i>Stilpon celertipes</i>)	304	fulvescens Miller (<i>Leptopeza</i>)	381
flavella Zett. (<i>Chelifera</i>)	260, 385	flavipes Zett. (<i>Edalea</i>)	64	fulvibarba Stro. (<i>Hilara</i>)	117
flavescens Miller (<i>Leptopeza fulvescens</i>)	381	flavipes Zett. (<i>Empis grisea</i>)	157	fulvida Bez. (<i>Drapetis fulvida</i>)	314
flavescens Rossi (<i>Platypalpus</i>)	354	flavipes Zett. (<i>Leptopeza borealis</i>)	67	fulvipes Hutt. (<i>Hilarempis</i>)	129
flavicans Oliv. (<i>Empis</i>)	156	flavipilosus Coq. (<i>Anthepeiscopus</i>)	107	fulvipes Macq. (<i>Hilara rufipes</i>)	125
flavicollis Beck. (<i>Drapetis</i>)	314	flavirostris Lw. (<i>Platypalpus</i>)	355	fulvipes Meig. (<i>Platypalpus flavipes</i>)	355
flavicornis Macq. (<i>Platypalpus</i>)	354	flavirostris Walk. (<i>Rhamphomyia</i>)	192	fulvipes Phil. (<i>Haplomera</i>)	131
flavicornis Meig. (<i>Platypalpus</i>)	354	flavitarsis Ros. (<i>Empis</i>)	156	fulvipes Wied.-Meig. (<i>Empis</i>)	156
flavicornis Mel. (<i>Drapetis</i>)	308	flaviventris Macq. (<i>Rhamphomyia</i>)	192	fulvithorax Wulp (<i>Drapetis</i>)	314
flavicoxa, nov. var. (<i>Proclinoptyga monogramma</i>)	223	flaviventris Stro. (<i>Empis pennaria</i>)	168	fulvolanata Frey (<i>Rhamphomyia</i>)	193
flavicoxa Scholz (<i>Rhamphomyia</i>)	192	flaviventris Stro. (<i>Platypalpus adencinus</i>)	362	fumida Coq. (<i>Empis</i>)	157
flavicoxa, nov. sp. (<i>Anthepeiscopus</i>)	107	flavobasalis Mats. (<i>Empis</i>)	156	fumipennis Meig. (var. <i>Hybos femoratus</i>)	21
flavicoxis Beck. (<i>Platypalpus</i>)	354	flavohalterata Stro. (<i>Hilara bistrigata</i>)	114	fumipennis Stro. (<i>Drapetis pusilla</i>)	309
flavida Brun. (<i>Chelipoda</i>)	267	flavus Macq. (<i>Platypalpus</i>)	355	fumipennis Zett. (<i>Rhamphomyia villosa</i>)	193, 385
flavida Will. (<i>Drapetis</i>)	314	flexicauda Zett. (<i>Rhamphomyia</i>)	192	fumosa Hutt. (<i>Timaphes</i>)	98
flavimana Zett. (<i>Leptopeza</i>)	67	flexuosa Coq. (<i>Rhamphomyia</i>)	192	fumosa Lw. (<i>Empis</i>)	157
flavinceris Miller (<i>Hilara</i>)	380	fuscicornis Lw. (<i>Drapetis ambigua</i> , <i>not mortella</i>)	311, 384	fumosa Lw. (<i>Rhamphomyia</i>)	193
flavinervis Phil. (<i>Empis</i>)	156	florisomma Lw. (<i>Empis</i>)	156	funebris Fall. (<i>Hybos culiciformis</i>)	20
flavipalpis Meig. (<i>Platypalpus pallipes</i>)	363	* florissantana Ckll. (<i>Empis</i>)	156	funebris Meig. (<i>Empis</i>)	157
flavipalpis Zett. (<i>Platypalpus maculatus</i>)	359	fluvialis Brun. (<i>Clinocera</i>)	236	funebris Meig. (<i>Hybos grossipes</i>)	22
flavipalpus Macq. (<i>Platypalpus luteocornis</i>)	357	fontinalis Miller (<i>Chelifera</i>)	378	funesta Meig. (<i>Empis</i>)	157
flavipennis Walk. (<i>Platypalpus</i>)	354	fontinalis Hal. (<i>Clinocera</i>)	237	furcatus Zett. (<i>Anthepeiscopus adalinus</i>)	107
flavipes auct. (<i>Platypalpus coarcatus</i>)	354, 381	forcipata Linn. (<i>Rhamphomyia</i>)	192	furcifer, nov. sp. (<i>Bicellaria</i>)	75
flavipes auct. (<i>Platypalpus interstinctus</i>)	354, 381	formalis Walk. (<i>Platypalpus</i>)	355	furcifer Wh-Mel. (<i>Rhamphomyia</i>)	193
flavipes Bez. (<i>Leptopeza setigera</i>)	68	* formosa Lw. (<i>Rhamphomyia</i>)	193	fusca Brun. (<i>Chelipoda</i>)	267
flavipes Brun. (<i>Hybos</i>)	22	formosae Bez. (<i>Drapetis</i>)	314	fusca Brun. (<i>Trichopoda</i>)	219
flavipes Brun. (<i>Parahybos</i>)	46	fortis Beck. (<i>Hilara</i>)	117	fusca Steph. (<i>Rhamphomyia</i>)	193
flavipes Brun. (<i>Platypalpus brunetti</i>)	349	fortis Bez. (<i>Drapetis</i>)	310	fuscescens Bez. (<i>Syneches</i>)	42
flavipes Cress. (<i>Mythicomysa</i>)	376	fossalis Miller (<i>Hilara</i>)	380	fuscicornis Bez. (<i>Chelipoda</i>)	266
flavipes Fab. (<i>Platypalpus</i>)	354			fuscicornis Frey (<i>Platypalpus lastadinorum</i>)	357
				fuscicornis Zett. (<i>Platypalpus</i>)	355
				fuscimana Bez. (<i>Chelifera</i>)	262

Pages	Pages	Pages			
<i>fuscimanus</i> Zett. (<i>Platypalpus albisetosus</i>)	346	<i>geniculata</i> Meig. (<i>Rhamphomyia</i>)	193	<i>gracilis</i> Lw. (<i>Hilara</i>)	118
<i>fuscinervis</i> Frey (<i>Tachydromia</i>)	285	<i>geniculata</i> Ros. (<i>Hilara litorea</i>)	118, 383	<i>gracilis</i> Lw. (<i>Rhamphomyia</i>)	194
<i>fuscipennis</i> Bonsd. (<i>Rhamphomyia coracina</i>)	189	<i>geniculata</i> Zett. (<i>Iteaphila macquarti</i>)	104	* <i>gracilis</i> Meun. (<i>Palæoleptopeza</i>)	371
<i>fuscipennis</i> Brun. (<i>Parahybos</i>)	46	<i>geniculata</i> Zett. (<i>Rhamphomyia squamigera</i>)	206	* <i>gracilis</i> Meun. (<i>Timalphes</i>)	98
<i>fuscipennis</i> Brun. (<i>Syueches</i>)	42	<i>geniculatus</i> Fall. (<i>Platypalpus albocapillatus</i>)	346	<i>gracilis</i> Scholz (<i>Platypalpus flavipes</i>)	355
<i>fuscipennis</i> Fall. (<i>Tachypeza</i>)	277	<i>geniculatus</i> Fall. (<i>Platypalpus exiguus</i>)	353	<i>gracilitarsis</i> Stro. (<i>Empis</i>)	157
<i>fuscipennis</i> Frey (<i>Iteaphila obscura</i>)	104	<i>geniculatus</i> Meig. (<i>Platypalpus</i>)	356	<i>grallatrix</i> Bez. (<i>Rhamphomyia</i>)	194
<i>fuscipennis</i> Lw. (<i>Clinocera</i>)	236	<i>geniculatus</i> Wulp (<i>Hybos</i>)	22	<i>graminum</i> auct. (<i>Stilpon nubila</i>)	304, 383
<i>fuscipennis</i> Macq. (<i>Ocydromia</i>)	69	<i>genitivus</i> , nov. sp. (<i>Euhybos</i>)	28	<i>graminum</i> Fall. (<i>Stilpon</i>)	304
<i>fuscipennis</i> Macq. (<i>Platypalpus luteicornis</i>)	357	<i>gentilis</i> Brun. (<i>Platypalpus</i>)	356	<i>grammoptera</i> Frey (<i>Rhamphomyia</i>)	194
<i>fuscipennis</i> Meig. (<i>Gloma</i>)	102	<i>gentilis</i> Lw. (<i>Rhamphomyia</i>)	194	<i>grandis</i> Lw. (<i>Empis crassa</i>)	153
<i>fuscipennis</i> Ros. (<i>Heleodromia immaculata</i>)	225	<i>genualis</i> Coq. (<i>Clinocera</i>)	237	<i>granditarsis</i> Curr. (<i>Hilara</i>)	380
<i>fuscipennis</i> Zett. (<i>Rhamphomyia</i>)	193	<i>genualis</i> Stro. (<i>Empis decora</i>)	153	<i>gratiosa</i> Beck. (<i>Chersodromia</i>)	296
<i>fuscipes</i> Bez. (<i>Micrempis</i>)	299	<i>gesticularis</i> , nov. sp. (<i>Platypalpus</i>)	328	<i>gravidus</i> Mel. (<i>Platypalpus</i>)	356
<i>fuscipes</i> Bonsd. (<i>Hilara griseola</i>)	118	<i>ghigiana</i> Bez. (<i>Empis</i>)	157	<i>gravipes</i> Lw. (<i>Empis</i>)	157
<i>fuscipes</i> Brullé (<i>Empis</i>)	157	<i>gibba</i> Fall. (<i>Rhamphomyia</i>)	194	<i>gravis</i> Wied.-Meig. (<i>Empis</i>)	157
<i>fuscipes</i> Fab. (<i>Hilara</i>)	117, 385	<i>gibbifera</i> Stro. (<i>Rhamphomyia</i>)	194	<i>grisea</i> Fall. (<i>Empis</i>)	157
<i>fuscipes</i> Gmel. (<i>Empis</i>)	157	<i>gibbipes</i> Stro. (<i>Empis</i>)	157	<i>grisea</i> Fall. (<i>Empis maculipes</i>)	163
<i>fuscipes</i> Zett. (<i>Microphorus velutinus</i>)	94	<i>gibbosa</i> Gmel. (<i>Empis</i>)	157	<i>grisea</i> Wied.-Meig. (<i>Empis hoffmannseggii</i>)	158
<i>fuscipes</i> Zett. (<i>Trichina elongata</i>)	56	<i>gilvihirta</i> Coq. (<i>Authalia</i>)	61	<i>griseiceps</i> Beck. (<i>Platypalpus alpinus</i>)	346
<i>fuscisetata</i> Bez. (<i>Chelipoda</i>)	266	<i>gilvipes</i> Coq. (<i>Platypalpus xanthopodus</i>)	367	<i>griseifrons</i> Coll. (<i>Hilara</i>)	380
<i>fuscitarsis</i> Zett. (<i>Platypalpus</i>)	356	<i>gilvipes</i> Lw. (<i>Rhamphomyia</i>)	194	<i>griseiventris</i> Phil. (<i>Hilarempis</i>)	129
<i>fuscohalteratus</i> Mel. (<i>Platypalpus hians</i>)	356	<i>gilvipes</i> Meig. (<i>Platypalpus</i>)	356	<i>griseola</i> Zett. (<i>Hilara</i>)	118
<i>fuscula</i> Zett. (<i>Rhamphomyia</i>)	193	<i>gilvipilosa</i> Coq. (<i>Rhamphomyia</i>)	194	<i>griseola</i> Zett. (<i>Rhamphomyia dispar</i>)	191
<i>fuscula</i> Zett. (<i>Rhamphomyia dispar</i>)	191	<i>glaber</i> Meig. (<i>Platypalpus</i>)	356, 384	<i>griseonigra</i> Brun. (<i>Empis</i>)	158
<i>fuscum</i> Phil. (<i>Brachystoma</i>)	13	<i>glaber</i> Zett. (<i>Platypalpus luteus</i>)	358	<i>griseonigra</i> Brun. (<i>Rhamphomyia</i>)	194
<i>fusitibia</i> Stro. (<i>Hilara</i>)	118	<i>glabra</i> Lw. (<i>Rhamphomyia</i>)	194	<i>grossipes</i> Linn. (<i>Hybos</i>)	22
<i>gagatina</i> , nov. sp. (<i>Tachyempis</i>)	289	<i>glabratulus</i> Meig. (<i>Platypalpus</i>)	356	Grossopalpus Scud. (see <i>Drapetis</i>)	305
<i>gagatinus</i> Big. (<i>Hybos</i>)	22	<i>glabricula</i> Fall. (<i>Ocydromia</i>)	69	<i>gryphus</i> , nov. sp. (<i>Euhybos</i>)	30
<i>gaigeri</i> Gerke (<i>Empis</i>)	157	<i>glacialis</i> , nov. sp. (<i>Platypalpus</i>)	331	<i>gubernans</i> , nov. sp. (<i>Clinocera</i>)	234
<i>galactodes</i> Bez. (<i>Rhamphomyia</i>)	193	<i>gladiator</i> Mel. (<i>Empis</i>)	157	<i>guifilar</i> Frey (<i>Rhamphomyia</i>)	194
<i>galactoptera</i> Stro. (<i>Hilara</i> , not <i>pseudosartrix</i>)	124, 380, 385	<i>glaucia</i> Coq. (<i>Rhamphomyia</i>)	194	<i>gulosa</i> Coq. (<i>Empis</i>)	158
<i>galactoptera</i> Stro. (<i>Rhamphomyia</i>)	193	<i>glaucella</i> Frey (<i>Rhamphomyia subglauccella</i>)	206	<i>guttata</i> Hal. (<i>Dolichocephala</i>)	247
<i>gallica</i> Meig. (<i>Hilara</i>)	118	<i>glaucella</i> Zett. (<i>Rhamphomyia</i>)	194	<i>gyllenhalli</i> Zett. (<i>Euthyneura</i>)	53
<i>garretti</i> Curr. (<i>Hilara</i>)	380	<i>glaucus</i> Beck. (<i>Heterotropus</i>)	375	<i>gymnaspis</i> Bez. (<i>Hilarempis</i>)	129
<i>gayi</i> Macq. (<i>Haplomera</i>)	131	<i>globifera</i> Stro. (<i>Rhamphomyia longipes</i>)	197	<i>gymnopoda</i> Bez. (<i>Empis</i>)	158
<i>gemmea</i> Bez. (<i>Lamprempis</i>)	136	<i>globulipes</i> Meig. (<i>Hilara maura</i>)	122	<i>gymnopoda</i> Bez. (<i>Haplomera</i>)	131
<i>geneatis</i> Mel. (<i>Empimorpha</i>)	179	Gloma Meig. (gen.)	101	<i>hæmi</i> Lw. (<i>Empis</i>)	158
* <i>generosa</i> Meun. (<i>Ragas</i>)	109	<i>gonatopus</i> Speis. (<i>Hemerodromia</i>)	256	<i>hæmorrhoidalis</i> Beck. (<i>Clinocera</i>)	239
<i>geniculata</i> Big. (<i>Rhamphomyia</i>)	193	<i>gracilipes</i> Bohem. (<i>Hilara flavipes</i>)	117	<i>halidayi</i> Coll. (<i>Euthyneura</i>)	379
<i>geniculata</i> Bonsd. (<i>Rhamphomyia plumipes</i>)	202	<i>gracilipes</i> Lw. (<i>Rhamphomyia squamigera</i>)	206	<i>halidayi</i> Coll. (<i>Tachydromia</i>)	383
<i>geniculata</i> Fall. (<i>Drapetis moriella</i>)	311	<i>gracilipes</i> Phil. (<i>Empis</i>)	157	Halsanalotes Beck. (gen.)	298
<i>geniculata</i> Kby. (<i>Iteaphila luctuosa</i>)	104	<i>gracilipes</i> Phil. (<i>Tenontomyia</i>)	133	<i>halteralis</i> Bez. (<i>Hybos</i>)	22
<i>geniculata</i> Macq. (<i>Empis macquarti</i>)	157,	<i>gracilis</i> Bez. (<i>Drapetis</i>)	312	* <i>halteralis</i> Ckll. (<i>Burmitempis</i>)	368
	379	<i>gracilis</i> Curt. (<i>Empis</i>)	157	<i>halteralis</i> Lw. (<i>Bicellaria</i>)	76
				<i>halterata</i> Beck. (<i>Gloma</i>)	102
				<i>halterata</i> , nov. sp. (<i>Tachyempis</i>)	292

DIPTERA

	Pages		Pages		Pages
hambergi Frey (<i>Rhamphomyia</i>)	194	Hilaropus White (<i>Hilarempis</i> Bez.)	128	humilis Frey (<i>Drapetis</i>)	310
hamifera Bez. (<i>Drapetis</i>)	312	himalayana Brun. (<i>Rhamphomyia</i>)	194	huttoni Bez. (<i>Hilarempis fulvipes</i>)	129
hamifera, nov. sp. (<i>Clinocera</i>)	233	hirsuta Beck. (<i>Empis</i>)	158	hyænoides, nov. var. (<i>Platypalpus juvenis</i>)	332
hamiltoni Miller (<i>Brachystoma</i>)	378	hirsuta Beck. (<i>Hilara</i>)	119	hyalina Bez. (<i>Lactistomyia</i>)	36
Haplocyrtoma , nov. gen.	78	hirsuta Old. (<i>Rhamphomyia</i>)	382	hyalina Brullé (<i>Rhamphomyia</i>)	195
Haplomera Macq. (gen.)	130	hirsutipes Beck. (<i>Ocydromia</i>)	70	hyalinata Meig. (<i>Empis vernalis</i>)	177
Haplomerinx Bez. (subgen. <i>Empis</i>)	142	hirsutipes Coll. (<i>Rhamphomyia</i>)	382	hyalinipennis Brethes (<i>Mythicomysia</i>)	376
Harpamerus Big. (<i>Syneches Walk.</i>)	39	hirsutitibia Meij. (<i>Drapetis</i>)	314	hyalinus Coq. (<i>Syneches</i>)	42
harpiger Mel. (<i>Platypalpus</i>)	356	hirsutus, nov. sp. (<i>Anthepiscopus</i>)	106	hyalipennis Fall. (<i>Empis</i>)	158
harti Mall. (<i>Tachydromia</i>)	285	*hirta Lw. (<i>Gloma</i>)	102	hyalipennis Macq. (<i>Tachyempis</i>)	292
* Hasmona Giebel (gen.)	369	hirta Str. (<i>Hilara</i>)	119	hyalipennis Schin. (<i>Empis albivenis</i>)	146
hastata Mik (<i>Wiedemannia</i>)	243	hirta Walk. (<i>Coloboneura</i>)	294	hyalogyne Bez. (<i>Empis</i>)	158
hastatus Mel. (<i>Platypalpus</i>)	356	hirtella Coll. (<i>Hilara</i>)	380	hyalopterus Bez. (<i>Euhybos</i>)	34
heeri Zett. (<i>Tachypeza</i>)	277	hirticrus, n. n. (<i>Empis</i>)	158	hyalopterus Bez. (<i>Syneches</i>)	45
Heleodromia auct. (<i>Clinocera</i> Meig.)	227	hirtiloba Speiser (<i>Clinocera</i>)	378	Hybos Meig. (gen.)	18
Heleodromia Hal. (gen.)	224	hirtimana Old. (<i>Rhamphomyia</i>)	382	<i>hybos</i> Lam. (<i>Syneches muscarius</i>)	43
helleni Frey (<i>Rhamphomyia</i>)	194	hirtipes Bez. (<i>Atrichofleura</i>)	134	hybotina Fall. (<i>Œdalia</i>)	64
helophila Lw. (<i>Empis</i>)	158	hirtipes Coq. (<i>Empis hirticrus</i>)	158	hybotina Zett. (<i>Rhamphomyia</i>)	195
helvolus Frey (<i>Syneches</i>)	45	hirtipes Coll. (<i>Hilara</i>)	380	HYBOTINÆ (subfam.)	15
Hemerodromia auct. (<i>Chelipoda</i> Macq.)	258, 259	hirtipes Lw. (<i>Rhamphomyia</i>)	194	<i>hybrida</i> Zett. (<i>Rhamphomyia sciarina</i>)	204
Hemerodromia auct. (<i>Chelipoda</i> Macq.)	264	hirtipes, nov. sp. (<i>Tachydromia</i>)	282	<i>hybridus</i> Frey (<i>Platypalpus agilis</i>)	345
Hemerodromia Meig. (gen.)	251, 254	hirtipes Wied. (<i>Empis</i>)	158	Hydromeria Macq. (subgen. <i>Clinocera</i>)	229
HEMERODROMIINÆ (sub-fam.)	248	hirtula Zett. (<i>Hilara</i>)	119	hygrobia Lw. (<i>Wiedemannia</i>)	244
hercynica Old. (<i>Rhamphomyia</i>)	382	hirtula Zett. (<i>Rhamphomyia</i>)	194	hyperborea Frey (<i>Hilara</i>)	380
herschelli Mall. (<i>Rhamphomyia</i>)	194	hispanica Lw. (<i>Empis morio</i>)	164	*hypolitha Ckll. (<i>Rhamphomyia</i>)	195
Hesperiempis Mel. (gen.)	111	hispanicus Stro. (<i>Platypalpus cur-</i> <i>sitanus</i>)	352	hystrichopyga Bez. (<i>Empis</i>)	158
Heterempis Brethes (<i>Hilarempis</i> Bez.)	128	hoffmannseggii Lw. (<i>Empis</i>)	158	hystrix Lw. (<i>Empis</i>)	158
Heterhybos Brethes (<i>Mythicomyia</i> Coq.)	375	holmgreni Zett. (<i>Œdalia</i>)	64	hystrix Stro. (<i>Hilara</i>)	119
heterochroma Bez. (<i>Rhamphomyia</i>)	194	Holeclora Mel. (<i>Microphorus</i> Macq.)	90	ibericus Arias (<i>Pistainia</i>)	287
heterogamus, nov. sp. (<i>Oreogeton</i>)	100	Holeclora Schin. (subg. <i>Rhamphomyia</i>)	182	ignobilis Zett. (<i>Rhamphomyia</i>)	195, 385
heterogastra Lw. (<i>Hilarempis</i>)	129	holosericea Meig. (<i>Rhamphomyia gibba</i>)	194	ignota Meig. (<i>Empis punctata</i>)	171
heterogastra Nov. (<i>Hilara</i>)	119	holosericea Thom. (<i>Hilarempis</i>)	129	immaculata Hal. (<i>Heleodromia</i>)	225
Heterophlebus Phil. (<i>Brachystoma</i> Meig.)	12	holosericea Walk. (<i>Empis</i>)	158	immaculatus Beck. (<i>Platypalpus</i>)	356
heteroptera Macq. (<i>Empis</i>)	158	holosericeus Meig. (<i>Microphorus velutinus</i>)	94	immaculatus Brun. (<i>Syneches</i>)	42
Heterotropus Lw. (gen.)	375, 377	holosericus Mel. (<i>Platypalpus</i>)	356	immatura Zett. (<i>Euthyneura</i>)	53
hians Mel. (<i>Platypalpus</i>)	356	Hemalecnemis Phil. (gen.)	14	impedita Lw. (<i>Rhamphomyia</i>)	195
hiemalis White (<i>Microphorus</i>)	93	Heplopoza Zett. (gen.)	70	impennis Stro. (<i>Empis albivenis</i>)	146
Hilara Meig. (gen.)	112	Hermopeza Zett. (gen.)	94	imperfecta Lw. (<i>Microsanita stigma-</i> <i>talis</i>)	374
hilariformis Bez. (<i>Hilarempis</i>)	129	houghii Mel. (<i>Chersodromia</i>)	296	impexus Mel. (<i>Platypalpus</i>)	356
Hilarempis Bez. (gen.)	128	hovgaardii Holmgr. (<i>Rhamphomyia</i>)	195	implicata Coll. (<i>Hilara</i>)	380
hilariformis Frey (<i>Rhamphomyia</i>)	194	Howlettia Brun. (<i>Platypalpus</i> Macq.)	315	impudica Mik. (<i>Wiedemannia fallaci-</i> <i>ciosa</i>)	244
Hilarimorpha Schin. (gen.)	373	hudsoni Hutt. (<i>Hilara</i>)	119	inæqualis Brun. (<i>Syneches</i>)	42
* Hilarites Heer (gen.)	369	humeralis, nov. sp. (<i>Tachypeza</i>)	276	inca Bez. (<i>Chelifera</i>)	262
		humilis Coq. (<i>Empis</i>)	158		

Pages	Pages	Pages
<i>incana</i> Beck. (<i>Chersodromia beckeri</i>) 296	integer Walk. (<i>Euhybos</i>) 34	<i>labiata</i> Lw. (<i>Empis</i>) 159
<i>incana</i> Walk. (<i>Chersodromia</i>) 296	<i>intercepta</i> , n. n. (<i>Empis</i>) 159	<i>lacertosus</i> , nov. sp. (<i>Platypalpus</i>) 319
<i>incertus</i> Bez. (<i>Parahybos</i>) 46	* <i>interfectoris</i> Meun. (<i>Platypalpus</i>) 356	<i>lacteipennis</i> , nov. sp. (<i>Anthalia</i>) 59
<i>incertus</i> Coll. (<i>Platypalpus</i>) 381	<i>interjectus</i> Lundb. (<i>Platypalpus</i>) 356	<i>lacteipennis</i> Stro. (<i>Hilara</i>) 119
<i>inchoata</i> Mel. (<i>Thinodromia</i>) 297	<i>intermedia</i> Fall. (<i>Hilara</i> , not <i>fuscipes</i>) 118, 385	<i>lacteipennis</i> Stro. (<i>Rhamphomyia</i> <i>galactoptera</i>) 193
<i>incisa</i> Brun. (<i>Tachypeza</i>) 277	<i>intermedia</i> Fall. (<i>Hilara quadri-</i> <i>vittata</i>) 125	<i>lacteipennis</i> Zett. (<i>Rhamphomyia</i> <i>niveipennis</i>) 200
<i>inclinata</i> Bez. (<i>Empis</i>) 159	<i>intermedia</i> Frey (<i>Rhamphomyia</i>) 195	<i>lacteiseta</i> Coll. (<i>Platypalpus</i>) 357
<i>incompleta</i> Beck. (<i>Tachydromia</i>) 285	<i>intermedia</i> Lundb. (<i>Bicellaria</i>) 76	Lactistomyia Mel. (gen.) 35
<i>incompleta</i> Coll. (<i>Drapetis</i>) 378	<i>interrupta</i> Beck. (<i>Empis intercepta</i>) 159	<i>laestadianorum</i> Frey (<i>Platypalpus</i>) 357
<i>incompleta</i> Lw. (<i>Rhamphomyia</i>) 195	<i>interrupta</i> Lw. (<i>Tachydromia</i>) 285	<i>laeta</i> Lw. (<i>Empis</i>) 159
<i>incompleta</i> Macq. (<i>Empis</i>) 159	<i>interrupta</i> , nov. sp. (<i>Anthalia</i>) 60	<i>laeta</i> Mel. (<i>Drapetis</i>) 314
<i>incompleta</i> Stro. (<i>Euthyneura myr-</i> <i>tilli</i>) 54	<i>interstincta</i> Fall. (<i>Hilara</i>) 119	<i>laetabilis</i> Coll. (<i>Empis</i>) 379
<i>incompta</i> Wahlb. (<i>Empis maculipes</i>) 163	<i>interstincta</i> Meig. (<i>Hilara pilosa</i>) 124	<i>laetabilis</i> , nov. sp. (<i>Platypalpus</i>) 324
<i>inconspicua</i> Brun. (<i>Empis</i>) 159	<i>interstinctus</i> Coll. (<i>Platypalpus</i>) 381	<i>laetus</i> Lw. (<i>Platypalpus</i>) 357
<i>inculta</i> Coq. (<i>Drapetis</i>) 310	<i>inusitata</i> Mel. (<i>Coloboneura</i>) 294	<i>lævigata</i> Lw. (<i>Empis</i>) 159
<i>incurvus</i> Mel. (<i>Platypalpus</i>) 356	<i>inusta</i> Mel. (<i>Tachypeza</i>) 277	<i>lævigata</i> Lw. (<i>Rhamphomyia</i>) 195
<i>indica</i> Brun. (<i>Chelipoda</i>) 266	* <i>involuta</i> Meun. (<i>Rhamphomyia</i>) 195	<i>lævipes</i> Fall. (<i>Rhamphomyia</i>) 195
<i>induta</i> Bez. (<i>Empis</i>) 159	<i>iridipennis</i> Kert. (<i>Parahybos</i>) 46	<i>lævipes</i> Beck. (<i>Drapetis</i>) 309
<i>inermis</i> Beck. (<i>Edalea</i>) 64	Ironomyia White (gen.) 48, 373	<i>lævipes</i> Lw. (<i>Empis</i>) 159
<i>inermis</i> Lw. (<i>Clinocera</i>) 239	<i>irregularis</i> Lw. (<i>Rhamphomyia</i>) 195	<i>lagunæ</i> Beck. (<i>Wiedemannia</i>) 244
<i>inermis</i> Mel. (<i>Drapetis</i>) 309	<i>irritans</i> Bez. (<i>Hilara</i>) 119	Lamachella (nov. gen.) 71
<i>infans</i> Zett. (<i>Hilara</i>) 119	<i>irrorata</i> Fall. (<i>Dolichocephala</i>) 247	<i>lamellata</i> Coll. (<i>Rhamphomyia</i>) 382
<i>infector</i> Coll. (<i>Platypalpus</i>) 381	<i>isommatus</i> , nov. sp. (<i>Microphorus</i>) 92	<i>lamellata</i> Lw. (<i>Wiedemannia</i>) 244
<i>inferialis</i> , nov. sp. (<i>Platypalpus</i>) 321	Iteaphila Zett. (gen.) 102	<i>lamellicornis</i> Beck. (<i>Empis</i>) 159, 379
<i>inflexa</i> Mel. (<i>Drapetis</i>) 314	<i>italica</i> Lw. (<i>Iteaphila</i>) 103	<i>lamellicornis</i> Lundb. (<i>Empis gym-</i> <i>nopoda</i>) 385
<i>infumata</i> Beck. (<i>Empis liosoma</i>) 160	<i>jacobsoni</i> Meig. (<i>Empis</i>) 159	<i>laminata</i> Coll. (<i>Empis pennaria</i>) 379
<i>infumata</i> Coq. (<i>Empis</i>) 159	<i>jardinei</i> White (<i>Syneches</i>) 383	Lamposoma Beck. (gen.) 248
<i>infumata</i> Mel. (<i>Drapetis</i>) 309	<i>javana</i> Meij. (<i>Leptopeza</i>) 67	Lamprempis Wh.-Mel. (gen.) 135
<i>infumata</i> Wied. (<i>Rhamphomyia</i>) 195	<i>jeanneli</i> Beck. (<i>Drymodromia</i>) 263	<i>lanceolata</i> , nov. sp. (<i>Edalea</i>) 63
<i>infuscata</i> Brullé (<i>Hilara infans</i>) 119	<i>johnsoni</i> Coq. (<i>Hilara</i>) 119	<i>landbecki</i> Phil. (<i>Empis</i>) 159
<i>infuscata</i> Lw. (<i>Edalea</i>) 64	<i>johnsoni</i> Mel. (<i>Empis</i>) 159	<i>laniventris</i> Esch. (<i>Empis</i>) 159
<i>infuscata</i> Meig. (<i>Rhamphomyia</i>) 195	<i>jugorum</i> Stro. (<i>Wiedemannia beckeri</i>) 242	<i>lanuginosa</i> Bez. (<i>Drapetis</i>) 314
<i>infuscatus</i> Meig. (<i>Platypalpus annu-</i> <i>lipes</i>) 356, 384	<i>jugulator</i> , nov. sp. (<i>Hemeroedromia</i>) 256	<i>lasiochira</i> Stro. (<i>Hilara</i>) 119
<i>infuscatus</i> Meij. (<i>Parahybos</i>) 46	<i>juncta</i> Coq. (<i>Ræderioides</i>) 226	<i>lasionota</i> Lw. (<i>Empis</i>) 159
<i>infuscatus</i> Old. (<i>Platypalpus enga-</i> <i>dinicus</i>) 381	<i>juvenilis</i> Zett. (<i>Wiedemannia bis-</i> <i>tigma</i>) 245	<i>lasiopyga</i> Lundb. (<i>Hilara mucinata</i>) 119,
<i>infuscatus</i> Zett. (<i>Hybos culiciformis</i>) 20	<i>juvenis</i> , nov. sp. (<i>Platypalpus</i>) 332	385
<i>ingenuus</i> Coll. (<i>Platypalpus</i>) 381	kaijensis Frey (<i>Rhamphomyia</i>) 195	<i>lata</i> Coq. (<i>Drapetis</i>) 310
<i>inops</i> Mel. (<i>Platypalpus</i>) 356	<i>kaiteriensis</i> Mill. (<i>Hilarempis</i>) 129	<i>lateralis</i> Beck. (<i>Platypalpus beckeri</i>) 348
<i>inornata</i> Lw. (<i>Empis</i>) 159	<i>kamtschatica</i> Frey (<i>Rhamphomyia</i>) 195	<i>lateralis</i> Coq. (<i>Platypalpus colla-</i> <i>teralis</i>) 350
<i>inornata</i> , nov. sp. (<i>Anthalia</i>) 61	<i>kerteszi</i> Bez. (<i>Drapetis</i>) 314	<i>lateralis</i> Lw. (<i>Platypalpus</i>) 357
* <i>inscita</i> Meun. (<i>Empis</i>) 159	<i>kerteszi</i> Old. (<i>Rhamphomyia</i>) 382	<i>latericia</i> Beck. (<i>Platypalpus</i>) 357
<i>insecta</i> Coq. (<i>Rhamphomyia</i>) 195	<i>kerteszii</i> Bez. (<i>Empis</i>) 159	<i>laticinctus</i> Walk. (<i>Platypalpus</i>) 357
<i>insignis</i> Brun. (<i>Syneches</i>) 42	<i>kjellmanii</i> Holmgr. (<i>Rhamphomyia</i>) 195	<i>laticornis</i> Big. (<i>Hilarempis</i>) 129
<i>insignis</i> Lw. (<i>Rhamphomyia</i>) 195	<i>klekovacensis</i> Stro. (<i>Rhamphomyia</i>) 195	<i>latifascipennis</i> Brun. (<i>Tachydro-</i> <i>mia</i>) 285
<i>insolita</i> Mel. (<i>Lactistomyia</i>) 36	Kowarzia Mik (subg. <i>Clinocera</i>) 229	<i>latifrons</i> Frey (<i>Rhamphomyia</i>) 196
* <i>insolita</i> Meun. (<i>Rhamphomyia</i>) 195	Kritempis Coll. (subg. <i>Empis</i>) 379	<i>latipennis</i> Bez. (<i>Euhybos</i>) 34
<i>insularis</i> Brethes (<i>Hilarempis</i>) 381	<i>kuntzei</i> Beck.-Bez. (<i>Empis</i>) 159	<i>latipennis</i> Meig. (<i>Rhamphomyia</i> <i>marginata</i>) 198
<i>insularis</i> Coll. (<i>Wiedemannia rhyn-</i> <i>chops</i>) 434		
<i>insularis</i> Mel. (<i>Tachyempis apicis</i>) 292		

DIPTERA

Pages	Pages	Pages			
latipennis Mel. (<i>Drapetis</i>)	309	lineatus Meig. (<i>Platypalpus</i>)	357	longipennis Lw. (<i>Rhamphomyia</i>)	196
latistrigatus Meij. (<i>Platypalpus</i>)	381	lineola Meij. (<i>Drapetis</i>)	314	longipennis Lw. (<i>Tachydromia cal-</i> <i>ceana</i>)	284
laudatoria Coll. (<i>Hemerocephala</i>)	380	liodes Bez. (<i>Empis</i>)	160	longipennis Mel. (<i>Syneches</i>)	42
laureæ Beck. (<i>Hilara</i>)	119	liosoma Bez. (<i>Empis</i>)	160	longipennis Mik. (<i>Wiedemannia</i> <i>escheri</i>)	243
leachianus Steph. (<i>Hybos</i>)	22	Lissomplis Bez. (subgen. <i>Empis</i>)	143	longipes Lw. (<i>Bicellaria</i>)	76
lecta Mel. (<i>Clinocera</i>)	239	lissonota Bez. (<i>Trichina clavipes</i>)	56	longipes Lw. (<i>Empis</i>)	161
Lemtopeza Macq. (see <i>Lepto-</i> <i>peza</i>)	65	Litanomyia Mel. (<i>Chelipoda</i> Macq.)	263	longipes Meig. (<i>Rhamphomyia</i>)	197
*leo Giebel (<i>Hasmona</i>)	369	*litigiosa Meun. (<i>Hilara</i>)	120	longipes Walk. (<i>Clinocera</i>)	237
lepidia Mel. (<i>Wiedemannia</i>)	243	litorea Fall. (<i>Hilara</i>)	120	longipes Walk. (<i>Rhamphomyia</i>)	207
Lepidomyia Big. (<i>Chelipoda</i> Macq.)	264	littoralis Frey (<i>Rhamphomyia sim-</i> <i>plex</i>)	205	longirostris Bez. (<i>Platypalpus</i>)	357
lepidopus Meig. (<i>Empis</i>)	159	littoralis Miller (<i>Hilara</i>)	380	longirostris Macq. (<i>Hilara</i>)	120
lepidopus Walk. (<i>Empis</i>)	160	liturata Lw. (<i>Rhamphomyia</i>)	196	longirostris Meig. (<i>Empis pennipes</i>)	168
Leptempis Coll. (subgen. <i>Empis</i>)	379	livida Fab. (<i>Empis tessellata</i>)	175	longirostris Meig. (<i>Empis serotina</i>)	172
leptogaster, nov. sp. (<i>Euhybos</i>)	33	livida Linn. (<i>Empis</i>)	160	*longirostris Meun. (<i>Euthyneu-</i> <i>rella</i>)	369
leptogastra Lw. (<i>Empis</i>)	160	lividiventris Zett. (<i>Rhamphomyia</i>)	196	longiseta Zett. (<i>Platypalpus pallidi-</i> <i>ventris</i>)	362
lepidium Phil. (<i>Brachystoma</i>)	13	lividus, nov. sp. (<i>Syneches</i>)	42	longispina, nov. sp. (<i>Tachyempis</i>)	289
Leptometopella , nov. gen. (<i>Oropezaella</i> Coll.)	70, 381	lobalis Thom. (<i>Empis</i>)	161	longitarsis, nov. sp. (<i>Micropho-</i> <i>rilla</i>)	89
leptomorion Bez. (<i>Empis</i>)	160	loewi Dahl (<i>Drapetis</i>)	311	longiventris, nov. sp. (<i>Syneches</i>)	42
leptoneura, nov. sp. (<i>Euhybos</i>)	30	loewi Now. (<i>Rhamphomyia anthra-</i> <i>cina</i>)	186	loripedis Coq. (<i>Empis</i>)	161
Leptopeza Macq. (gen.)	65	loewiana Bez. (<i>Empis</i>)	161	loripedis Coq. (<i>Rhamphomyia</i>)	197
leptopus Lw. (<i>Rhamphomyia</i>)	196	longeciliata Stro. (<i>Hilara quadri-</i> <i>faria</i>)	125	lota Haliday (<i>Wiedemannia</i>)	243
Leptoscelos Hal. (<i>Dolichoce-</i> <i>phala</i> Macq.)	246	longefilata Stro. (<i>Rhamphomyia</i>)	196	lucida Zett. (<i>Empis</i>)	161
lesinensis Stro. (<i>Platypalpus</i>)	357	longesetosa Stro. (<i>Hilara</i>)	120	lucidifrons Beck. (<i>Hilara</i>)	120
leucocephalus Ros. (<i>Platypalpus</i>)	357	longestylata Frey (<i>Rhamphomyia</i> <i>candata</i>)	196, 385	lucidilabris Bez. (<i>Empis</i>)	161
leucosticta Beck. (<i>Platypalpus ob-</i> <i>scura</i>)	357, 384	longevittata Zett. (<i>Hilara</i>)	120, 385	lucidula Zett. (<i>Rhamphomyia</i>)	197
leucopexa Lw. (<i>Empis flavitarsis</i>)	156	longicauda Lw. (<i>Rhamphomyia</i>)	196	luctator, nov. sp. (<i>Platypalpus</i>)	335
leucophenga Bez. (<i>Rhamphomyia</i>)	196	longicornis Brun. (<i>Platypalpus ne-</i> <i>palensis</i>)	360	luctifer Bez. (<i>Syneches</i>)	45
leucoptera Gmel. (<i>Empis</i>)	160	longicornis Lw. (<i>Rhamphomyia</i>)	196	luctifera Lw. (<i>Rhamphomyia</i>)	197
leucoptera Lw. (<i>Hilara</i>)	119	longicornis Macq. (<i>Empis</i>)	161	luctuosa Kirby (<i>Itapheila</i>)	104
leucoptera Lw. (<i>Rhamphomyia</i>)	196	longicornis Meig. (<i>Platypalpus</i>)	357	luctuosa Lw. (<i>Rhamphomyia</i>)	197
leucoptera Macq. (<i>Empis</i>)	160	longicornis Meig. (<i>Trichopoda</i>)	219	luctuosa, nov. sp. (<i>Gloma</i>)	102
leucoptera Meig. (<i>Empis vernalis</i>)	177	*longicornis Meun. (<i>Trichopoda</i> <i>succina</i>)	220	lugens Lw. (<i>Rhamphomyia luctuosa</i>)	197
leucostigma Bez. (<i>Empis</i>)	160	longicornis Mik. (<i>Wiedemannia</i>)	243	lugubrina Zett. (<i>Rhamphomyia fus-</i> <i>cipennis</i>)	193
leucothrix Stro. (<i>Platypalpus</i>)	357	longicornis, nov. sp. (<i>Ceratempis</i>)	218	lugubris Lw. (<i>Empis</i>)	162
levicosta White (<i>Leptopeza</i>)	67	longicornis Stro. (<i>Hilara</i>)	120	lugubris Meig. (<i>Hilara cornicula</i>)	116
levicula Coq. (<i>Empis</i>)	160	longifurca, nov. sp. (<i>Clinocera</i>)	231	lugubris, nov. sp. (<i>Bicellaria</i>)	76
limata Coll. (<i>Empis</i>)	379	longifurca Stro. (<i>Hilara clypeata</i>)	116	lugubris Zett. (<i>Hilara</i>)	120
limata Coq. (<i>Rhamphomyia</i>)	196	longimana Lw. (<i>Empis</i>)	161	lunata Walk. (<i>Stilpon nubila</i>)	304, 383
limbata Lw. (<i>Rhamphomyia</i>)	196	longimanus Stro. (<i>Platypalpus</i>)	357	lundbecki Frey (<i>Hilara</i>)	120
limbipennis Bez. (<i>Rhamphomyia</i>)	196	longipalpis, nov. sp. (<i>Anthepisco-</i> <i>pus</i>)	106	lundstroemi Frey (<i>Tachydromia</i>)	285
lindneri Old. (<i>Empis</i>)	379	longipennæ Miller (<i>Trichopoda</i>)	383	Lundstroemiella Frey (sub-	
lineata Fab. (<i>Empis livida</i>)	161	longipennis Bez. (<i>Wiedemannia</i> <i>czernyi</i>)	244	gen. <i>Rhamphomyia</i>)	182
lineata Gimm. (<i>Chelifera precatoria</i>)	261	longipennis Lw. (<i>Empis</i>)	161	lupatus Mel. (<i>Platypalpus</i>)	357
lineata Lw. (<i>Clinocera</i>)	236			lurida Fall. (<i>Hilara</i>)	121
lineata Meig. (<i>Empis</i>)	160				
lineata Scop. (<i>Empis maculata</i>)	162				
lineata Vill. (<i>Empis</i>)	160				

Pages	Pages	Pages
luridipennis Now. (<i>Rhamphomyia</i>) 197	maculifemur Meij. (<i>Platypalpus</i>) 358	maura Fab. (<i>Hilara</i>) 121
luridus Bez. (<i>Hybos</i>) 22	maculimanus Zett. (<i>Platypalpus arti-</i> <i>catus</i>) 347	maura Macq. (<i>Empis</i>) 163
lutea Bez. (<i>Phlebotoma</i>) 128	maculipennis Coq. (<i>Tachydromia</i> <i>bimaculata</i>) 284	medetera Mel. (<i>Drapetis</i>) 311
lutea Lw. (<i>Hilara</i>) 121	maculipennis Frey (<i>Tachydromia</i> <i>punctifera</i>) 286	medetera Mel. (<i>Drapetis scissa</i>) 311
lutea Meig. (<i>Empis</i>) 162	maculipennis Walk. (<i>Tachydromia</i>) 285	media Coll. (<i>Hilara</i>) 380
lutea Meij. (<i>Drapetis</i>) 314	maculipennis Zett. (<i>Rhamphomyia</i>) 197	* media Meun. (<i>Rhamphomyia</i>) 198
luteicollis Mel. (<i>Drapetis</i>) 314	maculipes Big. (<i>Boreodromia</i>) 216	mediocris Beck. (<i>Empis</i>) 163
luteicornis Frey (<i>Parahybos</i>) 46	maculipes Meig. (<i>Platypalpus</i>) 358	mediterranea Lw. (<i>Empis</i>) 163
luteicornis Meig. (<i>Platypalpus</i>) 357	maculipes Zett. (<i>Empis pictipes</i>) 163, 385	Megacyttarus Big. (subgen. <i>Rhamphomyia</i>) 183
luteipalpis Macq. (<i>Platypalpus</i>) 357	maculithorax White (<i>Syneches</i>) 383	Megagrapha (nov. gen.) 301
luteipes Lep. (<i>Drapetis</i>) 311	maculus Zett. (<i>Platypalpus</i>) 359	* Meghyperella Meun. (gen.) 370
luteiventris Lw. (<i>Rhamphomyia</i>) 197	mærens Lw. (<i>Empis</i>) 163	Meghyperus Lw. (gen.) 47
luteolus Coll. (<i>Platypalpus</i>) 381	magellanica Big. (<i>Hilarempis</i>) 129	meigeniana Steph. (<i>Hemerodromia</i>) 256
lutescens Bez. (<i>Chersodromia</i>) 297	magellenensis Bez.-Frey (<i>Rham-</i> <i>phomyia</i>) 197	meigenianus Gimm. (<i>Platypalpus</i>) 359
luteus Meig. (<i>Platypalpus glaber</i>) 358, 384	magica Mik (<i>Hilara thoracica</i>) 127	melæna Bez. (<i>Empis</i>) 163
luteus Wied. (<i>Syneches</i>) 42	major Bez. (<i>Hybos</i>) 23	melæna Hal. (<i>Bicellaria spuria</i>) 76, 384
luteus Zett. (<i>Platypalpus analis</i>) 346	major Bonsd. (<i>Platypalpus candicans</i>) 350	melanderi Cole (<i>Parathalassius</i>) 87
lyristes, nov. sp. (<i>Platypalpus</i>) 338	major Frey (<i>Rhamphomyia simplex</i>) 205	melangyna Coll. (<i>Hemerodromia</i>) 434
maackii Lw. (<i>Iteaphila</i>) 104	major Stro. (<i>Hilara longegettata</i>) 120	melania Beck. (<i>Rhamphomyia</i>) 198
mabelæ Mel. (<i>Hesperempis</i>) 111	major Stro. (<i>Hilara pinetorum</i>) 124	melania Stro. (<i>Rhamphomyia stig-</i> <i>mosa</i>) 206
macerrima Bez. (<i>Rhamphomyia</i>) 197	major Zett. (<i>Platypalpus</i>) 359	melanocephala Fab. (<i>Chelipoda</i>) 267
macilenta Lw. (<i>Rhamphomyia</i>) 197	* mala Meun. (<i>Empis</i>) 163	melanocephala Hal. (<i>Chelifera</i>) 260, 385
* macilenta Meun. (<i>Empis</i>) 162	* malefica Meun. (<i>Empis</i>) 163	melanocephala Lw. (<i>Dysaletria atri-</i> <i>pes</i>) 288
macquarti Beck. (<i>Empis geniculata</i>) 157	malleola Beck. (<i>Empis</i>) 163	melanocephala Lundb. (<i>Chelifera</i> <i>flavella</i>) 260, 385
macquarti Zett. (<i>Iteaphila</i>) 104	mallos Walk. (<i>Rhamphomyia</i>) 197	Malthacotricha Beck. (<i>Hete-</i> <i>rotropus</i> Lw.) 375
macra Lw. (<i>Empis</i>) 162	maluinensis End. (<i>Hilara</i>) 121	melanocerus, nov. var. (<i>Platypalpus</i> <i>crassifemoris</i>) 337
macrocera Big. (<i>Hilarempis</i>) 129	manca Coq. (<i>Empis</i>) 163	melanochatus Bez. (<i>Platypalpus</i> <i>flavipes</i>) 355
macropalpa Egg. (<i>Empis</i>) 162	manca Coq. (<i>Rhamphomyia</i>) 197	melanochira Bez. (<i>Hilara</i>) 122
macropalpus Stro. (<i>Platypalpus</i>) 358	mandalota, nov. sp. (<i>Anthalia</i>) 58	melanogaster, nov. sp. (<i>Platypalpus</i>) 323
* macrophthalma Först. (<i>Empis</i>) 162	manicata Meig. (<i>Hilara</i>) 121, 385	melanogastra Phil. (<i>Brachystoma</i>) 13
Macroptera Beck. (<i>Symball-</i> <i>ophthalmus</i> Beck.) 300	mantiformis Cuv. (<i>Chelifera preca-</i> <i>toria</i>) 261	melanopa Steph. (<i>Empis livida</i>) 161
macroptera Lw. (<i>Hilara</i>) 121	Mantipeza Rond. (<i>Chelifera</i> Macq.) 258	melanopleura Lw. (<i>Ocydromia gla-</i> <i>bricula</i>) 69
macropus Lw. (<i>Empis</i>) 162	mantispa Macq. (<i>Chelipoda voca-</i> <i>toria</i>) 267	melanotricha Lw. (<i>Empis</i>) 163
macrorrhyncha Phil. (<i>Empis</i>) 162	mantispa Panz. (<i>Chelipoda melano-</i> <i>cephala</i>) 267	melanura Bez. (<i>Drapetis</i>) 315
Macrostomus Wied. (<i>Rham-</i> <i>phomyia</i> Meig.) 180	marginalis Bez. (<i>Drapetis</i>) 315	melas Bez. (<i>Parahybos</i>) 46
Macroura Zett. (<i>Heleodromia</i> Hal.) 224	marginata Brun. (<i>Empis</i>) 163	* melia Heyden (<i>Empis</i>) 163
macrura Big. (<i>Empis</i>) 162	marginata Fab. (<i>Rhamphomyia</i>) 197	melleus, nov. sp. (<i>Platypalpus</i>) 321
macrura Coq. (<i>Rhamphomyia cla-</i> <i>uator</i>) 188	marginata Meig. (<i>Drapetis</i>) 311	mellipes Wh.-Mel. (<i>Hybos</i>) 23
macrura Lw. (<i>Rhamphomyia</i>) 197	marginipennis Stro. (<i>Hilara</i>) 121	memorandus, nov. sp. (<i>Philetus</i>) 110
maculata auct. (<i>Empis confusa</i>) 163	marshamanus Steph. (<i>Hybos</i>) 23	mendoza Brethes (<i>Hilarempis</i>) 130
maculata Fab. (<i>Empis</i>) 162	matrona Hal. (<i>Hilara</i>) 121	meridionalis Beck. (<i>Iteaphila ni-</i> <i>tidula</i>) 104
maculata Lw. (<i>Clinocera</i>) 237	matroniformis Stro. (<i>Hilara</i>) 121	meridionalis Meig. (<i>Empis</i>) 163
maculata Macq. (<i>Dolichocephala</i> <i>irrorata</i>) 247	matura, nov. sp. (<i>Euthyneura</i>) 53	merula Coll. (<i>Hilara</i>) 380
maculata Macq. (<i>Drapetis ephippiata</i>) 314		mesogramma Lw. (<i>Empis dimidiata</i>) 154
maculata Mats. (<i>Hilarimorpha</i>) 373		
maculifemoratus, nov. sp. (<i>Platy-</i> <i>palpus</i>) 358		

DIPTERA

Pages	Pages	Pages	
mesogrammus Lw. (<i>Platypalpus</i>)	359	minor Stro. (<i>Platypalpus cursitans</i>)	352
mesomelæna Bez. (<i>Hemerodromia</i>)	256	minuta Fall. (<i>Edalea flavipes</i>)	64
* Mesomyites Ckll. (gen.)	370	minuta Fall. (<i>Trichina clavipes</i>)	56
Metacheila Coq. (subgen. <i>Chelifera</i>)	259	minuta Greene (<i>Mythicomyia</i>)	376
metapleuralis Bez. (<i>Empis</i>)	163	minuta Mel. (<i>Micrempis</i>)	299
metatarsalis, nov. sp. (<i>Euhybos</i>)	28	minuta Walk. (<i>Trichina elongata</i>)	56
metatarsata Bez. (<i>Drapetis</i>)	315	minuta Will. (<i>Drapetis divergens</i>)	308
metatarsata Zett. (<i>Rhamphomyia</i>)	198	minutissima Stro. (<i>Platypalpus</i>)	359
mexicana Mel. (<i>Chelifera</i>)	262	minutus Brun. (<i>Syneches</i>)	42
mexicana Mel. (<i>Drapetis septentrionalis</i>)	311	minutus Meig. (<i>Platypalpus</i>)	359, 384
mexicana Wh.-Mel. (<i>Chelipoda</i>)	266	minutus Walk. (<i>Rhamphomyia</i>)	198
micans Old. (<i>Rhamphomyia</i>)	198	* <i>miocenica</i> Ckll. (<i>Empis</i>)	164
micans Schin. (<i>Empis</i>)	163	mira Big. (<i>Empis</i>)	164
micrargyra Bez. (<i>Rhamphomyia</i>)	198	mirifica Frey (<i>Rhamphomyia</i>)	198
Micrempis (nov. gen.)	298	mitrephorus, nov. sp. (<i>Oreogeston</i>)	99
Microcera Zett. (<i>Heleodromia Hal.</i>)	224	modesta Meig. (<i>Empis</i>)	164
<i>microcerus</i> , nov. var. (<i>Platypalpus flavirostris</i>)	325	modesta Meig. (<i>Hilara interstincta</i>)	119
Microcypta Big. (<i>Microsania</i> Zett.)	374	modesta Phil. (<i>Haplomera</i>)	131
Microdromia Big. (<i>Hemerodromia Meig.</i>)	254	modesta Wahlb. (<i>Rhamphomyia</i>)	198
<i>microphona</i> , nov. sp. (<i>Charadromia</i>)	293	mollicella White (<i>Hilara</i>)	122
Microphorella Beck. (gen.)	87	<i>mollis</i> , nov. var. (<i>Platypalpus crassifemoris</i>)	337
Micropherus Macq. (gen.)	89	monacha, nov. sp. (<i>Tachydromia</i>)	283
<i>microptera</i> Lw. (<i>Tachydromia brevipennis</i>)	286, 384	monochæta Bez. (<i>Drapetis</i>)	312
<i>micropyga</i> Bez. (<i>Empis</i>)	164	monedula Coll. (<i>Hilara</i>)	380
<i>micropyga</i> Coll. (<i>Rhamphomyia</i>)	382	monogramma Meig. (<i>Empis dimidiata</i>)	154
<i>micropyga</i> Mel. (<i>Drapetis</i>)	309	monogramma, nov. sp. (<i>Proclinoxyga</i>)	222
Microsania Zett. (gen.)	374	monostigma Meig. (<i>Chelifera preatoria</i>)	261
<i>microstigma</i> Bez. (<i>Wiedemannia</i>)	244	monostigma Zett. (<i>Chelifera concin-nicanda</i>)	378
<i>migrata</i> Walk. (<i>Hilara</i>)	122	<i>monserratensis</i> Stro. (<i>Tachydromia</i>)	286
<i>mikiana</i> Bez. (<i>Wiedemannia</i>)	243	<i>monstrosa</i> Bez. (<i>Rhamphomyia</i>)	198
<i>mikii</i> Beck. (<i>Platypalpus</i>)	359	<i>montana</i> Old. (<i>Rhamphomyia</i>)	198
<i>mikii</i> Stro. (<i>Empis</i>)	164	<i>montanus</i> Beck. (<i>Platypalpus</i>)	360
<i>mikii</i> Will. (<i>Hilarimorpha</i>)	373	<i>montanus</i> Frey (<i>Platypalpus ater</i>)	348
<i>mimus</i> , nov. sp. (<i>Platypalpus</i>)	324	<i>montanus</i> Mel. (<i>Platypalpus splendens</i>)	365
<i>mixopolia</i> Mel. (<i>Empis</i>)	164	<i>montezuma</i> Wh.-Mel. (<i>Empis</i>)	164
<i>minima</i> Beck. (<i>Tachydromia</i>)	286	<i>monticola</i> Lw. (<i>Empis</i>)	164
<i>minima</i> Meig. (<i>Drapetis</i> , not assi-milis)	308, 384	<i>monticola</i> , nov. sp. (<i>Brochella</i>)	109
<i>minima</i> Zett. (<i>Drapetis assimilis</i> , not <i>pussilla</i>)	309, 384	<i>monticola</i> Mel. (<i>Platypalpus</i>)	360
<i>minor</i> Bez. (<i>Syneches</i>)	45	<i>monticolus</i> White (<i>Platypalpus ceylonensis</i>)	381
<i>minor</i> Macq. (<i>Chelipoda melancephala</i>)	267	<i>morata</i> Coll. (<i>Hilara</i>)	380
<i>minor</i> , nov. sp. (<i>Clinocera</i>)	233	* <i>mordax</i> Meun. (<i>Empis</i>)	164
<i>minor</i> Old. (<i>Rhamphomyia</i>)	382	<i>moreirai</i> Brethes (<i>Hilarempis</i>)	381
		<i>morenæ</i> Stro. (<i>Empis</i>)	164
		<i>morenæ</i> Stro. (<i>Hilara cingulata</i>)	116

Pages	Pages	Pages			
<i>nervosa</i> Meig. (<i>Tachyepes nubila</i>)	277	<i>nigripennis</i> Meig. (<i>Ocydromia glabricula</i>)	69	<i>nitens</i> Mel. (<i>Drapetis</i>)	311
<i>nervosa</i> , nov. sp. (<i>Tachyempis</i>)	290	<i>nigripennis</i> Phil. (<i>Homalocnemis</i>)	14	<i>nitens</i> nov. var. (<i>Platypalpus politus</i>)	334
<i>nervosus</i> Steph. (<i>Hybos</i>)	23	<i>nigripes</i> Fab. (<i>Rhamphomyia</i>)	199	<i>nitida</i> Bez. (<i>Atrichopleura</i>)	134
Nioenla Walk. (<i>Empis</i> Linn.)	138	<i>nigripes</i> Meig. (<i>Platypalpus</i>)	361	<i>nitida</i> Lw. (<i>Syndyas</i>)	37
<i>niger</i> Brun. (<i>Hybos</i>)	23	<i>nigripes</i> Mel. (<i>Drapetis dividua</i>)	308	<i>nitida</i> Macq. (<i>Rhamphomyia</i>)	200
<i>niger</i> Macq. (<i>Platypalpus</i>)	360	<i>nigripes</i> , nov. sp. (<i>Euhybos</i>)	28	<i>nitida</i> Meig. (<i>Empis</i>)	165
<i>niger</i> Meig. (<i>Platypalpus</i>)	360	<i>nigripes</i> Stro. (<i>Anthepiscopus ribesii</i>)	108	<i>nitida</i> Mel. (<i>Drapetis unipila</i>)	312
<i>niger</i> Zett. (<i>Anthepiscopus oedalinus</i>)	107	<i>nigripes</i> Stro. (<i>Rhamphomyia aterrima</i>)	186	<i>nitida</i> , nov. sp. (<i>Trichina</i>)	56
<i>nigerrima</i> Bez. (<i>Tachydromia</i>)	286	<i>nigripes</i> Stro. (<i>Trichina</i>)	56	<i>nitida</i> Zett. (<i>Rhamphomyia nitidula</i>)	200
<i>nigerrima</i> Lw. (<i>Empis</i>)	164	<i>nigripes</i> Zett. (<i>Drapetis exilis</i>)	309, 384	<i>nitidicollis</i> Curt. (<i>Empis nigratarsis</i>)	165
<i>nigra</i> Fall. (<i>Drapetis curvipes</i> , not <i>flavipes</i>)	309, 384	<i>nigripes</i> Zett. (<i>Leptopeza</i>)	67	<i>nitidicollis</i> Frey (<i>Rhamphomyia</i>)	200
<i>nigra</i> Macq. (<i>Bicellaria spuria</i>)	78	<i>nigripes</i> Zett. (<i>Syndyas</i>)	37	<i>nitidipleura</i> nov. sp. (<i>Platypalpus</i>)	329
<i>nigra</i> Meig. (<i>Bicellaria</i>)	77	<i>nigripennis</i> Fab. (<i>Rhamphomyia</i>)	198	<i>nitidissima</i> Stro. (<i>Empis</i>)	166
<i>nigra</i> Meig. (<i>Clinocera</i>)	236	<i>nigrita</i> Big. (<i>Rhamphomyia</i>)	199	<i>nitidula</i> Zett. (<i>Empis</i>)	166
<i>nigra</i> Meig. (<i>Drapetis assimilis</i>)	308	<i>nigrita</i> Coll. (<i>Bicellaria</i>)	378	<i>nitidula</i> Zett. (<i>Hilara</i>)	123
<i>nigra</i> Mel. (<i>Drapetis aliternigra</i>)	307	<i>nigrita</i> Zett. (<i>Rhamphomyia</i>)	199	<i>nitidula</i> Zett. (<i>Iteaphila</i>)	104
<i>nigra</i> Miller (<i>Hilarempis</i>)	381	<i>nigritarsis</i> Fall. (<i>Platypalpus</i>)	361	<i>nitidula</i> Zett. (<i>Rhamphomyia</i>)	200
<i>nigra</i> Phil. (<i>Iteaphila</i>)	104	<i>nigritarsis</i> Meig. (<i>Empis</i>)	165	<i>nitidus</i> Macq. (<i>Platypalpus</i>)	361
<i>nigra</i> , nov. sp. (<i>Tachyempis</i>)	288	<i>nigritarsis</i> Zett. (<i>Hilara griseola</i>)	118	<i>nitidus</i> Mel. (<i>Meghyperus</i>)	47
<i>nigra</i> Vill. (<i>Empis</i>)	164	<i>nigritella</i> Zett. (<i>Drapetis not aterrima</i>)	310, 384	<i>nitidiventris</i> Lw. (<i>Empis</i>)	166
<i>nigra</i> Zett. (<i>Bicellaria intermedia</i>)	76	<i>nigritibialis</i> Stro. (<i>Empis</i>)	165	<i>nitidivittata</i> Macq. (<i>Rhamphomyia</i>)	200
<i>nigra</i> Zett. (<i>Drapetis curvipes</i> , not <i>morella</i>)	311, 384	<i>nigriventris</i> Lw. (<i>Chelifera</i>)	263	<i>nivalis</i> Zett. (<i>Clinocera</i>)	238
<i>nigricans</i> Mel. (<i>Drapetis</i>)	315	<i>nigriventris</i> Lw. (<i>Hilara</i>)	122	<i>niveipennis</i> Frey (<i>Rhamphomyia albipennis</i>)	184
<i>nigrescens</i> Beck. (<i>Schistostoma</i>)	94	<i>nigriventris</i> Macq. (<i>Rhamphomyia</i>)	199	<i>niveipennis</i> Zett. (<i>Hilara albipennis</i>)	123, 383
<i>nigrescens</i> Schum. (<i>Empis</i>)	165	<i>nigrofemoratus</i> Stro. (<i>Empis grisea</i>)	157	<i>niveipennis</i> Zett. (<i>Hilara braueri</i>)	114
<i>nigricans</i> Lw. (<i>Hormopeza</i>)	97	<i>nigrofemoratus</i> Stro. (<i>Platypalpus exiguus</i>)	353	<i>niveipennis</i> Zett. (<i>Rhamphomyia</i>)	200
<i>nigricans</i> Lw. (<i>Rhamphomyia</i>)	198	<i>nigrofemoratus</i> Stro. (<i>Platypalpus fascipes</i>)	354	<i>albissima</i>)	184
<i>nigricans</i> Macq. (<i>Empis</i>)	165	<i>nigrogeniculatus</i> Bez. (<i>Platypalpus baldensis</i>)	348	<i>niveiseta</i> Zett. (<i>Platypalpus</i>)	361
<i>nigricans</i> Meig. (<i>Empis</i>)	165	<i>nigrohalterata</i> Stro. (<i>Dolichocephala guttata</i>)	247	<i>nodipes</i> Fall. (<i>Rhamphomyia</i>)	200
<i>nigricans</i> , nov. sp. (<i>Chelipoda</i>)	266	<i>nigrohirta</i> Coll. (<i>Hilara</i>)	380	<i>nodipes</i> Mel. (<i>Empis</i>)	166
<i>nigricauda</i> Beck. (<i>Rhamphomyia</i>)	198	<i>nigrolineata</i> Ros. (<i>Hemerodromia</i>)	256	<i>nodosa</i> Beling (<i>Empis grisea</i>)	158
<i>nigricollis</i> Frey (<i>Platypalpus sahlbergi</i>)	365	<i>nigromaculata</i> Ros. (<i>Drapetis ephippata</i> , not <i>Platypalpus</i>)	361, 384	Noeza Meig. (<i>Hybos</i> Meig.)	18
<i>nigricoma</i> Lw. (<i>Empis</i>)	165	<i>nigromaculata</i> Ros. (<i>Rhamphomyia</i>)	199	<i>nonstriatus</i> Stro. (<i>Platypalpus pectoralis</i>)	363
<i>nigricorne</i> Phil. (<i>Brachystoma</i>)	13	<i>nigromaculata</i> White (<i>Ironomyia</i>)	48	<i>nordquistii</i> Holmg. (<i>Rhamphomyia</i>)	200
<i>nigricoxus</i> Mik (<i>Platypalpus</i>)	360	<i>nigronitidus</i> Brun. (<i>Hybos</i>)	23	<i>notata</i> Lw. (<i>Chelifera</i>)	260
<i>nigricrus</i> Gmel. (<i>Empis</i>)	165	<i>nigropunctata</i> White (<i>Drapetis</i>)	378	<i>notatithorax</i> White (<i>Drapetis</i>)	378
<i>nigrifemoratus</i> Macq. (<i>Platypalpus</i>)	361	<i>nigrosetosus</i> Stro. (<i>Platypalpus maculipes</i>)	359	<i>notatus</i> Meig. (<i>Platypalpus fascipes</i>)	353
<i>nigrifemur</i> Stro. (<i>Platypalpus baldensis</i>)	348	<i>nigroterminatus</i> Stro. (<i>Platypalpus exilis</i>)	353	<i>notogramma</i> Bez. (<i>Haplomera</i>)	131
<i>nigrimana</i> Beck. (<i>Empis</i>)	165	<i>nimia</i> White (<i>Hilara</i>)	122	<i>novakii</i> Mik (<i>Hilara</i>)	123
<i>nigrimana</i> Lw. (<i>Brachystoma</i>)	13	Niphogenia , nov. gen.	217	<i>novakii</i> Stro. (<i>Platypalpus</i>)	361
<i>nigrimana</i> Phil. (<i>Chelifera</i>)	262	<i>nitens</i> Brun. (<i>Hybos</i>)	23	<i>novecarolina</i> Beut. (<i>Rhamphomyia</i>)	200
<i>nigrimana</i> White (<i>Hilarempis</i>)	130	<i>nitens</i> Macq. (<i>Hilara chorica</i>)	115	<i>novemguttata</i> Stro. (<i>Dolichocephala ocellata</i>)	247
<i>nigrimanus</i> Ros. (<i>Platypalpus</i>)	361			<i>nox</i> Old. (<i>Rhamphomyia</i>)	200
<i>nigrimanus</i> Stro. (<i>Platypalpus</i>)	361			<i>nubecula</i> Beck. (<i>Chelifera</i>)	260
<i>nigrina</i> Fall. (<i>Hilara</i>)	122			<i>nubifera</i> Coq. (<i>Coloboneura</i>)	294
<i>nigrinus</i> Meig. (<i>Platypalpus</i>)	361				
<i>nigrinus</i> Schin. (<i>Platypalpus</i>)	361				
<i>nigripalpis</i> Big. (<i>Platypalpus</i>)	361				

DIPTERA

Pages	Pages	Pages			
nubigena Bez. (<i>Rhamphomyia</i>)	200	obscurus Coq. (<i>Microphorus</i>)	93	ornatipes, nov. sp. (<i>Microphorella</i>)	88
nubila Coll. (<i>Stilpon</i>)	383	obscurus Lw. (<i>Oreogeton</i>)	100	Oreopezella Coll. (gen.)	381
nubila Meig. (<i>Tachypeza</i>)	277	<i>obsecratoria</i> Walk. (<i>Chelipoda</i> <i>melanocephala</i>)	267	orthoneura Bez. (<i>Rhamphomyia</i>)	201
nubila White (<i>Hilara</i>)	123	<i>obsoleta</i> Lw. (<i>Chelifera</i>)	260	Orthorhynchium Mik. (<i>Wiede-</i> <i>mannia</i> Zeit.)	240
nuda Lw. (<i>Empis</i>)	166	<i>obsoleta</i> Stro. (<i>Tachydromia inter-</i> <i>rupta</i>)	285	ossicula Beck. (<i>Gloma</i>)	102
nudafacies Bez. (<i>Hilarempis</i>)	130	* <i>obtusa</i> Meun. (<i>Rhamphomyia</i>)	201	ostiorum Beck. (<i>Platypalpus</i>)	362
nudipes Lw. (<i>Clinocera</i>)	236	occidens Coq. (<i>Meghyperus</i>)	47	otakouensis Mill. (<i>Empis</i>)	166
nudipes Macq. (<i>Haplomera</i>)	131	occidentale Mel. (<i>Brachystoma</i>)	13	otiosa Coq. (<i>Empis</i>)	166
nudipes Old. (<i>Rhamphomyia</i>)	382	<i>occipitalis</i> Bez. (<i>Empis concolor</i>)	152	otiosa Coq. (<i>Rhamphomyia</i>)	201
<i>nudinacula</i> Stro. (<i>Hilara minuta</i>)	122	<i>ocellata</i> Costa (<i>Dolichocephala</i>)	247	otiraensis Mill. (<i>Chelipoda</i>)	378
nugax Mel. (<i>Hilara</i>)	123	<i>ochracea</i> Big. (<i>Hilarempis</i>)	130	* Oustatetimyia Meun. (gen.)	370
nuntia Meig. (<i>Empis</i>)	166	<i>ochracea</i> Curt. (<i>Empis lutea</i>)	162	oxystoma Bez. (<i>Wiedemannia</i>)	245
nuptus, nov. sp. (<i>Anthepiscopus</i>)	106	<i>ochraceus</i> Beck. (<i>Platypalpus</i>)	361	ozernajensis Frey (<i>Rhamphomyia</i>)	201
nura Mel. (<i>Trichina</i>)	57	<i>ochricollis</i> , nov. sp. (<i>Platypalpus</i>)	323	pachycnemus Lw. (<i>Platypalpus</i>)	362
nympha, nov. sp. (<i>Hemerodromia</i>)	256	<i>ochropus</i> Phil. (<i>Empis</i>)	166	pachymera Big. (<i>Rhamphomyia</i>)	201
obeliscus Bez. (<i>Syneches</i>)	43	<i>ocreatus</i> , nov. sp. (<i>Euhybos</i>)	26	pachymera Macq. (<i>Empis</i>)	167
obesa Lw. (<i>Empis</i>)	166	<i>oculata</i> Lw. (<i>Dolichocephala ocellata</i>)	247	Pachymeria Steph. (subgen. <i>Empis</i>)	143
<i>obesa</i> Verr.-Coll. (<i>Hilara platynura</i>)	380	<i>oculeus</i> , nov. sp. (<i>Platypalpus</i>)	342	Pachymerina Macq. (<i>Empis</i> Linn.)	139
obliqua, nov. sp. (<i>Micrempis</i>)	299	Ocydromia Meig. (gen.)	68	pachymerina Schin. (<i>Empis</i>)	167
obliquinervis Meij. (<i>Drapetis</i>)	315	OXYDROMIINAE (subfam.)	49	Pachyopeza Lioy (<i>Microsania</i> Zett.)	374
obliterata Zett. (<i>Hormopeza</i>)	97	Œdalem Meig. (gen.)	62	pachypodiata Big. (<i>Empis</i>)	167
<i>oblongoguttata</i> Dale (<i>Dolichocephala</i> <i>guttata</i>)	247	<i>œdalinus</i> Zett. (<i>Anthepiscopus</i>)	107	pachystoma Phil. (<i>Empis</i>)	167
<i>obscura</i> Big. (<i>Hilarimorpha</i>)	373	* <i>œdalooides</i> Meun. (<i>Rhamphomyia</i>)	201	* Palaeodalea Meun. (gen.)	370
<i>obscura</i> Brun. (<i>Clinocera</i>)	237	<i>œdicnemus</i> Stro. (<i>Platypalpus</i>)	362	* Palaeoleptopeza Meun. (gen.)	371
<i>obscura</i> Eversm. (<i>Rhamphomyia</i>)	200	<i>œdimera</i> Mel. (<i>Drapetis</i>)	311	* Palaeoparamesia Meun. (gen.)	371
<i>obscura</i> Lw. (<i>Rhamphomyia</i>)	200	<i>ohioensis</i> Mel. (<i>Œdalea</i>)	65	<i>pallens</i> Macq. (<i>Hilara manicata</i>)	121
<i>obscura</i> Macq. (<i>Empis serotina</i>)	172	<i>oldenberghi</i> Eng. (<i>Wiedemannia</i>)	243	<i>pallens</i> Ros. (<i>Empis, not aequalis</i>)	145, 383
<i>obscura</i> Meig. (<i>Hilara flavipes</i>)	117	<i>oldenberghi</i> Frey (<i>Rhamphomyia</i>)	201	<i>palliatia</i> Coq. (<i>Porphyrochroa</i>)	137
<i>obscura</i> Phil. (<i>Apalocnemis</i>)	101	<i>olivacea</i> , nov. sp. (<i>Clinocera</i>)	233	<i>pallida</i> Lw. (<i>Empis</i>)	167
<i>obscura</i> Ros. (<i>Platypalpus, not</i> <i>Tachypeza winthemi</i>)	278, 384	<i>ollius</i> Walk. (<i>Empis</i>)	166	<i>pallida</i> Lw. (<i>Synamphotera</i>)	217
<i>obscura</i> Zett. (<i>Iteaphila</i>)	104	<i>omissionis</i> Beck. (<i>Rhamphomyia</i>)	201	<i>pallida</i> Phil. (<i>Chelifera</i>)	262
<i>obscura</i> Zett. (<i>Rhamphomyia</i>)	201	<i>opaca</i> Meig. (<i>Empis</i>)	166	<i>pallida</i> Phil. (<i>Hilarempis</i>)	130
<i>obscurata</i> Phil. (<i>Haplomera</i>)	131	<i>opaca</i> Lw. (<i>Syndyas</i>)	37	<i>pallida</i> Zett. (<i>Allanthalia</i>)	62
<i>obscurella</i> Zett. (<i>Rhamphomyia</i>)	201	<i>opaca</i> Lw. (<i>Trichina</i>)	57	<i>pallidicornis</i> Coll. (<i>Platypalpus</i>)	381
<i>obscurior</i> Stro. (<i>Platypalpus annuli-</i> <i>pes</i>)	347	<i>opacitorax</i> Mall. (<i>Rhamphomyia</i>)	382	<i>pallidicoxa</i> Frey (<i>Platypalpus jas-</i> <i>cipes</i>)	354
<i>obscurior</i> Stro. (<i>Platypalpus fascipes</i>)	354	Opeatocerata (nov. gen.)	134	<i>pallidifurca</i> White (<i>Hilarempis</i>)	130
<i>obscripennis</i> Bez. (<i>Drapetis phaeo-</i> <i>tera</i>)	311	<i>optimus</i> Coll. (<i>Platypalpus</i>)	381	<i>palliditarsis</i> Brun. (<i>Syneches</i>)	43
<i>obscripennis</i> Meig. (<i>Rhamphomyia</i> <i>nigripennis</i>)	199	<i>oratoria</i> Fall. (<i>Hemerodromia</i>)	257	<i>palliditibiæ</i> Brun. (<i>Tachypeza</i>)	278
<i>obscripennis</i> Phil. (<i>Drapetis</i>)	311	<i>orchestris</i> Mel. (<i>Iteaphila</i>)	104	<i>pallidiventris</i> Beck. (<i>Platypalpus</i> <i>fasciatus</i>)	353
<i>obscripennis</i> Phil. (<i>Haplomera</i>)	131	Oreogeton Schin. (gen.)	98	<i>pallidiventris</i> Bonsd. (<i>Platypalpus</i> <i>bicolor</i>)	348
<i>obscripennis</i> Stro. (<i>Empis serotina</i>)	172	Oreothalla Mel. (gen.)	223	<i>pallidiventris</i> Bonsd. (<i>Rhampho-</i> <i>myia tenuirostris</i>)	207
<i>obscuripes</i> Adams (<i>Stilpon</i>)	304	<i>oriens</i> , nov. sp. (<i>Platypalpus</i>)	362	<i>pallidiventris</i> Fall. (<i>Rhamphomyia</i>)	201
<i>obscuripes</i> Lw. (<i>Brachystoma</i>)	13	<i>orientalis</i> Bez. (<i>Hilara</i>)	123	<i>pallidiventris</i> Meig. (<i>Platypalpus</i>)	362
<i>obscuripes</i> Lw. (<i>Empis</i>)	166	<i>orientalis</i> Brun. (<i>Platypalpus</i>)	362		
<i>obscuripes</i> Stro. (<i>Platypalpus minutus</i>)	360	<i>orientalis</i> Meij. (<i>Hemerodromia</i>)	257		
<i>obcuritarsis</i> Zett. (<i>Hilara</i>)	123	<i>orientalis</i> Meij. (<i>Parahybos</i>)	46		
		<i>ornata</i> Eng. (<i>Wiedemannia bicus-</i> <i>pidata</i>)	242		
		<i>ornatipes</i> Beck. (<i>Tachydromia</i>)	286		
		<i>ornatipes</i> Big. (<i>Coloboneura</i>)	294		
		<i>ornatipes</i> Meij. (<i>Parahybos</i>)	46		

Pages	Pages	Pages
<i>pallidus</i> Meig. (<i>Platypalpus lutens</i>) 358	<i>pectinator</i> Mel. (<i>Platypalpus</i>) 363	Phæoballa Mik. (subgen. <i>Clinocera</i>) 229
<i>pallidus</i> Stro. (<i>Platypalpus pollinosus</i>) 364	<i>pectinicauda</i> Coll. (<i>Chelifera</i>) 378	<i>phænomerus</i> Lw. (<i>Empis</i>) 168
<i>pallipes</i> Bez. (<i>Parahybos</i>) 46	<i>pectiniger</i> Mel. (<i>Stilpon</i>) 305	<i>phænomerus</i> Lw. (<i>Empis eumera</i>) 155
<i>pallipes</i> Fall. (<i>Platypalpus</i>) 362	<i>pectinipennis</i> Zett. (<i>Microsania pectinipennis</i>) 374	<i>phæoptera</i> Bez. (<i>Drapetis</i>) 311
<i>pallipes</i> Meij. (<i>Hybos</i>) 23	<i>pectinipes</i> Stro. (<i>Hilara</i>) 123	<i>phæopterus</i> Bez. (<i>Syneches</i>) 43
<i>pallipes</i> Meig. (<i>Microsania</i>) 374	<i>pectinipes</i> Zett. (<i>Microsania pectinipes</i>) 374	<i>phanerostigma</i> Frey (<i>Rhamphomyia</i>) 202
<i>pallipes</i> Meig. (<i>Platypalpus fuscicornis</i>) 356	<i>pectinulata</i> Stro. (<i>Heleodromia</i>) 225	<i>phantasma</i> Mik. (<i>Wiedemannia</i>) 242
<i>pallipes</i> Oliv. (<i>Empis</i>) 167	<i>pectipennis</i> Meig. (<i>Microsania</i>) 374	<i>phemius</i> Walk. (<i>Rhamphomyia</i>) 202
<i>pallipes</i> Zett. (<i>Platypalpus stigmatellus</i>) 365	<i>pectoralis</i> Fall. (<i>Platypalpus</i>) 363	<i>phengites</i> , nov. sp. (<i>Tachydromia</i>) 282
<i>pallipes</i> Zett. (<i>Trichina</i> , not <i>Edalea</i>) 56, 65, 384	<i>pectoris</i> Coq. (<i>Rhamphomyia</i>) 201	Philetus (nov. gen.) 110
<i>pallistigma</i> Ros. (<i>Rhamphomyia</i>) 201	<i>pedestris</i> Beck. (<i>Platypalpus</i>) 363	<i>philina</i> Speiser (<i>Hilara</i>) 123
<i>palloris</i> Coq. (<i>Chelifera</i>) 260	<i>pegasus</i> O.-S. (<i>Empis</i>) 167	<i>philippii</i> Bez. (<i>Hilarempis fulva</i>) 129
<i>palmarum</i> Stro. (<i>Hilara</i>) 123	<i>pellucida</i> Coq. (<i>Empis</i>) 167	<i>philippii</i> Big. (<i>Ocydromia</i>) 70
<i>palmeni</i> Frey (<i>Rhamphomyia</i>) 201	<i>pelops</i> Mel. (<i>Oreothalia</i>) 224	Philolutra Mik. (subgen. <i>Wiedemannia</i>) 241
<i>palparis</i> Egg. (<i>Empis scotica</i>) 167, 385	<i>penicillata</i> Bez. (<i>Rhamphomyia</i>) 201	<i>philpotti</i> Mill. (<i>Hilarempis</i>) 130
* <i>palpata</i> Lw. (<i>Timalphes</i>) 98	<i>peniscissa</i> Beck. (<i>Clinocera</i>) 239	Phlebotena Bez. (gen.) 127
<i>palpata</i> Meig. (<i>Drapetis</i>) 315	<i>pennaria</i> auct. (<i>Empis nuntia</i>) 167, 385	Phoneutrisca Lw. (<i>Tachydromia</i>) 279
<i>paludosa</i> Beck. (<i>Hilara</i>) 123	<i>pennaria</i> Fall. (<i>Empis</i>) 167, 379	Phoroxypha Rond. (<i>Platypalpus</i> Macq.) 316
<i>paludosus</i> Perris (<i>Platypalpus</i>) 363	<i>pennaria</i> Fall. (<i>Empis vernalis</i>) 178	<i>phthia</i> Walk. (<i>Syneches</i>) 43
<i>palustris</i> Scop. (<i>Empis</i>) 167	<i>pennata</i> Gimm. (<i>Edalea</i>) 65	Phyllodromia Zett. (subgen. <i>Chelipoda</i>) 263
<i>papuana</i> Bez. (<i>Empis</i>) 167	<i>pennata</i> Macq. (<i>Rhamphomyia barbata</i>) 201, 384	<i>physoprocta</i> Frey (<i>Rhamphomyia</i>) 202
<i>papuanus</i> Kert. (<i>Hybos</i>) 23	<i>pennata</i> Panz. (<i>Empis ciliata</i>) 151	<i>picea</i> Wied. (<i>Lactistomyia</i>) 36
<i>paradoxa</i> Wahlb. (<i>Rhamphomyia</i>) 201	<i>pennata</i> Schrank (<i>Empis pennipes</i>) 168	<i>picena</i> Bez. (<i>Empis</i>) 168
<i>paradoxus</i> Phil. (<i>Ceratomerus</i>) 215	<i>pennata</i> Stro. (<i>Empis albinervis</i>) 146	<i>picipes</i> Meig. (<i>Empis</i>) 169, 385
Parahybos Kert. (gen.) 45	<i>pennescens</i> Mel. (<i>Drapetis</i>) 311	<i>picipes</i> Zett. (<i>Drapetis curvipes</i> , not <i>moriella</i>) 311, 384
parallelia, nov. sp. (<i>Thinodromia</i>) 297	<i>pennipes</i> Linn. (<i>Empis</i>) 168	<i>picipes</i> Zett. (<i>Platypalpus</i>) 363
Paramesia Macq. (<i>Clinocera</i> Meig.) 227	<i>pennipes</i> Meig. (<i>Empis ciliata</i>) 151	<i>picipes</i> Zett. (<i>Rhamphomyia albipennis</i>) 184
Pararhamphomyia Frey (subgen. <i>Rhamphomyia</i>) 183	<i>penniventris</i> Bez. (<i>Empis</i>) 168	<i>picta</i> Lw. (<i>Empis spiloptera</i>) 173
* Parathalassia Meun. (gen.) 371	<i>pentophora</i> Wied. (<i>Empis atra</i>) 148	<i>picta</i> Stro. (<i>Clinocera dimidiata</i>) 239
Parathalassius Mik (gen.) 86	<i>peradeniæ</i> White (<i>Syneches</i>) 383	<i>pictipennis</i> Bez. (<i>Cephalodromia</i>) 268
<i>parilis</i> Coll. (<i>Drapetis</i>) 378	* <i>perdita</i> Coq. (<i>Empis</i>) 168	<i>pictipennis</i> Bez. (<i>Platypalpus</i>) 363
<i>parva</i> Coq. (<i>Rhamphomyia</i>) 201	<i>peregrina</i> Mel. (<i>Empis</i>) 168	<i>pictipennis</i> Bez. (<i>Rhamphomyia</i>) 202
<i>parva</i> , nov. sp. (<i>Chelipoda</i>) 266	<i>peregrina</i> White (<i>Hilarempis</i>) 130	<i>pictipes</i> Beck. (<i>Symballophthalmus dissimilis</i>) 300
<i>parvicauda</i> Coll. (<i>Platypalpus</i>) 382	<i>peregrinata</i> Walk. (<i>Ocydromia glabricula</i>) 70	<i>pictipes</i> Coq. (<i>Mythicomyia</i>) 376
<i>parvicellulata</i> Bez. (<i>Syndyas</i>) 37	<i>perimerus</i> , nov. var. (<i>Platypalpus vittatus</i>) 334	<i>pictipes</i> , nov. sp. (<i>Tachyempis</i>) 290
<i>parvicellulata</i> Frey (<i>Rhamphomyia</i>) 201	<i>periscelis</i> , nov. sp. (<i>Syneches</i>) 43	<i>pictitarsis</i> Beck. (<i>Platypalpus</i>) 363
<i>parvicornis</i> Mel. (<i>Drapetis</i>) 311	<i>perpendicularis</i> Lw. (<i>Empis</i>) 168	<i>pictithorax</i> Bez. (<i>Drapetis</i>) 315
<i>parvicornis</i> Zett. (<i>Platypalpus</i>) 363	<i>perplexa</i> Bez. (<i>Hilara</i>) 123	Pieltainia Arias (gen.) 287
<i>parvula</i> Egg. (<i>Empis pallens</i>) 167, 383	<i>perpulchra</i> Bez. (<i>Rhamphomyia</i>) 202	<i>pilatus</i> , nov. sp. (<i>Platypalpus</i>) 340
<i>parvula</i> Stro. (<i>Empis loewiana</i>) 161	* <i>personata</i> Meun. (<i>Empis</i>) 168	<i>pilicornis</i> Lw. (<i>Empis</i>) 169
<i>patagiata</i> Bez. (<i>Empis</i>) 167	<i>perspicuus</i> Hutt. (<i>Fraudator</i>) 379	<i>pilifer</i> Meig. (<i>Rhamphomyia</i>) 202
<i>pauciseta</i> , nov. sp. (<i>Stilpon</i>) 304	<i>pertinax</i> Harr. (<i>Empis stercorea</i>) 174	<i>piligeronis</i> Coq. (<i>Rhamphomyia</i>) 202
<i>paulseni</i> Phil. (<i>Platypalpus</i>) 363	<i>perturbans</i> Bez. (<i>Hilara</i>) 123	<i>pilimana</i> Lw. (<i>Empis</i>) 169
<i>pavesii</i> Bez. (<i>Empis</i>) 167	<i>perversa</i> Old. (<i>Hilara</i>) 123	<i>pilimanus</i> Stro. (<i>Microphorus</i>) 93
<i>pectinata</i> Lw. (<i>Rhamphomyia</i>) 201	<i>peshawarensis</i> Brun. (<i>Hilara</i>) 123	
<i>pectinata</i> , nov. sp. (<i>Bicellaria</i>) 76	<i>petiolata</i> , nov. var. (<i>Anthalia stigmatis</i>) 60	
	<i>petulans</i> Beck. (<i>Empis</i>) 168	

DIPTERA

	Pages		Pages		Pages
pilipes Lw. (<i>Bicellaria</i>)	77	plumipes Fall. (<i>Rhamphomyia squamigera</i>)	206	præputiata Lw. (<i>Empis</i>)	170
pilipes Lw. (<i>Drapetis</i>)	311	plumipes Mats. (<i>Empimorpha</i>)	179	præstans Bez. (<i>Syneches</i>)	43
pilipes Meig. (<i>Empis grisea</i>)	158	plumipes Meig. (<i>Rhamphomyia</i>)	202	præstans Frey (<i>Rhamphomyia</i>)	203
pilipes Meig. (<i>Hybos grossipes</i>)	22	plumipes Mel. (<i>Drapetis</i>)	311	prævia Coll. (<i>Empis</i>)	379
pilipes Stro. (<i>Hilara tanychira</i>)	126	plumipes Zett. (<i>Empis</i>)	169, 385	prasinata, nov. sp. (<i>Clinocera</i>)	232
pilipes Zett. (<i>Hilara</i>)	123	plurivittata Bez. (<i>Chelifera</i>)	262	pratincola Phil. (<i>Chelifera</i>)	262
pilipes Zett. (<i>Hilara lundbecki</i>)	120	pluto Mel. (<i>Platypalpus</i>)	364	precatoria Fall. (<i>Chelifera</i>)	260
pilitibia Macq. (<i>Empis</i>)	169	podagra Mel. (<i>Empis</i>)	169	precatoria Zett. (<i>Chelifera concinnata</i>)	378
pilosa Bonsd. (<i>Hilara interstincta</i>)	119	podagraria Meig. (<i>Empis</i>)	169	* predatoris Meun. (<i>Platypalpus</i>)	364
pilosa Lw. (<i>Empis</i>)	169	podocarpi Beck. (<i>Platypalpus</i>)	364	priapulus Lw. (<i>Rhamphomyia</i>)	203
pilosa Lundb. (<i>Bicellaria</i>)	77	pœcila Phil. (<i>Empis</i>)	169	primula Coll. (<i>Hilara</i>)	380
pilosa Mel. (<i>Drapetis</i>)	309	pœcioloptera Beck. (<i>Rhamphomyia paradoxo</i>)	201	* problematica Meun. (<i>Parathalassia</i>)	371
pilosa Zett. (<i>Hilara</i>)	124	pœcioloptera Lw. (<i>Empis</i>)	169	procera Lw. (<i>Empis</i>)	170
pilosiformis Bez. (<i>Enhybos</i>)	34	* poeppigi Giebel (<i>Empis</i>)	169	procera Lw. (<i>Haplocyrtoma</i>)	78
pilosipectinata Stro. (<i>Hilara</i>)	124	pokornyi Bez. (<i>Rhamphomyia</i>)	202	Preclinoptyga , nov. gen.	220
pilosus Schin. (<i>Enhybos</i>)	34	pokornyi Mik (<i>Clinocera</i>)	239	procurrrens, nov. sp. (<i>Drapetis</i>)	309
pinetorum auct. (<i>Hilara brevistyla</i>)	380	politata Lw. (<i>Rhamphomyia</i>)	202	prodromus Lw. (<i>Empis</i>)	170
pinetorum Beck. (<i>Hilara clypeata</i>)	116	politata Lw. (<i>Syndyas</i>)	37	productipes Stro. (<i>Tachydromia arrogans</i>)	284
pinetorum Beck. (<i>Hilara diversipes</i>)	117	politata Macq. (<i>Empis</i>)	169	* proosti Meun. (<i>Palaoparamesia</i>)	371
pinetorum Schin. (<i>Hilara nitidula</i>)	123	politata, nov. sp. (<i>Lactistomyia</i>)	36	propinquata Meij. (<i>Rhamphomyia subata</i>)	203, 384
pinetorum Zett. (<i>Hilara clypeata</i>)	124, 385	politellus, n. n. (<i>Platypalpus</i>)	382	Prerates Mel. (gen.)	376
pirata Mik (<i>Wiedemannia</i>)	243	polititus Coll. (<i>Platypalpus</i>)	382	prorsus, nov. sp. (<i>Platypalpus</i>)	328
pittoprocta Lw. (<i>Empis</i>)	169	polititus, nov. sp. (<i>Platypalpus politellus</i>)	333, 382	proserpina Bez. (<i>Platypalpus</i>)	364
planetica Coll. (<i>Empis lamelligornis</i>)	379	pollinosus Bez. (<i>Hybos</i>)	23	protarsalis Coll. (<i>Empis lamellicornis</i>)	379
planiceps, nov. sp. (<i>Syneches</i>)	43	pollinosus Stro. (<i>Platypalpus</i>)	364	* Protodalea Ckll. (gen.)	371
platycnemis Frey (<i>Rhamphomyia</i>)	202	poplitea Lw. (<i>Empis</i>)	169	proxima Meig. (<i>Empis hyalipennis</i>)	158
Platypalpus Macq. (gen.)	315	poplitea Wahlb. (<i>Rhamphomyia</i>)	202	pruinosa Coq. (<i>Edalea</i>)	65
Platyptera auct. (<i>Rhamphomyia</i> Meig.)	180	poppiusi Frey (<i>Platypalpus</i>)	364	pruinosa Coq. (<i>Tachypeza</i>)	278
Platyptera Meig. (subg. <i>Empis</i>)	143	Porphyrechrea (nov. gen.)	137	pruinosa Wied. (<i>Empis</i>)	170
platyptera Panz. (<i>Rhamphomyia marginata</i>)	198	* porphyropoides Meun. (<i>Meghyperella</i>)	370	pruinosa Wied.-Meig. (<i>Hilara</i>)	124
platyptera Stro. (<i>Empis</i>)	169	* porrecta Meun. (<i>Rhamphomyia</i>)	203	pseliophora, nov. sp. (<i>Tachydro-</i>	
Platypterygia Steph. (<i>Empis</i> Linn.)	139	porrectus Mel. (<i>Platypalpus</i>)	364	<i>mia</i>)	281
Platytelma Rond. (<i>Microsania</i> Zett.)	374	portacula Walk. (<i>Tachypeza corticalis</i>)	276	pseudobicolor Stro. (<i>Platypalpus</i>)	364
platytra Lw. (<i>Hilara griseola</i>)	118, 380	porteri Brethes (<i>Rhamphomyia</i>)	382	pseudochorica Stro. (<i>Hilara</i>)	124
plebeia Walk. (<i>Hilara</i>)	124	postica Walk. (<i>Tachypeza</i>)	278	pseudociliaris Stro. (<i>Platypalpus</i>)	364
plebeja Lw. (<i>Empis</i>)	169	postpositus nov. sp. (<i>Platypalpus</i>)	343	pseudocornicula Stro. (<i>Hilara</i>)	124
plectrum Mik (<i>Clinocera</i>)	239	Polyblepharis Bez. (subgen. <i>Empis</i>)	144	pseudocrinita Stro. (<i>Rhamphomyia</i>)	203
pleuralis Mel. (<i>Drapetis</i>)	315	polychaeta Bez. (<i>Hilara empis</i>)	130	pseudodecora Stro. (<i>Empis</i>)	170
pleuritica, nov. sp. (<i>Stilpon</i>)	302	Polydromia Big. (<i>Chelifera</i> Macq.)	258	pseudoexiguus Stro. (<i>Platypalpus</i>)	364
plorans Bez. (<i>Empis</i>)	169	polygynus, nov. sp. (<i>Anthepis copus</i>)	105	Pseudofulvipes Frey (<i>Platypalpus flavipes</i>)	355
plumbea Fab. (<i>Hilara fuscipes</i>)	118	* polymorpha Lw. (<i>Rhamphomyia</i>)	202	pseudogibba Stro. (<i>Rhamphomyia</i>)	203
plumea Mel. (<i>Drapetis</i>)	315	præcinctus Coll. (<i>Platypalpus</i>)	382	pseudomaculipes Stro. (<i>Platypal-</i>	
plumicornia White (<i>Drapetis</i>)	378	præcox Lw. (<i>Empis</i>)	170	<i>pus</i>)	364
plumicornis Bez. (<i>Hybos</i>)	23	præcox Lw. (<i>Microphorella</i>)	89	pseudomalleola Stro. (<i>Empis</i>)	170
plumifera Bonsd. (<i>Rhamphomyia lucidula</i>)	197	prælusio Walk. (<i>Tachypeza</i>)	278	pseudosartrix Stro. (<i>Hilara</i>)	124, 385
plumifera Zett. (<i>Rhamphomyia</i>)	202			<i>psuedotrislineata</i> Stro. (<i>Rhamphomyia lavipes</i>)	196
plumipes auct. (<i>Rhamphomyia geniculata</i>)	202, 384				

Pages	Pages	Pages			
pseudounguiculatus Stro. (<i>Platypalpus</i>)	364	<i>pusilla</i> Lw. (<i>Wiedemannia lota</i>)	243	raptoria Meig. (<i>Hemerodromia</i>)	257, 385
Pseudowiedemannia Eng. (subgen. <i>Wiedemannia</i>)	241	<i>pusilla</i> Scop. (<i>Rhamphomyia sulcata</i>)	207	rava Lw. (<i>Empis</i>)	171
Pterempis Bez. (<i>Empis</i> , s. str.)	139	<i>pusilla</i> Zett. (<i>Rhamphomyia</i>)	203	rava Lw. (<i>Rhamphomyia</i>)	203
*pteropa Lw. (<i>Rhamphomyia</i>)	203	<i>pusillus</i> Lw. (<i>Syneches</i>)	43	ravida Coq. (<i>Empis obesa</i>)	166
pteropoda Egg. (<i>Empis</i>)	170	<i>pusillus</i> Macq. (<i>Microphorus anomalis</i>)	93	ravidus Coq. (<i>Microphorus</i>)	93
Pterospilus Rond. (<i>Syneches</i> Walk.)	38	<i>pusillus</i> Meij. (<i>Parahybos</i>)	46	recedens Hal. (<i>Hilara</i>)	125
ptilocnemis Lw. (<i>Empis</i>)	170	<i>pusio</i> Egg. (<i>Empis</i>)	171	reciproca Walk. (<i>Empis</i>)	171
*ptilopa Lw. (<i>Rhamphomyia</i>)	203	<i>pusio</i> Lw. (<i>Rhamphomyia</i>)	203	rectineura Mel. (<i>Drapetis</i>)	315
Ptilophyllodromia Bez. (gen.)	268	*putidus Meun. (<i>Microphorus</i>)	93	recurvus, nov. sp. (<i>Platypalpus</i>)	339
ptilopoda Wied.-Meig. (<i>Empis</i>)	170	<i>pygmæa</i> Ros. (<i>Drapetis</i> , not assimilis)	308, 384	reflexa Zett. (<i>Rhamphomyia</i>)	203
pubescens Lw. (<i>Megagrapha</i>)	301	<i>pygmæus</i> Macq. (<i>Platypalpus</i>)	364	rejecta Coll. (<i>Hilara</i>)	380
pubescens, nov. sp. (<i>Platypalpus</i>)	331	<i>pygmæus</i> Meig. (<i>Platypalpus</i>)	364	*remitarsis Lw. (<i>Rhamphomyia</i>)	203
pubicornis Bez. (<i>Drapetis</i>)	311	<i>pygmæus</i> Zett. (<i>Platypalpus ciliaris</i>)	350	repletus Bez. (<i>Syneches</i>)	44
pubicornis Zett. (<i>Platypalpus longicornis</i>)	357	<i>pyramidatus</i> Bez. (<i>Syneches</i>)	43	reticulata Four. (<i>Empis livida</i>)	161
pubipes Lw. (<i>Hilara intermedia</i>)	124, 385	Pyrrempis , nov. subg. (<i>Empis</i>)	144	reversus Walk. (<i>Hybos</i>)	23
pudens, nov. sp. (<i>Platypalpus</i>)	329	quadrangularis Wh.-Mel. (<i>Syneches</i>)	44	rhabdoptera, nov. sp. (<i>Chelipoda</i>)	266
pudica Lw. (<i>Empis</i>)	170	quadricinctum Fab. (<i>Brachystoma</i>)	14	Rhadinempis Coll. (subgen. <i>Empis</i>)	379
puerinus, nov. var. (<i>Platypalpus juvenis</i>)	332	quadriclavata Stro. (<i>Hilara</i>)	125	Rhagas auct. (see <i>Ragas</i>)	109
pulchellus Walk. (<i>Platypalpus</i>)	364	quadrifaria Beck. (<i>Hilara empis</i>)	130	Rhamphomyia Meig. (gen.)	180
pulcherrima Bez. (<i>Leptopeza</i>)	67	quadrifaria Coll. (<i>Hilara subpolinosa</i>)	380	rhynchos Now. (<i>Wiedemannia</i>)	245, 434
pulchra Egg. (<i>Rhamphomyia umbripennis</i>)	209	quadrifaria Stro. (<i>Hilara</i>)	125	ribesii Beck. (<i>Anthepiscopus</i>)	108
pulchra Lw. (<i>Rhamphomyia</i>)	203	quadrilineata Gmel. (<i>Empis</i>)	171	rileyi Coq. (<i>Mythicomyia</i>)	376
pulchripes Frey (<i>Hilara</i>)	124	quadrilineata Macq. (<i>Hilara quadrivittata</i>)	125	riparia Rob. (<i>Clinocera</i>)	237
pulchripes Lw. (<i>Empis</i>)	170	quadripilosa Beck. (<i>Hilara</i>)	125	rivosa Big. (<i>Leptopeza</i>)	67
pulchriventris Bez (<i>Rhamphomyia</i>)	203	quadriseta Coll. (<i>Hilara</i>)	380	robertii Macq. (<i>Clinocera nigra</i>)	237
pulicaria Lw. (<i>Empis</i>)	170	quadrisetosa Mel. (<i>Drapetis</i>)	311	robertsonii Coq. (<i>Brachystoma</i>)	14
pulicarius Fall. (<i>Atelestus</i>)	384	quadrivittata Arrib. (<i>Empis</i>)	171	*robusta Meun. (<i>Edalea</i>)	65
pulicarius Meig. (<i>Platypalpus</i>)	364	quadrivittata Beck. (<i>Hilara carinthiaca</i>)	115	robustior Frey (<i>Rhamphomyia</i>)	203
pulla Bez. (<i>Syneches</i>)	45	quadrivittata Meig. (<i>Hilara</i>)	125	robustus, nov. sp. (<i>Microphorus</i>)	91
pulla Lw. (<i>Rhamphomyia</i>)	203	quadrivittata Zett. (<i>Hilara beckeri</i>)	114	robustus Walk. (<i>Platypalpus</i>)	364
pullata Mel. (<i>Heleodromia</i>)	225	quercifolia Eng. (<i>Wiedemannia</i>)	242	Roederella Eng. (subgen. <i>Wiedemannia</i>)	242
pullata, nov. sp. (<i>Trichina</i>)	57	quinquelineata Say (<i>Rhamphomyia</i>)	203	Roedereria Mik. (<i>Wiedemannia</i> Zett.)	240
pulverulentus, nov. sp. (<i>Platypalpus</i>)	341	quinquevittata Macq. (<i>Empis femorata</i>)	156	Roederioides Coq. (gen.)	225
*pulvillata Lw. (<i>Empis</i>)	170	Ragas Mel. (<i>Hesperempis Mel.</i>)	111	rogatoris Coq. (<i>Hemerodromia</i>)	257
punctata Fab. (<i>Empis digramma</i>)	154	Ragas Walk. (gen.)	109	rondaensis Stro. (<i>Platypalpus</i>)	364
punctata Fall (<i>Empis testacea</i>)	175	rapax Lw. (<i>Tachypeza fenestrata</i>)	276	rostellatus Lw. (<i>Microphorus</i>)	93
punctata Fall. (<i>Empis trigramma</i>)	176	rapax Pallas-Wied. (<i>Empis sternocoreata</i>)	174	rostrata Brun. (<i>Empis</i>)	171
punctata Meig. (<i>Empis</i>)	170	rapida Meig. (<i>Empis dasyprocta</i>)	153	rostrata Lw. (<i>Tachypeza</i>)	278
punctata Scop. (<i>Empis</i>)	171	rapidus Meig. (<i>Platypalpus</i>)	364	rostrata Zett. (<i>Euthyneura myrtilli</i> , not <i>Anthepiscopus</i>)	107, 384
punctifera Beck. (<i>Tachydromia</i>)	286	raptor Latr. (<i>Chelipoda melanoleuca</i>)	267	rostrata Zett. (<i>Heleodromia immaculata</i>)	225
punctipennis Wulp (<i>Syneches phthisia</i>)	43	raptor Macq. (<i>Chelifera precatoria</i>)	261	rostrifera Bez. (<i>Rhamphomyia</i>)	204
purpureus Walk. (<i>Euhybos</i>)	34	raptoria Bez. (<i>Empis</i>)	171	rotundicornis Brun. (<i>Drapetis</i>)	309
pusilla Lw. (<i>Drapetis exilis</i>)	309, 384			rotundipennis Bez. (<i>Rhamphomyia</i>)	204
pusilla Lw. (<i>Drapetis pygmæa</i>)	309, 384			rubefactus, nov. sp. (<i>Platypalpus</i>)	322
pusilla Lw. (<i>Tachydromia maculipennis</i>)	286			rubida Wh.-Mel. (<i>Opeatocerata</i>)	135
				rubripes Phil. (<i>Haplomera</i>)	132

DIPTERA

Pages	Pages	Pages			
rubrithorax Bez. (<i>Drapetis</i>)	312	sabulosa Meig. (<i>Tachydromia</i>)	286	Sciadromia Hal. (<i>Hileodromia</i> Hal.)	224
rubrithorax White (<i>Leptopeza</i>)	67	sachem, nov. sp. (<i>Tachydromia</i>)	283	scissa Mel. (<i>Drapetis</i>)	311
rufa Meig. (<i>Bicellaria</i>)	77	sahlbergi Frey (<i>Platypalpus</i>)	364	scitula Frey (<i>Rhamphomyia</i>)	204
rufescens Lw. (<i>Empis</i>)	171	salicina Lioy (<i>Empis</i>)	172	scolopacea Say (<i>Rhamphomyia</i>)	204
rufescens, nov. sp. (<i>Colabris</i>)	258	salictorum Lw. (<i>Rhamphomyia stigmatica</i>)	206	scoparia Coq. (<i>Empis</i>)	172
ruficollis Macq. (<i>Ocydromia glabricula</i>)	70	*samlandica Meun. (<i>Palaesdalea</i>)	370	scopifer Coq. (<i>Oreogeton</i>)	100
ruficollis Meig. (<i>Leptopeza</i>)	67	sancti-mauritii Beck. (<i>Rhamphomyia</i>)	204	scopulifera Bez. (<i>Empis</i>)	172
ruficollis Walk. (<i>Syneches</i>)	44	sancti-mauritii Stro. (<i>Rhamphomyia discoidalis</i>)	190	scotica Curt. (<i>Empis</i>)	167, 385
ruficornis Lw. (<i>Empis</i>)	171	sancti-mauritii Stro. (<i>Rhamphomyia villosa</i>)	210	scrobiculata Lw. (<i>Hilara</i>)	125
ruficornis Macq. (<i>Platypalpus flavicornis</i>)	354	sanduca, nov. sp. (<i>Hesperempis</i>)	111	scutellaris Bez. (<i>Empis</i>)	172
ruficornis, nov. sp. (<i>Tachyempis</i>)	291	sanguineolentus, n. n. (<i>Platypalpus</i>)	365	scutellaris Coq. (<i>Rhamphomyia</i>)	204
ruficornis Ros. (<i>Platypalpus</i>)	364, 384	sanguineus Arrib. (<i>Platypalpus</i>)	365	scutellaris, nov. sp. (<i>Anthalia</i>)	60
rufipes Bez. (<i>Clinocera nigra</i>)	237	sanguinensis Meij. (<i>Platypalpus sanguineolentus</i>)	365	scutellata Coq. (<i>Mythicomya</i>)	376
rufipes Brun. (<i>Drapetis</i>)	309	saniculae Curt. (<i>Rhamphomyia</i>)	204	scutellata Curt. (<i>Empis</i>)	172
rufipes Fab. (<i>Empis opaca</i>)	166	sapporensis Mats. (<i>Rhamphomyia</i>)	204	scutellata Meig. (<i>Ocydromia glabricula</i>)	70
rufipes Gmel. (<i>Empis</i>)	171	sartor Beck. (<i>Hilara</i>)	125	securigera Eng. (<i>Clinocera barbatula</i>)	239
rufipes Macq. (<i>Hilara</i>)	125	sartrix Handl. (<i>Hilara sartor</i>)	125	sellata Lw. (<i>Rhamphomyia</i>)	204
rufipes Macq. (<i>Edalea flavipes</i>)	64	satyriacus, nov. sp. (<i>Platypalpus</i>)	327	semaptera Gil (<i>Ariasella</i>)	377
rufipes Meig. (<i>Microphorus velutinus</i>)	94	sauteri Bez. (<i>Parahybos</i>)	46	semibrunneus Brun. (<i>Syneches</i>)	44
rufipes Meig. (<i>Ocydromia glabricula</i>)	70	sauteri Bez. (<i>Rhamphomyia</i>)	204	seminerea Lw. (<i>Empis</i>)	172
rufipes Meig. (<i>Platypalpus</i>)	364	sauteriana Bez. (<i>Empis</i>)	172	semifasciata Stro. (<i>Tachydromia styriaca</i>)	286
rufipes Meig. (<i>Rhamphomyia culicina</i>)	190	scapularis Lw. (<i>Chelisora</i>)	262	semifulvus Big. (<i>Microphorus</i>)	93
rufipes Meig. (<i>Rhamphomyia tenuirostris</i>)	207	scapuliferus Big. (<i>Microphorus</i>)	93	semihyalipennis Gimm. (<i>Platypalpus</i>)	365
rufipes Old. (<i>Wiedemannia cernyi</i>)	244	scatophagina Mel. (<i>Empis</i>)	172	semilugens Phil. (<i>Chelisora</i>)	262
rufipes Scop. (<i>Empis</i>)	171	saura Lw. (<i>Empis</i>)	172	seminigra Mel. (<i>Drapetis</i>)	315
rufipes Staeg. (<i>Empis tessellata</i>)	175	scaurissima Wh. (<i>Rhamphomyia</i>)	204	senator, nov. sp. (<i>Hormopeza</i>)	95
rufipes Wied. (<i>Empis</i>)	171	Scelolabes Reed (<i>Scelolabes Phil.</i>)	71	senilis Panz. (<i>Hilara pilosa</i>)	124
rufipes Zett. (<i>Rhamphomyia</i>)	204	Scelolabes Phil. (gen.)	71	septembris White (<i>Rhamphomyia</i>)	204
rufirostris Say (<i>Rhamphomyia</i>)	204	schildi, nov. sp. (<i>Eshybos</i>)	31	septemnotata Brun. (<i>Dolichocephala</i>)	247
rufitarsis Ros. (<i>Hybos femoratus</i>)	22	schildi, nov. sp. (<i>Tachyempis</i>)	289	septentrionalis Mel. (<i>Drapetis</i>)	311
rufitarsis Zett. (<i>Hybos culiciformis</i>)	21	schistaceus Meig. (<i>Rhamphomyia cassia</i>)	187	*sepulta Ckll. (<i>Rhamphomyia</i>)	204
rufithorax Brun. (<i>Empismorpha</i>)	179	Schistostoma Beck. (<i>Microphorus Macq.</i>)	90	sequens Mel. (<i>Lactistomyia dimidiata</i>)	35
rufithorax Brun. (<i>Rhamphomyia</i>)	204	schizophorus, nov. sp. (<i>Philetus</i>)	110	serena Pok. (<i>Empis</i>)	172
rufiventris Meig. (<i>Empis</i>)	171	schnabli Beck. (<i>Clinocera</i>)	239	seriata Lw. (<i>Hilara</i>)	126
rufiventris Mel. (<i>Platypalpus</i>)	364	schnaeusei Bez. (<i>Atrichopleura</i>)	134	sericans Brullé (<i>Empis</i>)	172
rufiventris Stro. (<i>Empis grisea</i>)	158	schnaeusei Bez. (<i>Carathybos</i>)	36	sericata White (<i>Empis</i>)	172
rufomaculata White (<i>Sciadocera</i>)	48	schoenherri Zett. (<i>Anthalia</i>)	61	sericatus, nov. sp. (<i>Platypalpus</i>)	320
rufopunctata Curr. (<i>Hilara</i>)	380	schrottkyi Bez. (<i>Haplomera</i>)	132	sericea Oliv. (<i>Empis femorata</i>)	156
rufus Lw. (<i>Oreogeton</i>)	100	schwarzii Coq. (<i>Tachydromia</i>)	286	sericeipalpis Frey (<i>Tachypeza</i>)	278
rufus Lw. (<i>Syneches</i>)	44	schwarzii Wh.-Mel. (<i>Tachydromia ciliata</i>)	285	sericeus Macq. (<i>Platypalpus</i>)	365
rugicollis Meig. (<i>Rhamphomyia sulcata</i>)	204, 384	Soladeocera White (gen.)	48, 373	serotina Lw. (<i>Empis</i>)	172
rugosiventris Stro. (<i>Leptopeza</i>)	68	sciapterus, nov. sp. (<i>Hybos</i>)	23	serotina Old. (<i>Rhamphomyia</i>)	205
rupestris Old. (<i>Rhamphomyia</i>)	383	sciarina Fall. (<i>Rhamphomyia</i>)	204	serperastrorum Mel. (<i>Empis poplitea</i>)	170
ruralis Meig. (<i>Empis femorata</i>)	156			serrata Bez. (<i>Lactistomyia</i>)	36
rustica Fall. (<i>Empis nigricans</i>)	165			serrata Schrank (<i>Empis</i>)	172
rustica Lw. (<i>Rhamphomyia</i>)	204			serraticosta White (<i>Leptopeza</i>)	68
*rustica Meun. (<i>Chelipoda</i>)	266				
rusticus Brun. (<i>Syneches</i>)	44				

Pages	Pages	Pages			
serratula Lw. (<i>Brachystoma</i>)	13	soeculatus Zett. (<i>Platypalpus cothurnatus</i>)	351	stenoptera Lw. (<i>Empis</i>)	173
setacea Beck. (<i>Iteaphila macquarti</i>)	104	sociabilis Will. (<i>Rhamphomyia</i>)	205	stentor Mel. (<i>Antheffiscopus</i>)	108
serpentata Lw. (<i>Rhamphomyia</i>)	205	solivaga, nov. sp. (<i>Procliuopyga</i>)	223	stercoraria Macq. (<i>Empis stercorea</i>)	174
setifrons Bez. (<i>Micrempis</i>)	299	sordida Lw. (<i>Empis</i>)	172	stercorea Linn. (<i>Empis</i>)	173
setigera Bez. (<i>Leptopeza</i>)	68	sordida Lw. (<i>Hilarempis</i>)	130	starcorea Zett. (<i>Empis univittata</i>)	177
setigera Coq. (<i>Lamprempis</i>)	136	sordida Lw. (<i>Rhamphomyia</i>)	205	stigma Coll. (<i>Platypalpus</i>)	382
setigera Lw. (<i>Drapetis, not moriella</i>)	311, 384	sordidus Zett. (<i>Platypalpus</i>)	365	stigma Meig. (<i>Empis stercorea</i>)	174
setigera Lw. (<i>Empis lepidopus</i>)	160	sordipes, nov. sp. (<i>Euhybos</i>)	28	stigma Walk. (<i>Syneches</i>)	45
setigera Lundb. (<i>Drapetis curvipes</i>)	311, 384	sorex Eng. (<i>Wiedemannia</i>)	243	stigmatis Coq. (<i>Anthalia</i>)	61
setigera Stein (<i>Rhamphomyia amana</i>)	185	sparsa Beck. (<i>Dolichocephala</i>)	247	stigmatellus Zett. (<i>Platypalpus</i>)	365
setosa Bez. (<i>Ragas</i>)	382	spectabilis Frey (<i>Rhamphomyia</i>)	205	stigmatica Lundb. (<i>Chelifera cinnicauda</i>)	378
setosa Coll. (<i>Hilara</i>)	380	spectabilis Lw. (<i>Empis</i>)	173	stigmatica Schin. (<i>Chelifera</i>)	261
setosa Coq. (<i>Rhamphomyia</i>)	205	spectabilis Mel. (<i>Drapetis</i>)	311	stigmatical Bez. (<i>Syneches</i>)	44
setosa Lw. (<i>Empis</i>)	172	specularis Bez. (<i>Empis</i>)	173	stigmatical Coq. (<i>Edalea ohioensis</i>)	65
setosa Scop. (<i>Empis</i>)	172	speculifera Walk. (<i>Chersodromia</i>)	297	stigmatical Zett. (<i>Edalea</i>)	65
setosus Meij. (<i>Hybos</i>)	23	sphenoptera Lw. (<i>Leptometopiella</i>)	70	stigmatical Zett. (<i>Microsania</i>)	374
setulosa Mel. (<i>Drapetis</i>)	309	sphenoptera Lw. (<i>Rhamphomyia</i>)	205	stigmatical Zett. (<i>Wiedemannia escheri</i>)	243
sexsetosa Frey (<i>Trichina clavipes</i>)	56	Sphicosa Phil. (<i>Iteaphila</i> Zett.)	103	stigmaticum Phil. (<i>Brachystoma</i>)	14
sibillina Bez. (<i>Empis</i>)	172	spiloptera Wied. (<i>Empis</i>)	173	stigmaticus Schin. (<i>Euhybos</i>)	34
sicula Lw. (<i>Empis chioptera</i>)	151	spinicosta Wh.-Mel. (<i>Hybos</i>)	23	stigmosa Macq. (<i>Rhamphomyia</i>)	206
Sicus Latr. (<i>Tachydromia</i> Meig.)	279	spinifera Bez. (<i>Dipsomyia</i>)	220	*stilicornis Lw. (<i>Empis</i>)	174
siebecki Stro. (<i>Rhamphomyia</i>)	205	spinifera Bez. (<i>Empis</i>)	173	Stillpon Lw. (gen.)	301
signatus Big. (<i>Syneches</i>)	45	spiniger, nov. sp. (<i>Euhybos</i>)	32	*stilpon Meun. (<i>Tachydromia</i>)	286
similata Mall. (<i>Rhamphomyia</i>)	205	*spinigera Lw. (<i>Leptopeza</i>)	68	storchi Mik (<i>Clinocera appendiculata</i>)	236
similis Beck. (<i>Empis</i>)	172	spinigera Stro. (<i>Hilara spinimana</i>)	126	straminipes Stro. (<i>Platypalpus sahlbergi</i>)	364
similis Walk. (<i>Tachypeza fenestrata</i>)	277	spinimana Zett. (<i>Hilara</i>)	126	straminipes Zett. (<i>Platypalpus pectoralis</i>)	363
simillima Stro. (<i>Euthyneura</i>)	54	spinipes Fall. (<i>Rhamphomyia</i>)	205	striatellus Vill. (<i>Hybos</i>)	23
simplex Lw. (<i>Wiedemannia</i>)	244	spinipes Macq. (<i>Hilara pilosa</i>)	124	striaticollis Beck. (<i>Hilara</i>)	126
simplex, nov. var. (<i>Euhybos triplex</i>)	30	spinipes Mel. (<i>Drapetis</i>)	311	striatipennis Stro. (<i>Tachydromia catalonica</i>)	284
simplex Walk. (<i>Syneches phthisia</i>)	43	spinipes, nov. sp. (<i>Euthyneura</i>)	54	strigata Lw. (<i>Empis</i>)	174
simplex Wied. (<i>Hilara maura</i>)	122	spinipes, nov. sp. (<i>Silpon</i>)	303	strigifera Meij. (<i>Drapetis</i>)	312
simplex Zett. (<i>Rhamphomyia</i>)	205	spinosa Bez. (<i>Hilarempis</i>)	130	strigifrons Frey (<i>Platypalpus fascipes</i>)	354
simplicinervis Frey (<i>Clinocera apendiculata</i>)	236	spinopipes Old. (<i>Rhamphomyia</i>)	206	strigifrons Zett. (<i>Platypalpus</i>)	365
simplicinervis, nov. sp. (<i>Helemordromia</i>)	257	spinopus, nov. sp. (<i>Platypalpus</i>)	319	strobliana Bez. (<i>Hilara</i>)	126
simplicior Wh.-Mel. (<i>Tachyempis</i>)	292	spinosula Phil. (<i>Hilarempis</i>)	130	stroblianus, n. n. (<i>Platypalpus</i>)	365
simplicipes Bez. (<i>Parahybos</i>)	46	*spinosulum Lw. (<i>Brachystoma</i>)	14	strollii Mik (<i>Platypalpus engadinius</i>)	353
simplicipes Lw. (<i>Empis</i>)	171	spirifera Bez. (<i>Empis</i>)	173	strumaticus, nov. sp. (<i>Euhybos</i>)	26
simplicipes Mel. (<i>Drapetis</i>)	315	spuria Fall. (<i>Bicellaria</i>)	77, 384	sturmii Meig. (<i>Hilara cingulata</i>)	116
simplicipes, nov. sp. (<i>Platypalpus</i>)	331	squalens Zett. (<i>Hilara manicata</i>)	121	stylata Coq. (<i>Rhamphomyia</i>)	206
simplices Stro. (<i>Hilara</i>)	126	squamata Brun. (<i>Empis</i>)	173	stylifera Mik (<i>Wiedemannia</i>)	245
simplices Zett. (<i>Bicellaria melæna</i>)	76	squamigera Lw. (<i>Rhamphomyia</i>)	206	styriaca Stro. (<i>Empis</i>)	174
similium Now. (<i>Rhamphomyia costata</i>)	189	squamipes Coq. (<i>Empis</i>)	173	styriaca Stro. (<i>Hilara longevittata</i>)	120
singularis Egg. (<i>Hilarimorpha</i>)	373	squamossima Stro. (<i>Rhamphomyia squamigera</i>)	206	styriaca Stro. (<i>Tachydromia</i>)	286
slossonæ Coq. (<i>Hybos reversus</i>)	23	stagnalis Hal. (<i>Clinocera</i>)	238	styriaca Stro. (<i>Tachydromia interrupta</i>)	285
smithi Wh.-Mel. (<i>Euhybos</i>)	34	Steleocheta Beck. (<i>Iteaphila Zett.</i>)	103		
smithii Hutt. (<i>Hilara</i>)	126	Stenoprocetus Lw. (gen.)	72		
soccata Lw. (<i>Rhamphomyia</i>)	205				
soccatus, nov. sp. (<i>Platypalpus</i>)	341				

DIPTERA

Pages	Pages	Pages
<i>styriensis</i> Beck. (<i>Iteaphila nitidula</i>) 104	Symballeophthalmus Beck.	tenuirostris Fall. (<i>Rhamphomyia</i>) 207
<i>suavis</i> Lw. (<i>Lamprempis</i>) 136	(gen.) 300	tenuirostris Thom. (<i>Empis</i>) 174
<i>subalpina</i> Old. (<i>Rhamphomyia</i>) 383	Synamphotera Lw. (gen.) 216	* <i>tenuis</i> Meun. (<i>Hybos</i>) 24
<i>subbrevis</i> Frey (<i>Platypalpus brevicornis</i>) 349	Synamphotera Mel. (<i>Boreodromia Coq.</i>) 215	<i>tenuis</i> , nov. sp. (<i>Platypalpus</i>) 326
<i>subbrunipennis</i> Old. (<i>Empis</i>) 370	Syndyas Lw. (gen.) 36	<i>tenuiterfilata</i> Beck. (<i>Rhamphomyia</i>) 207
<i>subciliata</i> Lw. (<i>Empis</i>) 174	Syneches Walk. (gen.) 38	<i>tephraea</i> Meig. (<i>Rhamphomyia</i>) 208
<i>subciliipes</i> Brun. (<i>Empis</i>) 174	<i>tabascensis</i> Wh.-Mel. (<i>Euhybos</i>) 35	<i>tephrodes</i> Phil. (<i>Atrichopleura</i>) 134
<i>subcinerascens</i> Coll. (<i>Rhamphomyia</i>) 383	Tachista Lw. (<i>Tachydromia Meig.</i>) 279	<i>tephrodes</i> Phil. (<i>Hilarempis</i>) 130
<i>subclavata</i> Lw. (<i>Empis</i>) 174	Tachydromia auct. (<i>Platypalpus Macq.</i>) 316	<i>teres</i> Mel. (<i>Empis</i>) 174
<i>suberis</i> Beck. (<i>Empis</i>) 174	Tachydromia Meig. (gen.) 278	<i>tergestinus</i> Egg. (<i>Platypalpus</i>) 366
<i>subglaucea</i> Frey (<i>Rhamphomyia</i>) 206	<i>tachydromiaeformis</i> Bez. (<i>Leptopeza</i>) 68	<i>ternovensis</i> Stro. (<i>Hilara</i>) 126
<i>subinfumata</i> Mall. (<i>Empis</i>) 370	TACHYDROMIINAE (subfam.) 269	<i>terrictola</i> Zett. (<i>Tachydromia</i>) 286
<i>subjectus</i> Walk. (<i>Euhybos</i>) 34	Tachyempis (nov. gen.) 288	<i>tersa</i> Coq. (<i>Rhamphomyia</i>) 208
<i>sublineata</i> Brullé (<i>Hilara</i>) 126	Tachypeza Meig. (gen.) 272	<i>tersa</i> Mel. (<i>Empis</i>) 174
<i>submaura</i> Coll. (<i>Hilara</i>) 380	<i>tachystiformis</i> , nov. sp. (<i>Platypalpus</i>) 320	<i>tersus</i> Coq. (<i>Platypalpus</i>) 366
<i>subnitida</i> Beck. (<i>Empis</i>) 174	<i>tacoma</i> , nov. sp. (<i>Tachydromia</i>) 281	<i>tessellata</i> Fab. (<i>Empis</i>) 174
<i>subpennata</i> Macq. (<i>Empis</i>) 174	<i>tacomæ</i> , nov. sp. (<i>Anomalempis</i>) 15	<i>testacea</i> Curt. (<i>Empis punctata</i>) 171
<i>subpilosa</i> Coll. (<i>Bicellaria</i>) 378	<i>tæniatus</i> Meig. (<i>Platypalpus</i>) 366	<i>testacea</i> Fab. (<i>Empis</i>) 175
<i>subpollinosa</i> Coll. (<i>Hilara</i>) 380	<i>talalis</i> , nov. sp. (<i>Platypalpus</i>) 342	<i>testacea</i> Lw. (<i>Hilara</i>) 126
<i>subsltans</i> Frey (<i>Rhamphomyia</i>) 206	<i>tantalus</i> Coll. (<i>Platypalpus</i>) 382	<i>testacea</i> Lw. (<i>Rhamphomyia</i>) 208
<i>subtilis</i> Coll. (<i>Platypalpus</i>) 382	<i>tanychira</i> Kow. (<i>Hilara</i>) 126	<i>testacea</i> , nov. sp. (<i>Micrempis</i>) 299
<i>subulifer</i> Meij. (<i>Platypalpus</i>) 365	<i>tanyphryra</i> Lw. (<i>Empis</i>) 174	<i>testacea</i> , nov. var. (<i>Edalea lanceolata</i>) 64
* <i>succinorum</i> Meun. (<i>Oustalietomyia</i>) 370	<i>tanythrix</i> Frey (<i>Hilara</i>) 126	<i>testacea</i> Walk. (<i>Empis scutellata</i>) 172
* <i>sucina</i> n. n. (<i>Trichopeza</i>) 220	<i>taos</i> Mel. (<i>Clinocera</i>) 238	<i>testacea</i> Zett. (<i>Empis digramma</i>) 154
<i>sudeticus</i> Lw. (<i>Meghyperus</i>) 47	* <i>tarda</i> Meun. (<i>Hilara</i>) 126	<i>testaceum</i> Phil. (<i>Brachystoma</i>) 14
<i>sudigeronis</i> Coq. (<i>Rhamphomyia</i>) 206	<i>tarsata</i> Meig. (<i>Rhamphomyia</i>) 207	<i>testaceus</i> Beck. (<i>Platypalpus engadensis</i>) 352
<i>sufasciatus</i> , nov. var. (<i>Platypalpus porrectus</i>) 336	<i>tarsata</i> Siebke (<i>Hilara</i>) 126	<i>testaceus</i> , nov. sp. (<i>Syneches</i>) 44
<i>sulcata</i> auct. (<i>Rhamphomyia profinqua</i>) 203	<i>tarsellus</i> Zett. (<i>Microphorus anomalous</i>) 93	<i>testaceus</i> Phil. (<i>Platypalpus</i>) 366
<i>sulcata</i> Meig. (<i>Rhamphomyia</i>) 206, 384	<i>tectifrons</i> Beck. (<i>Platypalpus</i>) 366	<i>testiculata</i> Bez. (<i>Empis</i>) 175
<i>sulcata</i> Meig. (<i>Rhamphomyia sulcata</i>) 206, 383	<i>tenax</i> , nov. sp. (<i>Platypalpus</i>) 339	<i>tetrachæta</i> , nov. sp. (<i>Tachyempis</i>) 291
<i>sulcata</i> Zett. (<i>Bicellaria, not spuria</i>) 77.	<i>tenebrosa</i> Coq. (<i>Empis hirsuterns</i>) 158	<i>tetragramma</i> Lw. (<i>Hilara</i>) 126
	<i>tenella</i> Fall. (<i>Hilara</i>) 126	Tetraneurella Dahl (<i>Stilpon Lw.</i>) 301
<i>sulcatella</i> Coll. (<i>Rhamphomyia</i>) 383	<i>tenella</i> Wahlb. (<i>Clinocera bipunctata</i>) 239	<i>thalhammeri</i> Stro. (<i>Empis</i>) 175
<i>sulcipes</i> Meig. (<i>Empis pictipes</i>) 164	<i>tenellus</i> Mel. (<i>Platypalpus</i>) 366	Thamnoredmia Mik (<i>Chelipoda Macq.</i>) 264
<i>sulcitarsis</i> Stro. (<i>Hilara</i>) 126	<i>tenera</i> Mel. (<i>Drapetis</i>) 315	<i>thermophila</i> Wied. (<i>Empis</i>) 175
<i>superba</i> Lw. (<i>Lamprempis</i>) 136	<i>teneriffensis</i> Beck. (<i>Platypalpus</i>) 366	Thinoredmia Mel. (gen.) 297
<i>superstitiosa</i> Say (<i>Hemeroedromia</i>) 257	Tenontomyia White (gen.) 133	* <i>Thirza</i> Giebel (gen.) 372
<i>supplicatoria</i> Meig. (<i>Chelipoda</i>) 266	<i>tenuicornis</i> Zett. (<i>Rhamphomyia hybotina</i>) 195	<i>thoracica</i> Eversm. (<i>Empis</i>) 175
<i>surata</i> Kuntze (<i>Empis</i>) 174	<i>tenuinervis</i> Bez. (<i>Empis</i>) 174	<i>thoracica</i> Macq. (<i>Hilara</i>) 126
* <i>suspiciosa</i> Meun. (<i>Micrempis</i>) 299	<i>tenuinervis</i> Zett. (<i>Hilara</i>) 126	<i>thoracica</i> Phil. (<i>Brachystoma</i>) 13
<i>sutor</i> Mel. (<i>Platypalpus</i>) 365	<i>tenuipes</i> Beck. (<i>Rhamphomyia</i>) 207	<i>thoracicus</i> Lundb. (<i>Platypalpus ru-</i>
<i>sycophantor</i> Mel. (<i>Microphorus</i>) 93	<i>tenuipes</i> Brun. (<i>Hybos</i>) 24	<i>ficornis</i>) 366, 384
<i>sydneyensis</i> Schin. (<i>Hybos</i>) 24	<i>tenuipes</i> Lw. (<i>Empis</i>) 174	<i>thoracicus</i> Say (<i>Syneches</i>) 44
<i>syletor</i> , nov. sp. (<i>Charadrodromia</i>) 293		<i>thrixothrix</i> , nov. sp. (<i>Euhybos</i>) 31
<i>sylvaticus</i> Beck. (<i>Stenoproctus</i>) 73		<i>tibialis</i> Bez. (<i>Hybos</i>) 24
<i>sylvicola</i> Coll. (<i>Platypalpus</i>) 382		<i>tibialis</i> Coq. (<i>Mythicomyia</i>) 376
		* <i>tibialis</i> Lw. (<i>Empis</i>) 176
		<i>tibialis</i> Macq. (<i>Edalea flavipes</i>) 64
		<i>tibialis</i> Macq. (<i>Tachyfeza nubila</i>) 277

Pages	Pages	Pages			
tibialis Meig. (<i>Rhamphomyia</i>)	208	trivittata Lw. (<i>Hilara</i>)	127	unimaculata Lw. (<i>Rhamphomyia</i>)	209
tibialis Ros. (<i>Atelostus pulicarius</i>)	78, 384	trivittata Macq. (<i>Empis</i>)	176	unipila Lw. (<i>Drapetis</i>)	311
tibialis Steph. (<i>Platypalpus</i>)	366	trunca, nov. sp. (<i>Clinocera</i>)	230	unipunctatus Lw. (<i>Stenoproctus</i>)	73
tibialis Zett. (<i>Leptopeza ruficollis</i>)	68	truncata Frey (<i>Rhamphomyia</i>)	208	uniseta Mel. (<i>Drapetis</i>)	315
tibiella Mik (<i>Clinocera</i>)	239	truncata Lw. (<i>Empis bistortæ</i>)	148	unistriata Beck. (<i>Empis lutea</i>)	162
tibiella Zett. (<i>Rhamphomyia costata</i>)	189	truncata Wied.-Meig. (<i>Empis</i>)	176	universalis Mel. (<i>Tachyempis</i>)	292
tieff Stro. (<i>Hilara</i>)	127	truncatus Lw. (<i>Schistostoma</i>)	94	univittata Lw. (<i>Empis</i>)	177
Timalphes (nov. gen.)	97	truncorum Fall. (<i>Tachypeza</i>)	278	univittata Meig. (<i>Hilara litorea</i>)	120
tipularia Fall. (<i>Rhamphomyia</i>)	208	truncorum Meig. (<i>Tachypeza nubila</i>)	277	univittata, nov. sp. (<i>Lamachella</i>)	72
tipulariata Zett. (<i>Rhamphomyia</i>		tuberculata Lw. (<i>Tachydromia</i>)	287	univittata Stro. (<i>Rhamphomyia fla-</i>	
<i>sciarina</i>)	204	tuberifera Bez. (<i>Lamprempis</i>)	136	<i>viventris</i>)	192
tipuloides Linn. (<i>Empis tessellata</i>)	175	tubifera nov. sp. (<i>Microphorella</i>)	88	univittatus Lw. (<i>Platypalpus</i>)	366
tolteca Wh.-Mel. (<i>Rhamphomyia</i>)	208	tumida Lw. (<i>Empis pudica</i>)	170	upsilon Mel. (<i>Drapetis</i>)	315
tonsa Lw. (<i>Rhamphomyia</i>)	208	tumida Meig. (<i>Empis</i>)	176, 385	uralensis Beck. (<i>Rhamphomyia</i>)	209
Toreus Mel. (gen.)	132	tumiditarsis Old. (<i>Rhamphomyia</i>)	208	<i>ursina</i> Mall. (<i>Rhamphomyia ursi-</i>	
torrentalis Mill. (<i>Empimorpha</i>)	378	turbida Meig. (<i>Empis volucris</i>)	178	<i>nella</i>)	209
totipennis Bell. (<i>Empis</i>)	176	turgidus Beck. (<i>Platypalpus</i>)	366	<i>ursina</i> Old. (<i>Rhamphomyia</i>)	209
transfuga Walk. (<i>Hilara</i>)	127	tympanica Bez. (<i>Rhamphomyia</i>)	208	<i>ursinella</i> , n. n. (<i>Rhamphomyia</i>)	209
trapezina Zett. (<i>Chelifera melan-</i>		typicus Wh.-Mel. (<i>Hybos</i>)	24	uruguayensis Arrib. (<i>Empis</i>)	177
<i>cephala</i>)	261, 385	tyrolensis Stro. (<i>Hilara</i>)	127	uvens, nov. sp. (<i>Bicellaria</i>)	74
triangula Coq. (<i>Iteaphila</i>)	104	umbilicata Lw. (<i>Rhamphomyia</i>)	208	*vaga Meun. (<i>Chelipoda</i>)	267
trianguligera Stro. (<i>Empis</i>)	176	umbrarum Hal. (<i>Tachydromia annu-</i>		vagans Lw. (<i>Hybos</i>)	24
Trichina Meig. (gen.)	54	<i>limana</i>)	283	vaginifer Mel. (<i>Empis</i>)	177
Trichopeza Rond. (gen.)	218	umbrina Wied.-Meig. (<i>Empis</i>)	176	valdiviana Phil. (<i>Drapetis</i>)	312
trichura Mel. (<i>Drapetis</i>)	309	umbripennis Eversm. (<i>Empis</i>)	176	valdiviana Phil. (<i>Empis</i>)	177
tricuspidata Bez. (<i>Wiedemannia</i>)	245	umbripennis Frey (<i>Rhamphomyia</i>		valens, nov. sp. (<i>Platypalpus</i>)	366
tridentata Coq. (<i>Empis</i>)	176	<i>nigripennis</i>)	199	valentis Coq. (<i>Empis</i>)	177
trigemina Old. (<i>Rhamphomyia</i>)	383	umbripennis Meig. (<i>Rhamphomyia</i>)	209	valga Coq. (<i>Rhamphomyia</i>)	209
trigemina Stro. (<i>Hilara cinereomi-</i>		umbripennis Meig. (<i>Tachypeza</i>)	278	valgus, nov. sp. (<i>Platypalpus</i>)	338
<i>cans</i>)	115	umbripes Beck. (<i>Rhamphomyia</i>)	209	valida Lw. (<i>Chelifera</i>)	261
trigramma Meig. (<i>Hilara lugubris</i>)	120	umbrosa Lw. (<i>Hilara</i>)	127	validis Adams (<i>Empis</i>)	177
trigramma Wied.-Meig. (<i>Empis</i>)	176	umbrosa Lw. (<i>Rhamphomyia</i>)	209	vallium Verr.-Coll. (<i>Hilara galac-</i>	
trilineata Gmel. (<i>Empis</i>)	176	uncicauda Bez. (<i>Hilara</i>)	127	<i>toptera</i>)	380
trilineata Zett. (<i>Rhamphomyia</i>)	208	uncinatus, nov. sp. (<i>Platypalpus</i>)	324	vanellus Schin. (<i>Hilaraempis</i>)	130
trinotata Mik (<i>Clinocera</i>)	240	undulata, nov. sp. (<i>Clinocera</i>)	230	vara Lw. (<i>Rhamphomyia</i>)	209
tripes Beck. (<i>Rhamphomyia para-</i>		undulata Stro. (<i>Tachydromia</i>)	287	variabilis Big. (<i>Empis</i>)	177
<i>doxa</i>)	201	unguiculata Frey (<i>Rhamphomyia</i>)	209	variabilis Fall. (<i>Rhamphomyia tenui-</i>	
triplex Walk. (<i>Euhybos</i>)	35	unguiculatus Zett. (<i>Platypalpus</i>)	366	<i>rostris</i>)	207
tristis Egg. (<i>Hilarimorpha</i>)	373	ungulata Lw. (<i>Rhamphomyia umbi-</i>		varians Big. (<i>Empis</i>)	177
tristis Lw. (<i>Empis</i>)	176	<i>licata</i>)	208	variata Mel. (<i>Drapetis</i>)	315
tristis Lw. (<i>Hilara</i>)	127	ungulifera Mel. (<i>Drapetis lata</i>)	314	varicolor Beck. (<i>Platypalpus</i>)	367
*tristis Meun. (<i>Empis tritava</i>)	176	*ungulina Lw. (<i>Rhamphomyia</i>)	209	variegata Bez. (<i>Apalocnemis</i>)	101
tristis Scholz (<i>Edalea</i>)	65	unica Walk. (<i>Ragas</i>)	109	variegata Brun. (<i>Drapetis</i>)	315
tristis Walk. (<i>Rhamphomyia</i>)	208	unicolor Brullé (<i>Empis</i>)	177	variegata Brun. (<i>Drapetis variata</i>)	315
tristriolata Now. (<i>Rhamphomyia</i>)	208	unicolor Curt. (<i>Clinocera nigra</i>)	237	variegata Meig. (<i>Empis</i>)	177
*tritava, n. n. (<i>Empis</i>)	176	unicolor Lw. (<i>Acarterus</i>)	38	varipennis Now. (<i>Clinocera</i>)	240
trivialis Lw. (<i>Platypalpus</i>)	366	unicolor Lw. (<i>Hilara</i>)	127	varipes Lw. (<i>Empis</i>)	177
trivialis Mel. (<i>Platypalpus coquill-</i>		unicolor Stro. (<i>Leptopeza ruficollis</i>)	68	varipes Lw. (<i>Stilpon</i>)	305
<i>letti</i>)	351	unicolor Walk. (<i>Empis</i>)	177	varipes Meig. (<i>Platypalpus</i>)	367
trivialis Wh.-Mel. (<i>Platypalpus</i>		unicolor Zett. (<i>Rhamphomyia</i>)	209	varipes, nov. sp. (<i>Micrempis</i>)	299
<i>caligaris</i>)	349	unifasciata Brun. (<i>Rhamphomyia</i>)	209	varipes White (<i>Acarterus</i>)	377
trivialis Wh.-Mel. (<i>Platypalpus</i>		unilineata Zett. (<i>Hemerodromia</i>)	257	varius Bez. (<i>Platypalpus oedicnemus</i>)	362

DIPTERA

	Pages		Pages		Pages
varius Walk. (<i>Platypalpus</i>)	367	virgata Coq. (<i>Rhamphomyia</i>)	210	winthemi Zett. (<i>Tachypeza</i>)	278
varus, nov. sp. (<i>Syneches</i>)	44	virgator, nov. sp. (<i>Hormopeza</i>)	96	woldstedti Frey (<i>Rhamphomyia</i>)	210
vector Mill. (<i>Hilara</i>)	380	viridis Coq. (<i>Lamprempis</i>)	137	woodi Coll. (<i>Empis</i>)	379
velox, nov. sp. (<i>Platypalpus</i>)	336	* vitiosa Meun. (<i>Drapetis</i>)	312	woodi Coll. (<i>Hilara</i>)	380
velutina Bez. (<i>Empis</i>)	177	vitripennis auct. (<i>Hybos grossipes</i>)	22	woodi Coll. (<i>Tachydromia</i>)	383
velutina Lw. (<i>Hilara</i>)	127	vitripennis Beck. (<i>Platypalpus</i>)	367	* wulpi Weyenb. (<i>Empidia</i>)	369
velutinus Macq. (<i>Microphorus</i>)	93	vitripennis Bez. (<i>Tachyempis</i>)	292	wuorentausi Frey (<i>Rhamphomyia</i>)	210
velutinus Meij. (<i>Syneches</i>)	45	vitripennis Brun. (<i>Leptopeza</i>)	68		
venaticus, nov. sp. (<i>Platypalpus</i>)	335	vitripennis Meig. (<i>Eupis volucris</i>)	178		
venosus Steph. (<i>Platypalpus</i>)	367	vitripennis Meig. (<i>Hybos culiciformis</i>)	21		
ventralis Meig. (<i>Platypalpus can-</i>		vittata Arrib. (<i>Chelipoda</i>)	266	Xanthempis Bez. (subgen. <i>Em-</i>	
<i>dicans</i>)	350	vittata Lw. (<i>Hemerodromia empiformis</i>)	256	<i>pis</i>)	145
ventralis Zett. (<i>Empis rufiventris</i>)	172	vittata Mel. (<i>Drapetis</i>)	315	xanthocephala Bez. (<i>Drapetis</i>)	315
verasi Brethes (<i>Haplomera</i>)	132	vittatus, nov. sp. (<i>Platypalpus</i>)	334	xanthocera Bez. (<i>Hilarempis</i>)	130
ernalis Meig. (<i>Empis pennaria</i>)	177, 385	vittatus Walk. (<i>Syneches</i>)	44	xanthochiton, nov. sp. (<i>Platypal-</i>	
verpus, nov. sp. (<i>Platypalpus</i>)	333	vittiger, nov. var. (<i>Platypalpus flavi-</i>		<i>pus</i>)	322
verralli Coll. (<i>Empis</i>)	379	<i>rostris</i>)	323	xanthopoda Will. (<i>Drapetis</i>)	312
verralli Coll. (<i>Platypalpus</i>)	382	vittigerum Felt. (<i>Brachystoma</i>)	14	xanthopoda Will. (<i>Drapetis flavida</i>)	314
verruciferus, nov. sp. (<i>Euhyas</i>)	26	vittipennis Walk. (<i>Tachypeza</i>)	278	xanthopodus, nov. sp. (<i>Hybos</i>)	24
versipes, nov. sp. (<i>Platypalpus</i>)	318	vividus Meig. (<i>Platypalpus albisetosa</i>)	346	xanthopodus, n. n. (<i>Platypalpus</i>)	367
versutus Mel. (<i>Platypalpus</i>)	367	vocatoria Fall. (<i>Chelipoda</i>)	267	vanhofus Bez. (<i>Platypalpus longirostris</i>)	357
vesiculosus Fall. (<i>Rhamphomyia</i>)	200	vocatoria Meig. (<i>Chelipoda albisetosa</i>)	265	xanthopyga Bez. (<i>Drapetis</i>)	309
vesiculosum Fab. (<i>Brachystoma</i>)	14	volucris Wied.-Meig. (<i>Empis</i>)	178	xanthopyga Schin. (<i>Empis</i>)	178
vespertilio Zett. (<i>Rhamphomyia</i>)	210	volucris Zett. (<i>Empis astiva</i>)	146	xiphias Bez. (<i>Hemerodromia</i>)	257
vicana Hart. (<i>Rhamphomyia nigripes</i>)	199	* voracis Meun. (<i>Tachypeza</i>)	278	Xiphidiocera Macq. (<i>Edalea</i> Meig.)	62
vicarius Walk. (<i>Platypalpus</i>)	367	vulgaris Meig. (<i>Platypalpus flavipes</i>)	355	Xiphidiocera Blanch. (<i>Edalea</i> Meig.)	62
vicina Arrib. (<i>Empis</i>)	178	vulneratus Schin. (<i>Hilara pruinosa</i>)	124	zochitl Wh.-Mel. (<i>Empis</i>)	178
vicinus Mik (<i>Microphorus</i>)	94	vulnificus, nov. sp. (<i>Platypalpus</i>)	339	Xiphidiocera Big. (<i>Edalea</i> Meig.)	62
vierecki Mel. (<i>Platypalpus</i>)	367	wachtlii Mik (<i>Wiedemannia</i>)	244	yakimensis, nov. sp. (<i>Schistostoma</i>)	94
villeneuvei Beck. (<i>Platypalpus</i>)	367	walkeri Bez. (<i>Empis</i>)	178	yucatanus Wh.-Mel. (<i>Euhyas</i>)	35
villipes Coq. (<i>Rhamphomyia</i>)	210	weedi Coq. (<i>Neocota</i>)	180	zaitzevi Beck. (<i>Rhamphomyia</i>)	210
villosa Macq. (<i>Empis</i>)	178	wesmaelii Macq. (<i>Clinocera</i>)	238	zetterstedti Fall. (<i>Wiedemannia</i>)	243
villosa Zett. (<i>Rhamphomyia</i>)	210, 385	wesmaelii Zett. (<i>Clinocera fontinalis</i>)	237	zetterstedti Walk. (<i>Clinocera bipunctata</i>)	239
vilosipes Bez. (<i>Rhamphomyia</i>)	210	westmanni Zett. (<i>Clinocera stag-</i>		zeylanicus White (<i>Platypalpus</i>)	382
vilosipes Stro. (<i>Rhamphomyia ur-</i>		<i>nalis</i>)	238	zontaki Now. (<i>Anthropiscopus</i>)	108
<i>sina</i>)	200	wheeleri Mel. (<i>Hilara</i>)	127		
vilosula Big. (<i>Empis</i>)	178				
violacea Lw. (<i>Lamprempis</i>)	137				
virgata Coq. (<i>Empis</i>)	178				

EXPLANATION OF THE PLATES

PLATE I

- Fig. 1. *Prorates claripennis* Melander, ♂.
 — 2. *Brachystoma vesiculosum* Fabricius, ♂.
 — 3. *Euhybos purpureus* Walker, ♂.
 — 4. *Syneches phthisia* Walker, ♂.
 — 5. *Hybos reversus* Walker, ♂.

- Fig. 6. *Syneches (Harpamerus) Bakeri*, nov. sp., ♂.
 — 7. *Parahybos chiragra* Bezzi, ♂.
 — 8. *Trichina clavipes* Meigen, ♂.
 — 9. *Œdalea tristis* Scholz, ♂.
 — 10. *Ocydromia glabricula* Fallen, ♂.

PLATE 2

- Fig. 11. *Hoplocyrtoma procera* Loew, ♂.
 — 12. *Porphyrochroa palliata* Coquillett, ♂.
 — 13. *Lamprempis chichimeca* Wheeler & Melander, ♀.
 — 14. *Deuteragonista breviventris* Philippi, ♂.

- Fig. 15. *Phleboctena lutea* Bezzi, ♀.
 — 16. *Brochella monticola*, nov. sp., ♂.
 — 17. *Oreothalia pelops* Melander, ♂.
 — 18. *Clinocera nigra* Meigen, ♂.
 — 19. *Roederioides juncta* Coquillett, ♀.

PLATE 3

- Fig. 20. *Boreodromia bicolor* Loew, ♂.
 — 21. *Niphogenia eucera*, nov. sp., ♂.
 — 22. *Ceratempis longicornis*, nov. sp., ♂.
 — 23. *Synamphotera pallida* Loew, ♂.
 — 24. *Dolichocephala irrorata* Fallen, ♂.

- Fig. 25. *Chelipoda vocatoria* Fallen, ♂.
 — 26. *Hemerodromia oratoria* Fallen, ♂.
 — 27. *Colabris coxalis*, nov. sp., ♂.
 — 28. *Chelifera precatoria* Fallen, ♂.

PLATE 4

- Fig. 29. *Tachydromia connexa* Meigen, ♂.
 — 30. *Platypalpus cursitans* Fabricius, ♂.
 — 31. *Tachypeza discifera*, nov. sp., ♂.
 — 32. *Micrempis nana*, nov. sp., ♂.
 — 33. *Megagrapha pubescens* Loew, ♀.

- Fig. 34. *Drapetis (Eudrapetis) spectabilis* Melander, ♂.
 — 35. *Tachyempis pictipes*, nov. sp., ♂.
 — 36. *Drapetis (Ctenodrapetis) aristalis* Melander, ♂.
 — 37. *Stilpon graminum* Fallen, ♂.

PLATE 5

- Fig. 38. *Anomalempis tacoma*, nov. sp. ♀.
 — 39. *Syndyas polita* Loew, ♂.
 — 40. *Meghyperus occidens* Coquillett, ♂.
 — 41. *Homalocnemis nigripennis* Philippi (from Philippi).
 — 42. *Acarterus unicolor* Loew (from Loew).
 — 43. *Scelolabes bivittatus* Philippi (fr. Philippi).

- Fig. 44. *Stenoproctus unipunctatus* Loew (fr. Loew).
 — 45. *Lactistomyia insolita* Melander, ♂.
 — 46. *Bicellaris spuria* Meigen, ♂.
 — 47. *Empis (Pachymeria) femorata* Fabricius, ♂.
 — 48. *Empis pennipes* Linnæus, ♂.
 — 49. *Empis pennipes* Linnæus, ♀.
 — 50. *Oreogoton basalis* Loew ♂.

PLATE 6

- Fig. 51. *Oreogoton mitraphorus*, nov. sp. ♂.
 — 52. *Gloma fuscipennis* Meigen, ♂.
 — 53. *Timalphes fumosa* Hutton, ♂.
 — 54. *Hilarempis nudifacies* Bezzi, ♂.
 — 55. *Hilara maura* Fabricius, ♂.

- Fig. 56. *Apalocnemis obscura* Philippi (fr. Philippi).
 — 57. *Neocota Weedii* Coquillett, ♂.
 — 58. *Haplomera gymnopaoda* Bezzi, ♀.
 — 59. *Iteaphila Macquarti* Zetterstedt, ♀.
 — 60. *Atrichopleura Schnusei* Bezzi, ♂.

PLATE 7

- Fig. 61. *Hesperempis Mabela* Melander, ♂.
 — 62. *Microphorus (Schistostoma) eremita* Beck., ♂.
 — 63. *Microphorus velutinus* Macquart, ♂.
 — 64. *Lamposoma cavaticum* Becker (from Beck.).
 — 65. *Proclinopyga amplectens*, nov. sp., ♂.
 — 66. *Philetus memorandus*, nov. sp., ♂.

- Fig. 67. *Ceratomerus paradoxus* Philippi (from Philippi).
 — 68. *Heleodromia pullata* Melander, ♂.
 — 69. *Trichopeza longicornis* Meigen, ♂.
 — 70. *Coloboneura inusitata* Melander, ♂.
 — 71. *Thinodromia inchoata* Melander, ♂.

PLATE 8

- Fig. 72. *Bicellaria*. Ventral forks of pygidium.
 A. *furcifer*, nov. sp.; B. *longipes* Loew;
 C. *pectinata*, nov. sp.; D. *lugubris* n. sp.;
 E. *filipes* Loew; F. *uvens*, nov. sp.;
 G. *brevifurca*, nov. sp.
 — 73. *Gloma luctuosa*, nov. sp. Pygidium.
 — 74. *Gloma fuscipennis* Meigen. Pygidium.
 — 75. *Proclinopyga fistulator*, nov. sp. Pygidium.
 — 76. *Hormopesa brevicornis* Loew. Wing.
 — 77. *Blepharoprocta serratula* Loew. Wing. ♂.
 — 78. *Philetus memorandus*, nov. sp. Pygidium.

- Fig. 79. *Leptopeza ruficollis* Meigen. Wing.
 — 80. *Anthepisopus stenor* Melander. Pygidium.
 — 81. *Philetus schizophorus*, nov. sp. Pygidium.
 — 82. *Euthyneura Myrtilli* Macquart. Wing.
 — 83. *Tenontomyia gracilipes* White. Wing (from White).
 — 84. *Ragas unica* Walker. Pygidium.
 — 85. *Rhamphomyia furcifer* Wheeler & Melander. Wing.
 — 86. *Ironomyia nigromaculata* White. Wing (from White).


The manuscript of this fascicle was sent to the Genera Insectorum on January 10, 1925, from the State College of Washington. M. P. Wytsman completed the editing of the printer's copy just prior to his death. The presswork has been supervised by Miss H. Van Hamme, as successor to M. Wytsman.

A. L. Melander,
 College of the City of New York.
 11 May, 1927.

POSTSCRIPT. — In the interim since May 11, the conclusion of Collin's article on British Empididae has appeared in the Entomologists' Monthly Magazine for May, 1927, pages 97 and 98. This section adds *Hemerodromia melangyna* Collin, p. 97, and *Wiedemannia rhynchos*, subsp. *insularis* Collin, p. 98, both from the British Isles. Descriptions have also appeared of **Hilara royo*, p. 98 (Oligocene, Spain) Collado, Nota sobre algunos insectos fósiles de Ribesalbes, Castellón (Bol. Inst. Geol. Esp. Madrid, Vol. 46, p. 89-107 [1926]); and *Empis deterra*, p. 96, and *Rhamphomyia bigelowi*, p. 97 (Ontario) Walley, Two new species of Empididae from Ontario (Canad. Ent. Vol. 59, p. 96-98 [1927]). (Final proofs returned to the printer June 21, 1927.)





GENERA INSECTORUM

DIPTERA




FAM. EMPIDIDÆ

SUBFAM. BRACHYSTOMATINÆ, HYBOTINÆ, OCYDROMIINÆ

*Eupogonius procera* Loew, ♂*Porphyrochroa palliata* Coquill., ♂*Lampromyia chichimeca* Wheel & Mel., ♀*Deuteragonista breviventris* Phil., ♂*Phlebotoma lutea* Bezz., ♀*Brochella monticola*, n.sp., ♂*Oreothalia pelope* Mel., ♂*Clinocera nigra* Meig., ♂*Roederioides juncta* Coquill., ♀



FAM. EMPIDIDÆ

SUBFAM. OCYDROMIINÆ, EMPIDINÆ, CLINOCERATINÆ



FAM. EMPIDIDÆ


SUBFAM. CLINOCERATINÆ, HEMERODROMIINÆ

*Tachydromia connexa* Meig. ♂*Platypalpus cursitans* Fabr. ♂*Megagrapha pubescens* Loew. ♂*Tachyempis pictipes* n.sp. ♂*Tachypora discifera*, n.sp., ♂*Drapetis (Eudrapetis) spectabilis* Mel. ♂*Drapetis (Ctenodrapetis) aristalis* Mel. ♂*Meromips nana*, n.sp., ♂*Sutpon graminum* Fall. ♂

FAM. EMPIDIDÆ








SUBFAM. TACHYDROMIINÆ






FAM. EMPIDIDÆ

SUBFAM. BRACHYSTOMATINÆ, HYBOTINÆ, OCYDROMIINÆ, EMPIDINÆ


*Oreogenes mitrophorus*, n. sp., ♂*Glossa trucipennis* Meig., ♂*Timaipes sonosa* Bull., ♂*Apalocnemis obscura* Phil.*Hilarempis nudifacies* Bezzi, ♂*Mura maura* Fadr., ♂*Neocoila Weedii* Coq., ♂*Haplomera gymnopoda* Bezzi, ♀*Itapilia Macquarti* Zell., ♀*Atrichopleura sohnensei* Bezzi, ♂

FAM. EMPIDIDÆ
SUBFAM. EMPIDINÆ



FAM. EMPIDIDÆ

SUBFAM. EMPIDINÆ, CLINOCERATINÆ, TACHYDROMIINÆ





BINDING LIST FEB 19 1939

QL Wytsman, Philogene (ed.)
468 Genera insectorum
W87
Fasc.184-185

Biological
& Medical

PLEASE DO NOT REMOVE
CARDS OR SLIPS FROM THIS POCKET

UNIVERSITY OF TORONTO LIBRARY
